Incorporating Social Determinants of Health into the Mathematical Modeling of HIV/AIDS

Robson Oliveira (✉ robson.bruniera@gmail.com)

Federal University of Bahia

Felipe Rubio

Federal University of Bahia

Rodrigo Anderle

Federal University of Bahia

Mauro Sanchez

University of Brasília

Luis de Souza

Federal University of Bahia

James Macinko

University of California, Los Angeles

Ines Dourado

Federal University of Bahia

Davide Rasella

University of Barcelona

Research Article

Keywords: HIV/AIDS, Epidemiology, Mathematical Modeling, Compartmental models, Social determinants of health

Posted Date: March 2nd, 2022

DOI: <https://doi.org/10.21203/rs.3.rs-1367356/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Incorporating Social Determinants of Health into the Mathematical Modeling of HIV/AIDS

*Robson Bruniera de Oliveira¹, Felipe Alves Rubio¹, Rodrigo Anderle¹, Mauro Sanchez², Luis Eugenio de Souza¹, James Macinko³, Ines Dourado¹, Davide Rasella^{1,4}

1. Institute of Collective Health, Federal University of Bahia (UFBA), Salvador, Brazil.
2. Department of Public Health, University of Brasilia, Brasilia, Brazil.
3. Departments of Health Policy and Management and Community Health Sciences, UCLA Fielding School of Public Health, Los Angeles, California.
4. ISGlobal, Hospital Clínic - Universitat de Barcelona, Barcelona, Spain.

* Corresponding author:
Robson.bruniera@gmail.com
Email Address: Robson.bruniera@gmail.com
Phone: +55 12988887885

Abstract

Background

Currently, it is estimated that 37.6 million people are living with the HIV/AIDS virus worldwide, placing HIV/AIDS among the 10 leading causes of death, mostly among low- and lower-middle-income countries. Despite the effective intervention in the prevention and treatment, this reduction did not occur equally among populations, subpopulations and geographic regions. This difference in the occurrence of the disease is associated with the social determinants of health (SDH), which could affect the transmission and maintenance of HIV. With the recognition of the importance of SDH in HIV transmission, the development of mathematical models that incorporate these determinants could increase the accuracy and robustness of the modeling. This article aims to propose a theoretical and conceptual way of including SDH in the mathematical modeling of HIV/AIDS.

Methods

The theoretical mathematical model with the Social Determinants of Health has been developed in stages. For the selection of SDH that were incorporated into the model, a narrative literature review was conducted. Secondly, we proposed an extended model in which the population (N) is divided into Susceptible (S), HIV-positive (I), Individual with AIDS (A) and individual under treatment (T). Each SDH had a different approach to embedding in the model.

Results

A total of 31 SDH were obtained in the review, divided into 4 groups: Individual Factors, Socioeconomic Factors, Social Participation, and Health Services. In the end, 4

determinants were selected for incorporation into the model: Education, Poverty, Use of Drugs and Alcohol abuse, and Condoms Use.

Conclusions

The equations presented with the chosen SDH exemplify some approaches that we can adopt when thinking about modeling social effects on the occurrence of HIV. The recognition of the importance of including the SDH in the modeling and studies on HIV/AIDS is evident, due to its complexity and multicausality. Models that do not take into account in their structure, will probably miss a great part of the real trends, especially in periods, as the current one, of economic crisis and strong socioeconomic changes.

Key Words: HIV/AIDS, Epidemiology, Mathematical Modeling, Compartmental models, Social determinants of health

Background

Currently, it is estimated that 37.6 million people are living with the HIV/AIDS virus worldwide, with an associated mortality of 1.5 million people per year (UNAIDS, 2021). These numbers place HIV/AIDS among the 10 leading causes of death, mostly among low- and lower-middle-income countries (WHO, 2020). In 2002, the World Health Organization (WHO) implemented the Getting to Zero (GTZ) initiative, with the aim of making 90% of people living with HIV aware of their status, 90% of diagnosed people under adequate antiretroviral treatment (ART), and 90% of those receiving ART be virologically suppressed (with a 95-95-95 coverage by 2030) (UNAIDS, 2014a; UNAIDS, 2014b; UNAIDS, 2020; Eisinger et al, 2019; End Inequalities, 2021)

Despite the effective interventions implemented for the prevention and treatment of HIV/AIDS, with a considerable reduction in new cases in the world, this reduction did not occur equally among populations, subpopulations and geographic regions (Dean and Fenton, 2010). This fact triggered the complexity of the challenge of controlling the disease, whose dynamics are closely related to behavioral factors, but also to social determinants to which populations are subjected (McMahan et al, 2021; UNAIDS, 2015, Leport et al., 2012). It became clear that responding to HIV is not just a biomedical issue, but is also a social challenge.

Social Determinants of Health (SDH) are conditions in the environments in which people are born, live and work that influence the occurrence of health problems and their risk factors (Healthy People, 2020). Not unlike other diseases, HIV/AIDS suffers the direct and indirect effect of these determinants, with an influence on its incidence and mortality. Socioeconomic determinants, for example, affect the dynamics of the disease, to the point that low-income populations have unequal rates in terms of HIV incidence, mortality and treatment success compared to individuals with better purchasing power

(Burch et al., 2016). Thus, low-income individuals are disproportionately affected by HIV, with an associated higher mortality (An Q et al, 2013). Even in countries with high incomes, such as the USA, impoverished urban areas are observed to have HIV prevalence rates equivalent to those of many low-income countries with generalized epidemics (Buot et al, 2014). In this context, a better understanding of the main social determinants of health (SDH), whether in the social, cultural and economic spheres, which interact with the risk factors for transmission and maintenance of HIV, gained space for better health outcomes and reduction of health inequities (WHO, 2008).

Mathematical models, in the medical field, are used to study, among other objectives, the behavior of infectious diseases in the population, the planning and evaluation of prevention and control programs, clinical trials and cost-benefit analysis of interventions. These models can even project how infectious diseases progress over time, with the ability to predict the evolution patterns of the dynamics of a disease in populations (Verma et al., 1981; Tolles and Luong, 2020). These models have been used to study the dynamics of HIV infection for decades. These models provide information about the potential impacts of interventions that are difficult to measure empirically for reasons of time, logistics or ethics (Cassels and Goodreau, 2011). With the recognition of the importance of SDH in HIV transmission, the development of mathematical models that incorporate these determinants in their scope, brings robustness to the results, with a significant improvement in the power and accuracy of the modeling (Galanis and Hanieh, 2021). This article aims to propose a theoretical and conceptual compartmental model for the inclusion of social determinants of health in the structure of the mathematical models for HIV/AIDS.

Methods

The theoretical mathematical model with the Social Determinants of Health has been developed in stages.

Social Determinants of Health.

For the selection of SDH that were incorporated into the model, a narrative literature review was conducted focused on peer-reviewed articles, in PubMed/Medline and Scopus databases between 2010 and 2020. This review included key words HIV, AIDS and Social Determinants. A total of 31 SDH were obtained in the review, divided into 4 groups: Individual Factors, Socioeconomic Factors, Social Participation and Health Services (Figure 1). In the end, 4 determinants were selected for incorporation into the model: Education, Employment, Use of Drugs and Alcohol abuse and Condoms Use. The choice of these SDH was based on the recurrence; recognized importance in the literature and the capacity for measurement and quantification, in order to be incorporated into the equations of the Compartmental Model. Although these 4 chosen determinants are among the most studied and important in the field of HIV/AIDS, they do not have the capacity and robustness to represent all other determinants and groups.

Education

Education is recognized as an important social determinant of health, being directly associated with the socioeconomic development and well-being of individuals. Different educational levels in a society lead to economic disparities and especially health inequities. Individuals with higher educational levels, in general, have better job opportunities, a lower unemployment rate, better economic conditions and, consequently, better psychosocial conditions for decision-making about their own health. Thus,

education, within society, has an influence and impact on life expectancy and its morbidities (The Lancet Public Health, 2020).

For HIV/AIDS, education plays an important role in reducing its incidence and prevalence, especially in low- and low-income countries. As many of the young people in developing countries who attend school have not yet had sexual intercourse, education for these young people is critical to the success of HIV/AIDS prevention programs. At this time, it is possible to guide young people to delay the beginning of their sexual activities, as well as encourage the use of protective methods such as condoms (Kirby et al, 2006). Petifor et al. (2016) noted that social protection programs, such as conditional cash transfer to school attendance, made young women stay in school longer, reducing the risk of acquiring HIV. Furthermore, education helps people to access and adhere to ART, as well as helping to reduce stigma, discrimination and gender inequality (UNESCO & WHO, 2005).

Poverty and Employment

Between the years 2020 and 2021, largely as a result of the COVID-19 pandemic, the number of unemployed people in the world increased from 187.3 million to 220.3 million, the largest annual increase in unemployment in this period of time, reaching the rate average of 6.5% (Statista, 2021). Many of these people who have lost their jobs have lost their only source of income, both personal and family, resulting in a drastic reduction in labor income and a consequent increase in poverty. Compared to the year 2019, 108 million more workers are now considered to be living in poverty or extreme poverty (ILO, 2021). Several studies have documented the association between unemployment and poor health status (DiClementi et al., 2004; Jin, Shah and Svoboda, 1995; Kasl et al., 1998.) Unemployed individuals, have their physical and mental health affected, more likely to suffer from depression, anxiety, low self-esteem, demoralization, worry and physical pain

(Avendano and Kawachi, 2014). In addition, these individuals end up having their perception of health reduced, exposing themselves more to risk of HIV infection, with risk behaviors such as exchanging sex for money, often with multiple partners and without condoms; low demand for care and health services, and consequently delay in detection and initiation of treatment for HIV/AIDS and increase considerable in mortality (Joy et al., 2008; Maruthappu et al., 2017)

Drug and Alcohol Abuse

Alcohol and drug abuse, whether injectable or non-injectable, are intrinsically associated with an increased risk of HIV infection. The main factor of injecting drugs is the sharing of syringes and needles, while alcohol and other types of drugs favor the increase in risky behavior due to the exchange of sex for drugs or money, sexual disinhibition (Pence et al., 2008). Approximately 15.6 million people inject drugs worldwide, while around 2.8 million of those are living with the HIV virus (Degenhardt et al., 2017). Sharing needles and syringes is the second risk behavior for HIV infection, second only to receptive anal sex. One in 160 people becomes infected every time they share a syringe, with 10% of HIV cases in the USA being attributed to this risky behavior. In this way, injecting drug users (IDUs) are exposed to the double route of contamination, sexual and intravenous. Another relevant factor is that substance abuse, alcohol consumption, as well as unemployment are factors related to reduced adherence to antiretroviral treatment (Costa et al., 2018). Therefore, the use of alcohol and drugs is a SDH that must be considered in the mathematical modeling of HIV/AIDS.

Condom Use

Approximately 95% of cases, worldwide, of people being infected by HIV are attributed to sexual practices without using condoms. Condoms are the best known, most

accessible and effective method to prevent HIV infection and other sexually transmitted infections, such as syphilis, gonorrhea and also some types of hepatitis (Pinkerton and Abramson, 1997). Consistent condom use can reduce HIV transmission among serodiscordant individuals by up to 80% (Weller and Davis, 2002).

Figure 1 - Social Determinants of Health for HIV/AIDS derived from Narrative review.

Mathematical Modeling

Compartmental model is a type of mathematical model that simulates the disease status of individuals within populations, which are divided into different compartments. Within each compartment, people are considered homogeneous in terms of their behavior

and risk factors (Verma et al., 1981; Tolles and Luong, 2020). The most generic and widely used model is the SIR model, in which individuals are classified into 3 types of compartments: Susceptible (S), Infected (I) and Recovered (R). Susceptible individuals are those who have never had the disease or are likely to become infected. After infection, these individuals migrate to the infected compartment and can spread the disease to susceptible individuals. The recovered ones can develop lifetime immunity or return to the Susceptible compartment, depending on the pathophysiology of the disease (Beckley 2020). In the case of HIV, the Recovered compartment does not exist, as this disease has no cure. However, a variation of the SIR model for HIV is the replacement of the Recovered compartment by the Treated compartment (T), where individuals in these compartments have an undetectable viral load and their contribution to transmission is almost zero (Huo et al., 2016). For this study, we propose an extended model in which the population (N) is divided into Susceptible (S), HIV-positive (I), Individual with AIDS (A) and individual under treatment (T). Thus, we consider the total population (N), where $N=S+I+A+T$. One way to model the dynamics of virus transmission is given by the system of equations:

$$\frac{dS}{dt} = \kappa - \mu S - \beta \frac{I}{N} S$$

$$\frac{dI}{dt} = \beta \frac{I}{N} S + \alpha_1 T - \rho I - \gamma_1 I - \mu I$$

$$\frac{dA}{dt} = \rho I + \alpha_2 T - \gamma_2 A - \delta_1 A - \mu A$$

$$\frac{dT}{dt} = \gamma_1 I + \gamma_2 A - \delta_2 T - \alpha_1 T - \alpha_2 T - \mu T$$

where κ represents the population's natality rate and μ the natural mortality rate (inverse of the average life expectancy). β represents the effective contact rate, I/N the fraction of

infected individuals. Thus, the term $\beta \frac{I}{N}$ represents the HIV transmission rate. After a period ρ^{-1} , the infected individual becomes “full-blown aids”. We consider that the individual in class I, upon adhering to treatment, after a period γ_1^{-1} it goes to compartment T, that is, reaches undetectable viral load levels, the same applies to the individual in compartment A, in this case at a rate γ_2 . However, if this individual with an undetectable viral load ceases treatment, the viral load may increase, which may lead the individual to class I (infectious) or even to compartment A, with rates α_1 and α_2 , respectively. Figure 2 shows the epidemiological scheme of the HIV/AIDS transmission model.

Figure 2 – Epidemiological scheme of the HIV/AIDS transmission model.

Education

One way to include the education factor would be to incorporate a new compartment, denoted here by (R-removed), representing individuals who through educational campaigns change their sexual behavior so as not to be susceptible to HIV transmission, as shown by (Huo et al., 2016). Parameter θ_1 represents this behavior. However, these individuals can change their behaviors again and return to infection susceptibility at a rate θ_2 . The dynamics of this class is given by the equation,

$$\frac{dR}{dt} = \theta_1 S - \theta_2 R - \mu R$$

Bearing in mind that as we incorporated a new compartment into the model, the total population is now given by $N=S+I+A+T+R$ and the susceptible dynamics is modified by including the term $-\theta_1 S$.

Condom use

To incorporate the effect of condom use, we can include this factor through the parameters ε and ν , which represent the condom efficacy and compliance, respectively. So, the product $c = \varepsilon\nu$ represents the level of protection against HIV through condom use (Nyabadza, 2006). We modified the effective contact rate, which goes as follows:

$$\beta = q(1 - c)\beta$$

where $1 - c$ measures the failure to prevent transmission through condom use and q the rate of acquisition of new partners.

Drug and Alcohol abuse

One way to incorporate the drug abuse as SDH in a mathematical model is to restrict the study to this subpopulation, that is, consider a compartmental model in which all compartments are related to drug users (Greenhalgh and Hay, 1997). For instance, Burattini et al. (1998) showed the impact of crack-cocaine use on the prevalence of HIV/AIDS among drug users. Alcohol abuse can be incorporated in the model decreasing (or increasing) the parameter values, which can be influenced by alcohol use. For instance, in the rate of acquiring partners (q), the parameter become

$$q = \phi_q q$$

and in the level of protection against HIV through condom use (c) as

$$c = \phi_c c$$

with $\phi_q > 1$ and $\phi_q < 1$, representing the alcohol abuse effect. In the case of absence of alcohol abuse effect in the mathematical model, we assume $\phi_q = \phi_q = 1$.

Poverty and Unemployment

The inclusion of factors such as poverty and unemployment can be applied to several parameters in the model. Galanis and Hanieh (2021) explore the use of these factors by modifying the transmission rate by a linear approximation of

$$\beta(x_1, x_2) = \beta_0 + \beta_1 x_1 + \beta_2 x_2$$

where x_1 and x_2 are the poverty and unemployment rates, respectively.

We can apply this modification to parameters such as α_1 , α_2 , γ_1 and γ_2 , as they can also be influenced by social determinants. Thus, the model with the inclusion of the social determinants mentioned here has the following form:

$$\frac{dS}{dt} = \kappa - \mu S - q(1 - c)\beta(x_1, x_2) \frac{I}{N} S - \theta_1 S + \theta_2 R$$

$$\frac{dI}{dt} = q(1 - c)\beta(x_1, x_2) \frac{I}{N} S + \alpha_1 T - \rho I - \gamma_1 I - \mu I$$

$$\frac{dA}{dt} = \rho I + \alpha_2 T - \gamma_2 A - \delta_1 A - \mu A$$

$$\frac{dT}{dt} = \gamma_1 I + \gamma_2 A - \delta_2 T - \alpha_1 T - \alpha_2 T - \mu T$$

$$\frac{dR}{dt} = \theta_1 S - \theta_2 R - \mu R$$

Discussion

Although many advances have occurred in recent decades in the detection, treatment and prevention of HIV/AIDS in the world, there are still a large number of people at risk of infection or living with HIV without access to care and treatment (HIV.gov, 2021; Pandey and Galvani, 2019). The positive progress in controlling the disease, caused a drop in the number of deaths by 64% since 2010, going from 1.9 million in 2004 to 680,000 people in 2020. However, this progress is uneven within and between countries, varying widely with age, gender and especially economic status of people (UNAIDS, 2021). Thus, the African continent, being the poorest, remains the most severely affected, with almost 1 in every 25 adults (3.6%) living with HIV. The east of the African continent, one of the poorest regions in the world, even comprising only 6.2% of the world population, reports the highest numbers of HIV/AIDS cases in the world, with a prevalence of 54% of the population (Avert, 2019). It is clear that not only biomedical measures are sufficient for the control and total elimination of HIV/AIDS, but the understanding and incorporation of social, cultural and environmental factors play important roles in the dynamics of the disease (Donkoin et al., 2018).

Recognition of the importance of social determinants of health in the distribution and impact of a disease on a given population is a consolidated concept in the literature. Thus, the circumstances in which individuals grow, live and age play as important a role as biological factors in the occurrence of diseases (Lowcock et al., 2012; USDHHS,

2020). The incorporation of these determinants in the design of epidemiological studies, with the development of new methodologies and the adaptation of methods that are already well used, to adapt to the new conditions, is a challenge that researchers have faced. This article aimed to propose a compartmental mathematical model for HIV/AIDS that includes SDH in its structure. The equations presented with the chosen SDH exemplify some approaches that we can adopt when thinking about modeling social effects on the occurrence of HIV. The role of mathematical modeling, in this context, is to support the foundation of the debate on the impact that a certain action or implemented program can have on a population, without losing the effect and influence that social and environmental factors play to the point of compromising the success of a well-planned program (Cassels et al., 2008). Deepening the understanding of current SDH and the search for identification of new factors, as well as a full understanding of the interaction mechanisms, is something that must always be present when conducting studies and interpreting the results. The more factors are added to mathematical models, the more robust and capable of approximating reality they become.

This article has limitations, mainly because it is only a theoretical discussion of mathematical modeling, without data or empirical simulations. However, its innovation lies in the proposition of a new methodological approach and in the development of a model that includes SDH in its differential equations, working as framework and starting point for similar studies. Also, the inclusion of only 5 SDH of the 31 identified by the literature review can be considered a limitation. However, these have been selected based on their strongest effect on HIV/AIDS dynamics, also considering that the inclusion of all determinants in a single model would have made very complex its calibration with real data.

Conclusion

The recognition of the importance of including the SDH in the modeling and studies on HIV/AIDS is evident, due to its complexity and multicausality. Models that do not take into account SDH in their structure, will probably miss a great part of the real trends, especially in periods, as the current one, of strong socioeconomic changes and economic crisis.

List of abbreviations

GTZ: Getting to Zero

IDUs: Injecting drug users

SDH: Social Determinants of Health

WHO: World Health Organization

Declarations

Availability of data and materials

Not applicable.

Ethics declarations

Ethics approval and consent to participate

Note applicable.

Consent for publication

Note applicable.

Competing interests

The authors declare that they have no competing interests.

Funding

This study is funded by the National Institute of Allergy and Infectious Diseases - NAIDS/NIH, Grant Number:1R01AI152938.

Authors' contributions

Robson Oliveira: Investigation, and writing - original draft

Felipe Rubio: Writing - Investigation, and writing - review & editing

Rodrigo Anderle: review & editing

Mauro Sanchez: Writing - review & editing

Luis Eugenio de Souza: Writing - review & editing

Inês Dourado: Writing - review & editing

James Macinko: Writing - review & editing

Davide Rasella: Writing - review & editing, Supervision

Acknowledgements

None

Data availability

All data are within the manuscript and additional files

References

An Q, Prejean J, McDavid Harrison K, Fang X (2013) Association between community Socioeconomic position and HIV diagnosis rate among adults and adolescents in the United

Avendano M, Kawachi I. Why do Americans have shorter life expectancy and worse health than do people in other high-income countries? *Annu Rev Public Health*. 2014;35:307-25. doi: 10.1146/annurev-publhealth-032013-182411. Epub

Avert. Funding for HIV and AIDS. Avert, March 19, 2019. <https://www.avert.org/professionals/hiv-around-world/global-response/funding>. (accessed 10 November 2021).

Buot ML, Docena JP, Ratemo BK, Bittner MJ, Burlew JT, Nuritdinov AR, Robbins JR. Beyond race and place: distal sociological determinants of HIV disparities. *PLoS One*. 2014 Apr 17;9(4):e91711. doi: 10.1371/journal.pone.0091711. PMID: 24743728; PMCID: PMC3990614.

Burattini, MN, Massad E, Coutinho FAB, Azevedo-Neto RS, Menezes RX, Lopes LF, A mathematical model of the impact of crack-cocaine use on the prevalence of HIV/AIDS among drug users, *Mathematical and Computer Modelling*, Volume 28, Issue 3, 1998, Pages 21-29, ISSN 0895-7177, doi.org/10.1016/S0895-7177(98)00095-8.

Burch LS, Smith CJ, Anderson J, Sherr L, Rodger AJ, O'Connell R, Geretti AM, Gilson R, Fisher M, Elford J, Jones M, Collins S, Azad Y, Phillips AN, Speakman A, Johnson MA, Lampe FC. Socioeconomic status and treatment outcomes for individuals with HIV on antiretroviral treatment in the UK: cross-sectional and longitudinal analyses. *Lancet*

Public Health. 2016 Nov;1(1):e26-e36. doi: 10.1016/S2468-2667(16)30002-0. PMID: 28299369; PMCID: PMC5341147.

Cassels S, Clark SJ, Morris M. Mathematical models for HIV transmission dynamics: tools for social and behavioral science research. *J Acquir Immune Defic Syndr*. 2008;47 Suppl 1(Suppl 1):S34-S39. doi:10.1097/QAI.0b013e3181605da3

Cassels S, Goodreau SM. Interaction of mathematical modeling and social and behavioral HIV/AIDS research. *Curr Opin HIV AIDS*. 2011;6(2):119-123. doi:10.1097/COH.0b013e328343acad

Costa JM, Torres TS, Coelho LE, Luz PM. Adherence to antiretroviral therapy for HIV/AIDS in Latin America and the Caribbean: Systematic review and meta-analysis. *J Int AIDS Soc*. 2018 Jan;21(1):e25066. doi: 10.1002/jia2.25066. PMID: 29356390; PMCID: PMC5810329

Dean HD, Fenton KA. Addressing Social Determinants of Health in the Prevention and Control of HIV/AIDS, Viral Hepatitis, Sexually Transmitted Infections, and Tuberculosis. *Public Health Reports*. 2010;125(4_suppl):1-5. doi:10.1177/00333549101250S401

Degenhardt L, Peacock A, Colledge S, Leung J, Grebely J, Vickerman P, Stone J, Cunningham EB, Trickey A, Dumchev K, Lynskey M, Griffiths P, Mattick RP, Hickman M, Larney S. Global prevalence of injecting drug use and sociodemographic characteristics and prevalence of HIV, HBV, and HCV in people who inject drugs: a multistage systematic review. *Lancet Glob Health*. 2017 Dec;5(12):e1192-e1207. doi: 10.1016/S2214-109X(17)30375-3. Epub 2017 Oct 23. Erratum in: *Lancet Glob Health*. 2017 Nov 15;: PMID: 29074409; PMCID: PMC5683738.

Diclementi, J.D., Ross, M.K., Mallo, C., & Johnson, S. (2004). Predictors of successful return to work from HIV-related disability. *Journal of HIV/AIDS and Social Services*, 3 (3), 89-96.

Donkin A, Goldblatt P, Allen J, et al Global action on the social determinants of health *BMJ Global Health* 2018;3:e000603.

Eisinger RW, Folkers GK, Fauci AS. Ending the Human Immunodeficiency Virus Pandemic: Optimizing the Prevention and Treatment Toolkits. *Clin Infect Dis*. 2019 Nov 27;69(12):2212-2217. doi: 10.1093/cid/ciz998. PMID: 31646338; PMCID: PMC7188222.

End Inequalities. End AIDS. GLOBAL COMMITMENTS, LOCAL ACTION After 40 years of AIDS, charting a course to end the pandemic, 2021. https://www.unaids.org/sites/default/files/media_asset/global-commitments-local-action_en.pdf. (accessed 15 November 2021).

Galanis G, Hanieh A. Incorporating Social Determinants of Health into Modelling of COVID-19 and other Infectious Diseases: A Baseline Socio-economic Compartmental Model. *Soc Sci Med*. 2021 Apr;274:113794. doi: 10.1016/j.socscimed.2021.113794. Epub 2021 Feb 23. PMID: 33662772; PMCID: PMC7900756.

Greenhalgh D, Hay G. Mathematical modelling of the spread of HIV/AIDS amongst injecting drug users. *IMA J Math Appl Med Biol*. 1997 Mar;14(1):11-38. PMID: 9080686.

Healthy People. Social Determinants of Health, 2020. <https://www.healthypeople.gov/2020/about/foundation-health-measures/Determinants-of-Health>. (accessed 22 September 2021).

HIV.gov. <https://www.hiv.gov/federal-response/pepfar-global-aids/global-hiv-aids-overview>. (accessed 14 September 2021).

Huo, H. F., Chen, R., & Wang, X. Y. (2016). Modelling and stability of HIV/AIDS epidemic model with treatment. *Applied Mathematical Modelling*, 40(13-14), 6550-6559

International Labour Organization (ILO) .World Employment and Social Outlook – Trends 2021. <https://www.ilo.org/global/research/global-reports/weso/trends2021/lang--en/index.htm>. (accessed 22 September 2021).

Jin RL, Shah CP, Svoboda TJ. The impact of unemployment on health: a review of the evidence. *CMAJ*. 1995 Sep 1;153(5):529-40. Erratum in: *CMAJ*. 1995 Dec 1;153(11):1567-8. PMID: 7641151; PMCID: PMC1487417.

Joy R, Druyts EF, Brandson EK, Lima VD, Rustad CA, Zhang W, Wood E, Montaner JS, Hogg RS. Impact of neighborhood-level socioeconomic status on HIV disease progression in a universal health care setting. *J Acquir Immune Defic Syndr*. 2008 Apr 1;47(4):500-5. doi: 10.1097/QAI.0b013e3181648dfd. PMID: 18197117.

Kasl, S. V., Rodriguez, E., & Lasch, K. E. (1998). The impact of unemployment on health and well-being. In B. P. Dohrenwend (Ed.), *Adversity, stress, and psychopathology* (pp. 111–131). Oxford University Press.

Kirby D, Obasi A, Laris BA. The effectiveness of sex education and HIV education interventions in schools in developing countries. *World Health Organ Tech Rep Ser*. 2006;938:103-50; discussion 317-41. PMID: 16921919.

Leport, C., Guégan, J., Zylberman, P., Bitar, D., Bricaire, F., Cavallo, J., Eliasiewicz, M., Moatti, J., 2012. Proceedings of the seminar on Emerging Infectious

Diseases, November 9, 2011: Current trends and proposals. *Médecine et Maladies Infectieuses* (In press).

Lowcock EC, Rosella LC, Foisy J, McGeer A, Crowcroft N. The social determinants of health and pandemic H1N1 2009 influenza severity. *Am J Public Health*. (2012) 102:e51–8. doi: 10.2105/AJPH.2012.300814

Maruthappu M, Zhou C, Williams C, Zeltner T, Atun R. Unemployment and HIV mortality in the countries of the Organisation for Economic Co-operation and Development: 1981-2009. *JRSM Open*. 2017 Jul 6;8(7):2054270416685206. doi: 10.1177/2054270416685206. PMID: 28748096; PMCID: PMC5507389.

McMahan LD, Lombe M, Evans CBR, Enelamah NV, Chu Y, Simms S, Verkamp-Ruthven J, Martinez JR, Mweemba O, LaForest L, Weiss DJ, Wideman L. Getting to zero HIV/AIDS in sub-Saharan Africa: Understanding perceptions of locals using the social determinants of health framework. *Health Soc Care Community*. 2021 May 24. doi: 10.1111/hsc.13444. Epub ahead of print. PMID: 34028915.

Nyabadza, F. (2006). A mathematical model for combating HIV/AIDS in Southern Africa: will multiple strategies work?. *Journal of Biological Systems*, 14(03), 357-372.

Pandey A, Galvani AP. The global burden of HIV and prospects for control. *Lancet HIV*. 2019 Dec;6(12):e809-e811. doi: 10.1016/S2352-3018(19)30230-9. Epub 2019 Aug 19. PMID: 31439533.

Pence BW, Thielman NM, Whetten K, Ostermann J, Kumar V, Mugavero MJ. Coping strategies and patterns of alcohol and drug use among HIV-infected patients in the United States Southeast. *AIDS Patient Care STDS*. 2008;22(11):869-877. doi:10.1089/apc.2008.0022

Pettifor A, MacPhail C, Hughes JP, Selin A, Wang J, Gómez-Olivé FX, Eshleman SH, Wagner RG, Mabuza W, Khoza N, Suchindran C, Mokoena I, Twine R, Andrew P, Townley E, Laeyendecker O, Agyei Y, Tollman S, Kahn K. The effect of a conditional cash transfer on HIV incidence in young women in rural South Africa (HPTN 068): a phase 3, randomised controlled trial. *Lancet Glob Health*. 2016 Dec;4(12):e978-e988. doi: 10.1016/S2214-109X(16)30253-4. Epub 2016 Nov 1. Erratum in: *Lancet Glob Health*. 2017 Feb;5(2):e146. PMID: 27815148; PMCID: PMC5626439.

Pinkerton SD, Abramson PR. Effectiveness of condoms in preventing HIV transmission. *Soc Sci Med*. 1997 May;44(9):1303-12. doi: 10.1016/s0277-9536(96)00258-4. PMID: 9141163.

Statista. Number of unemployed persons worldwide from 1991 to 2021 and projections until 2022. <https://www.statista.com/statistics/266414/unemployed-persons-worldwide/> 2021. (accessed 13 March 2003).

The Lancet Public Health. Education: a neglected social determinant of health. *Lancet Public Health*. 2020;5(7):e361. doi:10.1016/S2468-2667(20)30144-4

Tolles J, Luong T. Modeling Epidemics With Compartmental Models. *JAMA*. 2020;323(24):2515–2516. doi:10.1001/jama.2020.8420

UNAIDS 2021 - UNAIDS (2021) [AIDSinfo.unaids.org](https://aidsinfo.unaids.org)

UNAIDS 2021. FACT SHEET – WORLD AIDS DAY 2021. https://www.unaids.org/sites/default/files/media_asset/UNAIDS_FactSheet_en.pdf. (accessed 05 November 2021).

UNAIDS. 90-90-90: an ambitious treatment target to help end the AIDS epidemic. Geneva, Switzerland: Joint United Nations Programme on HIV/AIDS (UNAIDS), 2014a Contract No.: JC2684.

UNAIDS. Fast-Track: ending the AIDS epidemic by 2030. Geneva, Switzerland: Joint United Nations Programme on HIV/AIDS (UNAIDS), 2014b Contract No.: JC2686.

UNAIDS. GUIDANCE UNAIDS TERMINOLOGY GUIDELINES, 2015. https://www.unaids.org/sites/default/files/media_asset/2015_terminology_guidelines_en.pdf (accessed 05 November 2021).

UNAIDS. Seizing the moment: tackling entrenched inequalities to end epidemics. Geneva, Switzerland; 2020. <https://www.philstar.com/opinion/2020/07/09/2026614/seizing-moment-tackling-entrenched-inequalities-end-epidemics> (accessed 10 November 2021).

UNESCO & WHO. (2005, November 22-23). HIV and AIDS treatment education. Technical Consultation Report. UNESCO: Paris. Retrieved August 11, 2011, from <http://unesdoc.unesco.org/images/0014/001461/146120e.pdf> (accessed 05 November 2021).

USDHHS. Social Determinants of Health. Washington, DC (2020).

Verma BI, Ray SK, Srivastava RN. Mathematical models and their applications in medicine and health. Health Popul Perspect Issues. 1981 Jan-Mar;4(1):42-58. PMID: 10260952.

Weller S, Davis K. Condom effectiveness in reducing heterosexual HIV transmission. Cochrane Database Syst Rev. 2002;(1):CD003255. doi: 10.1002/14651858.CD003255. PMID: 11869658.

WHO 2020. The top 10 causes of death. <https://www.who.int/news-room/fact-sheets/detail/the-top-10-causes-of-death>. (accessed 03 November 2021).

World Health Organization (WHO), Commission on Social Determinants of Health. Closing the gap in a generation: Health equity through action on the social determinants of health. Final report of the Commission on Social Determinants of Health. Geneva: WHO; 2008.