“Building Palliative Care Capacity in Cancer Treatment Centres: A Participatory Action Research”

Sushma Bhatnagar (✉ sushmabhatnagar1@gmail.com)

Institute Rotary Cancer Hospital, All India Institute of Medical Sciences

Seema Rajesh Rao

Bangalore Hospice Trust – Karunashraya

Naveen Salins

Manipal Academy of Higher Education

Cynthia Ruth Goh

Duke NUS Graduate Medical School

Research Article

Keywords: Capacity-building, palliative care, low-resource setting, participatory action research

Posted Date: March 7th, 2022

DOI: <https://doi.org/10.21203/rs.3.rs-1369490/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Introduction:

There is a significant lack of palliative care access and service delivery in the Indian cancer institutes. In this paper, we describe the development, implementation, and evaluation of a palliative care capacity-building program in Indian cancer institutes.

Methods:

Participatory action research method was used to develop, implement and evaluate the outcomes of the palliative care capacity-building program. Participants were healthcare practitioners from various cancer institutes in India. Researchers developed interventions towards capacity building, which was modified and further developed after each cycle of the capacity-building program. Qualitative content analysis was used to develop an action plan to build capacity. Descriptive statistics were used to measure the outcomes of the action plan.

Results:

Seventy-three healthcare practitioners from 31 cancer treatment centers in India were purposively recruited between 2016 and 2020. Training and education in palliative care, infrastructure for palliative care provision, and opioid availability were identified as key requisites for capacity-building. Individual and organizational change champions were identified for context-specific capacity-building. A training program was developed and delivered to the change champions for knowledge- and skill-development and peer-mentorship for knowledge-translation. The outcome indicators of the project were defined a priori, and were audited by an independent auditor. The three cycles of the program resulted in the development of palliative care services in 23 of the 31 institutes enrolled in the program.

Conclusion:

This program illustrates how a transformational change at the organizational and individual level can lead to the development of sustained provision of palliative care services in cancer institutes.

Introduction

It is estimated that about 2.25 million people are living with cancer in India, with onemillion new cases every year, and over 0.88 million deaths annually[1]. A majority of them present with advanced metastatic disease, experience moderate-to-severe pain, andrequire palliative care [2-4]. The modified National Cancer Control Program of India emphasized the need palliative care at the primary care level [5], and led to the establishment of outpatient pain clinics in cancer centers, government, and private hospitals, stand-alone hospices, outreach clinics, and homecare services[6]. However, the Narcotic Drugs and Psychotropic Substances Act (NDPS) that restricted access to and availability of opioids was a major

barrier for pain relief [7]. The NDPS Act was amended in 2014 to make opioids more accessible, but lacked effective implementation[8].

Every year over seven million new patients need palliative care in India, with less than 4% having access to these services[9, 10]. Lack of access to palliative care results in poor symptom control, poor quality of life, inappropriate end-of-life care and increased economic burden[11]. Over 3.5-6.2% of the population in India become poorer every year due to enhanced health expenditure at end of life[12]. Evidence indicates that referral to palliative care results in reduced healthcare spending in patients with cancer and other chronic illnesses [11].

Palliative care activities in India have been on-going for three decades from 1980s. The National Programmed for Palliative Care (NPPC) was launched in 2012 [13]. Lack of budget allocation, provider awareness, education and employment opportunities, difficulty in accessing opioids, and absence of legal framework or policies regarding end-of-life-care has impacted timely and effective implementation of the NPPC[14]. A recent country-wide survey of the National Cancer Grid (NCG)[15] centers in India highlighted poor integration of palliative care in oncology[16]. Although India has capacity for generalized palliative care provision as per the Global Atlas of Palliative Care[17], many parts of rural India still have limited access[13]. Bridging the gaps in the capacity to provide palliative care that is cost-effective and equitable necessitates development of these services within the institutions[16]. A national level Cancer Treatment Centers Palliative Care (CTC) program was conceptualized to bridge this gap. In this paper we describe how we used Participatory Action Research (PAR) to design, implement, and evaluate the development of palliative care services in cancer centers in India.

This study aimed to develop palliative care services in cancer centers in India by recognizing receptive organizations and individuals, identifying facilitators and barriers for development of palliative care services, formulating strategies to overcome the constraints, and creating mechanisms to assess outcomes of program implementation.

Methods

Participatory action research was used to develop a program to build capacity to provide palliative care in cancer treatment centers in India. In PAR, groups of individuals work together to bring about a change in social or institutional practices [18]. It is a value based, action-oriented, and participatory research [19]. The values of the researcher and the participants inform and drive the research, which leads to an action that brings about the desired change, through a collective process of knowledge generation [20]. Participatory action research proceeds in cyclical stages where the learning of each cycle informs the next [21–23].

The participatory action research was conducted over three cycles from 2015 to 2020 in India. Each cycle had four stages that involved a) developing a critically-informed action plan for social change, b) selecting and implementing the action plan, c) observing the consequences of the actions (evaluation), and d) reflective learning and taking corrective actions. This program was a collaborative partnership

between an international palliative care organization and two university teaching hospitals with specialized palliative care units.

Purposive sampling was used to select participants for this program. Medical institutes providing oncology services, both public and private, that were receptive for palliative care, from the states and Union Territories of India where palliative care was underdeveloped or absent were invited to participate in the program [24]. Healthcare practitioners (doctors and nurses) employed in the cancer treatment institutes, willing to take part in the training, ongoing mentoring, and audit were recruited. The three cycles of the CTC program enrolled 54 doctors and 52 nurses from 31 cancer treatment institutes. The physician participants were from the specialties of anesthesia, oncology, internal medicine, psychiatry, and critical care. Nursing participants were from both general and specialty nursing pools.

Multiple sources of data like survey findings, transcripts of focus group discussion, mentor visit notes, and audit findings enabled methodological triangulation and provided rich in-depth information for analysis. Descriptive statistics was used to analyze quantitative data and content analysis to analyze qualitative data. This study was approved by the Institutional Ethics Committee of Kasturba Medical College and Kasturba Hospital, IEC No: 330/2021

Developing And Implementing The Intervention:

First Cycle:

Plan:

A rapid review of literature identified the barriers and enablers to palliative care provision in oncology settings in India [16, 25]. In addition to the challenges posed by population, geographic density and poverty, lack of institutional interest, restrictive opioid policies, poor workforce development, poor implementation and utilization of available resources for palliative care were identified as major barriers [25]. A core group of national and international palliative care experts participated in a focus group discussion (FGD) to develop a strategy that would enable development of palliative care services in the cancer treatment centers in India [26]. Two experts from the two university teaching hospitals facilitated this FGD. Eight palliative care experts in leadership positions responded to a voluntary call and consented to be part of the FGD. The findings of the rapid review informed the discussion. This group of experts deliberated on the processes needed to build capacity for palliative care in cancer treatment centers in India. Data from the FGD was collected and analyzed from the moderator notes, recorded conversations, and memory.

The barriers for capacity building identified in FGD included lack of knowledge and skills in palliative care, lack of opioid access, and poor implementation and utilization of available resources. The FGD also deliberated on the strategies for overcoming these barriers. This provided the framework for the CTC

palliative care program which was structured focusing on the three components of the WHO Public Health Model [27], education, drug availability and implementation.

Epidemiological and behavioral studies have shown that the critical number of personnel needed to bring about a social or organizational change is 25–30% [28]. When this tipping point is reached, new behaviors are supported and change is inevitable, self-sustaining, and fuels further growth [28]. The FGD deliberated on the critical number needed to bring about this change in India. It was estimated that by establishing palliative care services in at least 100 of the 327 cancer treatment institutes in India, this tipping point would be achievable.

Developing capacity to provide palliative care in an oncology setting requires organizational change as well as individual technical expertise [29]. Organizational climate can either facilitate or impede efforts at capacity building [30]. The FGD deliberated on a bottoms-up approach (individual skill building) and top-down approach (changing agency-specific policies and practices, buy-in from organizational leaders) for sustainable change [31].

Transformative adult learning is enhanced through critical incident analysis, small group discussions, reflective practices, and clinical immersion rotations [32]. This program utilized a workshop model for knowledge development along with problem-based learning thereby enabling improvements in individual knowledge, skills and behavior [33]. This program fostered experiential learning through real-life or simulated scenarios to enhance critical thinking, leadership, team building and collaboration, perspective transformation, and change management competencies [34].

Despite adequate training, the implementation of clinical practices and culture change is a slow and disorganized process and many patients remain deprived of high-quality care that is recommended by the guidelines [35]. Knowledge translation strategies help in addressing this gap. Studies in healthcare sector have highlighted the role of mentorship in improving leadership, management and clinical competencies among the healthcare workers in low-and-middle income countries [36, 37], while being cost-effective, and context-specific [29, 36, 38]. Site-based mentoring was utilized for academic detailing to identify site-specific policies and practices that impede or facilitate organizational change. The FGD also identified the outcome indicators for measuring program implementation. The elements of the FGD are outlined in Table 1.

Table 1
Focus Group Discussion Results

Probes	Response	Recommendation
How can we improve capacity for palliative care within oncology institutes?	Identify change champions for palliative care	<ul style="list-style-type: none"> • A team of 2 doctors and 2 nurses from each oncology institute to drive the organizational change
	Improve Infrastructure	<ul style="list-style-type: none"> • Stand-alone outpatient palliative care department
	Space, staff, time, equipment	<ul style="list-style-type: none"> • Consultation liaison service for inpatients • Task shifting to overcome health workforce shortage • Prevent task overload
	Increase access to opioids	<ul style="list-style-type: none"> • Procure license to store and dispense opioids • Educate regarding safe practices • Ensure uninterrupted supply
How can we improve individual capacity to provide palliative care in oncology treatment institutes?	Initiate advocacy activities to raise awareness about PC	<ul style="list-style-type: none"> • Develop competency for in-house training • Conduct continuing medical education programs with external support
	<ul style="list-style-type: none"> • Improve knowledge and skills about palliative care • Help in knowledge translation 	<ul style="list-style-type: none"> • Attend and complete recommended training in palliative care • Mentoring activities • Academic detailing • Develop institutional policies and guidelines
What are the outcome indicators to measure the implementation of the program?	For organizational capacity building	<ul style="list-style-type: none"> • Number of dedicated staff for palliative care • Number of hours of OPD per month • Number of patients seen in OPD per month • Number of patients seen in CL per month • Number of new patients referred to palliative care

Probes	Response	Recommendation
	For individual capacity building	<ul style="list-style-type: none"> • 5-day face-to-face training in Palliative Care • 5-day clinical attachment at specialist palliative care institute • 2-day mentorship training by a visiting mentor • Completion of certificate course in essentials of palliative care (CCEPC) by Indian Association of Palliative Care
	For morphine availability	<ul style="list-style-type: none"> • Number of milligrams of morphine use per month
	For advocacy activities	<ul style="list-style-type: none"> • Number of trainings conducted in a year • Number of doctors trained • Number of nurses trained • Number of allied healthcare practitioners trained • Observance activities for World Hospice and Palliative Care Day in the institute

The first cycle was initiated in January 2016. Thirty healthcare practitioners (16 nurses and 14 doctors) from 10 cancer treatment institutes participated in the training program. The duration of the first cycle was 24 months.

Act:

In this stage all 30 participants were provided training in palliative care. All participants underwent a 5-day residential face-to-face training program along with a 5-day clinical attachment at a specialist palliative care institute. A senior palliative care expert mentored the palliative care activities at each participating institute. This stage took four months. Each team on completion of the training initiated palliative care services within their respective institute. Organizational restructuring and task-shifting helped institutes overcome the workforce shortage in palliative care. Physical space for an outpatient department was identified and established. Healthcare personnel trained in the CTC program staffed the palliative care OPD, initially part-time. Consultation liaison services were established for inpatient care. Engaging with senior healthcare leadership and opinion leaders for improving access to opioids and ensuring uninterrupted supply was one of the key focuses of the action plan. In addition, the local change champions organized training activities with external resource personnel for all cadres of healthcare personnel within their institutes thereby increasing the workforce available for palliative care. The action

plan also involved building partnerships by engaging with community and local leaders to advocate for palliative care. This process took about 12 months.

Observation and Evaluation:

Both qualitative and quantitative methods were used to evaluate the action plan, longitudinally at different stages of the cycle. Pre- and post-evaluation questionnaires; survey forms that explored institutional activities like current care practices, organizational culture, barrier and enablers; mentor visit notes; and audit findings were used for evaluation. At the end of each cycle, an external auditor evaluated the implementation of project objectives. Of the 10 cancer treatment institutes enrolled in the CTC program five centers were able to develop capacity to provide palliative care.

Reflections from the first cycle:

The evaluation results of the first cycle were shared with the participants and faculties. The first cycle was followed by group reflective sessions. Rolfe's reflective model was used to better understand the barriers and enablers and modify the action plan for the second cycle accordingly [39]. This process took about six months. Literature has shown that local change champions are crucial in supporting transformative change efforts within an organization [33, 36]. Senior faculties in leadership roles were enrolled into the training program. A 3-day face-to-face centralized refresher course was incorporated into the regular training. The focus on small group problem-based learning (PBL) and peer learning during the refresher course enhanced collaboration and communication within and between members of the group [40]. More structured mentorship visits were initiated.

The barriers and adaptations are outlined in Table 2.

Table 2
Reflections from the first cycle of the CTC Program:

Learnings from the first cycle	Adaptations in the second cycle
Organizational Barriers	
Lack of buy-in from administrators and decision-makers	<ul style="list-style-type: none"> • A more rigorous process of sample selection through interviews and personal judgement • Identifying senior clinical and non-clinical leaders within organizations and engaging them
Workforce shortage to initiate PC service	<ul style="list-style-type: none"> • A 2-day on-site structured mentor visit where mentors addressed site-specific issues with administrators/decision-makers
Lack of resources – space, funding, time	<ul style="list-style-type: none"> • Sensitization and training programs by experts on palliative care
Lack of awareness about PC among other healthcare providers	
Hierarchical structure in the healthcare system that impedes communication and collaboration	<ul style="list-style-type: none"> • Group brainstorming with the team on how to enhance team collaboration and communication • Team building activities during the mentor visit and refresher course
Individual Barriers:	
Lack of motivation towards PC	<ul style="list-style-type: none"> • Improving selection of change champions by interviews and personal judgment • Selecting those with some awareness, understanding and commitment towards palliative care • Identifying those who have had short-term training in palliative care in the cancer treatment institutes and involving them in the program
<ul style="list-style-type: none"> • Deficits in PC skills and knowledge • Lack of leadership skills 	<ul style="list-style-type: none"> • Structured training program that included a 3-day centralized refresher training which focused on: • Problem-based learning and peer learning techniques to foster a culture of continuous self-directed learning • Microlearning to reinforce previously acquired knowledge and skills, address gaps in knowledge and help in retention
Competing interests of the healthcare provider	<ul style="list-style-type: none"> • Appropriate selection of candidates who would be able to devote exclusive time to PC • Organizational buy-in from administrators ensured to smoothen this transition
Barriers for drug availability	

Learnings from the first cycle	Adaptations in the second cycle
Opioid access and use	<ul style="list-style-type: none"> • Liaising with opinion leaders, administrators, local change champions, and governmental agencies for better opioid access • Training for healthcare staff on safe use of opioids

Second Cycle

The second cycle was initiated in January 2018, with nine cancer treatment institutes and 32 participants; 16 doctors and 16 nurses. The duration of this cycle was 12 months. Palliative care services were established in eight of the nine centers.

Third Cycle

The third cycle was initiated in January 2019, with 12 cancer treatment institutes enrolled in this cycle, with 44 participants; 22 doctors and 22 nurses. The duration of this cycle was 12 months. The growing number of participants and centers needed more continuous engagement of change champions. A full-time PC consultant was appointed to engage and monitor the program activities. Palliative care services were successfully established in 10 of the 12 institutes that had participated in the training.

The three cycles and steps of the research process are depicted in Fig. 1.

Results

The overall impact of the interventions of CTC program is presented in the Table 3.

Table 3
Overview of the CTC Program

SI No	CTC Program	Year	Number of institutes enrolled	Number of centers who completed training	Number of HCP trained in palliative care
1	First cycle	2016–2018	10	5	13
2	Second cycle	2018–2019	9	8	27
3	Third cycle	2019–2020	12	10	33
Total			31	23	73

Outcome Indicators: The outcome of the CTC programs was measured by the indicators that were defined a priori and included development of infrastructure (OPD space, personnel, and time), drug availability and training. These are listed below:

- a) Establishment of Palliative care outpatient services in the cancer treatment institutes: While 12 cancer treatment institutes had independent palliative care outpatient departments before CTC program, the number increased to 23 after CTC training. The number of healthcare providers working either part-time or full-time increased from 36 to 174 after initiation of the program.
- b) Morphine availability and use: While 14 institutes had the license to procure, store, and dispense morphine before the program, it increased to 23 after the CTC program. Total morphine consumption per month in the cancer treatment institutes increased from 413408 mg to 917638 mg, indicating increased availability and use. The use of morphine is depicted in Fig. 2.
- c) Training and Advocacy Activities: None of the centers were engaged in palliative care training or advocacy activities before CTC program. After CTC training 20 of the centers have initiated both in-house training and advocacy activities.
- d) Secondary Outcomes: There were several meaningful outcomes of this program, which were not quantifiable. Palliative care services were established in medical colleges states and Union Territories in India which hitherto had no or minimal access to palliative care. The change champions have assumed leadership roles in developing palliative care policies and have initiated postgraduate training in palliative care.

Discussion

Relief from pain and serious health-related suffering is a human right [24]. The disparity between need and availability of palliative care continues to persist, especially in lower and middle-income countries (LMIC) [9]. The Quality of Death Index published by the Economist Intelligence Unit in 2015 that measured the quality of palliative care around the world, ranked India 67th out of 80 countries in palliative care provision [25]. The barriers to development of palliative care in the oncology setting are complex and traverse multiple domains including policy, education, drug availability, awareness, advocacy, and implementation.

The central aim of PAR is to bring about a social change where groups of individuals work together collaboratively to bring about the change [19]. PAR fosters capacity development in all those who participate in the process. The CTC Program was envisaged to bring about a transformational change in practice in the cancer treatment institutes in India through capacity building for palliative care. This program was able to help 23 cancer treatment institutes build capacity to provide palliative care. The PAR framework aided the development of action plan that adopted a) appropriate context-specific capacity-building strategies [41], b) adult transformative learning for knowledge-sharing and knowledge-translation [34, 42], and c) partnerships between organizations [31].

Research shows that when changes are planned and executed by those within the organization then capacity building is sustainable and successful [31]. The participants in the CTC program were the drivers of change, who identified with the common goal of developing capacity to provide palliative care within

their institutions, and developed an action plan along with the researchers to achieve this goal. The participants and researchers revisited, evaluated, and redefined the outcomes with each cycle. This reflective cycle enabled identification of components that facilitated and impeded palliative care capacity building in the cancer treatment institutes and facilitated corrective actions for the next cycle [39].

In the four-tier hierarchy of capacity building needs, this model focused on the individual and institutional level needs, that is, staff and facilities, skills and tools [41]. To initiate, develop and implement any change you need change champions, both individual and organizational [43]. Identifying and training the change champions was the first step. The champions through advocacy activities in the local, national, and regional level have enhanced palliative care service provision in many states and union territories. Our experience suggests that while the participating institutes were able to bring about a transformative change and establish palliative care services within their institution, they were able to facilitate changes beyond their institution. Some of the participating centers were able to enlist and train other institutions. This dependence on change champions was also responsible for the failure to capacity build in some institutes. When the change champions lacked skills in leadership and teamwork, they were ineffective in starting the services within their institutes. When the change leaders resigned from the institute, the established services collapsed. Developing a structured selection process to identify organizational and individual champions will help in preventing dropouts. Strengthening the fourth pillar of capacity-building, that is, structures, systems and roles will help in overcoming this weakness in the program.

Education and training are important components of capacity building as is knowledge translation [42]. Adult transformative learning strategies were applied in this program [34]. The structured residential training program was as per national standards and aided conceptual learning. The refresher course created the space for the newly formed palliative care teams to share their experiences and aided experiential and peer-assisted learning. The rapid feedback through quizzes enabled participant learning and knowledge durability. It is important for knowledge learned to be translated into practice, that is, the knowledge to action cycle needs to be completed. The peer-based mentorship model helped the individual champions in implementing the learning in their institutes [40].

Partnerships were integral for the implementation of this project. Partnerships with international and national organizations were crucial for this project. The international organization provided resources for the training and mentoring activities. The aim of this initiative was to ensure sustainability of this process. The PAR approach co-opted the participants as partners who brought about a change in the institution's culture and attitude towards palliative care. The buy-in by the institutional and administrative leaders ensured sustainability of the program in the 23 centers without external support. In addition, collaborative partnerships between individual team members and between other teams ensured sustainability. Doctors and nurses from individual institutes worked together for the first time as a team during the CTC training and developed a team and group identity with a shared mission and vision. The interactive environment of the residential face-to-face training aided collaboration, communication and networking between individuals and institutes fostering partnerships.

The experience with the three CTC cycles also showed that what was transformative and worked in one setting is not necessarily relevant and applicable in the other. For the process to be successful more attention needs to be paid to the contextual starting conditions, facilitation skills of the mentors, and the leadership skills of the change champions. Lack of a structured process of selection is one of the limitations of the program. The CTC program has primarily focused on personal and performance capacity without focusing on the structural and system capacity. This is another limitation of the program. The development of the program did not factor in the end user, the patient and the family experiences, and is another major limitation of this program. These would need to be incorporated in the next cycle of the program.

There were methodological limitations in the study. The purposive sampling could have led to selection bias, choosing those centers where programs were likely to succeed. The data collection was achieved through pre-existing self-reported data, survey questionnaires, field notes, and through interviews which could have led to response and researcher bias. However, these were mitigated through methodological and data triangulation.

The purpose of the CTC program was to increase the capacity to provide high quality palliative care to patients with terminal illnesses. A service evaluation of the project against national standards with validated evaluation tools, while incorporating patient and provider experience will help us to define the effectiveness of the project

Conclusion

Palliative care needs to be incorporated into the routine care of patients with cancer and other terminal illnesses. The challenges in implementing this can be mitigated if we develop a sustainable working model to build capacity to provide palliative care. The CTC program focused on capacity building, knowledge development and translation and partnership between national and international palliative care organizations, individual and organizational stakeholders, and change champions. This program aimed at bringing about a transformational change at the organizational as well as at an individual level, led to development of palliative care services in cancer treatment institutes in India. Developing the institution-based generalist palliative care model in cancer institutes might enhance capacity to provide palliative care in India and might bridge the healthcare inequities related to palliative care access.

Declarations

Acknowledgements:

The authors would like acknowledge the contributions of the following palliative care experts involved in the CTC program training and mentoring: Dr. Anjum Khan Joad, Dr. Charu Singh, Dr. Jeremy Johnson, Dr. Mayank Gupta, Dr. Priyadarshini Kulkarni, Dr. Raghavendra Ramanjulu, Dr. Roopesh Jain, Dr. Santosh Chaturvedi, Dr. Savita Butola, Dr. Saurabh Joshi, Dr. Shiv Pratap Singh Rana, Dr. Vidya Viswanath, Ms.

HanifeMcGamwell, Ms. Alice Stella, Ms. Govindi Chauhan, and Ms. Nileema Sharad Singhade. We would also like to acknowledge Mr. Giam Cheong Leong, Executive Director, Asia Pacific Hospice Palliative Care Network for the organizational support.

Funding:

No research grants or research funds were received for preparing the manuscript or conducting the study.

Competing Interests:

The authors have no relevant financial or non-financial interests to disclose

Author Contributions:

All authors contributed to the study conception and design. All authors participated in the material preparation, data collection, analysis, and manuscript writing. All authors have read and approved the final manuscript.

Ethics Approval:

The study was approved by the Institutional Ethics Committee, of Kasturba Medical College and Kasturba Hospital IEC No: 330/2021. The study was carried out in accordance with the ethical guidelines and regulations of Institutional Ethics Committee of Kasturba Medical College and Kasturba Hospital, Manipal Academy of Higher Education (Deemed University). Informed consent was obtained from the study participants. Data confidentiality was strictly maintained and all data was anonymized. Only the study investigators had access to the data. Data was stored in password protected computers.

Consent to participate:

The authors confirm that written informed consent was obtained from all participants.

Consent for publication:

Not applicable

Data Material and Availability:

The datasets used and analysed for this study can be accessed through the Kasturba Medical College Research Cell. It is available from the corresponding author on reasonable request at the following email: sushmabhatnagar1@gmail.com.

References

1. Dhillon PK, Mathur P, Nandakumar A, Fitzmaurice C, Kumar GA, Mehrotra R, et al. The burden of cancers and their variations across the states of India: the Global Burden of Disease Study 1990–

2016. *Lancet Oncol.* 2018;
2. Chatterjee S, Chattopadhyay A, Senapati S, Samanta D, Elliot L, Loomis D, et al. Cancer Registration in India - Current Scenario and Future Perspectives. *Asian Pac J Cancer Prev.* 2016;17(8):3687–96.
3. Centeno C, Arias-Casais N. Global palliative care: from need to action. *The Lancet Global Health.* 2019.
4. Swetz KM, Kamal AH. Palliative Care. *Annals of internal medicine.* 2018.
5. Khanna I, Lal A. Palliative Care - An Indian Perspective. *ARC J Public Heal Community Med.* 2016;1(4):27–34.
6. Mohanti BK. Research Focus in Palliative Care. *Indian J Palliat Care.* 2011;17(Suppl):S8–11.
7. Rajagopal MR, Joranson DE. India: Opioid Availability—An Update. *J Pain Symptom Manage.* 2007;33(5):615–22.
8. Jacob A, Mathew A. End-of-Life Care and Opioid Use in India: Challenges and Opportunities. *J Glob Oncol.* 2017;
9. Khosla D, Patel FD, Sharma SC. Palliative care in India: Current progress and future needs. *Indian Journal of Palliative Care.* 2012.
10. Knaul FM, Farmer PE, Krakauer EL, De Lima L, Bhadelia A, Jiang Kwete X, et al. Alleviating the access abyss in palliative care and pain relief—an imperative of universal health coverage: the Lancet Commission report. *Lancet.* 2018;391(10128):1391–454.
11. May, P, Garrido MM, Cassel JB, Kelley AS, Meier DE, Normand C, et al. Palliative Care Teams' Cost-Saving Effect Is Larger For Cancer Patients With Higher Numbers Of Comorbidities. *Health Aff (Millwood).* 2016;35(1):44–53.
12. Hooda SK. Out-of-pocket payments for healthcare in India. *J Health Manag.* 19(1):1–15.
13. Kumar S. Models of delivering palliative and end-of-life care in India. *Current Opinion in Supportive and Palliative Care.* 2013.
14. Salins N, Johnson J, Macaden S. Feasibility and acceptability of implementing the integrated care plan for the dying in the Indian setting: survey of perspectives of Indian palliative care providers. *Indian J Palliat Care.* 2017;23(1):3–12.
15. Pramesh CS, Badwe RA, Sinha RK. The national cancer grid of India. *Indian J Med Paediatr Oncol.* 2014;
16. Damani A, Salins N, Ghoshal A, Chowdhury J, Muckaden MA, Deodhar J, et al. Provision of palliative care in National Cancer Grid treatment centres in India: A cross-sectional gap analysis survey. *BMJ Support Palliat Care.* 2020;
17. Connor S. *Global Atlas of Palliative Care.* 2nd ed. London, UK: World Hospice and Palliative Care Alliance; 2020. 1–119 p.
18. Kemmis S, McTaggart R. Participatory action research: Communicative Action and Public Sphere. In: Denzin NK, Lincoln YS, editors. *The Sage handbook of qualitative research.* 3rd ed. Thousand Oaks, California: SAGE PUBLICATIONS INC; 2005. p. 559–604.

19. Reason P, Bradbury H. Action Research: Participative Inquiry and Practice, Second Edition. Sage Publication Ltd. 2008.
20. Apgar JM, Allen W, Albert J, Douthwaite B, Paz Ybarnegaray R, Lunda J. Getting beneath the surface in program planning, monitoring and evaluation: Learning from use of participatory action research and theory of change in the CGIAR Research Program on Aquatic Agricultural Systems. Action Res. 2017;15(1):15–34.
21. Gillis A, Jackson W. Action Research. In: Research methods for nurses: Methods and interpretation. Philadelphia: F. A. Davis Company; 2002. p. 264–9.
22. MacDonald C. Canadian journal of action research CJAR. Can J Action Res. 2012;13(2):34–50.
23. Maguire P. Doing participatory action research: A feminist approach. Massachusetts: University of Massachusetts Press; 1987.
24. McDermott E, Selman L, Wright M, Clark D. Hospice and Palliative Care Development in India: A Multimethod Review of Services and Experiences. J Pain Symptom Manage. 2008;35(6):583–93.
25. Bag S, Mohanty S, Deep N, Salins N, Bag S. Palliative and End of Life Care in India - Current Scenario and the Way Forward. J Assoc Physicians India. 2020;68(11):61–5.
26. Bhatnagar S. Institute-Based Sustainable Palliative Care Program: A Unique Initiative of “Cancer Treatment Center Program”. Indian J Palliat Care. 2019;25(4):485–6.
27. Callaway M V, Connor SR, Foley KM. World Health Organization Public Health Model: A Roadmap for Palliative Care Development. J Pain Symptom Manage. 2018;55(2):S6–13.
28. Centola D, Becker J, Brackbill D, Baronchelli A. Experimental evidence for tipping points in social convention. Science (80-). 2018;360(6393):1116–9.
29. Fassbender K, Watanabe SM. Early palliative care and its translation into oncology practice in Canada: barriers and challenges. Annals of palliative medicine. 2015.
30. Danley KS, Ellison ML. An Introduction to Participatory Action Research. In: A handbook for participatory action researchers. 1st ed. Boston, MA: Implementation Science and Practice Advances Research Center Publications; 1999. p. 4.
31. Crisp BR, Swerissen H, Duckett SJ. Four approaches to capacity building in health: Consequences for measurement and accountability. Health Promot Int. 2000;15(2):99–107.
32. Mezirow J. Perspective transformation. Adult Educ Q. 1978;28(2):100–10.
33. Schwerdtle P, Morphet J, Hall H. A scoping review of mentorship of health personnel to improve the quality of health care in low and middle-income countries. Global Health. 2017;13(1):77.
34. Chacko T. Emerging pedagogies for effective adult learning: From andragogy to heutagogy. Arch Med Heal Sci. 2018;6(2):278–83.
35. Eccles MP, Armstrong D, Baker R, Cleary K, Davies H, Davies S, et al. An implementation research agenda. Implement Sci. 2009;4(18).
36. Vasan A, Mabey DC, Chaudhri S, Epstein HAB, Lawn SD. Support and performance improvement for primary health care workers in low- and middleincome countries: A scoping review of intervention

- design and methods. *Health Policy Plan.* 2017;32(3):437–52.
37. Belrhiti Z, Booth A, Marchal B, Verstraeten R. To what extent do site-based training, mentoring, and operational research improve district health system management and leadership in low- and middle-income countries: A systematic review protocol. *Syst Rev.* 2016;
 38. McCloughen A, O'Brien L, Jackson D. Positioning mentorship within Australian nursing contexts: a literature review. *Contemp nurse a J Aust Nurs Prof.* 2006;23(1):120–34.
 39. Rolfe G. Reflective practice: Where now? *Nurse Educ Pract.* 2002;2(1):21–9.
 40. Ghani ASA, Rahim AFA, Yusoff MSB, Hadie SNH. Effective Learning Behavior in Problem-Based Learning: a Scoping Review. *Med Sci Educ.* 2021;31(1):1–13.
 41. Potter C, Brough R. Systemic capacity building: A hierarchy of needs. *Health Policy Plan.* 2004;19(5):336–45.
 42. Gagliardi AR, Perrier L, Webster F, Leslie K, Bell M, Levinson W, et al. Exploring mentorship as a strategy to build capacity for knowledge translation research and practice: Protocol for a qualitative study. *Implement Sci.* 2009;4(55).
 43. Shaw EK, Howard J, West DR, Crabtree BF, Nease DE, Tutt B, et al. The role of the champion in primary care change efforts. *J Am Board Fam Med.* 2012;25(5):676–85.

Figures

Figure 1

Participatory Action Research Cycle of the Cancer Treatment Centers Program

Figure 2

Comparison on morphine consumption in CTC centers