Temporal Analysis for Multi-hazard Risk Assessment of Rice Cultivation in Coastal Areas: A Case Study of Soc Trang, Vietnam

Thanh Quoc Vo

IHE Delft Institute for Water Education <https://orcid.org/0000-0001-5940-1798>

Trung Hieu Nguyen (✉ nhtrung@ctu.edu.vn)

Research Institute for Climate Change, Can Tho University, Vietnam

Vo Thi Phuong Linh

Can Tho University

Research

Keywords: risk assessment, tropical cyclones, salinity, Mekong Delta, rice cultivation

Posted Date: January 6th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-137318/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Temporal analysis for multi-hazard risk assessment of rice cultivation in coastal areas: a case study of Soc Trang, Vietnam

Vo Quoc Thanh¹, Nguyen Hieu Trung^{2*}, Vo Thi Phuong Linh¹

¹ College of Environment and Natural Resources, Can Tho University, Vietnam

² Research Institute for Climate Change, Can Tho University, Vietnam

*Correspondence: nhtrung@ctu.edu.vn

Abstract

Rice is an important human crop and rice cultivation is threatened due to natural disasters, leading to negative effects on national and global food security. The natural disasters, such as tropical cyclones and saline intrusion, have dramatic influences in coastal regions. To investigate possible impacts of these disasters on rice cultivation, it needs an efficient tool to assess potential disasters impacts and a risk index is highly applicable. Therefore, this study aims at establishing a risk assessment of rice production in coastal areas under effects of tropical cyclones and saline intrusion. We adopted risk definition introduced by IPCC (2014) in which risk is a function of hazard, exposure and vulnerability. Multiple hazards of tropical cyclones and saline intrusion were indicated by their frequency and severity at some critical levels of 25%, 50% and >50% rice yield reduction. Each hazard was weighted by its damage on rice yield. Exposure and vulnerability of rice crops are evaluated at different growing phases. Tropical cyclone hazard index was ranked *high* and *very high* in the wet season while salinity hazard index was ranked *very high* in the dry season. Due to the combined effects of tropical cyclones and salinity, rice crop is highly susceptible during the reproduction phase and at the panicle initiation stage particularly. Based on the cropping calendar of My Xuyen, the period of October-November was the *very high* vulnerability period since it had the largest rice cultivable area and rice crops were at the reproduction phase. This result shows that rice crops are at *high* risk in October and November. Noticeably, saline intrusion reaches the highest level in April and May, but no risk is at this period because of no rice crop cultivated. This can reflect a measure to reduce risk by adjusting the cropping calendar.

Keywords: risk assessment, tropical cyclones, salinity, Mekong Delta, rice cultivation.

1. Introduction

According to FAO (2017) natural disasters caused damages of agricultural productions, including crops and livestock, about USD 93 billion in developing countries in a decade (2005-2014). Among these natural disasters, tropical cyclones are one of the main disastrous phenomena causing considerable damages around the world which cause a damage of USD 26 billion annually (Mendelsohn et al. 2012). In addition, there is an annual loss of approximately USD 12 billion due to saline-affected land (Qadir et al. 2008). The mentioned damaged phenomena are the major factors affecting coastal areas.

The Vietnamese Mekong Delta (VMD) is located at the downstream of the Mekong River which is one of the largest rivers in the world (MRC 2010). The VMD is a crucial economic region in Vietnam. Moreover, it plays an important role in ensuring food security not only for Vietnam, but also for the world since the VMD contributes up to 90% to the annual rice export of Vietnam (GSOVN 2010). With the land resources of approximately 4 million ha, three-quarters of this area is used for agricultural production (Kakonen 2008). Therefore, rice production in the coastal VMD is facing challenges of tropical cyclones and saline intrusion. For example, Soc Trang is a coastal province, dealing with these challenges. Although Soc Trang is a coastal province, rice is the major agricultural products. Consequently, rice cultivation in Soc Trang is at risk of tropical cyclones and saline intrusion effects. Generally, to reduce risk of these events, it needs to conduct a risk assessment to evaluate how rice crops can be damaged. Risk is resulted from interaction of hazard and vulnerability (Merz et al. 2010). In order to assess risk, it requires hazard and vulnerability assessment. The outputs of these assessment support decision making

process.

There are some methods to assess hazard and vulnerability using indicators, field or lab experiments and numerical models (Buan et al. 1996; Zeng and Shannon 2000; Zeng et al. 2001; Hakim et al. 2010; Bauder et al. 2011; Kunitsugu 2012; Lee et al. 2012; Masutomi et al. 2012; Tri et al. 2013; Bates et al. 2014; Esteban et al. 2014; IRRI 2015; Kakar et al. 2019). In comparison, using indicators is a rapid and simple method compared to the others mentioned. For instance, Balica et al. (2012) developed a coastal city flood vulnerability index which is able to assess vulnerability to flood and evaluate adaptation options. Tri, Trung, and Thanh (2013) applied this index to evaluate impacts of the projected climate change and sea level rise on flood vulnerability in the VMD. It is determined that vulnerability index is an effective and rapid tool to assist managers and decision-makers in assessing impacts of predicted scenarios and improving decision-making procedure. The outputs resulted in applying this index enhance mitigation and adaptation strategies. Therefore, this study is conducted to establish a risk assessment of rice production due to effects of tropical cyclones and saline intrusion. We assessed impacts of tropical cyclones and saline intrusion on rice production in a coastal area (My Xuyen district, Soc Trang province, Vietnam, presented in Figure 1) by using the risk definition of IPCC (2014). Based on this definition, risk includes hazard, exposure and vulnerability components. Rice is an important food crop and is grown seasonally, so rice crops are not always available on the field. This suggests that a temporal analysis should be taken into account for the risk assessment. The outputs of the risk assessment helps to mitigate and reduce impacts of tropical cyclones and saline intrusion on rice cultivation.

Figure 1. Land-use map in 2010 of My Xuyen district.

2. Methods

2.1. Risk definition

To identify the natural hazard risk, we used the definition of IPCC (2014) in which risk is defined as a function of hazards and consequences which present the impacts of the hazardous events. In a hazardous event, its impacts are highly depended on vulnerability and exposure. Therefore, risk is defined by the three components of hazard, exposure and vulnerability.

$$\text{risk} = f(\text{hazard}, \text{exposure}, \text{vulnerability}) \quad (1)$$

Hazard describes probability of occurrence and intensity of natural or human-induced events such as storms, floods and saline intrusion. Exposure shows the presence of people,

infrastructure, livelihoods, environmental functions, services, etc. in the region that could be affected by the events. Vulnerability is identified as the extent of harm, including sensitivity/susceptibility and resilience. There is a broad set of elements of an system which is suffering from the natural disasters. This set can be characterized by hydrogeological, social and economic components (Tri et al. 2013; Balica et al. 2014).

2.2. Multiple natural hazard assessment

In order to assess natural hazards, the main hazardous events which cause considerable damages on agricultural production are selected for the assessment. Identifying and assessing hazardous events help to understand their nature and behaviour, suggesting awareness and planning for disaster mitigation strategies. The natural hazards was assessed in terms of frequency, affected areas and degree of severity. In reality, a number of natural events threaten the agricultural production in the coastal area, such as saline intrusion, tropical cyclones, tornadoes, etc. These threats were selected for multi-hazards assessment based on their damages on agricultural crops in the study area and this was observed by the Department of Agriculture and Rural Development of My Xuyen district. We assessed the impacts of multi-hazards on agricultural crops, focusing on rice crops. The agricultural crops are only influenced by the natural hazards during their cultivated durations which are represented by their cropping calendars. The major land-use types in My Xuyen district are agriculture and aquaculture, consisting of intensive rice farming, intensive shrimp farming and rice-shrimp farming (Thảo et al. 2017). Therefore, a temporal analysis is important for the hazard assessment and its temporal resolution should be fine enough to present seasonal variations of the cropping systems. In this study, the monthly interval are reasonable for presenting the cropping calendar of each farming system.

According to the recent reports on natural disaster prevention and control of My Xuyen district, saline intrusion and tropical cyclones are the most dangerous phenomena. These phenomena cause substantial damages to agricultural cultivation recently. Therefore, these two events are considered for multiple hazards assessment. Hazard indexes are calculated by the below equation.

$$HI = \sum_{i=1}^n w_i HI_i \quad (2)$$

where HI is multiple hazard index; HI_i is the hazard index of phenomenon i ; w_i is the weight of phenomenon i . The HI_i of each phenomenon is computed based on its frequency and intensity. The hazard index of each phenomenon should be normalized due to different range of the hazard index. We used the weights in calculation of hazard index because this help to prioritize the importance of each phenomenon. The weight of a phenomenon is simply identified as damage percentage caused by that phenomenon of the total damage.

Tropical cyclones

Tropical cyclones are of the most destructive natural hazards not only in Vietnam but also over the world. They affect large areas depending on their intensity and track of the tropical cyclones. Wang et al. (2020) defined the warning area due to the tropical cyclones within an 800 km radius. Therefore, we analysed the tropical cyclones occurred within a circle whose radius is 800 km from My Xuyen district. The tropical cyclones mentioned in this study are tropical storms or greater events (e.g. severe tropical storms, typhoons, or hurricanes). Data of tropical cyclones, which were recorded for the period of 1951 to 2019 by the Japan Meteorological Agency, were used for the hazard assessment. The data contain characteristic features of tropical cyclone tracks for every six hours. First, the eye's locations of tropical cyclones within the 800 km circle were

selected. To assess tropical cyclone impact, we used and modified the Tropical Cyclone Potential Index (TCI) which was introduced by (Xiao et al. 2011). Then monthly TCI was computed as follows:

$$TCI = \sum_{i=1}^n w_i v_i^2 \quad (3)$$

$$w_i = 1 - \frac{d_i}{800} \quad (4)$$

where n is the number of points of 6-hour tropical cyclones in the circle; v_i is the maximum wind speed near the tropical cyclone center i ; w_i is the weight of the tropical cyclone center i which was calculated by its inverse distance to the study area; and d_i is the distance between the tropical cyclone center and the study area centroid (km).

Saline intrusion

Saline intrusion is the most influential factor that impacts agricultural production in My Xuyen district. Saltwater intrudes into My Xuyen district through the My Thanh River which are directly connected to the East Sea (Bé et al. 2017). Salinity data from 2016 to 2019 at Nga Ba Vam Leo (located in Hoa Tu 2,

142)

143 was collected by the Irrigation Management Station of My Xuyen district. These data of salinity

144 were analysed monthly frequency of salt stress. The levels of salt stress are identified based on

145 the cultivated crops and their stages. Generally, salinity levels of higher than 2 dS/m can reduce

146 rice yield regarding to timing of salt stress and rice variety (Asch and Wopereis 2001).

147 According to Ayers and Westcot (1985), the agricultural crops are not affected by the electrical

148 conductivity (EC) of irrigation water which is lower than 0.7 dS/m. When the EC is higher than

149 0.7 dS/m, the agricultural crops begin to be damaged slightly. The crops are severely affected by

150 the EC of higher than 3.0 dS/m. This is a common classification for salt stress of irrigation water

151 for agriculture and this highly agrees with salinity hazard of irrigation water defined by Bauder

152 et al. (2011); Follett and Soltanpour (1914). Specifically, each agricultural crop has different

153 capacity to adapt to saline water. For instance, effects of saline water on rice yield were

evaluated by Ayers and Westcot (1985) and is presented in **Error! Reference source not found.** Total dissolved solids (TDS) are the common data of water salinity and they were estimated by using its relationship with EC. The average ratio of total dissolved solids and EC is 0.64 (Ali et al. 2012).

Table 1. Salinity hazard categories of irrigation water for rice crops (Ayers and Westcot 1985).

EC (dS/m)	TDS (g/l)	Category	Effects on rice crops
EC < 2.0	TDS < 1.3	None	Rare effect on rice yield
2.0 ≤ EC < 3.4	1.3 ≤ TDS < 2.2	Slight	Reduction of 25% rice yield
3.4 ≤ EC ≤ 4.8	2.2 ≤ TDS ≤ 3.1	Moderate	Reduction of 50% rice yield
EC > 4.8	TDS > 3.1	Severe	Reduction of over 50% rice yield or destruction of rice crops

The salinity data collected were used to analyse the monthly frequency of the selected salinity levels. To assess effects of saline water on rice crops, we defined the Salinity Potential Index (SPI) based on salinity levels and their frequency, presented as follows.

$$SPI = \sum_{i=1}^3 n_i w_i \quad (5)$$

where i is the number of salinity hazard categories, n_i is the number of salinity occurrence of the category i , w_i is the category weights which are equal to 0.25, 0.5, and 1 for *slight*, *moderate* and *severe* categories respectively.

2.3. Susceptibility of rice crops in the rice-based cropping system

Rice is the major agricultural crops in My Xuyen district and the rice-based farming systems are double rice crops and shrimp-rice crops. Rice crops are highly susceptible to the effects of tropical cyclones and salinity in irrigation water. Each type of these hazardous events has a different way to influence the rice crops.

Effects of tropical cyclones

The tropical cyclones damage the rice crops due to high wind velocities which leads to injury of plant organs. Masutomi et al. (2012) found that the highest rice vulnerability to tropical cyclones is the heading stage and assumed that the fragility curve of rice plants can be formulated by the Weibull distribution. Therefore, rice vulnerability to tropical cyclones was computed from the damage ratios introduced by Masutomi et al. (2012).

Effects of saline intrusion

The rice crops are highly vulnerable to salt stress and the damages are corresponded to their growth stages. The rice crops are really sensitive to salinity, particularly at the seedling stage (Grattan et al. 2002; Kakar et al. 2019). By another way, saline irrigation water increases salt accumulation in the crop root zone and this causes that the rice crops are not able to sufficient water. Moreover, the saline irrigation water can slow growth rate of rice plants, leading to a reduction or a loss of rice yield. Salinity in irrigation water differently influences rice growth and yield during the growing stages. Thus we used the cropping calendar of the farming systems in order to analyse susceptibility of the rice crops. During the seedling stage, effects of saline water on rice growth at different timing during the vegetative stage are relatively similar. These effects

in the reproductive phase are slightly higher in the vegetative phase (Zeng et al. 2001). In this study, the susceptibility of rice was indicated by reduction of rice yields. The reduction of rice yields at different timing of salinity was computed based on experimental data of Rad et al. (2011). These reductions are considerably changed under different salinity levels, illustrated in Table 2. Rice growth has similar patterns of responses to the different salinity levels during growing phases (Figure 2). The panicle initiation is the most susceptible stage in which rice yield can be reduced by a half due to effects of 4 dS/m salinity. We used average reduction of rice yield due to 2 and 4 dS/m salinity for indicating rice sensitivity during growing phases. At the seedling stage, rice survival is about 70.2% because of impacts of 3.3 dS/m salinity (Zeng et al. 2003). Response of the rice crops to salinity is highly various because of rice varieties. We assumed that the rice varieties cultivated in My Xuyen district has similar capacity to response to salinity and tropical cyclones to those in experiments of Masutomi et al. (2012) and Rad et al. (2011).

Figure 2. The growth stages of rice (IRRI 2015).

Table 2. Reduction of rice yield at different timing of salinity (Rad et al. 2011).

EC (dS/m)	Tillering	Panicle initiation	Heading	Ripening
2	26%	43%	9%	4%
4	33%	50%	13%	3%
Average	30%	46%	11%	4%

The decreased percentages of rice yields were normalized. Values of the vulnerability index were normalized on a scale from 0 to 1 where vulnerability is lowest or highest respectively. These vulnerability are grouped into five categories, presented in Table 3.

Table 3. Categories of vulnerability.

Category	Hazard and vulnerability	Risk
Very low	0 – 0.2	0 – 0.04
Low	0.2 – 0.4	0.04 – 0.16
Moderate	0.4 – 0.6	0.16 – 0.36
High	0.6 – 0.8	0.36 – 0.64
Very high	0.8 - 1	0.64 – 1

3. Results and discussion

Many regions are exposed to a number of natural hazards depending on their spatial

characteristics. There are several specializing hazards in coastal areas such as tropical cyclones and saline intrusion. It is evident that these natural hazards cause huge damage to agricultural production in the coastal areas. Therefore, the outputs of the hazard and vulnerability assessment assist to deal with risks of these phenomenon. In this section, we present results of hazard, vulnerability and risk assessment of the mentioned hazardous events, influencing the rice crops in My Xuyen district.

3.1. Temporal hazard assessment

Figure 3 shows the tropical cyclone tracks within the region of potential effects from 1951 to 2019. There are 136 tropical cyclones occurred in the region. The tropical cyclones originate from the eastern-oriented directions, developing in the Pacific Ocean. Generally, the tropical cyclones occur during the southwest monsoon season. The monthly numbers of tropical cyclones are presented Figure 4. My Xuyen has high potential effects of tropical cyclones from September to December, particularly in October and November which contribute to over a half of the total tropical cyclone number. The tropical cyclones are rarely approach the southern coast (My Xuyen district) while they hit the central and northern parts of Vietnam (Takagi et al. 2014). It is obviously evident that the tropical cyclones move to the south from November to February. This is indicated by the low latitudes of tropical cyclones.

226

227 Figure 3. Tropical cyclone tracks (1951-2019) within the region in which they can affect My
 228 Xuyen.

Figure 4. Monthly numbers and mean latitude of tropical cyclones from 1951 to 2019.

We calculated the monthly TCI of all tropical cyclones from 1951 to 2019 (Table 4) in order to identify stress timing due to tropical cyclones. The results clearly show that November has the highest TCI value, followed by October since these two months have the greatest occurrence of tropical cyclones. Although September and December have similar numbers of tropical cyclones, the TCI value of December is considerably higher than that of September. This happens because tropical cyclones in December tends to get closer to My Xuyen. Consequently, October and November are the most hazardous period, which are categorized as *high* and *very high* respectively.

Table 4. Hazard indexes of tropical cyclones and salinity.

Month	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec

Average												
TCI	49	482	131	173	336	242	160	11	1285	8646	13065	4709
HI_{Tropical} cyclones	0.00	0.04	0.01	0.01	0.03	0.02	0.01	0.00	0.10	0.66	1.00	0.36
HI_{Salinity}	0.60	0.76	0.96	1.00	1.00	0.98	0.68	0.34	0.43	0.40	0.46	0.48
HI	0.59	0.77	0.95	1.00	1.00	0.98	0.68	0.34	0.44	0.52	0.63	0.54
	Very low	Low	Moderate	High	Very high							

240 Saltwater intrudes into My Xuyen district through the My Thanh River (

241).

242 The My Thanh River is strongly dominated by the coastal processes and marginally controlled

by the fluvial process. Thus surface water resources become saline due to effects of the coastal processes. Figure 5 presents variations of daily maximum salinity at Nga Ba Vam Leo station from 2016 to 2019. It obviously depicts that saline intrusion are varied seasonally. Specifically, salinity reaches the highest level in April or May when river flows are low in the dry season; in contrast, salinity becomes lowest in September coinciding with high river flows in the rainy season in the Mekong Delta. However, salinity at Nga Ba Vam Leo is generally higher than the S1 level in which rice yield can be reduced by 25%. Therefore, surface water resources at Nga Ba Vam Leo need to be noticed to use for rice irrigation. This results in high values of salinity hazard index over the year. The salinity hazard indexes are usually categorized as *moderate*, *high* and *very high* (Table 4).

Based on the rates of saline intrusion and tropical cyclones influencing agricultural cropping systems in My Xuyen, saline intrusion is the main phenomenon damaging rice crops while tropical cyclones has much less impact. The ratio of saline intrusion to tropical cyclones impacts is approximately 85:15. This ratio is relatively reasonable because probability of tropical cyclones is low while saline intrusion frequently happens in My Xuyen. The multiple hazard index (illustrated in Table 4) indicates that May-June is the most hazardous period for rice production. This calculation agrees with the cropping calendar of the double rice cropping system in My Xuyen This period is free in the cropping calendar (Nguyen et al. 2019) due to *very high* hazard index. The lowest hazard appears in August, followed by a *moderate* hazard period. Thus rice is cultivated during this period even in the rice-shrimp cropping system.

Figure 5. Daily maximum salinity at Nga Ba Vam Leo. S1, S2 and S3 are critical levels that reduce rice yields of 25%, 50% and 75% respectively.

3.2. Vulnerability

Table 5 shows temporal variations of vulnerability index of a rice crop which indicates potential effects of tropical cyclones and salinity, damaging at different growth phases. We found that rice crops response to tropical cyclones and salinity differently. Rice is most vulnerable to tropical cyclones at the heading stage and to salinity at the panicle initiation phase. Noticeably, the first half of rice crop is highly vulnerable to salinity while the second half is considerably damaged due to tropical cyclones. The combined vulnerability index is averaged and normalized. It shows that the panicle initiation stage has the highest value of the combined vulnerability index, followed by the heading stage with an index of 0.66.

Table 5. Vulnerability index of a rice crop.

Growth phase	Seedling	Tillering	Panicle initiation	Heading	Ripening	Harvesting
Days after sowing	1	20	50	70	100	105
Vulnerability to tropical cyclones	0.00	0.33	0.87	1.00	0.77	0.67
Vulnerability to salinity	0.64	0.64	1.00	0.23	0.08	0.08
Combined vulnerability index	0.34	0.52	1.00	0.66	0.45	0.4

Rice cultivated in My Xuyen are usually short-duration varieties which mature in a range of 105-120 days. This is coincided with surveyed data in My Xuyen district recently (Nguyen et al. 2019). Table 6 presents cropping calendar or the double rice and rice-shrimp cropping system in My Xuyen. The double rice cropping system includes Summer-Autumn (from June to September) and Winter-Spring (October to January) crops. The rice crop in the rice-shrimp cropping system is cultivated from September to December. The cropping calendar of each system was used for computing vulnerability index which is depicted in Table 6. My Xuyen district has highly potential damage to rice in October and November because it has the largest area of rice production at the reproductive phase in the both cropping systems. In contrast, there are no damage of rice crops from February to May because of no rice grown during this period. This period was obviously coincided with the highest salinity in My Xuyen. It can be explained that the local community and farmers may have a measure to adapt to saline intrusion by adjusting the cropping calendar. Consequently, saline intrusion is the most hazardous events in

289 this area.

290 Table 6. Cropping calendar of rice-based farming systems in My Xuyen (after Nguyen et al.,
291 2019) and rice vulnerability index.

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Double rice						Summer-Autumn				Winter-Spring		
Rice-Shrimp					Shrimp				Winter-Spring			
Vulnerability index	0.17	0.00	0.00	0.00	0.00	0.25	0.55	0.44	0.43	0.81	1.00	0.64
Very low	Low	Moderate		High		Very high						

292 3.3. Multi-hazard risk

293 Based on the hazard and vulnerability identification, monthly risk indexes indicate possibility of
294 rice damage or loss due to effects of tropical cyclones and saline intrusion, are presented in Table
295 7. October and November belong to the *high* category while February-May have no risk. The
296 rice crop is at no risk from February to May and this is resulted from no rice crop cultivated
297 during this period. The *high* risk of rice production in October and November is occurred
298 because of *very high* vulnerability and *high* hazard. Results of the temporal analysis have a
299 reasonable agreement with Wassmann et al. (2019) outputs of risk analysis for the coastal VMD
300 in general.

301 Table 7. Monthly risk of rice to multiple hazards.

Month	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Risk	0.10	0.00	0.00	0.00	0.00	0.24	0.37	0.15	0.19	0.42	0.63	0.34
	Very low		Low		Moderate		High			Very high		

The nature of risks is considerably useful for assessing potential impacts of tropical cyclones and saltwater intrusion on rice production. It is noticed that the damages of agricultural crops vary with timing of the hazards and crop growth stages. Therefore, to reduce the risks, it needs to reduce hazards or/and vulnerability. First, the hazards can be reduced by improving early warning system. The hazardous phenomenon are tropical cyclones and salinity which are possibly predicted by their dominated factors. For example, tropical cyclones are predicted by several predominant factors (Safaripour et al. 2012; Camargo et al. 2019). The salinity is usually projected by the El Niño–Southern Oscillation indexes (Apel et al. 2020) or water discharge of the Mekong River (Dat et al. 2011). In addition, a structure such as sluice gates is a common measure to prevent saline intrusion in coastal areas. Second, there are some solutions to reduce vulnerability of rice-based cropping systems, including using salt-tolerant varieties, adjusting cropping calendar and changing to other agricultural crops (Renaud et al. 2015; Thảo et al. 2017). The results of risk assessment can help the local community become more resilient in rice cultivation to the natural hazardous events.

4. Conclusions

We assessed multiple natural hazards of tropical cyclones and saline intrusion. These hazardous events are the major factors, causing significant damages on rice crops in My Xuyen district. To assess risk of the multiple natural hazards, we used a framework of risk on rice crops in a coastal

area (My Xuyen, Soc Trang). The study primarily focused on evaluation of potential impacts of the mentioned hazards on rice crops. A temporal analysis was embedded in risk assessment because rice crops vary seasonally.

The study have shown that rice crop may face extreme salinity from March to June while it deals with tropical cyclones in October and November. These hazards have various effects on rice crops. Rice crops are considerably susceptible to salinity during the first half of the growing period and to tropical cyclones during the other half. In general, therefore, the results show that rice crops are sharply susceptible in the reproduction phase to the combination of salinity and tropical cyclones. For rice cultivation in My Xuyen, it is considerably vulnerable to these hazards in October and November based on the cropping calendar. We found that the rice crops are at *high* risk in October and November. Our findings are that saline intrusion reaches the highest level in April and May, but the rice crop was at no risk because of no rice crop cultivated during this period.

The multiple hazard assessment provides essential information for the case study in order to create an awareness for natural disaster mitigation. In addition, the integrated result of multiple hazards is much more simply than these of each hazard in providing information to planners and decision-makers because many numbers and scales can be confusing and cumbersome to them. If the results of multi-hazards and risk assessment are used efficiently, it can reduce damage or loss of rice crops/yield. However, this needs an effort of users who convey the information to the natural disaster mitigation procedure. An implication of these findings is to assist decision-makers, planners and managers in mitigation and adaptation strategies.

341 **List of abbreviations**

342 VMD: the Vietnamese Mekong Delta; TCI: Tropical Cyclone Potential Index; EC: Electrical
343 conductivity; SPI: Salinity Potential Index.

344 **Availability of data and materials**

345 The datasets generated and/or analysed during the current study are available from the
346 corresponding author on reasonable request.

347 **Competing interests**

348 The authors declare that they have no competing interests.

349 **Funding**

350 This study was funded in part by the Can Tho University Improvement Project VN14-P6,
351 supported by a Japanese ODA loan.

352 **Authors' contributions**

353 VQT and NHT were responsible for conceptualizing this study and formal analysis. VTPL was
354 responsible for data curation. VQT wrote the initial draft and all authors contributed to the
355 manuscript by providing comments and suggestions.

356 **Acknowledgements**

357 The authors would like to thank staff at Department of Agriculture and Rural Development of
358 My Xuyen district for their support.

359 **References**

360 Ali NS, Mo K, Kim M (2012) A case study on the relationship between conductivity and
361 dissolved solids to evaluate the potential for reuse of reclaimed industrial wastewater.

362 KSCE J Civ Eng 16:708–713. <https://doi.org/10.1007/s12205-012-1581-x>
 363 Apel H, Khiem M, Hong Quan N, Quang Toan T (2020) Brief communication: Seasonal
 364 prediction of salinity intrusion in the Mekong Delta. *Nat Hazards Earth Syst Sci* 20:1609–
 365 1616. <https://doi.org/10.5194/nhess-20-1609-2020>
 366 Asch F, Wopereis MCS (2001) Responses of field-grown irrigated rice cultivars to varying levels
 367 of floodwater salinity in a semi-arid environment. *F Crop Res* 70:127–137.
 368 [https://doi.org/10.1016/S0378-4290\(01\)00128-9](https://doi.org/10.1016/S0378-4290(01)00128-9)
 369 Ayers RS, Westcot DW (1985) Water quality for agriculture. FAO irrigation and drainage paper
 370 29 Rev 1, Rome, Italia
 371 Balica S, Dinh Q, Popescu I, et al (2014) Flood impact in the Mekong Delta, Vietnam. *J Maps*
 372 10:257–268. <https://doi.org/10.1080/17445647.2013.859636>
 373 Balica SF, Wright NG, Meulen F (2012) A flood vulnerability index for coastal cities and its use
 374 in assessing climate change impacts
 375 Bates PD, Pappenberger F, Romanowicz RJ (2014) Uncertainty in flood inundation modelling.
 376 In: *Applied Uncertainty Analysis for Flood Risk Management*. pp 232–269
 377 Bauder TA, Waskom RM, Sutherland PL, Davis JG (2011) Irrigation water quality criteria.
 378 Colorado State University Extension Publication. Crop series: Irrigation
 379 Bé NV, Trí VPĐ, Vũ PT, Vũ PH (2017) Challenges in agriculture in My Xuyen district, Soc
 380 Trang province in the context of saline intrusion. *Can Tho Univ J Sci* 2:187–196.
 381 <https://doi.org/10.22144/ctu.jsi.2017.067>
 382 Buan RD, Maglinao AR, Evangelista PP, Pajuelas BG (1996) Vulnerability of rice and corn to
 383 climate change in the Philippines. *Water Air Soil Pollut* 92:41–51.
 384 https://doi.org/10.1007/978-94-017-1053-4_4
 385 Camargo SJ, Camp J, Elsberry RL, et al (2019) Tropical cyclone prediction on subseasonal time-
 386 scales. *Trop Cyclone Res Rev* 8:150–165. <https://doi.org/10.1016/j.tcr.2019.10.004>
 387 Dat TQ, Likitdecharote K, Srisatit T, Trung NH (2011) Modeling the influence of river discharge
 388 and sea level rise on salinity intrusion in the Mekong Delta. In: *The 1st Environment Asia*
 389 *International Conference on Environmental Supporting in Food and Energy Security: Crisis*

390 and Opportunity. Thai Society of Higher Education Institutes on Environment, pp 685–701

391 Esteban M, Takagi H, Thao ND (2014) Tropical Cyclone Damage to Coastal Defenses: Future
 392 Influence of Climate Change and Sea Level Rise on Shallow Coastal Areas in Southern
 393 Vietnam. Elsevier Inc.

394 FAO (2017) The impact of disasters on agriculture: Addressing the information gap. Rome, Italia

395 Follett R, Soltanpour P (1914) Irrigation water quality criteria. Colorado State University
 396 Publication No. 0.506

397 Grattan SR, Zeng L, Shannon MC, Roberts SR (2002) Rice is more sensitive to salinity than
 398 previously thought. *Calif Agric* 56:189–198

399 GSOVN (GENERAL STATISTICS OFFICE of VIET NAM) (2010) Statistical yearbook of
 400 Vietnam. Hanoi

401 Hakim MA, Juraimi AS, Begum M, et al (2010) Effect of salt stress on germination and early
 402 seedling growth of rice (*Oryza sativa* L.). *African J Biotechnol* 9:1911–1918

403 IPCC (2014) Summary for policy makers. In: Field CB, Barros VR, Dokken DJ, et al. (eds)
 404 Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part A: Global and Sectoral
 405 Aspects. Contribution of Working Group II to the Fifth Assessment Report of the
 406 Intergovernmental Panel on Climate Change. Cambridge University Press, Cambridge,
 407 United Kingdom and New York, NY, USA, pp 1–32

408 IRRI (2015) IRRI Rice Production Manual: Steps to successful rice production. International
 409 Rice Research Institute, Los Baños (Philippines)

410 Kakar N, Jumaa SH, Redoña ED, et al (2019) Evaluating rice for salinity using pot-culture
 411 provides a systematic tolerance assessment at the seedling stage. *Rice* 12:.
 412 <https://doi.org/10.1186/s12284-019-0317-7>

413 Kakonen M (2008) Mekong Delta at the crossroads: More control or adaptation? *Ambio* 37:205–
 414 212

415 Kunitsugu M (2012) Tropical cyclone information provided by the RSMC Tokyo - Typhoon
 416 Center. *Trop Cyclone Res Rev* 1:51–59. <https://doi.org/10.6057/2012TCRR01.06>

417 Lee T-C, Knutson TR, Kamahori H, Ying M (2012) Impacts of climate change on tropical
 418 cyclones in the Western North Pacific Basin. Part I: Past observations. *Trop Cyclone Res*
 419 *Rev* 1:213–235. <https://doi.org/10.6057/2012TCRR02.08>

420 Masutomi Y, Iizumi T, Takahashi K, Yokozawa M (2012) Estimation of the damage area due to
 421 tropical cyclones using fragility curves for paddy rice in Japan. *Environ Res Lett* 7:.
 422 <https://doi.org/10.1088/1748-9326/7/1/014020>

423 Mendelsohn R, Emanuel K, Chonabayashi S, Bakkensen L (2012) The impact of climate change
 424 on global tropical cyclone damage. *Nat Clim Chang* 2:205–209.
 425 <https://doi.org/10.1038/nclimate1357>

426 Merz B, Hall J, Disse M, Schumann A (2010) Fluvial flood risk management in a changing
 427 world. *Nat Hazards Earth Syst Sci* 10:509–527. <https://doi.org/10.5194/nhess-10-509-2010>

428 MRC (2010) State of the Basin Report 2010. Vientiane, Laos

429 Nguyen MT, Renaud FG, Sebesvari Z, Can Nguyen D (2019) Resilience of agricultural systems
 430 facing increased salinity intrusion in deltaic coastal areas of Vietnam. *Ecol Soc* 24:19.
 431 <https://doi.org/10.5751/ES-11186-240419>

432 Qadir M, Tubeileh A, Akhtar J, et al (2008) Productivity enhancement of salt-affected
 433 environments through crop diversification. *L Degrad Dev* 19:429–453.
 434 <https://doi.org/10.1002/ldr>

435 Rad HE, Aref F, Khaledian M, et al (2011) The effects of salinity at different growth stage on
 436 rice yield. In: *ICID 21st International Congress on Irrigation and Drainage*. Tehran, Iran, pp
 437 155–165

438 Renaud FG, Le TTH, Lindener C, et al (2015) Resilience and shifts in agro-ecosystems facing
 439 increasing sea-level rise and salinity intrusion in Ben Tre Province, Mekong Delta. *Clim*
 440 *Change* 133:69–84. <https://doi.org/10.1007/s10584-014-1113-4>

441 Safaripour M, Monavari M, Zare M, et al (2012) Flood Risk Assessment Using GIS (Case
 442 Study : Golestan Province , Iran). *Polish J Environ Stud* 21:1817–1824

443 Takagi H, Thao ND, Esteban M (2014) Tropical Cyclones and Storm Surges in Southern
 444 Vietnam. In: *Coastal Disasters and Climate Change in Vietnam: Engineering and Planning*

Perspectives. Elsevier Inc., pp 3–16

Thảo NH, Trung NH, Trí LQ (2017) Establishing the model for supporting agricultural land use allocation - A case study in My Xuyen district, Soc Trang province. *Can Tho Univ J Sci* 2:166–177. <https://doi.org/10.22144/ctu.jsi.2017.065>

Tri VPD, Trung NH, Thanh VQ (2013) Vulnerability to flood in the Vietnamese Mekong Delta : mapping and uncertainty assessment. *J Environ Sci Eng B* 2:229–237

Wang L, Zhou Y, Lei X, et al (2020) Predominant factors of disaster caused by tropical cyclones in South China coast and implications for early warning systems. *Sci Total Environ* 726:138556. <https://doi.org/10.1016/j.scitotenv.2020.138556>

Wassmann R, Phong ND, Tho TQ, et al (2019) High-resolution mapping of flood and salinity risks for rice production in the Vietnamese Mekong Delta. *F Crop Res* 236:111–120. <https://doi.org/10.1016/j.fcr.2019.03.007>

Xiao F, Yin Y, Luo Y, et al (2011) Tropical cyclone hazards analysis based on tropical cyclone potential impact index. *J Geogr Sci* 21:791–800. <https://doi.org/10.1007/s11442-011-0880-3>

Zeng L, Lesch SM, Grieve CM (2003) Rice growth and yield respond to changes in water depth and salinity stress. *Agric Water Manag* 59:67–75. [https://doi.org/10.1016/S0378-3774\(02\)00088-4](https://doi.org/10.1016/S0378-3774(02)00088-4)

Zeng L, Shannon MC (2000) Salinity effects on seedling growth and yield components of rice. *Crop Breeding, Genet Cytol* 40:996–1003

Zeng L, Shannon MC, Lesch SM (2001) Timing of salinity stress affects rice growth and yield components. *Agric Water Manag* 48:191–206

Figures

Figure 1

Land-use map in 2010 of My Xuyen district.

Figure 2

The growth stages of rice (IRRI 2015).

Figure 3

Tropical cyclone tracks (1951-2019) within the region in which they can affect My Xuyen.

Figure 4

Monthly numbers and mean latitude of tropical cyclones from 1951 to 2019.

Figure 5

Daily maximum salinity at Nga Ba Vam Leo. S1, S2 and S3 are critical levels that reduce rice yields of 25%, 50% and 75% respectively.