Oral Administration of the Gut-bacterial Metabolite 5-hydroxyindole and Its Subsequent Acceleration of Gut Motility Have Marginal Effects on the Rat Microbiota

Barbora Waclawiková

University of Groningen

Markus Schwalbe

University of Groningen

Sahar EL Aidy (✉ sahar.elaidy@rug.nl)

University of Groningen

Short Report

Keywords: Gut bacterial metabolite, 16S rRNA analysis, total gut transit time, Prevotella

Posted Date: February 23rd, 2022

DOI: <https://doi.org/10.21203/rs.3.rs-1374818/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Abstract

Background: Intestinal microbiota and microbiota-derived metabolites play a key role in regulating the host physiology, including gut motility. We have previously shown that a wide variety of gut bacteria can produce 5-hydroxyindole from its precursor, dietary supplement and antidepressant, 5-hydroxytryptophan. When 5-hydroxyindole was administered orally in rats, it significantly accelerated the total gut transit time.

Results: Here we show, using 16S rRNA sequencing, that 5-hydroxyindole has no effect on the bacterial richness. No separation between the treated and untreated groups was detected based on their microbial composition as shown by principal component analysis. A significant increase in the abundance of only two genera, *Shuttleworthia* and *Prevotellaceae_UCG-001*, was detected in the 5-hydroxyindole treated group compared to the vehicle group. No significant association was found between the reduced total gut transit time as a result of 5-hydroxyindole administration and the cecal bacterial composition.

Conclusions: Together, our study infers a marginal effect of 5-hydroxyindole or its subsequent accelerated gut motility on the cecal microbiota composition of rats. The results urge further investigation of the impact of 5-hydroxyindole oral administration in humans, which will support its potential application as a treatment for slow intestinal motility disorders.

Background

The gastrointestinal tract is home to trillions of microbes. The gut microbiota produces a wide range of small bioactive molecules derived from various substrates, including dietary precursors and medications [1, 2]. Such microbial conversion represents a significant regulatory mechanism by which gut microbes can alter intestinal host physiology, including gastrointestinal motility [3–6]. Recently, we have identified 5-hydroxyindole, a product of gut microbial conversion of the dietary supplement and antidepressant 5-hydroxytryptophan, as a potent accelerator of the gastrointestinal motility via its activation of L-type calcium channels located on the colonic smooth muscle cells and possibly also through its induction of serotonin production from enterochromaffin cells [7]. These findings propose 5-hydroxyindole as a potentially safe treatment for gastrointestinal slow motility disorders. Slow gastrointestinal motility disorders, such as constipation, is a common, debilitating motility disorder affecting up to 27% of the population [8]. Constipation is also often associated with colorectal cancer, Parkinson disease, childhood attention-deficit/hyperactivity disorder, and autism spectrum disorder, as well as mood disorders [9–13].

Indole (tryptophan) metabolites, one of the most studied gut microbiota-produced metabolites [14], are known for their effect on, for example, the aryl hydrocarbon receptors (AhR) [6, 15, 16], which in turn affects gastrointestinal barrier function [17–19] and host gut motility [6]. Thus, the influence of 5-hydroxyindole, being an indole derivative, can modify the fluid flow within the intestinal lumen, which is directly responsible for the transport and dispersion of the nutrients, bacterial colonization and the growth [20]. This led us to hypothesize that since 5-hydroxyindole stimulates the gut motility it may impact the microbiota composition. Using cecal samples from rats treated orally with 5-hydroxyindole for 11 days, we show that, compared to the vehicle treated rats, 5-hydroxyindole has only a marginal effect on the gut microbiota composition.

Methods

Cecal samples collection

All animal procedures were approved by the Groningen University Committee of Animal experiments (approval number: AVD1050020197786) and were performed in adherence to the NIH Guide for the Care and Use of Laboratory Animals. Wild-type Groningen (WTG) male rats were orally administered either 30 mg/kg 5-hydroxyindole (H31859, Sigma) (n = 10) or vehicle (10% sucrose) (n = 6) for a period of 11 days and after measurement of total gut transit time (TGTT) (more detailed protocol can be found in [7]), the rats were anesthetized with isoflurane, killed and the whole cecum from every rat was collected and snap-frozen in a liquid nitrogen and stored in -80°C until further procedure.

DNA isolation

DNA isolation followed a previously proposed protocol by Yu and Morrison (2004). Briefly, about 500 mg of cecal material was resuspended in 750 µL of lysis buffer (500 mM NaCl, 50 mM Tris-HCl, 50 mM EDTA, 4% SDS) and transferred to a screw cap tube containing ~ 500 mg of 0.1 mm zirconium beads and four 3 mm glass beads. Samples were homogenized 3 × 1 min with 1-minute intervals on ice in a mini bead-beater (Biospec, Bartlesville, USA). Subsequently, samples were further incubated at 95°C for 15 min and centrifuged at 16000 x g for 15 min at 4°C. All subsequent centrifugation steps were conducted under the same conditions. The supernatant was transferred to a clean tube and 20 µL of 10 M ammonium acetate was added and incubated on ice for 10 min before centrifugation. The supernatant was again transferred to a clean tube and one volume of 100% ice cold isopropanol was added. Samples were incubated on ice for 1.5 - 2h and centrifuged. The supernatant was then aspirated and the pellet was washed with 200 µL of 70% ethanol with following centrifugation to collect the pellet. Ethanol was removed and tubes were left to air-dry for ~1h before resuspension of the pellet in 200 µL of TE buffer.

16S rRNA sequencing

Illumina 16S rRNA gene amplicon libraries were generated and sequenced at Novogene (Bioinformatics Technology Co., Ltd., Beijing, China). In short, 16S rRNA genes of distinct regions (16SV3-V4) were amplified using specific primers (341F and 806R) with the barcode. All PCR reactions were carried out with Phusion® High-Fidelity PCR Master Mix with GC Buffer (New England Biolabs). The PCR products were detected by 2% agarose gel electrophoresis, and the samples were mixed equally according to the concentration of PCR products. After full mixing, the 2% agarose gel electrophoresis was used for the detection again, and the target band was recovered by using the gel recovery kit provided by Qiagen company. The library was constructed by NEBNext® Ultra™ II DNA Library Prep Kit, and the constructed library was quantified by Qubit and Q-PCR. After the library was qualified, NovaSeq6000 was used for sequencing.

Microbiota analysis

Paired-end reads were trimmed of their barcodes and sequencing primers and subsequently merged using FLASH (v1.2.11) [22]. Fastp was used for quality control and read filtering [23], while further VSEARCH was employed to detect chimera sequences by searching them against the Greengenes database [24]. Filtered high-quality reads were then subjected to read denoising using DADA2 to obtain amplicon sequencing variants (ASVs), while making use of the QIIME2 software [25, 26]. Sequences with less than 5 counts were removed and the remaining ASVs were classified with Classify-sklearn in QIIME2 leveraging the Greengenes database.

For downstream analysis, QIIME2 was further used to assess alpha diversity. We used the *phyloseq* and *microbiome* packages in the statistical programming language R to process our data further [27, 28]. ASV absolute abundances were collapsed on genus level (**Table G in Supplementary Table 1**) and CSS normalized [29] and the resulting abundance table was used for principal component analysis (PCA) (supporting data for PCA analysis can

be found in **Table E and F in Supplementary Table 1**). Significance of the model was determined by an ANOVA-like permutation test implemented in the *vegan* package [30] (PERMANOVA results can be found in **Table B in Supplementary Table 1**). Differential abundance between control and 5-hydroxyindole-treated groups was assessed by unpaired *t* test with Welch's correction. Significance was evaluated using FDR < 0.05. Marginal effect was assessed by *P* value < 0.05 (see main text for more details and **Table 1**). Additionally, LEfSe was used with default parameters to investigate differentially abundant taxa [31]. Pairwise correlations between microbial taxa and TGTT were performed using the *associate* function of the *microbiome* package, using Spearman correlations. Significance was evaluated using FDR < 0.05. Marginal effect was assessed by *P* value < 0.05 (see main text for more details, **Table 2 and Table D in Supplementary Table 1**).

Statistical analysis

For alpha diversity, Mann Whitney test was used. For pairwise comparison (beta diversity) between groups, unpaired *t* test with Welch's correction was used. Data are presented as mean ± SEM. Data were evaluated with FDR < 0.05 for significance and *P* value < 0.05 for marginal effect (see main text for more details). For correlations between cecal microbiota, 5-hydroxyindole treatment and TGTT, Spearman correlation was used. Data were evaluated with FDR < 0.05 for significance and *P* value < 0.05 for marginal effect (see main text for more details).

Results And Discussion

5-hydroxyindole has a marginal effect on the richness and composition of the cecal microbiota in wild-type Groningen rats

Recently, we showed that a daily oral administration of the gut microbiota-produced 5-hydroxyindole (30 mg/kg) to wild-type Groningen (WTG) rats for 11 days results in a significant decrease of the total gut transit time (TGTT) [7]. The dose for 5-hydroxyindole (30 mg/kg) was chosen based on a previous report [32]. To investigate the possible effect of 5-hydroxyindole and the subsequent change in the gut motility on the microbiota composition, 16 cecal samples were collected (5-hydroxyindole-treated group (*n* = 10); vehicle-treated group (*n* = 6)) after the TGTT was measured and amplicon sequencing of the V3-V4 regions of the bacterial 16S gene was performed. Microbial richness, assessed by the Chao1 index and observed number of OTUs, showed a marginal but not significant (*P* value = 0.056) increase in 5-hydroxyindole-treated rats compared to the control group (Figure 1A; Table A in Supplementary Table 1). Next, the microbiota diversity was determined by Shannon's H and Simpson's index, both indices are used to measure similar parameters of alpha diversity [33, 34]. The diversity index did not differ between the treated and untreated groups (Figure 1A; Table A in Supplementary Table 1). The data highlight that 5-hydroxyindole has a negligible effect on the richness and no effect on the diversity of the cecal microbiota. Despite the considerable recent progress in describing the effects of the indole (tryptophan) metabolites on the composition and diversity of the intestinal microbiota [35–38], the overall impact of these metabolites on the host microbiota is contradicting and remains largely unknown. For example, Liang *et al.* showed that indole and its derivative, indole-3-acetic acid, significantly enhanced the richness, but not the diversity of the bacterial population in the cecal contents of piglets [35]. On the other hand, other studies showed no effect of high levels of indole metabolites produced from tryptophan and Mediterranean diet on the alpha diversity in mice and humans, respectively [37, 38].

As a general exploratory analysis, principal component analysis (PCA) was performed, explaining 21.3% and 16% of the variance respectively, and showed no significant difference between the 5-hydroxyindole and vehicle-treated groups (PERMANOVA: *P* value = 0.175, stratified *P* value = 1; Table B in Supplementary Table 1). Next, LEfSe (Linear

discriminant analysis Effect Size; [31]) was employed to complement our differential abundance analysis. The main discriminant feature separating the groups (5-hydroxyindole and vehicle group) in WTG rats was the *Allobaculum* genus (Figure 1C). To support this analysis and investigate whether we can identify individual bacterial taxa to be affected by the 5-hydroxyindole treatment, pairwise comparisons of bacterial abundances were performed between 5-hydroxyindole-treated and vehicle-treated groups. Focusing on the phylum level, no significant changes were observed. On the family level, 5-hydroxyindole treatment seemed to only increase the abundance of family Yersiniaceae (P value = 0.03; Table C in Supplementary Table 1). On the genus level, 5-hydroxyindole treatment was associated with an increase in the abundance of *Allobaculum*, *Prevotellaceae_UCG-001*, *Serratia*, *Prevotellaceae_NK3B31_group*, *Shuttleworthia*, *Rikenellaceae_RC9_gut_group*, *Tuzzerella*, *Eubacterium_eligens_group*, *Parvibacter*, *Lachnospiraceae_NK4B4_group*, while reduced the abundance of *Acetatifactor* (P value < 0.05, unpaired t test with Welch's correction) (Figure 1D; Supplementary Table 1). Nonetheless, after multiple comparison correction (false discovery rate; FDR), only *Prevotellaceae_UCG-001* and *Shuttleworthia* (FDR < 0.05) showed a significant increase in their abundance. In agreement with our results, tryptophan enriched diets, and the consequent higher concentrations of indole metabolites in the cecal or fecal samples of piglets and mice, respectively, have been shown to increase the abundance of *Prevotella* genus [35, 38]. Taken together, the data analysis infers a minimal impact of 5-hydroxyindole treatment on the composition of the microbiota in the cecal samples of rats.

Because the gut transit time was significantly affected in the 5-hydroxyindole-treated group [7], the TGTT was tested for its association with the abundance of genera using Spearman correlations. The correlation analysis revealed nine genera to be associated with the TGTT covariate (Spearman, P value < 0.05); Figure 1D; Table 2; Table D in Supplementary Table 1). Eight genera correlated negatively (*Lachnospiraceae_UCG-006*, *Lachnospiraceae_NK4B4_group*, *Tuzzerella*, *Prevotellaceae_NK3B31_group*, *Eubacterium_ventriosum_group*, *Prevotellaceae_UCG-001*, *Barnesiella*, *Bacteroides*) and one genus positively (*Anaerovibrio*). Nonetheless, none of these associations could be detected as significant after multiple comparison correction (FDR). Overall, the results indicate that the TGTT covariate, which was shown to be significantly enhanced by 5-hydroxyindole treatment [7], is not significantly associated with the rat cecal bacterial composition.

Conclusions

Together, our current study suggests a marginal, presumably beneficial effect, of 5-hydroxyindole or its subsequent acceleration of the gut motility on the cecal microbiota composition when orally administered for 11 consecutive days to the WTG rats. Slow intestinal motility disorders, such as constipation, are highly prevalent gastrointestinal disorders in humans [8]. We speculate that potential future application of 5-hydroxyindole as a targeted drug, may have a marginal impact on the host gut microbiota. This gives an advantage for this microbiota-produced metabolite over several other available medications against constipation, which have been linked to significant changes in the gut microbiota composition [39]. Additionally, *Prevotella* genus has been reported to exhibit significant reduction in abundance in the gut microbiome of the constipated patients, compared to the controls [40]. Thus, the observed significant increase in the abundance of *Prevotellaceae_UCG-001* upon the 5-hydroxyindole treatment, may provide another beneficial effect for the consideration of 5-hydroxyindole as a treatment for constipated patients. Nevertheless, human interventions aiming to investigate long-term impact of 5-hydroxyindole on the microbial composition and the gastrointestinal sensitization, are needed to support the present claim.

List Of Abbreviations

AhR, aryl hydrocarbon receptor; WTG, wild-type Groningen; TGTT, total gut transit time; ASVs, amplicon sequencing variants; PCA, principal component analysis; FDR, false discovery rate; 5-HI, 5-hydroxyindole-treated group; Veh, vehicle-treated group; LDA, linear discriminant analysis.

Declarations

Ethics approval and consent to participate

All animal procedures were approved by the Groningen University Committee of Animal experiments (approval number: AVD1050020197786) and were performed in adherence to the NIH Guide for the Care and Use of Laboratory Animals.

Consent for publication

Not applicable

Availability of data and material

All sequencing data is available at PRJNA800624 and at this link:

<https://dataview.ncbi.nlm.nih.gov/object/PRJNA800624?reviewer=c9ag5ptv7vabikefplu16t8idt>

Competing Interests Statement

The authors have declared that no competing interests exist.

Funding

S.E.A. is supported by Rosalind Franklin Fellowships, co-funded by the European Union and University of Groningen. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Authors' Contributions

B.W. and S.E.A. conceptualized and designed the study. B.W. and M.S. performed the analysis of the data. S.E.A. assisted with analysis of the data. B.W. and S.E.A. wrote the original manuscript that was reviewed by M.S. Funding was acquired by S.E.A.

Acknowledgments

We thank Amber Bullock, Host-microbe Interaction group, Groningen Biomolecular Sciences and Biotechnology Institute (GBB), University of Groningen, Groningen, The Netherlands and Prof. Dr. Gertjan van Dijk, Department of Behavioral Neurosciences, Cluster Neurobiology, Groningen Institute of for Evolutionary Life Sciences (GELIFES), University of Groningen, Groningen, The Netherlands, for their help with the collection of the cecal samples.

References

1. van Kessel SP, Frye AK, El-Gendy AO, Castejon M, van Dijk G, El Aidy S. Gut bacterial tyrosine decarboxylases restrict the bioavailability of levodopa, the primary treatment in Parkinson's disease. *Nat Commun.* 2019;**31**:1–31.

2. Donia MS, Fischbach MA. Small molecules from the human microbiota. *Science* (80-). 2015;**349**.
3. Yano JM, Yu K, Donaldson GP, Shastri GG, Ann P, Ma L, et al. Indigenous Bacteria from the Gut Microbiota Regulate Host Serotonin Biosynthesis. *Cell*. 2015;**161**:264–76.
4. Reigstad CS, Salmonson CE, Rainey JF, Szurszewski JH, Linden DR, Sonnenburg JL, et al. Gut microbes promote colonic serotonin production through an effect of short-chain fatty acids on enterochromaffin cells. *FASEB J*. 2015;**29**:1395–403.
5. Bhattarai Y, Williams BB, Battaglioli EJ, Whitaker WR, Till L, Grover M, et al. Gut Microbiota-Produced Tryptamine Activates an Epithelial G-Protein-Coupled Receptor to Increase Colonic Secretion. *Cell Host Microbe*. 2018;**23**:775-785.e5.
6. Obata Y, Castaño Á, Boeing S, Bon-Frauches AC, Fung C, Fallesen T, et al. Neuronal programming by microbiota regulates intestinal physiology. *Nature*. 2020;**578**:284–9.
7. Wacławiková B, Bullock A, Schwalbe M, Aranzamendi C, Nelemans SA, van Dijk G, et al. Gut bacteria-derived 5-hydroxyindole is a potent stimulant of intestinal motility via its action on L-type calcium channels. *PLOS Biol*. 2021;**19**:e3001070.
8. Sanchez MIP, Bercik P. Epidemiology and burden of chronic constipation. *Can J Gastroenterol*. 2011;**25**:11–5.
9. Hosseinzadeh ST, Poorsaadati S, Radkani B, Forootan M. Psychological disorders in patients with chronic constipation. *Gastroenterol Hepatol from Bed to Bench*. 2011;**4**:159–63.
10. Guérin A, Mody R, Fok B, Lasch KL, Zhou Z, Wu EQ, et al. Risk of developing colorectal cancer and benign colorectal neoplasm in patients with chronic constipation. *Aliment Pharmacol Ther*. 2014;**40**:83–92.
11. Fasano A, Visanji NP, Liu LWC, Lang AE, Pfeiffer RF. Gastrointestinal dysfunction in Parkinson's disease. *Lancet Neurol*. 2015;**14**:625–39.
12. Mckeown C, Hisle-Gorman E, Eide M, Gorman GH, Nylund CM. Association of constipation and fecal incontinence with attention-deficit/hyperactivity disorder. *Pediatrics*. 2013;**132**.
13. Pang KH, Croaker GDH. Constipation in children with autism and autistic spectrum disorder. *Pediatr Surg Int*. 2011;**27**:353–8.
14. Agus A, Planchais J, Sokol H. Gut Microbiota Regulation of Tryptophan Metabolism in Health and Disease. *Cell Host Microbe*. 2018;**23**:716–24.
15. Lamas B, Hernandez-Galan L, Galipeau HJ, Constante M, Clarizio A, Jury J, et al. Aryl hydrocarbon receptor ligand production by the gut microbiota is decreased in celiac disease leading to intestinal inflammation. *Sci Transl Med*. 2020;**12**.
16. Lamas B, Natividad JM, Sokol H. Aryl hydrocarbon receptor and intestinal immunity. *Mucosal Immunol*. 2018;**11**:1024–38.
17. Bansal T, Alaniz RC, Wood TK, Jayaraman A. The bacterial signal indole increases epithelial-cell tight-junction resistance and attenuates indicators of inflammation. *Proc Natl Acad Sci*. 2010;**107**:228–33.
18. Venkatesh M, Mukherjee S, Wang H, Li H, Sun K, Benechet AP, et al. Symbiotic Bacterial Metabolites Regulate Gastrointestinal Barrier Function via the Xenobiotic Sensor PXR and Toll-like Receptor 4. *Immunity*. 2014;**41**:296–310.
19. Shimada Y, Kinoshita M, Harada K, Mizutani M, Masahata K, Kayama H, et al. Commensal bacteria-dependent indole production enhances epithelial barrier function in the colon. *PLoS One*. 2013;**8**:1–10.
20. Wacławiková B, Codutti A, Alim K, El Aidy S. Gut microbiota-motility interregulation: insights from in vivo, ex vivo and in silico studies. *Gut Microbes*. 2022;**14**:e1997296.

21. Yu Z, Morrison M. Improved extraction of PCR-quality community DNA from digesta and fecal samples. *Biotechniques*. 2004;**36**:808–12.
22. Magoč T, Salzberg SL. FLASH: fast length adjustment of short reads to improve genome assemblies. *Bioinformatics*. 2011;**27**:2957–63.
23. Chen S, Zhou Y, Chen Y, Gu J. fastp: an ultra-fast all-in-one FASTQ preprocessor. *Bioinformatics*. 2018;**34**:i884–90.
24. Rognes T, Flouri T, Nichols B, Quince C, Mahé F. VSEARCH: a versatile open source tool for metagenomics. *PeerJ*. 2016;**4**.
25. Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. DADA2: High resolution sample inference from Illumina amplicon data. *Nat Methods*. 2016;**13**:581–3.
26. Bolyen E, Ram Rideout J, Dillon MR, Bokulich NA, Abnet CC, Al-Ghalith GA, et al. Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2. *Nat Biotechnol*. 2019;**37**:848–57.
27. Lahti L, Shetty S. microbiome [Internet]. 2012.
28. McMurdie PJ, Holmes S. Phyloseq: An R Package for Reproducible Interactive Analysis and Graphics of Microbiome Census Data. *PLoS One*. 2013;**8**.
29. Paulson JN, Stine OC, Bravo HC, Pop M. Differential abundance analysis for microbial marker-gene surveys. *Nat Methods*. 2013;**10**.
30. Dixon P. VEGAN, a package of R functions for community ecology. *J Veg Sci*. 2003;**14**:927–30.
31. Segata N, Izard J, Waldron L, Gevers D, Miropolsky L, Garrett WS, et al. Metagenomic biomarker discovery and explanation. *Genome Biol*. 2011;**12**:R60.
32. Mannaioni G, Carpenedo R, Moroni F. 5-Hydroxyindole causes convulsions and increases transmitter release in the CA1 region of the rat hippocampus. *Br J Pharmacol*. 2003;**138**:245–53.
33. Shannon CE. A Mathematical Theory of Communication. *Bell Syst Tech J*. 1948;**27**:379–423.
34. Simpson E. Measurement of Diversity. *Nature*. 1949;**163**:688.
35. Liang H, Dai Z, Liu N, Ji Y, Chen J, Zhang Y, et al. Dietary L-tryptophan modulates the structural and functional composition of the intestinal microbiome in weaned piglets. *Front Microbiol*. 2018;**9**:1736.
36. Menni C, Zhu J, Le Roy CI, Mompeo O, Young K, Rebholz CM, et al. Serum metabolites reflecting gut microbiome alpha diversity predict type 2 diabetes. *Gut Microbes*. 2020;**11**:1632–42.
37. Zhu C, Sawrey-Kubicek L, Beals E, Rhodes CH, Houts HE, Sacchi R, et al. Human gut microbiome composition and tryptophan metabolites were changed differently by fast food and Mediterranean diet in 4 days: a pilot study. *Nutr Res*. 2020;**77**:62–72.
38. Yusufu I, Ding K, Smith K, Wankhade UD, Sahay B, Patterson GT, et al. A tryptophan-deficient diet induces gut microbiota dysbiosis and increases systemic inflammation in aged mice. *Int J Mol Sci*. 2021;**22**.
39. Vich Vila A, Collij V, Sanna S, Sinha T, Imhann F, Bourgonje AR, et al. Impact of commonly used drugs on the composition and metabolic function of the gut microbiota. *Nat Commun*. 2020;**11**.
40. Zhu L, Liu W, Alkhouri R, Baker RD, Bard JE, Quigley EM, et al. Structural changes in the gut microbiome of constipated patients. *Physiol Genomics*. 2014;**46**:679–86.

Tables

Table 1

Bacterial taxa on the genus level that were affected by the 5-hydroxyindole treatment. Significance was assessed by multiple comparison correction (FDR < 0.05). Marginal effect was assessed by unpaired t-test (P value < 0.05).

Genus	Avg(5-HI)	Sd(5-HI)	Avg(Veh)	Sd(Veh)	p.value	q.value	Difference
<i>Prevotellaceae_UCG-001</i>	0.0222	0.0040	0.0074	0.0023	0.0000	0.0000	0.0148
<i>Shuttleworthia</i>	0.0016	0.0003	0.0003	0.0004	0.0002	0.0174	0.0013
<i>Tuzzerella</i>	0.0012	0.0003	0.0006	0.0003	0.0014	0.0850	0.0007
<i>Allobaculum</i>	0.0309	0.0238	0.0076	0.0030	0.0128	0.4308	0.0234
<i>Parvibacter</i>	0.0002	0.0002	0.0000	0.0000	0.0136	0.4308	0.0002
<i>Prevotellaceae_NK3B31_group</i>	0.0027	0.0010	0.0011	0.0012	0.0205	0.4332	0.0017
<i>Rikenellaceae_RC9_gut_group</i>	0.0026	0.0012	0.0015	0.0005	0.0197	0.4332	0.0012
<i>Lachnospiraceae_NK4B4_group</i>	0.0001	0.0002	0.0000	0.0000	0.0283	0.4889	0.0001
<i>Serratia</i>	0.0108	0.0071	0.0033	0.0054	0.0326	0.5174	0.0075
<i>Eubacterium_eligens_group</i>	0.0008	0.0005	0.0004	0.0002	0.0419	0.6133	0.0004
<i>Acetatifactor</i>	0.0004	0.0003	0.0010	0.0004	0.0163	0.4332	-0.0006
Abbreviations: 5-HI, 5-hydroxyindole-treated group; Veh, Vehicle-treated group; avg, average; sd, standard deviation							

Table 2

Spearman correlations of bacterial taxa on the genus level that were marginally affected by the 5-hydroxyindole treatment.

Genus	Feature	Correlation	p.value	q.value
<i>Lachnospiraceae_UCG-006</i>	TGTT	-0.6702	0.0045	0.7847
<i>Lachnospiraceae_NK4B4_group</i>	TGTT	-0.6696	0.0045	0.7847
<i>Prevotellaceae_NK3B31_group</i>	TGTT	-0.6415	0.0074	0.8313
<i>Tuzzerella</i>	TGTT	-0.5670	0.0220	0.8457
<i>Eubacterium_ventriosum_group</i>	TGTT	-0.5457	0.0288	0.8457
<i>Prevotellaceae_UCG-001</i>	TGTT	-0.5404	0.0307	0.8457
<i>Barnesiella</i>	TGTT	-0.5200	0.0389	0.8457
<i>Bacteroides</i>	TGTT	-0.5107	0.0432	0.8457
<i>Anaerovibrio</i>	TGTT	0.6292	0.0090	0.8450

Figures

Figure 1

5-hydroxyindole has a marginal effect on the cecal microbiota in wild-type Groningen rats. (A) Comparison of cecal microbiota alpha diversity between 5-hydroxyindole-treated group depicted by red bars, and vehicle-treated group depicted by grey bars, including species richness (represented by Chao1 and Observed OTUs) and diversity (represented by Shannon and Simpson index). Data were analyzed using the Mann Whitney test (P value is indicated above the box plots; ns = not significant). Error bars represent SEM. The alpha diversity data can be found in **Table A in Supplementary Table**. (B) Principal component analysis (PCA) indicating no separation of 5-hydroxyindole-treated and vehicle-treated group. Top 10 most contributing species are shown in the figure, the rest of the supporting data for the PCA analysis can be found in **Table E and F in Supplementary Table 1**. (C) LEfSe (Linear discriminant analysis Effect Size) for the 5-hydroxyindole-treated group. The length of the bar represents the $\log_{10}$ transformed Linear discriminant analysis (LDA) score for genera significantly changed in the 5-hydroxyindole-treated group, indicated by vertical dotted lines. (D) Difference in the abundance of the *Prevotellaceae_UCG-001* and *Shuttleworthia* genera in 5-hydroxyindole-treated group (red bars) and vehicle-treated group (grey bars). Left panel represents means in groups. Right panel represents differences between groups, where each dot is colored by its FDR value < 0.05. Significance was assessed by multiple comparison correction. Error bars represent SEM.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SupplementaryTable1.xlsx](#)