Gastrointestinal bleeding caused by metastatic testicular choriocarcinoma: a case report and literature review

Pengliang Zhang

First Affiliated Hospital of Sun Yat-sen University

Yu Wang

First Affiliated Hospital of Sun Yat-sen University

Lishou Xiong (✉ xionglishou@163.com)

First Affiliated Hospital of Sun Yat-sen University

Case Report

Keywords: Testicular choriocarcinoma, Metastatic, Gastrointestinal bleeding, Genital examination

Posted Date: February 23rd, 2022

DOI: <https://doi.org/10.21203/rs.3.rs-1374847/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background: Testicular tumor is one of the common solid tumors in young men. Testicular choriocarcinoma is a non spermatogonial germ cell tumor, which is the most rare of all testicular cancers. Choriocarcinoma usually shows bleeding at the metastatic site, while gastrointestinal involvement is rare.

Methods: Here we report a case of testicular choriocarcinoma with gastrointestinal bleeding as the first diagnosis, and summarize the similar cases all over the world in recent 20 years.

Results: A 28 year old male was treated with repeated black stools for 2 months. No bleeding foci of stomach, duodenum, colon and rectum were found in endoscopy, and no bleeding foci of digestive tract were found in selective angiography, but space occupying lesions of lung, liver and upper jejunum were found in chest and abdominal CT, Considering the possibility of metastatic tumor and the ineffectiveness of medical treatment, the patient was converted to surgical treatment. The postoperative pathology was consistent with testicular choriocarcinoma. The patient received a chemotherapy regimen of paclitaxel, ifosfamide and cisplatin. At present, the chemotherapy regimen is well tolerated.

Conclusions: The case report confirmed that even if we can't find the logical relationship between clinical manifestations and genital examination, genital examination should also be part of the patient's systematic examination.

Introduction

Gastrointestinal bleeding has many clinical manifestations, including hematemesis, black stool and hematochezia, and its common causes include peptic ulcer bleeding, esophageal and gastric variceal bleeding, gastrointestinal primary tumor bleeding and biliary tract bleeding, but bleeding caused by extragastrointestinal tumor metastasis to the digestive tract is very rare in clinic, Especially, bleeding caused by testicular choriocarcinoma metastasis to the gastrointestinal tract is more rare. In 1983, Teryn et al.^[1] Described the first case of jejunal bleeding caused by metastatic testicular choriocarcinoma. This paper summarizes clinical data of a patient with metastatic choriocarcinoma with gastrointestinal bleeding as the clinical manifestation in the First Affiliated Hospital of Sun Yat sen University, combined with the review of relevant literature in the past 20 years, so as to improve the diagnostic rate of the etiology of gastrointestinal bleeding and reduce misdiagnosis and missed diagnosis.

Case Presentation

Clinical data

A 28 year old male patient was admitted to the hospital mainly because of "repeated black defecation for more than 2 months". He denied the history of peptic ulcers, liver cirrhosis and portal hypertension. One year ago, he underwent a CT examination of his chest, upper and lower abdomen and basin due to "right testicular swelling". The results showed that there was a huge tumor on the right testicle with clear

boundary and protruding into the inguinal canal, about 91mm×86mm×119 mm in size(FIGURE 1), uneven density, medium enhancement in enhanced scanning. The left testicle is located in the inguinal canal. Imaging diagnosis: right testicular mass, considering the possibility of seminoma. Left cryptorchidism. No obvious lesions were found in the chest and abdomen. Improve the examination of tumor markers: AFP 86.57ng/ml, HCG 11439 mIU/ml. "Radical resection of the right testis" was performed. The postoperative pathology showed that some tumor cells in testicular tissue were nest like, adenoid or sieve reticular distribution, rich cytoplasm, large nucleus and obvious nucleolus. Syncytial cells were seen with massive bleeding; In addition, squamous epithelium, glandular epithelium, cartilage and small glandular tubular structure can be seen. The morphology is consistent with a malignant mixed germ cell tumor. It is suggested to add immunohistochemical detection to assist in the diagnosis. However, the patient did not undergo immunohistochemical detection, and did not undergo any follow-up treatment such as chemotherapy or radiotherapy. Physical examination after admission: poor spirit, severe anemia, pale eyelid conjunctiva, lips and nail bed, no obvious abnormalities in physical examination such as cardiopulmonary examination. Improve relevant laboratory tests after admission: HB 38g/L, normal cell anemia; Albumin, liver and kidney function and coagulation function were normal.

Diagnosis and treatment process

After admission, the gastroscopy was urgently improved. Gastroscopy showed that there was no bleeding focus in the stomach and duodenum; Colonoscopy showed that a large number of black feces and fecal water were seen in the intestinal cavity, no bright red blood was found, and no bleeding focus was found in the terminal ileum, ileocecal valve, cecum, appendix opening, colon and rectal mucosa(FIGURE 2); Improve selective angiography, see intrahepatic multiple tumor staining, not excluding the possibility of multiple liver metastases, but no clear gastrointestinal bleeding focus has been found. Perfect CT examination of chest and whole abdomen showed that there was a mass in the tongue segment of the upper lobe of the left lung, considering metastasis, and a nodule in the anterior basal segment of the lower lobe of the right lung, considering the possibility of metastasis; Abnormal enhancement of local intestinal tract in the upper jejunum did not rule out angiogenic lesions(FIGURE 3). After medical treatment, the symptoms of black stool were relieved and the hemoglobin increased to 87g/L.

The patient began to eat cold fluid on the 5th day after stopping defecation, but the symptoms of defecation occurred again on the 3rd day. The volume was about 1000ml, blood pressure decreased to 69/44mmhg, and the hemoglobin was about 59g/L, so he was given emergency laparotomy, jejunectomy and ileal diverticulectomy. It can be seen during operation: about 15 cm away from treiz ligament, the upper end of jejunum can be overlapped, and soft masses can be touched. Black intestinal fluid accumulates in this segment of the intestinal cavity. Combined with preoperative CT images, it is considered that the masses here are bleeding sites. Resection of jejunal tumor: the mesentery of the small intestine supplying the tumor segment was ligated and cut off during the operation, with length of about 5cm(FIGURE 4). Postoperative pathology showed that: (small intestine) there was tumor infiltration in the intestinal septum with massive bleeding. The tumor cells showed two forms, one was a rich and bright cytoplasm with a clear boundary, and the other was a multinucleated syncytial cell like; The tumor

cells are heterotypic, with mitotic images and focal necrosis (FIGURE 5). Immunohistochemistry showed: cancer cells CK7 (+), HCG diffuse (+), AFP weak (+), p57 part (+), p53 part (+), CD17 individual cells (+), PLAP focal (+), CD56 individual cells (+), CK20 (-), CDX-2 (-), SATB (-), TTF-1 (-), napsina (-), p40 (-), Sall4 (-), OCT3 / 4 (-), CD30 (-), syn (-). It is considered to be intestinal metastasis of malignant germ cell tumor, and the metastatic component is mainly choriocarcinoma. The patient stopped bleeding after the operation and is currently receiving further chemotherapy. The chemotherapy regimen is tip (paclitaxel, ifosfamide and cisplatin). At present, the patient tolerated the chemotherapy regimen well and was discharged for outpatient follow-up. During the one month follow-up, the patient showed no signs of recurrent gastrointestinal bleeding.

Literature Review

16 cases reported in other journals in the past 20 years were reviewed and analyzed (Table 1). We found that the age of the patients ranged from 17 to 60 years old. Most of them had melena as the first diagnosis symptom (12/16, 75%), and a few had anemia as the first diagnosis symptom (3/16, 18.75%). Only one patient had hematemesis as the first diagnosis symptom, which may be related to the fact that most of the bleeding sites were in the small intestine (10/16, 62.5%), of which duodenal bleeding was the most, with 6 cases. There were only 4 patients with gastric bleeding alone, and 2 patients with bleeding in stomach and colon. Almost all patients received chemotherapy, 6 patients received surgical treatment, but only 1 patient survived (survival rate 16.7%), while 5 patients survived (50%) in 10 patients who received chemotherapy only without surgery. This result may be related to the fact that patients undergoing surgery generally have a large amount of bleeding and serious condition, and surgical intervention must be taken.

Table 1
Clinical data of gastrointestinal bleeding caused by metastatic testicular choriocarcinoma

Case	Year	Age	Symptom	Position of bleeding	Treatment	Ending
1 ^[2]	2002	60	anemia	jejunum	surgical operation	death
2 ^[3]	2004	28	haematemesis	stomach	chemotherapy	death
3 ^[4]	2004	37	anemia	Stomach and colon	chemotherapy	death
4 ^[5]	2005	51	melenas	Stomach and colon	surgical operation+chemotherapy	death
5 ^[6]	2009	24	melenas	Intestinum tenue	surgical operation+chemotherapy	live
6 ^[7]	2010	17	melenas	duodenum	chemotherapy	live
7 ^[8]	2011	25	melenas	stomach	-	death
8 ^[9]	2012	24	melenas	duodenum	chemotherapy	death
9 ^[10]	2013	24	melenas	duodenum	chemotherapy	live
10 ^[11]	2013	20	anemia	stomach	surgical operation+chemotherapy	death
11 ^[12]	2015	18	melenas	stomach	chemotherapy	live
12 ^[13]	2019	30	melenas	duodenum	chemotherapy	death
13 ^[14]	2020	17	melenas	duodenum	chemotherapy	live
14 ^[15]	2021	32	melenas	jejunum	surgical operation+chemotherapy	death
15 ^[16]	2021	33	melenas	Intestinum tenue	surgical operation+chemotherapy	death
16 ^[17]	2021	40	melenas	duodenum	chemotherapy	live
Death refers to death during hospitalization						

Discussion

Testicular cancer is the most common tumor in men aged 15 to 44 ^[18, 19], which is generally divided into germ cell tumors and non germ cell tumors. Germ cell tumors include several cell types, roughly divided into seminoma and non seminoma. Among them, choriocarcinoma is a non seminoma germ cell tumor,

which is the most rare, accounting for 1% - 3% of all testicular tumors^[20]. At present, the cause of testicular choriocarcinoma is not clear, which may be related to a variety of risk factors. Cryptorchidism may be one of the important factors leading to testicular choriocarcinoma. It is reported that the probability of cryptorchidism patients with choriocarcinoma is 20-40 times higher than that of normal testis^[21] in this paper, the patient is left cryptorchidism, which may be one of the important factors leading to testicular choriocarcinoma.

Choriocarcinoma mainly metastasizes through blood. Because of its strong invasiveness to blood vessels and tissues, it leads to tissue bleeding and necrosis. Metastasis occurs early and widely. Therefore, when choriocarcinoma is diagnosed, a large number of cases have metastasized, so that most of the initial manifestations are metastasis related symptoms^[22]. Because the most common metastatic sites are lung, liver and brain^[23], patients usually show seizures, stroke like symptoms, blurred consciousness and-or hemoptysis. Gastrointestinal metastasis of choriocarcinoma is very rare. Gastrointestinal metastasis occurs in 5% of germ cell tumors. Gastrointestinal metastasis is considered to be the result of direct diffusion or hematogenous diffusion from adjacent retroperitoneal lymph nodes, and direct infiltration is more common than hematogenous diffusion. The small intestine, the most common duodenum, is the most common metastatic site (72%), followed by the esophagus, stomach and colon^[6, 24]. The involvement of the small intestine is characterized by intestinal obstruction or gastrointestinal bleeding, usually abdominal pain, black stool or anemia. In this paper, the patient was treated with repeated black stool as the first symptom, but in the follow-up examination, it was found that there were tumor metastasis in the lung and liver in addition to small intestinal metastasis, but did not show clinical symptoms.

The determination of serum tumor markers HCG and AFP may be helpful in the diagnosis of choriocarcinoma because they are elevated in about 80% of cases. The serum concentration of HCG can also be used to monitor the response to treatment. According to the international cooperative organization for germ cell cancer, HCG higher than 50000miu/ml indicates poor prognosis. However, in this paper, the monitoring of HCG level after radical resection of testicular cancer is ignored, which leads to multiple metastasis and poor prognosis. In addition, as a transcription factor, GATA3 is another immune tumor marker sensitive to choriocarcinoma^[25].

The imaging of testicular choriocarcinoma lacks characteristic changes that can be distinguished from other types of germ cell tumors. It is difficult to diagnose choriocarcinoma first. Most of the specimens obtained during surgical resection are confirmed by pathology. The typical histopathological feature of metastatic choriocarcinoma is the coexistence of cytotrophoblast and syncytiotrophoblast cells without mesenchymal cells, which is different from other germ cell tumors with only scattered syncytiotrophoblast cells.

The treatment of testicular choriocarcinoma depends on the stage of the disease. Radical orchiectomy and dissection of affected lymph nodes are the treatment of early diseases. The treatment of bleeding caused by gastrointestinal metastasis of choriocarcinoma is similar to that of other gastrointestinal

bleeding, including endoscopic intervention, embolization or surgical resection. Abdelkader et al reported a case of bleeding from duodenal choriocarcinoma. The exudation point was ablated by an endoscopic adrenalin injection and argon plasma coagulation system to finally stop bleeding ^[12]. Bain et al. ^[7] Reported a case of bleeding treated by angiography and embolization. Iglesias et al. ^[26] Used surgical hemostasis after endoscopic injection of adrenalin and argon plasma coagulation system ablation failure. In this paper, because there was no bleeding focus under endoscopy and selective angiography, the patient finally stopped bleeding by surgical means.

Chemotherapy consolidation is usually required after bleeding stops. For metastatic choriocarcinoma, platinum therapy is recommended as the first-line chemotherapy, but unfortunately, choriocarcinoma is not so sensitive to chemotherapy. Most patients' tumors progress so rapidly that they do not respond to the standard chemotherapy regimen of three to four cycles of BEP (bleomycin, etoposide and cisplatin) ^[27]. In recurrent cases, salvage chemotherapy with vincristine and ifosfamide may help to reduce the tumor burden, but these patients may finally have to choose palliative treatment. Simple testicular choriocarcinoma usually has a poor prognosis, with a five-year survival rate of less than 80% ^[20]. In some reports, the long-term survival rate is even lower ^[28, 29], while mixed choriocarcinoma is slightly better. In this paper, the patient was treated with paclitaxel, ifosfamide and cisplatin.

Testicular choriocarcinoma is a rare tumor with strong invasiveness and rapid growth in young men. It mainly metastasizes to the lung, liver and brain. Reports of metastasis to the gastrointestinal tract are rare, which makes it easy to ignore the existence of the disease in the clinical diagnosis and treatment of gastrointestinal bleeding. Therefore, we recommend a more detailed inquiry into medical history and systematic examination. It is very important for us to correctly distinguish and diagnose the etiology of gastrointestinal bleeding. The lack of accurate physical examination and laboratory examination will lead to waste of patients' diagnosis and treatment time, the increase of mortality, the extension of hospital stay, and the increase of patients' medical expenses. Therefore, for any young male patient, the most basic reproductive system examination is very important. Among the patients in this article, the reproductive system examination was omitted during the patient's repeated hospitalization outside the hospital, and it was not carried out when he was admitted to our hospital. In addition, the detection of HCG was also omitted. Although the final surgical pathology guided us to diagnose testicular choriocarcinoma, during the process, the patient once had hemorrhagic shock, which may have been life-threatening, and we don't know how many patients died because their doctors missed this article.

Conclusion

Testicular choriocarcinoma is a rare malignant tumor with early metastasis. Although it is extremely rare, gastrointestinal metastasis of choriocarcinoma should be a part of the differential diagnosis of upper gastrointestinal bleeding in young male patients. The earlier we diagnose the disease, the greater the opportunity for us to start treatment in the golden age, and the mortality, hospital stay and treatment cost will be greatly reduced.

Abbreviations

CT: Computed tomography

AFP: Alpha fetoprotein

HCG: Human chorionic gonadotropin

CK: Cytokeratin

CD: Cluster of Differentiation

PLAP: Placental alkaline phosphatase

CDX-2: Caudal-type homebox-gene transcription factor-2

SATB: Special AT rich sequence binding protein

TTF-1: Thyroid transcription factor-1

OCT3 / 4: Octamer Binding Transcription Factor 3/4

GATA3: GATA binding protein 3

Declarations

Availability of data and materials

The data that support the findings of this study are available on reasonable request from the corresponding author. The data are not publicly available due to privacy or ethical restrictions.

Acknowledgements

Not applicable

Funding

This manuscript did not receive any funding from any organization.

Author information

Affiliations

Department of Gastroenterology and Hepatology, First Affiliated Hospital of Sun Yat-sen University, Guangzhou, CN

Pengliang Zhang, Yu Wang and Lishou Xiong

Contributions

P.L.Z.: concept/design, data analysis/interpretation, drafting, critical revision. Y.W.: data analysis/interpretation. L.S.X.: concept/design, data interpretation, critical revision, approval. The authors read and approved the final manuscript.

Corresponding author

Correspondence to Lishou Xiong.

Ethics declarations

Ethics approval and consent to participate

Not applicable

Consent for publication

Informed consent for publication was obtained by the patient for this case report presentation.

Competing interests

We declare no support from any organisation for the submitted work; no financial relationships with any organisations that might have an interest in the submitted work; no other relationships or activities that could appear to have influenced the submitted work.

References

1. S Meryn, R Pötzi, W Graninger, T Egger, K Pirich, N Neuhold. **Endoscopic removal of choriocarcinoma metastasis in a patient with chronic intestinal hemorrhage.** Endoscopy. 1983 Jan; 15(1): 34-5.
2. P Zerbib, E Prieur, A Khoury-Helou, P Catala, F R Pruvot, J P Chambon. **Hemorrhagic digestive metastases from testicular choriocarcinoma.** Ann Chir. 2002 Apr; 127 (4): 300-01.
3. R Harikumar, K Harish, K P Aravindan, **Varghese Thomas. Testicular choriocarcinoma with gastric metastasis presenting as hematemesis.** Indian J Gastroenterol. Nov-Dec 2004;23(6):223-24.
4. J Molina Infante, I Beceiro Pedreño, C Ripoll Noiseux, I Marín Jiménez, **C González Asanza, P Menchén Fernández-Pacheco. Gastrointestinal hemorrhage due to metastatic choriocarcinoma with gastric and colonic involvement.** Rev Esp Enferm Dig. 2004 Jan;96(1):77-80.
5. Shahrokh F Shariat 1, David Duchene, Wareef Kabbani, **Zachary Mucher, Yair Lotan. Gastrointestinal hemorrhage as first manifestation of metastatic testicular tumor.** Urology. 2005 Dec;66(6):1319.
6. Irfan C, Filiz O, Hakan K, Fatma Y, Sernaz U, Mustafa K. **Massive upper gastrointestinal bleeding from pure metastatic choriocarcinoma in patient with mixed germ cell tumor with subclinical intestinal metastasis.** Urology. 2009 Feb;73(2):443.e15-17.

7. Andrew JB, David JO, Thomas JS. **Image of the month. Upper gastrointestinal bleeding caused by metastatic testicular choriocarcinoma.** Clin Gastroenterol Hepatol. 2010 Apr;8(4):A22.
8. M Fosado Gayosso, J L Pérez Hernández, F Bernal Sahagún, C Acevedo García, E L Aguilar Ayala, J Pérez Espinosa. **Upper gastrointestinal bleeding as an initial manifestation of metastasis, secondary to a choriocarcinoma in a patient suffering from testicular mixed cell tumour.** Rev Esp Enferm Dig. 2011 Jan;103(1): 46-47.
9. Shuang Fu, Boris Avezbakiyev, Wanqing Zhi, Sreenath Kodali, Kaleem Rizvon, Artur Alaverdian, Lester Freedman, Jose Mejia, Ghulamullah Shahzad, Vladimir Gotlieb. **Germ cell cancer presenting as gastrointestinal bleeding and developing brain metastases: case report and review of the literature.** Future Oncol. 2012 Nov: 8(11): 1487-93.
10. Magdalene Vardaros, Miral Subhani, Kaleem Rizvon, Vladimir Gotlieb, Paul Mustacchia, Lester Freedman, Vikas Garg, Jaspreet Singh, Ghulam Siddiqui. **A case of gastrointestinal bleeding due to duodenal metastasis from a testicular choriocarcinoma.** J Gastrointest Cancer. 2013 Jun;44(2):234-37.
11. Yusuf Yazgan, Kemal Öncü, Mustafa Kaplan, Alpaslan Tanoğlu, Irfan Küçük, Halil Onur Özari, Ufuk Berber, Levent Demirtürk. **Upper gastrointestinal bleeding as an initial manifestation of metastasis secondary to choriocarcinoma.** Turk J Gastroenterol. 2013;24(6):565-67.
12. Kirsty Lowe, Sharon Armstrong, Shaun Walsh, Max Groome, Craig Mowat. **Metastatic Testicular Choriocarcinoma: A Rare Cause of Upper GI Bleeding.** ACG Case Rep J. 2015 Oct; 3(1): 36-38.
13. Abdelkader C, Jason A M,Ahmed A,and Mohammed B. **Metastatic Testicular Choriocarcinoma: An Unusual Cause of Upper Gastrointestinal Bleed.** Cureus.2019 Jul;11(7):e5243.
14. Shahryar Zeighami, Nima Naghdi Sede, Fatemeh Khajeh, Ali Ariafar, Mohammadreza Soltani, Zahra Jahanabadi, Aliasghar Khaleghi, Zahra Bazargan. **Genital exam, a missed piece of the puzzle in medical diagnosis, can be lifesaving in men: A lesson from a case of a state of shock due to duodenal metastasis of testicular choriocarcinoma.** Qatar Med J. 2020 Nov 9:2020(2):31.
15. Jennifer Yoon, Steve Hu, Jessica Farrell, Kandarp K Shah, Jaya Krishna Chintanaboina. **An Exceptionally Rare Cause of Refractory Gastrointestinal Bleed: Choriocarcinoma Syndrome.** Cureus. 2021 Apr 20:13(4):e14599.
16. Cherihane Dassouli, Chama El Manjra, Adil Ait Errami, Sofia Oubaha, Zouhour Samlani, Khadija Krati. **Small bowel metastases from a testicular choriocarcinoma revealed by gastrointestinal bleeding: a case report.** Pan Afr Med J. 2021; 39: 259.
17. Ahmed Elfiky, Asmaa Mokhtar, Mira Alsheikh, Hassan Almoussawi, Stephen Mulrooney. **Gastrointestinal Bleeding Secondary to Metastatic Duodenal Choriocarcinoma in a Patient with Concomitant Peptic Ulcer Disease.** Case Rep Gastrointest Med. 2021; 2021: 6664147.
18. McGlynn KA, Cook MB. **Etiologic factors in testicular germ-cell tumors.** Future Oncol. 2009;5:1389-402.
19. Ghazarian AA, Trabert B, Devesa SS, McGlynn KA. **Recent trends in the incidence of testicular germ cell tumors in the United States.** Andrology. 2015; 1: 13-18.

20. Reilley MJ, Pagliaro LC. **Testicular choriocarcinoma: a rare variant that requires a unique treatment approach.** Curr Oncol Rep. 2015;**17**:2.
21. Richie JP. **Re: a meta-analysis of the risk of boys with isolated cryptorchidism developing testicular cancer in later life.** J Urol, 2013,**190**(3): 1045.
22. Smith ZL, Werntz RP, Eggener SE. **Testicular cancer: Epidemiology, diagnosis, and management.** Med Clin North Am. 2018;**102**:251-64.
23. Smith ZL, Werntz RP, Eggener SE. **Testicular cancer: epidemiology, diagnosis, and management.** Med Clin North Am, 2018,**102**(2): 251-64.
24. Chait M. M., Kurtz R. C., Hajdu S. I. **Gastrointestinal tract metastasis in patients with germ-cell tumor of the testis.** The American Journal of Digestive Diseases. 1978; **23**(10):925-28.
25. Miettinen M., McCue P. A., Sarlomo-Rikala M., *et al.* **GATA3: a multispecific but potentially useful marker in surgical pathology: a systematic analysis of 2500 epithelial and nonepithelial tumors.** American Journal of Surgical Pathology. 2014; **38**(1):13-22.
26. Iglesias García J., López García E., Macías García F., *et al.* **Severe gastrointestinal bleeding secondary to duodenal metastasis from a choriocarcinoma.** Revista Española de Enfermedades Digestivas. Oct 2009; **101**(10): 720-22.
27. Rajpert-De Meyts E, McGlynn KA, Okamoto K, Jewett MA, Bokemeyer C. **Testicular germ cell tumours.** Lancet. 2016;**387**:1762-74.
28. Papiani G, Einhorn LH. **Salvage chemotherapy with high-dose carboplatin plus etoposide and autologous peripheral blood stem cell transplant in male pure choriocarcinoma: a retrospective analysis of 13 cases.** Bone Marrow Transplant. 2007; **40**:235-37.
29. Batata MA, Chu FC, Hilaris BS, Papantoniou PA, Whitmore WF Jr, Golbey RB. **Therapy and prognosis of testicular carcinomas in relation to TNM classification.** Int J Radiat Oncol Biol Phys. 1982(**8**):1287–93.

Figures

Figure 1

CT images of patients. The red arrow indicates the right testicular tumor, and the yellow arrow indicates the left cryptorchidism.

Figure 2

The picture a and b indicates gastroscopy results, the picture c and d indicates colonoscopy results, and no active bleeding focus is found.

Figure 3

The picture a represents chest CT results, the picture b represents abdominal CT results, and red arrows represent metastases.

Figure 4

A gross specimen removed surgically (Specimen fixed by fixing liquid).

Figure 5

The tumor cells are heterotypic, with mitotic images and focal necrosis.