Langerhans cell histiocytosis of the uvea with a ciliochoroidal mass: a Case report of management with systemic therapy

Zahra Mahdizad

Tehran University of Medical Sciences

Hamid Riazi-Esfahani

Tehran University of Medical Sciences

Nazanin Ebrahimiadib

Tehran University of Medical Sciences

Abdulrahim Amini

Tehran University of Medical Sciences

Fariba Ghassemi (✉ fariba.ghassemi@gmail.com)

Tehran University of Medical Sciences

Short Report

Keywords: Langerhans cell histiocytosis, choroidal mass, choroidal Langerhans cell histiocytosis, case report

Posted Date: February 21st, 2022

DOI: <https://doi.org/10.21203/rs.3.rs-1374911/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Abstract

Background. This study aims to explore a case of choroidal Langerhans cell histiocytosis (LCH) successfully treated with systemic corticosteroid and immunosuppressant.

Case presentation. A 24-year-old man with known multisystem LCH developed loss of vision, ocular pain, conjunctival injection, panuveitis, and a ciliochoroidal mass. *After receiving an intravenous methylprednisolone pulse, oral high-dose corticosteroids, and methotrexate, the mass resolved quickly and completely without flare-ups during 6 months of follow-up.*

Conclusions. Intraocular involvement of LCH is rare and can present with or without a history of multisystemic disease. The diagnosis is challenging and the standard treatment is not established. Systemic anti-inflammatory and immunosuppressive therapy could be an effective treatment, as the LCH itself contains an essential element of inflammation and the symptoms may be mainly inflammatory.

Background

Langerhans cell histiocytosis (LCH) is the most common histiocytic disorder characterized by the accumulation of histiocytes in various tissues (1). It includes a broad spectrum of clinical presentations, ranging from self-healing lesions to life-threatening disseminated disease (1). The disorder is most common in children (peaks between ages 1 and 4), but anyone can be affected (2). Although there is strong evidence for an underlying inflammatory process (3), a recent mutation found in LCH patients of BRAF-V600E makes the cancerous etiology more clear [1, 4]. Therefore, some may consider LCH to be inflammatory myeloid neoplasia (5).

LCH can be diagnosed by clinical and paraclinical findings, but in particular by histopathological analyses demonstrating the presence of histiocytes with specific characteristics of Langerhans cells (LCs) (6). More frequently affected organs are the bones, skin, pituitary gland, liver, spleen, hematopoietic system, and lungs (1). Most cases of LCH present with ophthalmic involvement, namely orbital involvement, whereas intraocular involvement is rare [3, 7].

Case Presentation

A 24-year-old man was referred to our oncology clinic for evaluation of an amelanotic mass in his left eye. His symptoms have included pain, redness, proptosis, and visual loss in his left eye, as well as nasal congestion and posterior nasal drip over the past few months. About 3 months before, he underwent ethmoid and maxillary sinus biopsies, which were reported to be a non-specific orbital inflammation (NSOI) with extension to the sinuses based on the presence of only CD34 positive cells. While the patient was treated with systemic steroids, his symptoms (especially the loss of vision) did not improve significantly. The patient presented with a history of bilateral anterior uveitis in remission in addition to juvenile idiopathic arthritis (JIA). His history also included multisystemic LCH since childhood, which was diagnosed based on his symptoms (lymphadenopathy, skin eruptions, frontal bone lesions, sinusitis, epilepsy, and diabetes insipidus) and the presence of CD1a positive cells in lymph node specimens. He received systemic prednisolone and vinblastine, which were tapered upon improvement of his condition. At the time of presentation, he was taking oral prednisolone 50 mg daily, valproic acid 500 mg daily, eye drop betamethasone QID, and eye drop homatropine TDS.

His best-corrected visual acuity (BCVA) in the right eye (OD) was 10/10, while in the left eye (OS) it was 6/10. On the temporal side of OS, diffuse conjunctival injections with localized scleral and episcleral thickenings were found with dilated (sentinel) vessels on the temporal side. A 2+ anterior chamber (AC) and anterior vitreous (AV) reactions were also detected. In the far peripheral region of the superotemporal area of the retina, a subretinal creamy-white mass with a cuff of subretinal fluid (SRF) was found (Figure 1B). On ultrasonography, a choroidal mass with a diameter of 12x9x6.7 mm was seen with a homogenous internal reflectivity that extended anteriorly to the ciliary body (Figure1C-1D)

Fluorescein angiography (FA) in OD was unremarkable while in OS it showed posterior and peripheral vasculitis, as well as leakage from the optic disc. (Figure 1E)

The diagnosis of unilateral panuveitis with anterior scleritis and inflammatory choroidal mass was made. The systemic workup for rheumatologic disease and granulomatosis disease came back negative and the medical examination revealed no other systemic focus of disease activity. Ibuprofen (400 mg QID) and methotrexate (15 mg weekly) were administered. In the following follow-up sessions by 1 month, both his symptoms and the size of the choroidal lesion decreased substantially.

A flare-up occurred two months after the initial presentation. The BCVA of OS was counting finger at 3 meters and the examination revealed conjunctival injection, chemosis, hazy media, and AC, and AV inflammation (3+) along with a larger mass (15x11.5x9.7mm). He was admitted with the diagnosis of LCH associated panuveitis and choroidal mass. Intravenous methylprednisolone pulse was administered for 3 days, and he was discharged with prescription of oral prednisolone 50 mg daily, methotrexate 15 mg weekly, and Ibuprofen 400mg every 4 hours. The pain, conjunctival injection as well as SRF amount decreased in a few days and after two weeks his BCVA improved to 2/10, the AC and AV inflammation decreased to trace cells and the mass size decreased significantly as it could hardly be seen in the fundus examination and B-scan. Subretinal white deposits with trace SRF at the site of resolved mass were visible (Figure 2A-C). In addition to discontinuing the ibuprofen, the dosage of the

corticosteroid medication was tapered cautiously. The patient has been followed for the last six months with a stable condition, and his vision improved to 9/10.

Discussion And Conclusions

LCH is a rare systemic disorder characterized by the proliferation of clonal S-100, CD1a, and CD207 positive Langerhans cells in a wide range of organs (8). Due to its myeloid origin and inflammatory characteristics, it has been considered a myeloid neoplasm (5). A number of mutations, including BRAF (the most common one), MAP2K1, and ARAF have been identified in LCH patients, all of which activate MAPK/ERK signaling pathway (9).

A typical pathologic histiocyte in LCH consists of mononucleated cells with indented or kidney-shaped nuclei and a Birbeck granule; however, in recent years the search for CD1a and CD207 (langerin) expression has largely replaced Birbeck granule detection [6, 10]. Furthermore, BRAF V600E mutation analysis in plasma and urine has shown to be an effective diagnostic tool (9).

LCH may affect any organ of the body (individually or in conjunction). The disease is classified clinically as single-system Langerhans cell histiocytosis (SS-LCH) or multisystem Langerhans cell histiocytosis (MS-LCH) with multiple organ involvement. The hematopoietic system (bone marrow), spleen, and liver are also at risk (8). Isolated pulmonary LCH is the most common manifestation in adults, but diabetes insipidus (DI) is the most prevalent initial sign of central nervous system involvement in LCH.

Intraocular involvement has mostly been reported in case reports (7). The choroid is the most common intraocular site of manifestation and it may be infiltrated diffusely or as a solitary mass (11). Although there have been reports of abnormal LCs infiltrating the sclera, retina, vitreous, uveal tract, and optic nerve sheath in patients with and without a history of multisystemic LCH, this is a rare occurrence [10–12].

The treatment of LCH depends on its heterogeneous presentation; generally, patients with a single-system disease only need local treatment (e.g. radiotherapy) or observation, while those with a more extensive disease require systemic therapy (9). The standard treatment for MS-LCH has yet to be identified, but vinblastine and prednisolone therapy have been suggested (as with our patient); short-term chemotherapy with MTX, doxorubicin, cyclophosphamide, vincristine, and bleomycin, as well as allogeneic hematopoietic stem cell transplantation have shown efficacy in more aggressive cases (8). Clinical trials of targeted therapies with selective BRAF and MEK1/2 inhibitor medications are ongoing and seem to be effective [8, 9].

Our patient presented at the age of 24 with unilateral panuveitis, scleritis, and choroidal mass refractory to the treatment with oral prednisolone. Considering his unremarkable laboratory work-up and systemic disease, we concluded that MS-LCH was the cause of his ophthalmic symptoms. Similar cases of choroidal LCH presenting as unilateral choroidal masses, or panuveitis, or a combination of both have been reported (Table1)[7, 10, 12, 13, 14]. In one case, a diagnosis of choroidal melanoma was made, but immunohistochemical studies revealed LCH [10], and in the second eye, the iris portion of an iridociliochoroidal mass was found to be LCH after fine-needle aspiration (7). Some reported vitreous biopsy for diagnosis of intraocular LCH [17], whereas others made the diagnosis based on a systemic history of LCH [11, 15, 16], as we did for our patient.

Table 1
reported cases of choroidal Langerhans cell histiocytosis mass

No.	Authors	Year	Age	Sex	Presentation	Tumor size	Diagnosis method	Treatment	Outcome	Ref.
1	Shields et al.	2010	6	M	Iridociliochoroidal mass and neovascular glaucoma	A 14 mm in basal diameter and 8.4 mm in thickness	Fine needle aspiration biopsy	Low dose plaque radio-therapy of	Complete regression to a flat scar of 2 mm thickness and no sign of radiation retinopathy	7
2	Kim and Lee	2000	49	M	Choroidal mass resembling choroidal melanoma	6 DD	Tumor biopsy	Enucleation (with a diagnosis of choroidal melanoma)	-	10
3	Narita et al.	1993	61	F	A solitary massive tumor and recurrent anterior uveitis		biopsy	Enucleation		12
4	Patton et al.	2006	29	M	Choroidal mass	5 DD, 2mm thickness	Patient history of Central nervous system Langerhans cell histiocytosis lesion	Low-dose external fractionated radiotherapy	A rapid improvement in both visual symptoms and clinical appearance	13
5	Angell and Burton	1978	5	F	Choroidal mass and secondary retinal detachment	1.75mm	Patient history of MS-LCH	-	-	14
6	Mahdizad et al.	2021	24	M	Choroidal mass, panuveitis and scleritis	15x11.5x9.7mm	Patient history of MS-LCH	intravenous methylprednisolone pulse followed by oral high-dosage corticosteroid and methotrexate	Rapid and complete resolution	

Considering the rare incidence of intraocular involvement of LCH, there is no standard treatment. Patton et al.(13) reported the first case of choroidal involvement as a mass by LCH and documented successful outcomes of fractionated low-dose, whole-eye external beam radiotherapy for choroidal LCH. Shields and colleagues (7), used brachytherapy for treatment of an iridociliochoroidal tumor in a 6-year-old boy with known MS-LCH. The tumor responded rapidly with complete regression.

As LCH itself contains an essential element of inflammation and our patient had inflammatory symptoms and a first response to systemic corticosteroid, we decided to continue with high-dose corticosteroids (intravenous methylprednisolone pulse) in combination with immunosuppression (methotrexate), which proved to be effective. To the best of our knowledge, this is the first case of choroidal mass associated with LCH, successfully treated with systemic corticosteroid and immunosuppressant without radiotherapy.

Abbreviations

LCH: Langerhans cell histiocytosis; LCs: Langerhans cells; NSOI: non-specific orbital inflammation; JIA: juvenile idiopathic arthritis; CD: cluster of differentiation; QID: quater in die; BCVA: best-corrected visual acuity; OD: oculus dexter; OS: oculus sinister; AC: anterior chamber; AV: anterior vitreous; SRF: subretinal fluid; FA: Fluorescein angiography; SS-LCH: single-system Langerhans cell histiocytosis; MS-LCH: multisystem Langerhans cell histiocytosis; DI: diabetes insipidus.

Declarations

Ethics approval and consent to participate

“Not applicable”

Consent for publication

Written informed consent was obtained from the patient for publication of this case report and any accompanying images. A copy of the written consent is available for review by the Editor-in-Chief of this journal.

Availability of data and materials

The datasets used during the current study available from the corresponding author on reasonable request.

Competing interests

The authors declare that they have no competing interests.

Funding

No funding was obtained for this study.

Authors' contributions

"HR visited and followed up the patient and edited the manuscript. NE visited and admitted the patient for treatment. FGH made the final decision about the treatment approach and was a major contributor in writing the manuscript. ZM gathered the patient's documents and wrote the manuscript. AA edited the manuscript. All authors read and approved the final manuscript. "

Acknowledgements

"Not applicable"

References

1. Krooks J, Minkov M, Weatherall AG. Langerhans cell histiocytosis in children: History, classification, pathobiology, clinical manifestations, and prognosis. *J Am Acad Dermatol*. 2018 Jun;78(6):1035–44.
2. Hamre M, Hedberg J, Buckley J, Bhatia S, Finlay J, Meadows A, et al. Langerhans cell histiocytosis: an exploratory epidemiologic study of 177 cases. *Med Pediatr Oncol*. 1997 Feb;28(2):92–7.
3. Sitaula RK, Khatri A. Langerhans cell histiocytosis with hemorrhagic uveitis and exudative retinal detachment. Vol. 11, *International medical case reports journal*. 2018. p. 65–8.
4. Badalian-Very G, Vergilio J-A, Degar BA, MacConaill LE, Brandner B, Calicchio ML, et al. Recurrent BRAF mutations in Langerhans cell histiocytosis. *Blood* [Internet]. 2010;116(11):1919–23. Available from: <https://www.sciencedirect.com/science/article/pii/S0006497120329955>
5. Berres M-L, Merad M, Allen CE. Progress in understanding the pathogenesis of Langerhans cell histiocytosis: back to Histiocytosis X? *Br J Haematol*. 2015 Apr;169(1):3–13.
6. Emile J-F, Abila O, Fraitag S, Home A, Haroche J, Donadieu J, et al. Revised classification of histiocytoses and neoplasms of the macrophage-dendritic cell lineages. *Blood*. 2016 Jun;127(22):2672–81.
7. Shields CL, Hogarty MD, Kligman BE, Christian C, Ehya H, Shields JA. Langerhans cell histiocytosis of the uvea with neovascular glaucoma: diagnosis by fine-needle aspiration biopsy and management with intraocular bevacizumab and brachytherapy. *J AAPOS Off Publ Am Assoc Pediatr Ophthalmol Strabismus*. 2010 Dec;14(6):534–7.
8. Kobayashi M, Tojo A. Langerhans cell histiocytosis in adults: Advances in pathophysiology and treatment. *Cancer Sci*. 2018 Dec;109(12):3707–13.
9. Monsereenusorn C, Rodriguez-Galindo C. Clinical Characteristics and Treatment of Langerhans Cell Histiocytosis. *Hematol Oncol Clin North Am*. 2015 Oct;29(5):853–73.
10. Kim IT, Lee SM. Choroidal Langerhans' cell histiocytosis. *Acta Ophthalmol Scand*. 2000 Feb;78(1):97–100.
11. Thanos A, Vavvas D, Young LH, Foster CS. Choroidal neovascular membrane formation and retinochoroidopathy in a patient with systemic langerhans cell histiocytosis: a case report and review of the literature. Vol. 3, *Case reports in ophthalmology*. 2012. p. 128–35.
12. Narita K, Yoshida K, Yoshimoto H, Matsuyama S, Kamata Y. [A case of intraocular histiocytosis X]. *Nihon Ganka Gakkai Zasshi*. 1993 May;97(5):649–54.
13. Patton N, Lai T, Robbins P, Holthouse D, Barry C, Constable I. Presumed choroidal langerhans cell histiocytosis following a previously resected solitary central nervous system lesion in an adult. *Arch Ophthalmol (Chicago, Ill 1960)*. 2006 Aug;124(8):1193–5.
14. Heath P. The Ocular Features of a Case of Acute Reticuloendotheliosis (Letterer-Siwe Type). *Trans Am Ophthalmol Soc*. 1959;57:290–302.
15. Angell LK, Burton TC. Posterior choroidal involvement in Letterer-Siwe disease. *J Pediatr Ophthalmol Strabismus*. 1978;15(2):79–81.
16. Tsai JH, Galaydh F, Ching SST. Anterior uveitis and iris nodules that are associated with Langerhans cell histiocytosis. *Am J Ophthalmol*. 2005 Dec;140(6):1143–5.

Figures

Figure 1

A 24-year-old man presented with left conjunctival injection (A), pain, and visual loss. He was found to have an amelanotic choroidal mass (B) and panuveitis. The mass was located temporally, had medium homogenous internal reflectivity, and measured 12* 9*6.7 mm by ultrasonography (C) and spread anteriorly to the iris root (D). Fluorescein angiography revealed optic nerve head leakage, posterior and peripheral vasculitis, and staining of the lesion (E).

Figure 2

Two weeks following therapy, slit-lamp examination of the left eye revealed reduced conjunctival injection and scleral edema (A), fundus examination demonstrated significant decrease in mass size and subretinal white deposits with trace SRF at the site of resolved mass (B), b-scan also showed resolution of choroidal mass with small amount of subretinal fluid(C).