

# Performance of age-adjusted D-dimer values for predicting DVT before the knee and hip arthroplasty

**Jian Xiang Wu**

Nanjing University Medical School Affiliated Nanjing Drum Tower Hospital

**Jiang Hui Qing**

Nanjing University Medical School Affiliated Nanjing Drum Tower Hospital

**Yao Yao**

Nanjing University Medical School Affiliated Nanjing Drum Tower Hospital

**Dong Yang Chen**

Nanjing University Medical School Affiliated Nanjing Drum Tower Hospital

**Qing Jiang** (✉ [qingj@nju.edu.cn](mailto:qingj@nju.edu.cn))

Drum Tower Hospital, School of Medicine, Nanjing University <https://orcid.org/0000-0002-2552-9686>

---

## Research article

**Keywords:** preoperative D-dimer, age-adjusted D-dimer, joint arthroplasty, ultrasonography, screening DVT

**Posted Date:** December 1st, 2020

**DOI:** <https://doi.org/10.21203/rs.2.23200/v3>

**License:**  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---

**Version of Record:** A version of this preprint was published on January 25th, 2021. See the published version at <https://doi.org/10.1186/s13018-020-02172-w>.

# Abstract

**Purpose:** To compare the specificity and sensitivity of preoperative D-dimer and age-adjusted D-dimer value for predicting the incidence of the DVT preoperatively in total joint arthroplasty (TJA) patients.

**Methods:** We enrolled 406 patients finally above 50 years old. Everyone had done ultrasonography bedside, and D-dimer concentrations were collected before surgery. The D-dimer and age-adjusted D-dimer cut-off was calculated by multiple logistic regression and receiver operating curve (ROC) analyses.

**Results:** A total of 39 patients had found asymptomatic deep vein thrombosis (DVT) by ultrasonography. The age (odds ratio [OR]: 1.067;  $p=0.003$ ) and D-dimer (OR:1.331;  $p=0.025$ ) were related to the existence of DVT. For conventional D-dimer and age-adjusted D-dimer value, the area under the curves (AUCs) were 0.685 (0.499-0.696) and 0.795 (0.611-0.881), respectively.

**Conclusion:** Compared to traditional D-dimer, age-adjusted D-dimer showed better performance in screening DVT, which was useful clinically.

## Introduction

As is known to all, the total hip arthroplasty (THA) and total knee arthroplasty (TKA) are the effective methods to treat osteoarthritis, but it also can bring some complications, such as infection, pain and deep vein thrombosis (DVT), which was the most common complication. The incidence of DVT is about 11.9% in THA and 20.8% in TKA after surgery. [1] DVT can lead to Pulmonary thromboembolism (PTE), which is the most serious complication after arthroplasty. There are many researches focus on the incidence of DVT postoperatively, few studies about preoperative DVT were reported. [2] In clinical work, the removal of thrombus preoperatively may need the help of imaging examination such as ultrasound, contrast radiography and so on. However, there are inadequacies in all the above examinations, such as time-consuming, expensive and labor-consuming, so they are not suitable for large-scale use before operation. If patients have asymptomatic DVT preoperatively, there might cause severe consequence. Thus, a valuable method in detection of DVT or PTE before surgery is necessary.

In clinical work, the prompt diagnosis for thrombosis is the D-dimer test, which can give references to the surgeon. [3] D-dimer is produced by fibrin degradation. [4] The monitoring value of D-dimer is identified according to different methods. The cut off value of D-dimer is 0.5 $\mu$ g/ml by latex agglutination turbidimetry (LATEX) in our hospital, however, in the elderly patients, the D-dimer concentrations higher than the upper Limit of normal with age. [5] Thus, the value of D-dimer is discounted in the elderly patients. [6] Moreover, the pregnancy, infection, inflammation and cancer also can lead to the high D-dimer value. As a result, the only use of D-dimer to predict DVT is not appropriate.

Some studies showed that the use of age-adjusted D-dimer cut-off in the diagnostic strategy for the specificity and sensitivity of deep vein thrombosis is more valuable over 50 years old. [7] [8] Mads Nybo et al. argued that the use of an age-adjusted D-dimer in patients above 50 years of age for ruling out DVT

seems as safe as using a standard D-dimer cut-off. [9] Schouten et al. also described that the application of age-adjusted cut-off values to the D-dimer value ( $0.01 \times \text{age} [\text{years}]$ ) considerably elevates specificity without altering sensitivity in patients aged >50 years with suspected PTE. [10]

Therefore, the use of D-dimer concentrations for predicting deep vein thrombosis is controversial. [11] However, the application of age-adjusted cut-off values to the D-dimer value is relatively rare in orthopaedics. So, it is valuable for us to research. The method of this study is to compare the specificity and sensitivity of preoperative D-dimer and age-adjusted D-dimer value for predicting the incidence of the DVT preoperatively. Our aim is to explore the feasibility and effectiveness of this method in the field of joint replacement.

## Materials And Methods

We enrolled 834 patients, whose age was over 50 years old and underwent THA or TKA at Affiliate Drum Tower Hospital, Medical school of Nanjing University from September 1<sup>st</sup> 2015 to April 1<sup>st</sup> 2017. All patients signed the informed consent. The patients who underwent the pregnancy, infection, and cancer were excluded. In addition, patients were excluded if had a femoral neck fracture, whose concentration of d-dimer was above normal level. Also, the patients with cardiovascular disease, a previous history of thromboembolism were excluded. We removed the patients who had taken anticoagulant or antiplatelet agents previously. In the end, we included 406 patients in our study. All the patients drew blood for collecting the plasma D-dimer level when were admitted to hospital. The cut off value of D-dimer was  $0.5\mu\text{g/ml}$  by latex agglutination turbidimetry (LATEX). We divided patients aged >50 into 10 year age groups. Renee A Douma made three large cohorts to find the age-adjusted D-dimer value. They plotted the D-dimer cut-off level against age group and performed linear regression analysis to obtain the regression coefficient. They finally proved the multiplication factor for age in the new age-adjusted cut-off value was 0.01. Thus, the age-adjusted D-dimer value was calculated as  $\text{age (years)} \times 0.01\mu\text{g/mL}$ . The plasma D-dimer level, diagnosis, age, sex and body mass index (BMI) of the patients were recorded preoperatively. All the patients did not take anticoagulant or antiplatelet agents. The surgeries were done by four skilled orthopaedical surgeons. The patients accepted the same program of rehabilitation and postoperative management. We observed vein thrombosis 1 day preoperatively. The skilled radiologists performed thrombosis examinations for all patients. The double lower limbs were observed with bedside ultrasound machine with two mode (B-mode and color Doppler ultrasound examination) for the common femoral, popliteal, superficial and calf veins bilaterally. DVT was diagnosed that : (1) the vein could not be compressed. (2) there were abnormal vascular signals in doppler color ultrasound. Plasma D-dimer normal value was defined as  $0.5\mu\text{g/ml}$ . We wanted to analysis the age-adjusted D-dimer cut-off to predict the DVT event. At the same time, we also analyzed the risk factors for preoperative deep venous thrombosis. We predicted that the age-adjusted D-dimer was a more valuable index for monitoring the DVT before the TKA and THA.

All statistical analyses were from SPSS (version 25; SPSS Inc., Chicago, IL, USA). The qualitative data such as the number of females or males was analyzed by Fisher's exact test. The quantitative data such

as body height, body weight, age, body mass index, and D-dimer value was analyzed by unpaired Student's t-test. Multiple logistic regression analysis was used to discover the following risk factors related to the presence of DVT: body height, body weight, body mass index (BMI), age, sex, and D-dimer value. Receiver operating characteristic (ROC) curves were produced for age-adjusted D-dimer value and conventional D-dimer value. Also, the area under the curve (AUC) was calculated by ROC curves. In addition, we used the Youden index [12] to determine the cut-off value. We considered  $P < 0.05$  to be statistically significant.

## Results

In the end, we included 406 patients. There were 118 males and 288 females. The average BMI of all patients was  $25.3 \pm 3.9 \text{ kg/m}^2$ . The mean age was  $65.4 \pm 8.4$  years old. The patients' detailed preoperative basic data and related blood tests are shown in Table I.

**Table I Patients characteristics**

	Total patients	Patients with DVT	Patients without DVT	P value
Patient (n)	406	39	367	-
Sex(male/female)	118/288	9/30	109/258	0.386
Age(years)	$65.4 \pm 8.4$	$69.8 \pm 6.9$	$65.0 \pm 8.5$	$< 0.001$
Body weight(kg)	$66.0 \pm 10.8$	$67.5 \pm 11.6$	$65.7 \pm 10.7$	0.316
Body height(cm)	$161.4 \pm 7.0$	$160.5 \pm 6.3$	$161.5 \pm 7.0$	0.307
BMI ( $\text{kg/m}^2$ )	$25.3 \pm 3.9$	$27.2 \pm 3.2$	$25.0 \pm 3.9$	0.201
Preoperative D-dimer ( $\mu\text{g/ml}$ )	0.45(0.25,0.84)	0.91(0.61,1.12)	0.66(0.32,0.98)	0.045
APTT (s)	$28.1 \pm 3.8$	$28.1 \pm 4.6$	$28.1 \pm 3.7$	0.968
PT (s)	$12.0 \pm 0.9$	$12.0 \pm 0.8$	$12.0 \pm 0.9$	0.865
INR	$1.04 \pm 0.08$	$1.04 \pm 0.07$	$1.04 \pm 0.08$	0.927
Platelet count ( $10^9/\text{L}$ )	$206.9 \pm 61.7$	$194.8 \pm 55.8$	$208.2 \pm 62.2$	0.193
Hb(g/L)	$128.7 \pm 18.4$	$126.1 \pm 12.0$	$129.0 \pm 18.9$	0.091

DVT: deep vein thrombosis; BMI: body mass index; APTT: activated partial thromboplastin time  
PT: prothrombin time; INR: international normalized ratio; Hb: hemoglobin

We divided patients aged  $>50$  into 10 year age groups. We found that the incidence of thrombosis in the group older than 80 years old was higher than that in the other groups, and the difference was statistically significant. In addition, we compared conventional D-dimer and age-adjusted D-dimer cut-off by age group, more patients below cutoff values could be found in all age groups using age-adjusted D-dimer cut-off values. The specific information is in the Table II.

**Table II Comparison of different cut-off values stratified by age group**

	Total patients	Age range (years)			
		50-59	60-69	70-79	>80
No(%) of patients	406	103(25.4)	182(44.8)	99(24.4)	22(5.4)
Median age (years)	63.3	54.7	64.9	73.5	83.3
No(%) of patients with DVT	39	3(2.9)	16(8.8)	16(16.2)	4(18.2)
Conventional cut-off value					
No (%) of patients below cut-off value	226(55.7)	73(70.9)	99(54.4)	46(46.5)	8(36.4)
Age-adjusted cut-off value					
No (%) of patients below cut-off value	269	78(75.7)	118(64.8)	60(60.6)	12(54.5)

Total of 39 patients had found deep vein thrombosis of the lower extremities, all of which were asymptomatic. The incidence of thrombosis was 9.6%. Two of them were iliac vein thrombosis, and the rest were intramuscular venous thrombosis. The mean age of patients with thrombosis was  $69.8 \pm 6.9$  years, and the mean age of patients without thrombosis was  $65.0 \pm 8.5$  years, with a statistically significant difference ( $P < 0.001$ ). For preoperative D-dimer levels, patients with thrombosis had an average of  $0.91(0.61,1.12)$  mg/mL, compared with  $0.66(0.32,0.98)$  mg/mL for patients without thrombosis, and there was a statistically significant difference. ( $P = 0.045$ ).

By multivariate logistic regression analysis, when age, gender, BMI, body weight, and preoperative D-dimer levels were independent variables, we found that only age and D-dimer levels were associated with the development of deep vein thrombosis (Table III).

**Table III Multiple logistic regression analysis for preoperative deep vein thrombosis**

	OR	95% CI	p-value
Age (year)	1.067	1.023-1.113	0.003
Sex (female)	0.615	0.208-1.814	0.378
Body height (cm)	0.977	0.910-1.049	0.52
Body weight (kg)	1.022	0.988-1.057	0.209
D-dimer ( $\mu\text{g/ml}$ )	1.331	1.032-1.716	0.027

CI, confidence interval OR, odds ratio

With regard to preoperative D-dimer, a total of 180 patients had a D-dimer concentration greater than  $0.5\mu\text{g/mL}$ , of which only 18 had thrombosis. With the use of age-adjusted D-dimer, there were only 33 patients' D-dimer concentration above the age-adjusted D-dimer reference value. In addition, we plotted the ROC curve for the age-adjusted D-dimer and the traditional D-dimer (Figure I). The area under the curve was 0.795 and 0.685, respectively (Table IV). For DVT observations, the traditional D-dimer had a cut-off value of 0.845 with a sensitivity of 41% and a specificity of 0.658 (Table IV). The cut-off value of

age-adjusted D-dimer cut-off value was 0.655 which caused a sensitivity of 0.769 and a specificity of 0.768 (Table IV).

**Table IV AUC and cut-off**

	cut-off value	Sensitivity	Specificity	AUC	95% CI	p-value
Age-adjusted D-dimer	0.655	0.769	0.768	0.795	0.611-0.881	< 0.001
D-dimer	0.845	0.41	0.658	0.685	0.499-0.696	0.045

AUC, area under the curve; CI, confidence interval

## Discussion

Patients with joint replacement were generally older. If the traditional D-dimer value was used as a preoperative diagnosis method, the false positive patient would be greatly increased. [5]

We used both age-adjusted D-dimer and traditional D-dimer values to detect the effectiveness of deep vein thrombosis. The area under the ROC curve was 0.795 and 0.685, respectively, and the age-adjusted D-dimer had a better predictive value in predicting deep vein thrombosis. In addition, we found that the sensitivity of the age-adjusted D-dimer cut-off value (76.8%) was significantly improved compared to the traditional method (41%), while the specificity, age-adjusted D-dimer cut-off value (76.9%) was also slightly better than the traditional D-dimer (65.8%).

The basic D-dimer value increased with age. P. L. Harper et al. found that D-dimer concentration increased with age reducing the clinical value of the D-dimer assay in the elderly. [5]

Through the age-stratification study, we found that the elderly had higher incidence of thrombosis, which is consistent with the conclusion of our risk factor analysis, and by the age-adjusted D-dimer cut-off value in all age groups, it can be found that more samples are lower than the cut-off value, which improved the sensitivity of the model.

In our study, the age-adjusted D-dimer value was defined as age x 0.01 ( $\mu\text{g}/\text{mL}$ ), which was considered safe. Mads Nybo et al. had found that the age-adjusted D-dimer cut-off was safe to recommend the use of an age-adjusted D-dimer in a DVT setting as well as for PE from a systematic review. [9] In this study, we also made multivariate logistic regression analysis to find that age is the risk factor of DVT.

In previous reports, Norio Imai et al. demonstrated that age and D-dimer index can be useful in screening patients for DVTs before THA. [13] The conclusion was similar to us. Moreover, age-adjusted D-dimer value was calculated as patients' age times 0.01 ( $\mu\text{g}/\text{mL}$ ) by Henrike J Schouten et al. [10] They drew

conclusions that age-adjusted D-dimer value can increase specificity, whereas sensitivity was not. The sensitivity was 97%, but the specificity was below 60%. The results did not coincide with our research. In addition, some studies focused on the performance of age adjusted D-dimer cut-off values for reducing the use of ultrasounds. [3] [14] [15] Kelly Broen et al. inclined the view that age adjusted D-dimer cut-off values can lead to less duplex ultrasounds performed to find patients that suffers from DVT. [16]

In our study, the age-adjusted D-dimer value cut-off increased sensitivity, which was important to find patients with DVT. DVT was the most critical complication. Once we did not observe the asymptomatic DVT patients, it might cause serious consequences, such as pulmonary embolism (PE). Thus, the age-adjusted D-dimer could help us find more asymptomatic DVT patients. Actually, that was what we cared about more.

As far as we know, it was the first study to focus on the performance of age adjusted D-dimer cut-off values for the screening of deep vein thrombosis in both TKA and THA. We gave the lower extremity venous color ultrasound bedside to all patients admitted to the hospital. Thus, we had enough samples for observation.

However, there were several limitations in our study. First, we did not use Wills score to assess the sensitivity and specificity of age-adjusted D-dimer cut-off value. Wills score could help us predict DVT better. [17] Second, our hospital used latex agglutination turbidimetry (LATEX) to measure the concentrations of D-dimer. Compared to enzyme linked fluorescent assays (ELISA), LATEX may lead lower sensitivity and specificity in our study. Finally, we did not group patients to several level by age, which may get a better result in this study.

In conclusion, compared to traditional D-dimer, age-adjusted D-dimer showed better performance in screening DVT, which was useful clinically. Meanwhile, more clinical observations were needed to verify our ideas.

## List Of Abbreviations

Deep vein thrombosis: DVT

The knee arthroplasty: TKA

The hip arthroplasty: THA

The joint arthroplasty: TJA

Pulmonary thromboembolism: PTE

Body mass index: BMI

Receiver operating characteristic: ROC

Area under the curve: AUC

Activated partial thromboplastin time: APTT

Prothrombin time: PT

International normalized ratio: INR

Hemoglobin: Hb

Confidence interval: CI

Odds ratio: OR

Pulmonary embolism: PE

Enzyme linked fluorescent assays: ELISA

Latex agglutination turbidimetry: LATEX

## **Declarations**

### **Ethics approval and consent to participate**

This study was approved by the institutional review board (IRB) of Nanjing Drum Tower Hospital (No. 2012029).

### **Consent for publication**

Not applicable

### **Availability of data and materials**

The data used to support the findings of this study are available from the corresponding author upon request.

### **Competing interests**

The authors declare that they have no competing interests

### **Funding**

This work was supported by Key Program of NSFC (81730067), Jiangsu Provincial Key Medical Center Foundation, Jiangsu Provincial Medical Outstanding Talent Foundation, Jiangsu Provincial Medical Youth Talent Foundation, Jiangsu Provincial Key Medical Talent Foundation, National Natural Science Foundation of China (81972124) and Basic Research Program of Jiangsu Province (20170123).

## Authors' contributions

Qing Jiang contributed substantially to the conception and design of the study. Jian Xiang Wu, Jiang Hui Qing, Yao Yao and Dong Yang Chen collected and analyzed the data. Jian Xiang Wu wrote the manuscript with support from Yao Yao. All authors read and approved the final manuscript.

## Acknowledgement

This work was supported by Key Program of NSFC (81730067), Jiangsu Provincial Key Medical Center Foundation, Jiangsu Provincial Medical Outstanding Talent Foundation, Jiangsu Provincial Medical Youth Talent Foundation, Jiangsu Provincial Key Medical Talent Foundation, National Natural Science Foundation of China (81972124) and Basic Research Program of Jiangsu Province (20170123). We thank the patients and their families who donated their blood samples for this study.

## References

1. Yamaguchi, T., et al., *Incidence and time course of asymptomatic deep vein thrombosis with fondaparinux in patients undergoing total joint arthroplasty*. *Thromb Res*, 2010. 126(4): p. e323-6.
2. Dai, X., et al., *Associations of Serum Lipids and Deep Venous Thrombosis Risk After Total Knee Arthroplasty in Patients With Primary Knee Osteoarthritis*. *Int J Low Extrem Wounds*, 2019: p. 1534734619868123.
3. Sartori, M., et al., *D-dimer for the diagnosis of upper extremity deep and superficial venous thrombosis*. *Thromb Res*, 2015. 135(4): p. 673-8.
4. Ruiz-Gimenez, N., et al., *Rapid D-dimer test combined a clinical model for deep vein thrombosis. Validation with ultrasonography and clinical follow-up in 383 patients*. *Thromb Haemost*, 2004. 91(6): p. 1237-46.
5. Harper, P.L., et al., *D-dimer concentration increases with age reducing the clinical value of the D-dimer assay in the elderly*. *Intern Med J*, 2007. 37(9): p. 607-13.
6. Cini, M., et al., *D-dimer use for deep venous thrombosis exclusion in elderly patients: a comparative analysis of three different approaches to establish cut-off values for an assay with results expressed in D-dimer units*. *Int J Lab Hematol*, 2014. 36(5): p. 541-7.
7. Farm, M., et al., *Age-adjusted D-dimer cut-off leads to more efficient diagnosis of venous thromboembolism in the emergency department: a comparison of four assays*. *J Thromb Haemost*, 2018. 16(5): p. 866-875.
8. Li, J., et al., *The Diagnostic Efficacy of Age-Adjusted D-Dimer Cutoff Value and Pretest Probability Scores for Deep Venous Thrombosis*. *Clin Appl Thromb Hemost*, 2019. 25: p. 1076029619826317.
9. Nybo, M. and A.M. Hvas, *Age-adjusted D-dimer cut-off in the diagnostic strategy for deep vein thrombosis: a systematic review*. *Scand J Clin Lab Invest*, 2017. 77(8): p. 568-573.
10. Schouten, H.J., et al., *Diagnostic accuracy of conventional or age adjusted D-dimer cut-off values in older patients with suspected venous thromboembolism: systematic review and meta-analysis*. *BMJ*,

2013. 346: p. f2492.

11. Haenssle, H.A., et al., *Limited diagnostic value of Wells-score and D-dimer testing in hospitalized dermatologic patients with symptoms of deep vein thrombosis*. Eur J Dermatol, 2013. 23(6): p. 830-6.
12. Fluss, R., D. Faraggi, and B. Reiser, *Estimation of the Youden Index and its Associated Cutoff Point*. Biometrical Journal, 2005. 47(4): p. 458-472.
13. Imai, N., et al., *Usefulness of a novel method for the screening of deep vein thrombosis by using a combined D-dimer- and age-based index before total hip arthroplasty*. PLoS One, 2017. 12(2): p. e0172849.
14. Riva, N., et al., *Age-adjusted D-dimer to rule out deep vein thrombosis: findings from the PALLADIO algorithm*. J Thromb Haemost, 2018. 16(2): p. 271-278.
15. Prochaska, J.H., et al., *Age-related diagnostic value of D-dimer testing and the role of inflammation in patients with suspected deep vein thrombosis*. Sci Rep, 2017. 7(1): p. 4591.
16. Broen, K., B. Scholtes, and R. Vossen, *Predicting the need for further thrombosis diagnostics in suspected DVT is increased by using age adjusted D-dimer values*. Thromb Res, 2016. 145: p. 107-8.
17. Sartori, M., et al., *The Wells rule and D-dimer for the diagnosis of isolated distal deep vein thrombosis*. J Thromb Haemost, 2012. 10(11): p. 2264-9.

## Figures




Figure 1

the ROC curve for the age-adjusted D-dimer and the traditional D-dimer

## Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SupplementaryMaterial.pdf](#)