Fecal Metabolome and Bacterial Composition in Severe Obesity: Impact of Diet and Bariatric Surgery

Nuria Salazar

Instituto de Productos Lácteos de Asturias (IPLA-CSIC)

Manuel Ponce-Alonso

Servicio de Microbiología. Hospital Universitario Ramón y Cajal, & Instituto Ramón y Cajal de Investigación Sanitaria (IRYCIS), & CIBERINFECT

María Garriga

Hospital Universitario Ramón y Cajal, & Instituto Ramón y Cajal de Investigación Sanitaria (IRYCIS)

Sergio Sánchez-Carrillo

Consejo Superior de Investigaciones Científicas

Ana María Hernández-Barranco

Instituto de Productos Lácteos de Asturias (IPLA-CSIC)

Begoña Redruello

Instituto de Productos Lácteos de Asturias (IPLA-CSIC)

María Fernández

Instituto de Productos Lácteos de Asturias (IPLA-CSIC)

José Ignacio Botella-Carretero

Hospital Universitario Ramón y Cajal, & Instituto Ramón y Cajal de Investigación Sanitaria (IRYCIS), & CIBER Epidemiology and Public Health (CIBERESP)

Belén Vega-Piñero

Hospital Universitario Ramón y Cajal, & Instituto Ramón y Cajal de Investigación Sanitaria (IRYCIS), & CIBER Epidemiology and Public Health (CIBERESP)

Javier Galeano

Universidad Politécnica de Madrid

Javier Zamora

Hospital Universitario Ramón y Cajal, & Instituto Ramón y Cajal de Investigación Sanitaria (IRYCIS), & CIBER Epidemiology and Public Health (CIBERESP)

Manuel Ferrer

Consejo Superior de Investigaciones Científicas

Clara G. de los Reyes Gavilán

Instituto de Productos Lácteos de Asturias (IPLA-CSIC)

Rosa del Campo (✉ rosacampo@yahoo.com)

Servicio de Microbiología. Hospital Universitario Ramón y Cajal, & Instituto Ramón y Cajal de Investigación Sanitaria (IRYCIS), & CIBERINFECT

Research Article

Keywords:

Posted Date: February 23rd, 2022

DOI: <https://doi.org/10.21203/rs.3.rs-1378909/v1>

Abstract

Background: The aim of this study was to monitor the impact of a preoperative diet and bariatric surgery on the bacterial gut microbiota composition and functionality in severe obesity and to compare sleeve gastrectomy (SG) *versus* Roux-en-Y gastric bypass (RYGB). The study also aimed to incorporate big data analysis for the omics results and machine learning by a Lasso-based analysis to detect the potential markers for excess weight loss.

Methods: Forty patients who underwent bariatric surgery were recruited (14 underwent SG, and 26 underwent RYGB). Each participant contributed 4 fecal samples (baseline, post-diet, 1 month after surgery and 3 months after surgery). The bacterial composition was determined by 16S rDNA massive sequencing using MiSeq (Illumina). Metabolic signatures associated to fecal concentrations of short-chain fatty acids, amino acids, biogenic amines, gamma-aminobutyric acid and ammonium were determined by gas and liquid chromatography. Orange 3 software was employed to correlate the variables, and a Lasso analysis was employed to predict the weight loss at the baseline samples.

Results: A correlation between Bacillota (formerly Firmicutes) abundance and excess weight was observed only for the highest body mass indexes. The crash diet had little impact on composition and targeted metabolic activity. RYGB had a deeper impact on bacterial composition and putrefactive metabolism than SG, although the excess weight loss was comparable in the two groups. Significantly higher ammonium concentrations were detected in the feces of the RYGB group.

Conclusions: We detected individual signatures of composition and functionality, rather than a gut microbiota characteristic of severe obesity, with opposing tendencies for almost all measured variables in the two surgical approaches. The gut microbiota of the baseline samples was not useful for predicting excess weight loss after the bariatric process.

Background

Bariatric surgery is the most cost-effective treatment for reducing body mass index (BMI) in severe obesity (BMI > 40 or > 35 combined with comorbidities) and resolving endocrine-related dysfunctions such as type 2 diabetes mellitus (DM2) [1, 2]. The most common surgical approaches are Roux-en-Y gastric bypass (RYGB) and sleeve gastrectomy (SG), both by laparoscopy. SG is a restrictive method based on stomach reduction, whereas RYGB combines both restrictive and malabsorptive approaches through stomach reduction and the anatomical reorganization of the biliary and digestive transits. Selecting the appropriate surgical approach requires a consensus between the patient and surgeon based on BMI, dietary habits and comorbidities, with RYGB recommended for a BMI $\geq$ 50 and for patients with compulsive eating disorder [3]. Although the physical reduction of the stomach forces drastic diet changes, not all patients achieve complete loss of the baseline excess weight.

The gut microbiota's contribution to obesity has been repeatedly discussed in recent years, but no solid conclusions have been reached, particularly for severe obesity. Bariatric surgery not only affects bacterial gut composition [4, 5], with RYGB causing deeper changes, but also its overall functionality [1, 6]. Beyond the bacterial composition, scientific research is currently focused on microbial metabolism, particularly the mechanisms involved in DM2 remission [7], as well as on the microbiome functionality evolution [8], including short-chain fatty acids (SCFAs) [9] and other microbial products related to the excess weight loss. The aim of the present study was to monitor the impact of a preoperative crash diet and bariatric surgery on bacterial gut microbiota composition and functionality in severe obesity, to compare the RYGB and SG approaches, and to apply a Lasso predictive analysis based on multiple linear regressions to detect potential markers of excess weight loss.

Methods

Study design and participants. A prospective study was conducted with 40 participants (22 women and 18 men, all older than 18 years) who were recruited during 2015–2017 at the Bariatric Surgery Unit of Ramón y Cajal University Hospital in Madrid, Spain. None of the participants reported infectious gastrointestinal disorders or antibiotic consumption in the last 3 months. Our center's ethics committee approved the study (accession number 379/14), and all participants voluntarily signed an informed consent for participation in the study, which did not interfere with the bariatric surgical procedure.

Each participant provided 4 fecal samples at the following points: 1) baseline, the initial visit at the Bariatric Surgery Unit; 2) post-diet, after 1 month of the crash diet; 3) 1 month after surgery; and 4) 3 months after surgery. Immediately after collection, the feces were aliquoted and frozen at -80°C until processing.

The baseline visit took place 1 month prior to the surgery to provide nutritional instructions with a low-calorie, high-protein crash diet to reduce liver size in order to facilitate laparoscopic access (1400 calories for women and 1700 for men, distributed as 38% protein, 36% carbohydrates and 26% fat). Regardless of the surgical approach, the postoperative diet included only liquids for the first 2 days, followed by purées/soft food for the first month; thereafter, the patients received only general nutritional advice with no specific restrictions, with recommendations for a high protein intake (to prevent the loss of muscle mass) and multivitamin and mineral complexes. All participants were administered antibiotic prophylaxis during surgery with 2 g of intravenous amoxicillin/clavulanic acid.

Fecal metabolome. The SCFA concentration was determined in cell-free fecal supernatants prepared as previously described [10] using gas chromatography in a system composed of a 6890N injection module (Agilent Technologies Inc., Palo Alto, CA, USA) connected to a flame injection detector and a mass spectrometry 5973N detector (Agilent). Samples were analyzed in triplicate and the results were expressed as medians.

Quantification of amino acids, amines, ammonium and gamma-aminobutyric acid (GABA) was performed by ultra-high-performance liquid chromatography on cell-free fecal supernatants following diethyl ethoxymethylenemalonate (DEEMM, Sigma-Aldrich, USA) derivatization. Briefly, fecal supernatants were filtered through 3kDa centrifugal filters (Amicon Ultra-0.5, Merk KGaA, Germany), and DEEMM derivatization reactions were performed in 100 μl as previously described by Redruello et al. [11]; however, this was the first time that the procedure was performed on feces. The accuracy of the chromatographic method employed to quantify all of the studied amino compounds was tested using a standard addition procedure [12]. Thus, two known concentrations of analytes were independently added (in triplicate) to cell-free fecal supernatants. Non-spiked sample replicates (blanks) were used to determine the sample's initial analyte content. The percentage recovery at each concentration was calculated as $[(\text{amount found in the spiked sample}) - (\text{amount found in the blank}) / \text{amount added}] \times 100$. The recovery percentages for all analytes ranged from 80.1 to 104.4 when 0.2 mM was added (the accepted range for this concentration is 80–110) and from 90.1 to 104.3 when 1 mM was added (the accepted range is 90–107), reflecting the method's good accuracy [12]. Samples were also filtered through 0.22- μm -pore diameter polytetrafluoroethylene membranes (VWR International, USA) into conical vials (VWR) prior to injection into the ultra-high-performance liquid chromatography system [11].

Bacterial gut microbiota characterization. Samples were slowly thawed, first at -20°C for 24 h, followed by another 24 h at 4°C . After that, 0.5 g of the samples was solubilized in 5 ml of sterile water, centrifuged and extracted with the Speedtools tissue DNA extraction kit (Biotools, Madrid, Spain). The resulting total DNA was used for sequencing (2×300 bp) the V3 and V4 regions of the 16S rDNA gene on a MiSeq platform (Illumina, USA). To prepare the library, the 16S Metagenomic Sequencing Library Preparation protocol (Illumina; Cod. 15044223 Rev. A) was used with the following primers: Forward (5'TCGTCGGCAGCAGCGTCGTCAGATGTGTAAGACAGCCTACGGNGGCWGCAG), and Reverse (5'GTCTCGTGGCTCGAGGAGGTAAGAGAGACGACTACHVGGTATCTAATCC).

The quality of the raw sequences was measured according to the minimum length (250 bp), type of trimming quality measure (medium), quality value for trimming from the 3' end (30) and trimming quality window (10 bp). Taxonomic affiliations were assigned using the SILVA 119 database and reads with a Ribosomal Database Project score < 0.8 were assigned to the higher taxonomic rank, leaving the last rank as unidentified. The relative abundance and contingency tables of operational taxonomic units (OTUs) included taxa with very low representation. Sequences not assigned to any taxon or classified as non-bacterial were eliminated. Alpha and beta diversity studies were performed using the q2-diversity add-on of QIIME2 [13], after normalizing the samples by rarefaction (subsampling without replacement). In addition, a linear effect size discriminant analysis (LEfSe) [14] was performed to assess which taxa explained the differences between groups. Sequence data were deposited in Genbank (BioProject PRJNA639545).

Statistical analysis. For the statistical analysis of the bacterial diversity results, we employed the q2-diversity plugin for QIIME2. In those groups/variables with statistically significant differences, we further explored which taxa explained these differences,

using LEfSe, which employs the Kruskal–Wallis test and the pairwise Wilcoxon test. The alpha value for the Kruskal–Wallis factorial test was set at 0.05, and the linear discriminant analysis score threshold was set at 2.0.

The correlation between bacterial abundance and the various study metabolites was explored using a Big Data strategy, calculating the Spearman coefficient in Orange 3 software (<https://orangedatamining.com>) and the Wilcoxon test in GraphPad Prism 9 (GraphPad Software, USA, <http://www.graphpad.com>).

Lasso regression model. To predict the excess weight loss at 3 months after surgery using only numerical variables from the baseline sample, we employed a Lasso regression model, a machine learning-based analysis with multiple linear regressions. The independent variable was the percentage of excess weight loss (%EWL), defined as the remaining excess weight 3 months after surgery divided by the excess weight in the baseline sample. The excess weight at time 2 (post-diet) and time 3 (1 month after surgery) were used as the dependent variables. Given that many dependent variables were analyzed and to avoid overfitting, the most important variables were delimited using the correlation between the main variable and the dataset of bacterial composition, SCFAs, amino acids and biogenic amines. The resulting dataset was divided into a test dataset (25%) and a training dataset (75%) to generate the Lasso model. The advantage of using this model is that it allows for a new selection of functions to obtain a better relationship between the test score and the training score. To measure the goodness of the method, the root mean square error was calculated. All calculations were performed using Python scripts developed by the authors (https://github.com/galeanojav/Severe_Obesity).

Results

%EWL. Table 1 summarizes the participants' relevant clinical and demographic data, and Fig. 1 provides an illustration of the %EWL progression. The RYGB group ($n = 26$) had significantly higher excess weight than the SG group ($n = 14$), both at baseline (57.2 vs. 47.3, $p = 0.006$) and after the crash diet (55.6 vs. 47.8, $p = 0.02$). In fact, the crash diet resulted in immediate weight loss only in the RYGB group; 3 months after surgery, however, the two groups had a similar %EWL (50.1% for RYGB and 49.0% for SG).

Table 1
Main characteristic of recruited severe obese (M = male, F = female).

Main characteristic of recruited severe obese (M = male, F = female).							
Sex/Age (years)	T2DM/ Resolution	Basal BMI	Weight Excess in Sample (kg)				Final % EWL
SG (n = 14)							
F/58		50.4	55.7	53.2	47.2	38.7	69.4
M/58		40.4	45.7	48.4	34.6	27.4	59.9
F/34		42.8	41.2	38.7	27.4	17.2	41.7
F/66		47.8	48.7	97.7	25.7	16.5	33.8
M/64	+/-	36.4	32.9	28.7	20.4	14.1	38.7
F/37		39.1	38.9	39.1	26.8	18.7	48.0
M/65	+/+	40.8	41.4	47.3	31.4	24.9	60.1
F/48		46.3	57.3	57.7	46.6	35.8	62.4
F/45		45.0	53.3	51.5	40.0	29.3	54.9
M/39		44.3	69.1	66.5	52.5	39.0	56.4
M/59		41.5	52.4	52.7	49.7	28.5	54.3
F/68		42.1	45.5	46.5	33.8	25.9	56.9
F/49		47.2	58.3	58.3	47.9	30.8	52.8
F/40		43.7	44.9	38.1	28.9	18.1	40.3
Media		43.1	47.3	47.8	33.8	23.5	50.1
RYGB (n = 26)							
M/57		44.5	61.9	52.8	39.1	27.8	44.9
F/56	+/+	47.5	58.2	55.2	46.7	40.5	69.5
M/50		45.5	68.0	54.8	38.8	32.5	47.7
M/59	+/+	40.7	50.9	44.0	31.8	23.0	45.1
M/48		63.4	109.6	100.6	79.8	66.6	60.7
F/49		59.3	99.3	94.8	79.3	66.1	66.5
F/62		43.5	44.4	60.1	38.9	25.9	58.3
F/43	+/+	50.1	69.9	76.3	52.3	42.3	60.5
F/53		50.7	69.9	69.9	48.9	48.9	69.9
F/63	+/+	44.3	51.2	52.4	42.4	26.1	50.9
M/56	+/-	56.8	94.0	79.0	60.0	50.0	53.1
F/62	+/+	38.0	34.1	30.2	21.8	14.8	43.4
F/62	+/+	36.5	28.6	37.6	25.6	12.1	42.3
M/62		47.6	68.3	61.7	51.3	32.9	48.1
F/58		46.1	52.6	51.6	41.1	22.6	42.9

Sex/Age (years)	T2DM/ Resolution	Basal BMI	Weight Excess in Sample (kg)				Final % EWL
			1	2	3	4	
M/53		47.7	67.0	49.8	35.2	17.5	26.1
M/51		48.4	70.2	70.7	54.3	43.0	61.2
M/47		47.9	75.9	76.6	57.9	43.7	57.5
M/37		48.7	66.8	57.0	47.5	38.3	57.3
M/50	+/-	38.2	44.3	44.3	20.3	17.6	39.7
F/42		50.7	69.9	74.3	58.1	42.5	60.8
M/36	+/+	60.5	115.0	107.0	89.5	78.4	68.1
F/50		45.2	54.9	47.2	37.5	25.7	46.8
F/63		39.9	38.1	40.0	28.4	20.0	52.4
M/60	+/+	44.0	54.2	57.6	29.6	14.8	27.3
F/36		41.7	43.9	41.4	33.4	23.2	52.8
Media		46.3	57.2	55.6	39.9	27.3	49.0

Bacterial Composition and Distribution. The 160 fecal samples yielded a total of 11,889,557 reads corresponding to 250 OTUs. The data were normalized by establishing a sampling depth of 24,758 reads per sample. No statistically significant differences in the alpha diversity values were detected by sampling point or type of surgery (Fig. 2A). The Shannon index remained stable, but the Chao1 index experienced opposing trends for the SG (slight decrease) and RYGB (slight increase) groups, without reaching statistical significance in any case. There were no statistically significant correlations between alpha diversity and the BMI data (data not shown).

Beta diversity based on the Bray-Curtis dissimilarity index was comparable for all patients at baseline and post-diet. From this point on, however, the progression of this index in the microbiota diverged for the two groups (Fig. 2B), with significant changes in the SG group starting only after surgery ($p = 0.03$); these changes in SG patients were considerably less pronounced than for the RYGB group ($p = 0.0015$).

The taxonomic analysis demonstrated the dominance of Bacillota (formerly Firmicutes) in the samples from both surgical groups (Fig. 3A), with a total of 167 bacterial genera identified, 14 of which accounted for more than 80% of the overall abundance (Fig. 3B). The differential traits in the microbiota of the SG group included the increase in Bacillota abundance across the bariatric process and the unique representation of certain minority genera such as *Butyricoccus*, *Eggerthella*, *Gordonibacter* and *Intestinibacter* (data not shown).

In the SG group, there was a significant increase in certain genera of the phylum Bacillota (*Hungatella*, *Dorea* and *Eisenbergiella*) as the result of a decrease in the abundance of Bacteroidota after the crash diet. In contrast, the RYGB group showed a slight increase in the genera *Ruminococcus* and *Anaerococcus* and a decrease in the families *Lactobacillaceae* and *Bifidobacteriaceae*.

In terms of the surgical process, the intergroup differences were greater. Overall, bariatric surgery led to a decrease in the genera *Roseburia*, *Faecalibacterium*, *Ruminococcus* and *Bifidobacterium* and a significant increase in *Escherichia/Shigella* and *Akkermansia*. When stratifying according to type of surgery, the differences between the samples at baseline and at the end of the follow-up were much more profound in the RYGB group (Fig. 4A), in accordance with the Bray-Curtis dissimilarities indexes. Thus, the Proteobacteria, Bacteroidota, Verrucomicrobiota and Fusobacteriota phyla experienced a significant increase in number at 3 months after RYGB surgery, inversely to Bacillota and Actinomycetota. However, the changes were considerably

less marked in the SG group (Fig. 4B), with a slight enrichment of certain Bacillota, such as *Streptococcus*, *Parvimonas*, *Hungatella*, *Lactobacillus* and *Desulfovibrio*, together with a decrease in Bacteroidota and Negativicutes.

The dynamics of the Bacillota phylum during the entire bariatric surgical process was particularly noteworthy for its opposing trends: the SG group maintained or even increased the high initial relative proportions, whereas the RYGB group experienced a marked decrease (Fig. 3A). In the SG group, the sole correlation between weight and Bacillota abundance was the excess weight in the baseline sample (Fig. 5, $r = 0.29$), whereas in the RYGB group, the %EWL was directly correlated to the decrease in Bacillota during the follow-up ($r = 0.31$).

Short-Chain and Branched-Chain Fatty Acids. Table 2 and Fig. 6 show the absolute concentrations and relative abundances of these microbial metabolites. The relative abundance of the main SCFAs (acetate > 50%, propionate ~ 20% and butyrate 15–20%) was comparable for the two groups. In general, there were higher concentrations of all measured SCFAs in the feces of the SG group, reaching statistical significance only for valeric acid ($p = 0.04$). There were no significant differences in SCFA and branched-chain fatty acid (BCFA) abundance according to sex, age or excess weight in the baseline sample (data not shown). The concentrations of each SCFA experienced moderate variations in the subsequent samples. The crash diet promoted a significant increase in the concentration of isobutyric acid, a branched SCFA, whereas no statistically significant changes were observed for the samples from the RYGB group. The postoperative fecal samples from the SG group had significantly decreased acetic and propionic acid concentrations compared with the concentrations after the crash diet. There was also a statistically significant decrease in acetic acid levels in the RYGB samples and a statistically significant increase in isobutyric acid levels. Statistically significant correlations for SCFA and BCFA and specific bacterial taxa were not detected.

Table 2
Median, standard deviation and range of SCFA values (mM) measured at the 4 times for both groups of patients.

Median ± SD Range (mM)	Acetate	Propionate	Butyrate	Isobutyrate	Valerate	Isovalerate	Caproate
SG*	e	e		a			b
Basal	83.1 ± 28.8 43.6–132.4	29.9 ± 10.5 6.9–44.2	29.9 ± 10.5 6.9–44.2	2.3 ± 1.1 0.5–4.7	3.8 ± 2.3 0.7–10.3	2.8 ± 2.1 0.7–8.3	0.5 ± 0.9 0.08–3.4
Diet	92.1 ± 37.8 37.9–178.6	31.6 ± 17.9 8.3–82.8	31.6 ± 17.9 8.3–82.8	3.1 ± 1.0 0.7–4.5	4.0 ± 2.2 0.8–8.6	4.3 ± 1.5 1.4–6.2	0.5 ± 0.4 0.04–1.5
1 month	61.8 ± 42.2 10.4–168.5	25.0 ± 30.1 2.6–111.9	25.0 ± 30.1 2.6–111.9	2.5 ± 1.6 0.4–6.5	3.0 ± 2.6 0.4–10.4	3.9 ± 2.5 0.6–9.2	0.1 ± 0.3 0.0–0.8
3 months	62.6 ± 24.9 19.8–123.9	25.3 ± 9.6 3.9–40.7	25.3 ± 9.6 3.9–40.7	2.9 ± 1.3 1.0–5.7	3.6 ± 1.4 0.8–5.6	3.9 ± 2.3 1.5–8.6	0.3 ± 0.5 0.06–1.7
RYGB*	b,c,e			c		c,e	
Basal	69.1 ± 31.7 19.7–126.1	24.8 ± 12.3 9.7–52.3	17.2 ± 15.5 0.6–58.3	2.1 ± 1.0 0.2–4.3	3.2 ± 2.0 0.0–9.3	2.9 ± 1.5 0.9–6.8	0.7 ± 0.7 0.0–3.3
Diet	69.9 ± 33.0 22.8–158.2	23.5 ± 17.2 3.8–73.4	18.3 ± 14.8 0.0–66.7	2.1 ± 1.5 0.9–6.7	3.4 ± 2.8 0.4–14.2	2.8 ± 2.3 1.0–10.6	0.7 ± 0.6 0.0–2.4
1 month	53.6 ± 25.1 14.2–114.9	21.7 ± 13.1 3.1–51.2	10.5 ± 10.2 0.0–35.2	2.4 ± 1.1 0.5–5.7	3.3 ± 2.4 0.0–11.7	3.6 ± 1.7 1.6–9.0	0.5 ± 0.4 0.0–1.6
3 months	52.9 ± 22.9 16.0–91.1	26.5 ± 15.8 3.6–55.7	16.1 ± 14.9 0.9–56.4	3.0 ± 1.5 0.9–8.7	2.9 ± 2.1 0.6–7.8	4.7 ± 2.8 1.4–15.7	0.4 ± 0.6 0.0–3.1
*a: statistically significant difference determined by Wilcoxon test between baseline and diet; b: baseline and 1 month; c: baseline and 3 months; d: diet and 1 month; and e: diet and 3 months. Only values of $p \leq 0.05$ were considered.							

Amino acids, biogenic amines and GABA. Fecal concentrations of amino acids and biogenic amines are shown in Tables 3 and 4, respectively. Significant differences were observed between bariatric surgery groups, as well as different trends throughout the study period within each group. Globally, higher concentrations of amino acids were found in the SG samples both after the crash diet and after surgery, while the biogenic amines cadaverine, histamine and tyramine had significantly higher concentrations in fecal samples of the group of RYGB patients.

Table 3

Median, standard deviation and range of amino acid values (mM) measured at the 4 times for both groups of patients.

Median $\pm$ SD Range (mM)	Ala	Arg	Asn	Asp	Gln	Glu	Gly	His	Ile	Leu	Lys
SG*	a,c	a			a,c	a			c	c	
Basal	4.2 $\pm$ 3.8 0.8–15.5	0.3 $\pm$ 0.3 0.06–1.3	0.03 $\pm$ 0.2 nd–1.1	1.3 $\pm$ 0.9 0.4–3.7	0.1 $\pm$ 0.2 nd–0.9	3.5 $\pm$ 3.6 1.0–14.9	1.2 $\pm$ 1.5 0.2–6.5	0.1 $\pm$ 0.1 0.04–0.5	0.9 $\pm$ 1.0 0.05–3.6	1.5 $\pm$ 1.5 0.05–5.3	2.0 $\pm$ 1.3 0.05–5.0
Diet	8.2 $\pm$ 5.2 1.6–20.5	0.4 $\pm$ 0.3 0.08–1.2	0.02 $\pm$ 0.1 nd–0.7	1.0 $\pm$ 0.6 0.6–2.9	0.2 $\pm$ 0.1 0.01–0.4	7.3 $\pm$ 3.0 0.2–12.1	1.8 $\pm$ 1.9 0.4–7.1	0.2 $\pm$ 0.2 0.03–0.8	1.4 $\pm$ 1.3 0.2–5.1	3.2 $\pm$ 1.7 0.5–6.7	2.6 $\pm$ 1.5 0.8–5.8
1 month	6.5 $\pm$ 5.2 1.6–24.4	0.2 $\pm$ 0.2 0.02–0.7	0.02 $\pm$ 0.04 nd–0.1	1.1 $\pm$ 1.0 0.2–3.8	0.1 $\pm$ 0.2 0.08–1.1	4.9 $\pm$ 3.6 1.3–16.7	1.9 $\pm$ 2.3 0.4–10.5	0.1 $\pm$ 0.1 0.03–0.6	1.8 $\pm$ 1.1 0.3–4.6	3.2 $\pm$ 1.9 0.4–8.8	2.2 $\pm$ 1.3 0.1–5.5
3 months	8.3 $\pm$ 5.3 2.7–19.8	0.2 $\pm$ 0.2 0.03–0.8	0.03 $\pm$ 0.04 nd–0.1	1.1 $\pm$ 0.6 0.4–3.1	0.1 $\pm$ 0.1 nd–0.5	4.7 $\pm$ 2.4 2.2–10.2	2.4 $\pm$ 1.9 0.6–6.8	0.2 $\pm$ 0.1 0.01–0.6	2.2 $\pm$ 1.2 0.5–4.6	3.1 $\pm$ 1.8 1.1–7.3	2.6 $\pm$ 1.2 0.7–4.7
RYGB*		a,f			a,f	a,e,f					
Basal	4.3 $\pm$ 2.9 1.5–14.0	0.5 $\pm$ 0.2 0.09–0.9	0.02 $\pm$ 0.04 nd–0.1	1.1 $\pm$ 0.6 0.3–2.6	0.1 $\pm$ 0.1 0.03–0.5	5.2 $\pm$ 2.5 1.3–11.9	1.2 $\pm$ 2.3 0.2–10.9	0.1 $\pm$ 0.4 0.03–1.9	0.9 $\pm$ 0.7 0.3–3.5	1.5 $\pm$ 1.2 0.8–6.7	1.8 $\pm$ 0.7 0.4–3.6
Diet	5.2 $\pm$ 4.1 0.8–17.9	0.3 $\pm$ 0.2 0.02–0.9	nd $\pm$ 0.03 nd–0.1	1.3 $\pm$ 0.7 0.2–2.9	0.1 $\pm$ 0.1 nd–0.7	4.8 $\pm$ 3.0 0.6–12.2	1.7 $\pm$ 1.3 0.1–5.7	0.1 $\pm$ 0.1 0.03–0.4	1.3 $\pm$ 0.9 0.1–4.0	2.3 $\pm$ 1.4 0.2–6.1	1.8 $\pm$ 1.0 0.4–4.8
1 month	4.1 $\pm$ 3.8 0.9–15.6	0.2 $\pm$ 0.1 0.02–0.6	0.01 $\pm$ 0.2 nd–1.2	0.8 $\pm$ 0.6 0.2–3.1	0.1 $\pm$ 0.2 0.01–1.1	3.7 $\pm$ 2.5 nd–9.2	1.2 $\pm$ 0.9 0.2–3.2	0.1 $\pm$ 0.1 0.03–0.8	0.9 $\pm$ 1.4 0.1–7.4	1.8 $\pm$ 2.3 0.3–12.2	1.4 $\pm$ 1.0 0.5–4.1
3 months	3.0 $\pm$ 4.1 1.0–18.6	0.5 $\pm$ 0.4 0.04–2.1	nd $\pm$ 0.02 nd–0.07	0.9 $\pm$ 0.7 0.1–3.5	0.1 $\pm$ 0.08 0.01–0.2	2.2 $\pm$ 2.1 0.5–8.6	0.7 $\pm$ 1.1 0.2–5.5	0.1 $\pm$ 0.09 0.01–0.4	0.6 $\pm$ 1.0 0.1–3.9	1.3 $\pm$ 1.7 0.4–7.3	1.2 $\pm$ 1.0 0.5–4.6
Ala: alanine, Arg: arginine, Asn: asparagine, Asp: aspartic acid, Gln: glutamine, Glu: glutamate, Gly: glycine, His: histidine, Ile: isoleucine, Leu: leucine. *a: statistically significant difference determined by Wilcoxon test between baseline and diet; b: baseline and 1 month; c: baseline and 3 months; d: diet and 1 month; e: diet and 3 months; and f: 1 month and 3 months. nd: not detected. Only values of $p \leq 0.05$ were considered.											

Table 3
Cont.

Median ± SD	Met	Phe	Pro	Ser	Thr	Trp	Tyr	Val	Orn	BCAAs	TOTAL amino acids
Range (mM)											
SG*		c	c,e					c		c	
Basal	0.2 ± 0.3	0.7 ± 0.5	0.09 ± 0.5	1.0 ± 1.0	1.4 ± 1.9	0.1 ± 0.1	0.9 ± 0.7	1.2 ± 1.0	0.2 ± 0.3	3.5 ± 3.4	
	0.02– 1.0	0.05– 2.0	0–1.9	nd– 3.7	0.1– 8.5	0.05– 0.6	0.1– 2.7	0.1– 4.0	0.05– 1.5	0.4– 13.1	
Diet	0.6 ± 0.4	1.4 ± 0.6	0.1 ± 0.2	1.5 ± 1.1	1.6 ± 2.8	0.1 ± 0.1	1.8 ± 0.8	1.9 ± 1.2	0.3 ± 0.3	6.5 ± 4.3	
	0.01– 1.5	0.3– 2.4	0–0.6	nd– 3.9	0.3– 11.4	0.05– 0.4	0.4– 3.5	0.3– 4.8	0.1– 1.2	1.2– 16.7	
1 month	0.3 ± 0.4	1.1 ± 0.6	0.03 ± 0.2	1.8 ± 0.8	1.5 ± 3.0	0.2 ± 0.1	1.4 ± 0.8	2.0 ± 1.4	0.3 ± 0.2	6.7 ± 4.5	
	0.03– 1.8	0.1– 2.9	0.002– 0.9	0.3– 3.5	0.3– 10.5	0.03– 0.7	0.2– 3.6	0.3– 6.3	0.08– 0.9	1.1– 19.7	
3 months	0.4 ± 0.3	1.3 ± 0.5	0.04 ± 0.1	1.6 ± 0.8	1.7 ± 0.9	0.2 ± 0.08	1.2 ± 0.6	2.2 ± 1.3	0.4 ± 0.2	7.7 ± 4.3	
	0.02– 1.3	0.5– 2.6	0–0.5	0.7– 3.5	0.5– 3.7	0.08– 0.4	0.6– 3.0	0.6– 4.7	0.2– 1.0	2.5– 15.9	
RYGB*				a		a					
Basal	0.2 ± 0.2	0.6 ± 0.5	0.05 ± 0.1	0.9 ± 0.4	1.1 ± 0.6	0.1 ± 0.2	0.7 ± 0.2	0.9 ± 0.9	0.2 ± 0.6	3.5 ± 2.9	
	nd– 1.2	0.3– 3.1	nd–0.8	0.3– 2.6	0.4– 3.0	0.05– 1.0	0.4– 1.7	0.4– 4.5	0.08– 3.3	1.6– 14.7	
Diet	0.3 ± 0.2	0.9 ± 0.4	0.04– 0.1	1.3 ± 0.9	1.2 ± 0.8	0.1 ± 0.1	0.8 ± 0.5	1.5 ± 1.3	0.2 ± 0.2	5.5 ± 3.8	
	0.03– 1.0	0.1– 1.9	nd–0.6	0.1– 3.1	0.1– 2.6	0.01– 0.5	0.1– 1.9	0.1– 6.0	0.04– 0.7	0.4– 14.7	
1 month	0.2 ± 0.7	0.6 ± 0.8	0.1 ± 0.3	1.0 ± 0.7	0.8 ± 2.0	0.1 ± 0.3	0.6 ± 0.4	1.0 ± 2.1	0.3 ± 1.5	3.8 ± 5.9	
	0.02– 3.7	0.1– 4.3	nd–1.1	nd– 3.0	0.2– 10.5	0.02– 1.5	0.2– 2.0	0.2– 10.8	0.09– 7.6	0.7– 30.5	
3 months	0.2 ± 0.3	0.7 ± 1.4	0.07 ± 0.4	0.8 ± 0.7	0.7 ± 0.7	0.1 ± 0.1	0.6 ± 0.5	0.7 ± 1.3	0.2 ± 0.3	2.6 ± 4.0	
	1.0– 18.6	0.2– 6.0	nd–1.8	0.2– 3.0	0.2– 2.9	0.02– 0.5	0.2– 2.5	0.2– 5.5	0.08– 1.3	0.8– 16.5	

Lys: lysine, Met: methionine, Phe: phenylalanine, Pro: proline, Ser: serine, Thr: threonine, Trp: Tryptophan, Tyr: tyrosine, Val: valine, Orn: ornithine, BCAAs: Branched-Chain Amino Acids (leucine, isoleucine and valine). *a: statistically significant difference determined by Wilcoxon test between baseline and diet; b: baseline and 1 month; c: baseline and 3 months; d: diet and 1 month; e: diet and 3 months; and f: 1 month and 3 months. nd: not detected. Only values of $p \leq 0.05$ were considered.

Table 4

Median, standard deviation and range of the values of biogenic amines, phenylethylamine (PEA) and γ -aminobutyric acid (GABA), and ammonium (mM).

Median ± SD	Agmatine	Histamine	Tiramine	Putrescine	Tryptamine	Cadaverin	PEA	GABA	Ammonium
Range (mM)									
SG*			e		f			e,f	e,f
Basal	0.03 ± 0.09 nd–0.4	0.07 ± 0.09 nd–0.3	0.01 ± 0.02 nd–0.1	0.3 ± 0.5 0.03–1.7	nd ± 0.03 nd–0.1	0.5 ± 1.6 0.01–6.2	nd ± 0.01 nd–0.05	0.2 ± 1.9 0.02–6.6	32.7 ± 11.3 7.5–40.3
Diet	0.02 ± 0.1 nd–0.6	0.05 ± 0.1 nd–0.5	0.01 ± 0.04 nd–0.1	0.1 ± 0.3 0.02–1.3	nd ± 0.01 nd–0.04	0.6 ± 1.4 nd–4.0	nd ± nd nd–0.002	0.3 ± 1.2 0.03–3.5	34.0 ± 16.5 10.1–63.5
1 month	0.01 ± 0.03 nd–0.1	0.02 ± 0.3 nd–1.3	nd ± 0.1 nd–0.5	0.06 ± 0.9 nd–3.9	nd ± nd nd–0.03	0.5 ± 2.0 nd–7.1	nd ± nd nd–0.004	0.5 ± 0.7 0.03–2.3	20.7 ± 15.4 5.9–71.1
3 months	0.02 ± 0.04 nd–0.1	0.02 ± 0.1 nd–0.5	0.01 ± 0.02 nd–0.1	0.01 ± 0.4 0.01–1.6	nd ± 0.02 nd–0.1	0.4 ± 1.8 nd–6.3	0 ± 0 0–0.01	0.4 ± 1.4 0.04–5.0	24.6 ± 13.9 6.6–53.3
RYGB*					a			a	
Basal	0.02 ± 0.07 nd–0.3	0.07 ± 0.2 nd–1.1	0.03 ± 0.3 nd–1.5	0.2 ± 0.7 0.02–2.0	nd ± nd nd–0.004	3.0 ± 2.3 nd–7.7	nd ± nd nd–0.02	0.2 ± 1.5 0.02–7.2	27.9 ± 13.6 6.9–60.4
Diet	0.02 ± 0.02 nd–0.07	0.08 ± 0.1 nd–0.5	0.02 ± 0.1 nd–0.5	0.1 ± 0.9 nd–4.7	nd ± nd nd–0.01	1.4 ± 2.0 0.1–6.7	nd ± nd nd–0.01	0.1 ± 1.7 nd–6.8	28.6 ± 13.0 6.5–47.9
1 month	0.02 ± 0.04 nd–0.1	0.04 ± 0.1 nd–0.5	0.05 ± 0.4 nd–2.1	0.4 ± 0.8 nd–3.1	nd ± nd nd–0.03	2.9 ± 3.7 0.01–12.8	nd ± 0.04 nd–0.2	0.2 ± 0.2 0.03–0.7	30.7 ± 12.5 9.6–57.1
3 months	0.01 ± 0.05 nd–0.2	0.1 ± 0.1 nd–0.8	0.1 ± 0.5 nd–0.2	0.6 ± 0.7 nd–2.3	nd ± nd nd–0.01	3.1 ± 3.0 nd–11.4	nd ± nd nd–0.02	0.7 ± 1.8 0.0–8.0	40.5 ± 12.5 11.5–58.7
*a: statistically significant difference determined by Wilcoxon test between baseline and diet; b: baseline and 1 month; c: baseline and 3 months; d: diet and 1 month; e: diet and 3 months; and f: 1 month and 3 months. Only values of $p \leq 0.05$ were considered									

There was a statistically significant increase in alanine, arginine, glutamine and glutamic acid levels after the crash diet in the SG group, as well in the levels of the biogenic amine tryptamine and the neurotransmitter GABA. Regarding the impact of the surgical process, there was a statistically significant increase in alanine and branched amino acid levels and a decrease in

phenylalanine, proline and glutamine levels in the SG group when comparing the baseline samples and the samples taken 3 months after surgery.

By contrast, the crash diet produced an increase in the RYGB group in the concentrations of tyrosine and serine and a decrease in arginine, glutamine and glutamic acid levels. In addition, surgery was associated with a statistically significant decrease in glutamic acid levels and a statistically significant increase in levels of the biogenic amine tryptamine and GABA.

Ammonium. The progression of the fecal ammonium concentrations showed opposing trends over the sampling time for the two surgical groups, with a significant decrease for the SG group and a significant increase for the RYGB group (Fig. 7A). Given that the normal range for fecal ammonium has not yet been defined, we arbitrarily established three categories based on the following tertiles: low (0–19 mM), medium (20–39 mM) and high concentration (> 40 mM) (Fig. 7B). We compared the bacterial abundance in the baseline samples to the fecal ammonium profile, demonstrating the predominance of *Paraprevotella* to the detriment of other members of the Bacteroidales order (*Parabacteroides*, *Barnesiella* and *Odoribacter*) and *Lentisphaerae* in those patients with higher ammonium levels (Supplementary Fig. 1).

DM2 resolution after bariatric surgery. During enrollment, 12 participants (7 men; 10 of them undergoing RYGB) had insulin-dependent DM2. After the surgery, 9 (75%, 8 of them undergoing RYGB) of the 12 experienced complete remission without drug treatment, while the remaining 3 patients continued to require oral metformin and insulin.

The differential bacterial abundance was analyzed in the 8 patients who underwent RYGB with complete resolution of the diabetes (baseline vs. 3-month samples), showing that the phyla Verrucomicrobiota (*Akkermansia*) and Fusobacteriota (*Fusobacterium*) increased significantly after surgery, while the relative abundance of the phyla Bacillota (*Faecalibacterium*, *Erysipelotrichia*, *Gemmiger* and *Lactobacillus*) and Actinobacteriota (*Bifidobacterium*) decreased. When comparing these 8 patients with the remaining 18 RYGB patients at baseline, a significantly lower abundance of the Clostridia class was associated with DM2, although surgery resulted in a clear decrease of this taxon in both groups (Supplementary Fig. 2).

Lasso model. Analyses with the Lasso model failed to identify any metabolite able to predict %EWL after 100 repetitions; however, they were able to identify 8 minority bacterial genera (Table 5). The root mean square error values were always low (approximately 0.05). The model pointed to *Anaerofustis* and *Acetanaerobacterium* as the most discriminant genera; however, both are underrepresented in the global gut microbiota. The most promising finding was related to *Bifidobacterium*, a genus considerably more abundant in the microbiota than the others, despite its lower prediction capability. This genus decreased in both groups of surgery as directly correlated with weight loss.

Table 5
Median values of the 100 Lasso regression values obtained for the significant bacterial genera and their abundance at baseline in each group of patients.

Genera	Lasso	% Basal abundance	
	regression	SG	RYGB
<i>Anaerofustis</i>	4.910	0.001	0.001
<i>Acetanaerobacterium</i>	3.234	0.002	0.01
<i>Coproacter</i>	0.182	0.02	0.03
<i>Anaerophylum</i>	0.088	0.002	0.002
<i>Slackia</i>	0.024	0.07	0.05
<i>Gemmiger</i>	0.003	1.7	2.5
<i>Romboutsia</i>	0.002	0.3	0.1
<i>Bifidobacterium</i>	0.001	3.3	2.4

Discussion

Obesity is a complex and multifactorial disease that goes far beyond simple excess weight, requiring an individualized study to understand its nature, causes, and treatment, particularly in severe obesity. The gut microbiota's contribution to obesity is gradually being recognized, although the lack of established normal ranges for the microbiota's composition and functionality precludes reaching solid conclusions in certain cases. Most of the available studies have been designed to identify significant differences in bacterial gut microbiota composition between lean individuals and those with obesity or to decipher the impact of each surgical approach on bacterial composition, employing a wide range of methodologies and analytical strategies.

The present study aimed to monitor fecal microbiota composition and functionality fluctuations during the bariatric process in severe obesity and included robust statistical analysis with machine learning approaches for prediction. Our main findings are the fecal determination of SCFAs, amino acids and biogenic amines in severe obesity during the bariatric process, highlighting the comparatively high levels of ammonia detected in the patients who underwent RYGB compared with those who underwent SG and, relevantly, the suggestion that a particular baseline gut microbiota might be necessary to reach the highest BMI values, as the dynamics of Bacillota suggests.

Previous studies have emphasized the low microbial gene richness, low alpha diversity values and higher abundance of Bacillota in severe obesity [6, 15–18]. The utility of the Bacillota/Bacteroidota ratio (formerly the Firmicutes/Bacteroidetes ratio) in obesity is controversial, and recent data point to its lack of validity as a microbiota marker, given that an expansion of Bacillota leads to the proportional reduction of the remaining phyla [19, 20]. The most noticeable feature we found was that, although the Bray-Curtis analysis found no significant difference in phyla abundance in the baseline samples, there was a direct correlation between Bacillota abundance and baseline BMI occurring in SG, whereas in RYGB, the extremely high proportions of Bacillota in the baseline samples experienced a proportional decrease in accordance with the %EWL. Surprisingly, the abundance of Bacillota increased in the postoperative SG samples with no correlation to weight loss.

The dissimilar impact of the two bariatric surgical approaches on gut microbial composition has been exhaustively reviewed, with consistent results [9, 21–27]. Alpha diversity dynamics appear not to be reproducible across studies; however, beta diversity analyses consistently find deeper changes for RYGB, including an expansion of the Pseudomonadota (formerly Proteobacteria) phyla [28–33]. Biliary acid circuit redistribution, whose antibacterial activity limits the expansion of gamma-Proteobacteria in the small intestine, is likely the major cause of the increase in members from this phylum in samples from patients who undergo RYGB. Despite the differential trends in bacterial composition, weight loss was uniformly reached in both groups.

Acetic, propionic and butyric acids were the most abundant SCFAs in our fecal samples, as other authors have reported [34]. SCFAs are derived mainly from carbohydrate bacterial fermentation in the small intestine; however, a small amount is also produced by the colonic fermentation of the branched-chain amino acids valine, leucine, and isoleucine, leading to the production of the BCFAs isobutyric, isovaleric and 2-methyl butyric acid [9, 35]. A number of authors have recently reported an increase in fecal BCFAs after bariatric surgery [33, 36]. In our study, only the increase in isobutyric acid levels was statistically significant in the RYGB group after surgery with the concomitant decrease of leucine and valine levels. In general, we observed modest variations in SCFA concentrations, highlighting the decrease in acetic acid levels in both groups; however, the bariatric process implies drastic changes in food intake, particularly after surgery. SCFAs can represent up to 10% of the total energy intake, and it has been postulated that there is a significant overproduction of these microbial metabolites in obesity [18, 37], particularly propionic acid [38]. In general, the major limitation to reaching solid conclusions is the lack of defined cut-offs for SCFA, as well as measurement standardization for food intake and the subsequent time required for SCFA production.

We performed an exhaustive analysis to decipher the relationship between bacterial taxonomic levels and SCFA concentrations but did not obtain relevant results. In that sense, certain metabolic routes of SCFA production could be shared by unrelated bacterial taxa, which, together with the possible absorption by the intestinal epithelium of formed compounds, makes it difficult to establish clear correlations between fecal SCFA levels and bacterial abundance [34, 35, 39]. Several authors have inferred this by shotgun metagenomics, but the existence of a metabolic pathway does not always lead to their expression, and the final on/off control is largely modulated by bacterial population densities.

Most dietary proteins are absorbed during digestion, and only a small proportion of undigested proteins (approximately 5%), together with the rest of the luminal apoptotic host cells or cells from dead microorganisms, reach the colon. There is currently a lack of data on the normal concentrations of amino acids in feces, even more so in the case of bariatric surgery, which again limits our analysis; significant increases in glutamate levels have been reported only in rats subjected to RYGB [40]. We found noticeable variations in the levels of certain amino acids associated with the preoperative crash diet and the type of bariatric surgery. Notably, levels of total amino acids were higher in the samples from the SG group than from the RYGB group after the crash diet and surgery, whereas conversely, certain biogenic amines had higher levels in the RYGB samples. Bariatric surgery, particularly RYGB by malabsorption, could lead to increased protein concentrations in the lumen of the colon. Amino acids from these proteins could be metabolized by anaerobic bacteria, increasing the potential production of toxic metabolites. The higher levels of biogenic amines found in RYGB support this conclusion, and although the low fecal concentrations of these compounds are not suggestive a priori of endogenous intoxication, higher microbial production, particularly in diseases involving intestinal dysmotility, cannot be ruled out [41].

One of the most striking results of our study was the opposing tendency in fecal ammonium concentrations of the two approaches. RYGB provoked an abrupt increase (almost doubling the baseline values), while SG caused a marked decrease. High variability in intestinal ammonium levels has been previously reported in healthy volunteers [42], with Martínez-Cuesta *et al.* demonstrating higher concentrations in severe obesity [18]. Once again, normal cut-offs need to be established, as well as determining the correlation between serum and feces and their production after food intake. Ammonium is a normal metabolite of amino acid digestion and is one of the most significant toxicants for humans. Hyperammonemia due to defective ammonium detoxification of urea occurs in liver injury, causing severe neurological disorders [43], which are successfully treated with antibiotics to reduce the bacterial load in the small intestine. The contribution of gut microbiota to ammonium production is likely underestimated in healthy individuals because it is rapidly converted to urea and excreted. The capacity of intestinal bacteria to overproduce ammonium should, however, not be neglected in various health conditions, as well as in digestive tract surgery. Other metabolites such as cresol, indole, trimethylamine N-oxide and even bile salts can contribute to intestinal and systemic toxicity in patients who undergo bariatric surgery and should be determined in further studies. In fact, the bacterial overgrowth has been previously described after RYGB [44, 45], being the SG approach recommended for hepatopathy as cirrhosis, but in Non-Alcoholic SteatoHepatitis (NASH), RYGB and SG have similar results [46–48].

We aimed to assess the impact of the preoperative crash diet, given that postoperative dietary changes cannot be properly evaluated without the intrinsic effect of the surgery. Bacterial rearrangements were detected in both groups. Al Assal *et al.* reported a significant decrease in *Ruminococcaceae/F.prausnitzii* after a crash diet [15]. In our series, however, *F. prausnitzii* decreased only in the SG group, in which, contrary to expected, Bacillota considerably increased, and these variations were not followed by weight loss. Individuals with severe obesity are often undernourished, particularly for essential micronutrients, and recent data have increased the suspicion that the gut microbiota can contribute to this state [49], which is a target of interest for future studies to improve postoperative diets.

The DM2 resolved with no drug treatment after the bariatric surgery in 75% of the patients who had this metabolic disorder, which was accompanied by defined changes in intestinal microbial patterns, as previously reported [7, 15, 50, 51]. The contribution of the intestinal microbiota in resolving DM2 is still controversial, whereas biliary salts appear to play a relevant role [32]. We were unable to identify a correlation between bacterial genera abundance and DM2 resolution in 75% of our patients (9/12). The baseline status was related to a lower abundance of the Clostridia class, although bariatric surgery promoted a significant decrease in this taxon in all participants.

A robust bioinformatics analysis was incorporated to detect correlations between microbiota composition and functional variables, as were machine learning models to predict weight loss from the baseline samples. Despite our efforts, the integrative analysis provided no statistically significant results, and although the Lasso model identified 8 bacterial genera for predicting weight loss, their low representation in the entire ecosystem hinders their use, except for *Bifidobacterium*, which decreased in both surgical groups in direct correlation with weight loss.

At the design stage, all participants were equally considered in their baseline sampling when the surgical technique had not yet been assigned. However, given that each group displayed significant particularities, we decided to analyze the results by considering RYGB and SG separately. Our results strongly suggest different types/degrees of obesity, particularly differentiated by BMI and bacterial metabolism, as has been previously suggested by other authors [52]. Certainly, a BMI ≥ 50 implies differential bacterial characteristics compared with lower BMIs, although we cannot ignore the fact that almost all the participants with the highest BMI values underwent RYGB.

Further studies should address the characterization of fecal and serum metabolites from gut microbiota, focusing on their normal ranges to decipher their actual role in obesity and determine the impact of bariatric surgery on the host's health. Our sampling ended 3 months after surgery, so the long-term effects could not be determined. Although most authors follow patients for a year, the most drastic changes related to gut microbiota appear to occur during the first three months [30].

In summary, rather than a gut microbiota characteristic of severe obesity, we have detected individual signatures of composition and functionality, with opposing trends for almost all determined variables in the two surgical approaches. RYGB provokes a deeper impact on gut microbiota composition and a more putrefactive metabolism, probably related to biliary acid redistribution. SCFA production is similar in the two groups, with minimal effects from the bariatric process. Lastly, significantly higher ammonium production occurs in RYGB, although its clinical significance is uncertain. Standardized determinations of microbiota metabolites in feces and serum, as well as normality cut-offs, are needed to reach solid conclusions in future studies.

Abbreviations

GABA: gamma-aminobutyric acid; OTU: operating taxonomic unit; SCFAs: short-chain fatty acids; NASH: non-alcoholic steatohepatitis.

Declarations

Acknowledgments

We are indebted to all the patients who have participated in the study, and we would like to thank all the people who have contributed to this work, especially Marta Cobo and Yolanda Gutiérrez.

Authors' contributions

Conceptualization—NS, JIBC, CGRG, RdC; Methodology—NS, MPA, SSC, AMHB, BR; Validation—NS, BR; Formal analysis—MF, JZ, JG, MF; Investigation—NS, MPA, MG, CGRG, RdC; Data curation—MG, JIBC, BVP, JG; Writing original draft—NS, MPA, CGRG, RdC; Writing-review and editing—All authors. All authors read and approved the final manuscript.

Funding

This work was supported by the Spanish Ministry of Science and Innovation, Instituto de Salud Carlos III (ICI21/00012), and co-funded by the NextGenerationEU and PRTR. NS was financed by a postdoctoral Juan de la Cierva contract (Ref. IJCI-2014-19885) from Ministry of Economy and Competitiveness, Spain and is the recipient of a postdoctoral contract awarded by the Fundación para la Investigación y la Innovación Biosanitaria del Principado de Asturias (FINBA). MPA was supported by a Rio Hortega contract (CM19/00069) from the Instituto de Salud Carlos III and co-funded by the European Social Fund (ESF, "Investing in your future"). MF acknowledge the financial support by the Instituto de Salud Carlos III, and Fundación Agencia Española contra el Cáncer (projects ERA NET TRANSCAN-2 AC17/00022) and co-financed by the European Regional Development Fund (ERDF).

Availability of data and materials

<https://www.ncbi.nlm.nih.gov/bioproject/?term=PRJNA639545>

Ethics approval and consent to participate

Our local ethical committee approved the project, and all participants signed the informed consent.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

References

1. Alsumali A, Al-Hawag A, Samnaliev M, Eguale T. Systematic assessment of decision analytic models for the cost-effectiveness of bariatric surgery for morbid obesity. *Surg Obes Relat Dis*. 2018;14:1041–1059. doi: 10.1016/j.soard.2018.03.005.
2. Castellana M, Procino F, Biacchi E, Zupo R, Lampignano L, Castellana F, Sardone R, Palermo A, Cesareo R, Trimboli P, Giannelli G. Roux-en-Y gastric bypass *versus* sleeve gastrectomy for remission of type 2 diabetes. *J Clin Endocrinol Metab*. 2020:dga737. doi: 10.1210/clinem/dga737.
3. Arakawa R, Febres G, Cheng B, Krikhely A, Bessler M, Korner J. Prospective study of gut hormone and metabolic changes after laparoscopic sleeve gastrectomy and Roux-en-Y gastric bypass. *PLoS One*. 2020;15(7):e0236133. doi: 10.1371/journal.pone.0236133.
4. Gutiérrez-Repiso C, Moreno-Indias I, de Hollanda A, Martín-Núñez GM, Vidal J, Tinahones FJ. Gut microbiota specific signatures are related to the successful rate of bariatric surgery. *Am J Transl Res*. 2019;11:942–952. eCollection 2019.
5. Luijten JCHBM, Vugts G, Nieuwenhuijzen GAP, Luyer MDP. The importance of the microbiome in bariatric surgery: a systematic review. *Obes Surg*. 2019;29:2338–2349. doi: 10.1007/s11695-019-03863-y.
6. Sánchez-Alcoholado L, Gutiérrez-Repiso C, Gómez-Pérez AM, García-Fuentes E, Tinahones FJ, Moreno-Indias I. Gut microbiota adaptation after weight loss by Roux-en-Y gastric bypass or sleeve gastrectomy bariatric surgeries. *Surg Obes Relat Dis*. 2019;15:1888–1895. doi: 10.1016/j.soard.2019.08.551.
7. Insenser M, Vilarrasa N, Vendrell J, Escobar-Morreale HF. Remission of diabetes following bariatric surgery: plasma proteomic profiles. *J Clin Med*. 2021;10:3879. <https://doi.org/10.3390/jcm10173879>.
8. Sanchez-Carrillo S, Ciordia S, Rojo D, Zubeldia-Varela E, Méndez-García C, Martínez-Martínez M, Barbas C, Ruiz-Ruiz S, Moya A, Garriga M, Salazar N, Botella-Carretero JI, Vega-Piñero B, de Los Reyes-Gavilán CG, del Campo R, Ferrer M. A body weight loss- and health-promoting gut microbiota is established after bariatric surgery in individuals with severe obesity. *J Pharm Biomed Anal*. 2021;193:113747. doi: 10.1016/j.jpba.2020.113747.
9. Gutiérrez-Repiso C, Moreno-Indias I, Tinahones FJ. Shifts in gut microbiota and their metabolites induced by bariatric surgery. Impact of factors shaping gut microbiota on bariatric surgery outcomes. *Rev Endocr Metab Disord*. 2021;22:1137–1156. <https://doi.org/10.1007/s11154-021-09676-8>.
10. Nogacka AM, Salazar N, Arbolea S, Ruas-Madiedo P, Mancabelli L, Suarez A, Martinez-Faedo C, Ventura M, Tochio T, Hirano K, Endo A, G de Los Reyes-Gavilán C, Gueimonde M. *In vitro* evaluation of different prebiotics on the modulation of gut microbiota composition and function in morbid obese and normal-weight subjects. *Int J Mol Sci*. 2020;21:906. doi: 10.3390/ijms21030906.
11. Redruello B, Ladero V, Cuesta I, Álvarez-Buylla JR, Martín MC, Fernández M, Álvarez MA. A fast, reliable, ultra-high performance liquid chromatography method for the simultaneous determination of amino acids, biogenic amines and ammonium ions in cheese, using diethyl ethoxymethylenemalonate as a derivatising agent. *Food Chem*. 2013;139:1029–1035. doi: 10.1016/j.foodchem.2013.01.071.
12. Taverniers I, De Loose M, Van Bockstaele E. Trends in quality in the analytical laboratory. II. Analytical method validation and quality assurance. *Trends Anal Chem*. 2004;23:535e552. doi: 10.1016/j.trac.2004.04.001.

13. Bolyen E, Rideout JR, Dillon MR, Bokulich NA, Abnet CC, Al-Ghalith GA, Alexander H, Alm EJ, Arumugam M, Asnicar F, Bai Y, Bisanz JE, Bittinger K, Brejnrod A, Brislawn CJ, Brown CT, Callahan BJ, Caraballo-Rodríguez AM, Chase J, Cope EK, Da Silva R, Diener C, Dorrestein PC, Douglas GM, Durall DM, Duvallet C, Edwardson CF, Ernst M, Estaki M, Fouquier J, Gauglitz JM, Gibbons SM, Gibson DL, Gonzalez A, Gorlick K, Guo J, Hillmann B, Holmes S, Holste H, Huttenhower C, Huttley GA, Janssen S, Jarmusch AK, Jiang L, Kaehler BD, Kang KB, Keefe CR, Keim P, Kelley ST, Knights D, Koester I, Kosciulek T, Kreps J, Langille MGI, Lee J, Ley R, Liu YX, Loftfield E, Lozupone C, Maher M, Marotz C, Martin BD, McDonald D, McIver LJ, Melnik AV, Metcalf JL, Morgan SC, Morton JT, Naimy AT, Navas-Molina JA, Nothias LF, Orchanian SB, Pearson T, Peoples SL, Petras D, Preuss ML, Priesse E, Rasmussen LB, Rivers A, Robeson MS 2nd, Rosenthal P, Segata N, Shaffer M, Shiffer A, Sinha R, Song SJ, Spear JR, Swafford AD, Thompson LR, Torres PJ, Trinh P, Tripathi A, Turnbaugh PJ, Ul-Hasan S, van der Hooft JJJ, Vargas F, Vázquez-Baeza Y, Vogtmann E, von Hippel M, Walters W, Wan Y, Wang M, Warren J, Weber KC, Williamson CHD, Willis AD, Xu ZZ, Zaneveld JR, Zhang Y, Zhu Q, Knight R, Caporaso JG. Reproducible. interactive. scalable and extensible microbiome data science using QIIME 2. *Nat Biotechnol.* 2019;37:852-857. doi: 10.1038/s41587-019-0209-9.
14. Segata N, Izard J, Waldron L, Gevers D, Miropolsky L, Garrett WS, Huttenhower C. Metagenomic biomarker discovery and explanation. *Genome Biol.* 2011;12(6):R60. doi: 10.1186/gb-2011-12-6-r60.
15. Al Assal K, Prifti E, Belda E, Sala P, Clément K, Dao MC, Doré J, Levenez F, Taddei CR, Fonseca DC, Rocha IM, Balmant BD, Thomas AM, Santo MA, Dias-Neto E, Setubal JC, Zucker JD, Belarmino G, Torrinhas RS, Waitzberg DL. Gut microbiota profile of obese diabetic women submitted to Roux-en-Y Gastric Bypass and its association with food intake and postoperative diabetes remission. *Nutrients.* 2020;12:278. doi: 10.3390/nu12020278.
16. Crovesy L, Masterson D, Rosado EL. Profile of the gut microbiota of adults with obesity: a systematic review. *Eur J Clin Nutr.* 2020;74:1251-1262. doi: 10.1038/s41430-020-0607-6.
17. Debédat J, Clément K, Aron-Wisniewsky J. Gut microbiota dysbiosis in human obesity: impact of bariatric surgery. *Curr Obes Rep.* 2019;8:229-242. doi: 10.1007/s13679-019-00351-3.
18. Martínez-Cuesta MC, Del Campo R, Garriga-García M, Peláez C, Requena T. Taxonomic characterization and short-chain fatty acids production of the obese microbiota. *Front Cell Infect Microbiol.* 2021;11:598093. doi: 10.3389/fcimb.2021.598093.
19. Magne F, Gotteland M, Gauthier L, Zazueta A, Pessoa S, Navarrete P, Balamurugan R. The firmicutes/bacteroidetes ratio: a relevant marker of gut dysbiosis in obese patients? *Nutrients.* 2020;12:1474. doi: 10.3390/nu12051474.
20. Palmisano S, Campisciano G, Silvestri M, Guerra M, Giuricin M, Casagrande B, Comar M, de Manzini N. Changes in gut microbiota composition after bariatric surgery: a new balance to decode. *J Gastrointest Surg.* 2020;24:1736-1746. doi: 10.1007/s11605-019-04321-x.
21. Cook J, Lehne C, Weiland A, Archid R, Ritze Y, Bauer K, Zipfel S, Penders J, Enck P, Mack I. Gut microbiota, probiotics and psychological states and behaviors after bariatric surgery-a systematic review of their interrelation. *Nutrients.* 2020;12:2396. doi: 10.3390/nu12082396.
22. Ejtahed HS, Angoorani P, Hasani-Ranjbar S, Siadat SD, Ghasemi N, Larijani B, Soroush AR. Adaptation of human gut microbiota to bariatric surgeries in morbidly obese patients: A systematic review. *Microb Pathog.* 2018 Mar;116:13-21. doi: 10.1016/j.micpath.2017.12.074.
23. Guo Y, Huang ZP, Liu CQ, Qi L, Sheng Y, Zou DJ. Modulation of the gut microbiome: a systematic review of the effect of bariatric surgery. *Eur J Endocrinol.* 2018;178:43–56. doi: 10.1530/EJE-17-0403
24. Koulas SG, Stefanou CK, Stefanou SK, Tepelenis K, Zikos N, Tepetes K, Kapsoritakis A. Gut microbiota in patients with morbid obesity before and after bariatric surgery: a ten-year review study (2009-2019). *Obes Surg.* 2021;31:317-326. doi: 10.1007/s11695-020-05074-2.
25. Morales-Marroquin E, Hanson B, Greathouse L, de la Cruz-Munoz N, Messiah SE. Comparison of methodological approaches to human gut microbiota changes in response to metabolic and bariatric surgery: A systematic review. *Obes Rev.* 2020;21:e13025. doi: 10.1111/obr.13025.
26. Paganelli FL, Luyer M, Hazelbag CM, Uh HW, Rogers MRC, Adriaans D, Berbers RM, Hendrickx APA, Viveen MC, Groot JA, Bonten MJM, Fluit AC, Willems RJL, Leavis HL. Roux-Y Gastric Bypass and Sleeve Gastrectomy directly change gut

- microbiota composition independent of surgery type. *Sci Rep.* 2019;9:10979. doi: 10.1038/s41598-019-47332-z.
27. Tabasi M, Ashrafi F, Khezerloo JK, Eshghjoo S, Behrouzi A, Javadinia SA, Poursadegh F, Eybpoosh S, Ahmadi S, Radmanesh A, Soroush A, Siadat SD. Changes in gut microbiota and hormones after bariatric surgery: a bench-to-bedside review. *Obes Surg.* 2019;29:1663-1674. doi: 10.1007/s11695-019-03779-7.
 28. Kong LC, Tap J, Aron-Wisniewsky J, Pelloux V, Basdevant A, Bouillot JL, Zucker JD, Doré J, Clément K. Gut microbiota after gastric bypass in human obesity. increased richness and associations of bacterial genera with adipose tissue genes. *Am J Clin Nutr.* 2013;98:16–24. doi: 10.3945/ajcn.113.058743.
 29. Graessler J, Qin Y, Zhong H, Zhang J, Licinio J, Wong ML. Metagenomic sequencing of the human gut microbiome before and after bariatric surgery in obese patients with type 2 diabetes. correlation with inflammatory and metabolic parameters. *Pharmacogenom J.* 2013;13:514–522. doi: 10.1038/tpj.2012.43.
 30. Palleja A, Kashani A, Allin KH, Nielsen T, Zhang C, Li Y, Brach T, Liang S, Feng Q, Jørgensen NB, Bojsen-Møller KN, Dirksen C, Burgdorf KS, Holst JJ, Madsbad S, Wang J, Pedersen O, Hansen T, Arumugam M. Roux-en-Y gastric bypass surgery of morbidly obese patients induces swift and persistent changes of the individual gut microbiota. *Genome Med.* 2016;8:67. doi: 10.1186/s13073-016-0312-1.
 31. Medina DA, Pedreros JP, Turiel D, Quezada N, Pimentel F, Escalona A, Garrido D. Distinct patterns in the gut microbiota after surgical or medical therapy in obese patients. *PeerJ.* 2017;5:e3443. doi: 10.7717/peerj.3443.
 32. Murphy R, Tsai P, Jüllig M, Liu A, Plank L, Booth M.. Differential changes in gut microbiota after gastric bypass and sleeve gastrectomy bariatric surgery vary according to diabetes remission. *Obes Surg.* 2017;27:917-925. doi: 10.1007/s11695-016-2399-2.
 33. Tremaroli V, Karlsson F, Werling M, Ståhlman M, Kovatcheva-Datchary P, Olbers T, Fändriks L, le Roux CW, Nielsen J, Bäckhed F. Roux-en-Y gastric bypass and vertical banded gastroplasty induce long-term changes on the human gut microbiome contributing to fat mass regulation. *Cell Metab.* 2015;22: 228–238. doi: 10.1016/j.cmet.2015.07.009.
 34. Ríos-Covián D, Ruas-Madiedo P, Margolles A, Gueimonde M, de Los Reyes-Gavilán CG, Salazar N. Intestinal short chain fatty acids and their link with diet and human health. *Front Microbiol.* 2016;7:185. doi: 10.3389/fmicb.2016.00185.
 35. Rios-Covian D, González S, Nogacka AM, Arbolea S, Salazar N, Gueimonde M, de Los Reyes-Gavilán CG. An overview on fecal branched short-chain fatty acids along human life and as related with body mass index: associated dietary and anthropometric factors. *Front Microbiol.* 2020;11:973. doi: 10.3389/fmicb.2020.00973.
 36. Farup PG, Maseng MG. Are faecal microbiota analyses on species-level suitable clinical biomarkers? A pilot study in subjects with morbid obesity. *Microorganisms* 2021;9:664. Doi: 10.3390/microorganisms9030664.
 37. Patil DP, Dhotre DP, Chavan SG, Sultan A, Jain DS, Lanjekar VB, Gangawani J, Shah PS, Todkar JS, Shah S, Ranade DR, Patole MS, Shouche YS. Molecular analysis of gut microbiota in obesity among Indian individuals. *J Biosci.* 2012;37:647-57. doi: 10.1007/s12038-012-9244-0.
 38. Schwiertz A, Taras D, Schäfer K, Beijer S, Bos NA, Donus C, Hardt PD. Microbiota and SCFA in lean and overweight healthy subjects. *Obesity (Silver Spring).* 2010;18:190-5. doi: 10.1038/oby.2009.167.
 39. Amaretti A, Gozzoli C, Simone M, Raimondi S, Righini L, Pérez-Brocal V, García-López R, Moya A, Rossi M. Profiling of protein degraders in cultures of human gut microbiota. *Front Microbiol.* 2019;10:2614. doi: 10.3389/fmicb.2019.02614.
 40. Haange SB, Jehmlich N, Krügel U, Hintschich C, Wehrmann D, Hankir M, Seyfried F, Froment J, Hübschmann T, Müller S, Wissenbach DK, Kang K, Buettner C, Panagiotou G, Noll M, Rolle-Kampczyk U, Fenske W, von Bergen M. Gastric bypass surgery in a rat model alters the community structure and functional composition of the intestinal microbiota independently of weight loss. *Microbiome* 2020;8:13. doi: 10.1186/s40168-020-0788-1.
 41. Li JV, Ashrafi F, Bueter M, Kinross J, Sands C, le Roux CW, Bloom SR, Darzi A, Athanasiou T, Marchesi JR, Nicholson JK, Holmes E. Metabolic surgery profoundly influences gut microbial–host metabolic cross-talk. *Gut.* 2011;60:1214-23. doi: 10.1136/gut.2010.234708.
 42. Delgado S, Ruas-Madiedo P, Suárez A, Mayo B. Interindividual differences in microbial counts and biochemical-associated variables in the feces of healthy Spanish adults. *Dig Dis Sci.* 2006;51:737-43. doi: 10.1007/s10620-006-3200-5.

43. Holeček M. Branched-chain amino acids and branched-chain keto acids in hyperammonemic states: metabolism and as supplements. *Metabolites*. 2020;10:324. doi: 10.3390/metabo10080324.
44. Dolan RD, Baker J, Harer K, Lee A, Hasler W, Saad R, Schulman AR. Small intestinal bacterial overgrowth: clinical presentation in patients with Roux-en-Y gastric bypass. *Obes Surg*. 2021;31:564-569. doi: 10.1007/s11695-020-05032-y.
45. Sollier C, Barsamian C, Bretault M, Poghosyan T, Rahmi G, Chevallier JM, Bouillot JL, Carette C, Czernichow S, Rives-Lange C. Diagnostic and therapeutic management of post-gastric bypass chronic diarrhea: a systematic review. *Obes Surg*. 2020;30:1102-1111. doi: 10.1007/s11695-019-04253-0.
46. Aminian A, Al-Kurd A, Wilson R, Bena J, Fayazzadeh H, Singh T, Albaugh VL, Shariff FU, Rodriguez NA, Jin J, Brethauer SA, Dasarathy S, Alkhouri N, Schauer PR, McCullough AJ, Nissen SE. Association of bariatric surgery with major adverse liver and cardiovascular outcomes in patients with biopsy-proven nonalcoholic steatohepatitis. *JAMA*. 2021;326:2031-2042. doi: 10.1001/jama.2021.19569.
47. Miller A, Jalil S, Hussan H, Kelly S, Hinton A, Hanje J, Michaels A, Conteh L, Black S, Mumtaz K. Apprising the choice of bariatric surgery in patients with compensated cirrhosis with morbid obesity: results of a national analysis. *Updates Surg*. 2021;73:1899-1908. doi: 10.1007/s13304-021-01142-z.
48. Ahmed S, Pouwels S, Parmar C, Kassir R, de Luca M, Graham Y, Mahawar K; Global bariatric research collaborative. outcomes of bariatric surgery in patients with liver cirrhosis: a systematic review. *Obes Surg*. 2021;31:2255-2267. doi: 10.1007/s11695-021-05289-x.
49. Ciobârcă D, Cătoi AF, Copăescu C, Miere D, Crișan G. Bariatric surgery in obesity: effects on gut microbiota and micronutrient status. *Nutrients*. 2020;12:235. doi: 10.3390/nu12010235.
50. Pucci A, Batterham RL. Mechanisms underlying the weight loss effects of RYGB and SG: similar yet different. *Obes Surg*. 2021;31:327-336. doi: 10.1007/s11695-020-04974-7.
51. Wang M, Li L, Chen Y, Lian G, Wang J, Zhang J, Shan K, Shang L, Tian F, Jing C. Role of gut microbiome and microbial metabolites in alleviating insulin resistance after bariatric surgery. *Obes Surg*. 2021;31:327-336. doi: 10.1007/s11695-020-04974-7.
52. Aron-Wisnewsky J, Prifti E, Belda E, Ichou F, Kayser BD, Dao MC, Verger EO, Hedjazi L, Bouillot JL, Chevallier JM, Pons N, Le Chatelier E, Levenez F, Ehrlich SD, Dore J, Zucker JD, Clément K. Major microbiota dysbiosis in severe obesity: fate after bariatric surgery. *Gut*. 2019;68:70-82. doi: 10.1136/gutjnl-2018-316103.

Supplementary Figures

Supplementary Figures 1 and 2 are not available with this version.

Figures

Figure 1

% Weight excess in the 40 patients along the 4 samples of each one. *: significant differences between SG and RGYB groups at basal and diet samples.

Figure 2

Bacterial diversity analysis in the two groups of patients throughout the 4 fecal samples. A: alpha diversity values (Shannon and Chao1 indices). B: PCoA based on beta diversity values (Bray Curtis index), calculated from the bacterial profiles of stool samples. Each point represents a sample, and the color refers to sampling time.

A

B

Figure 3

Bacterial distribution in feces. A: at phylum level. B: at genus level, showing only the most abundant (80% of the total abundance).

Figure 4

Bacterial taxa with differential abundance according to linear effect size discriminant analysis (LEfSe). The left side of each graph shows a cladogram, in which the yellow circles represent bacterial taxa that show no significant differences between the groups analyzed, while the green and red circles represent taxa whose abundance is significantly differential between the groups analyzed. The other side of each graph shows the significant taxa ordered according to the magnitude of the differences (LDA score). A: Comparison between baseline and 3 months after surgery in SG. B: Comparison between baseline and 3 months after surgery in RYGB (only taxa with LDA>4 are shown, to simplify the figure).

Figure 5

Correlation study between Firmicutes abundance and weight excess, distinguishing according to the type of surgery (SG: green; RYGB: red) and sex. Left: basal state, right: difference between 3 months after surgery and basal state.

Figure 6

Evolution of the proportion (left) and absolute concentration (right, expressed in mM) of the majority (acetate, propionate and butyrate) and minority (caproate, isobutyrate, isovalerate and valerate) SCFA measured at the 4 times (from left to right: baseline, diet, 1 month after surgery, 3 months after surgery) for both groups. The horizontal line represents the mean of each distribution.

A **B** **Figure 7**

Results of fecal ammonium determination. A: ammonium concentration (mM) at the 4 fecal samples, for each group of patients. B: Percentage distribution of patients according to their fecal ammonium levels at the basal sample and at the end (3 months) of follow-up, for each type of surgery.