Surgical Treatment on Infective Endocarditis: Impact of Diabetes on Mortality

Alexander Kogan (✉ alexanderkogan140@hotmail.com)

Sheba Medical Center

Anat Wieder

Sheba Medical Center

Jonathan Frogel

Sheba Medical Center

Yael Peled-Potashnik

Sheba Medical Center

Eilon and Leonid Sternik^{1,5}. Ram

Sheba Medical Center

Ehud Raanani

Sheba Medical Center

Leonid Sternik

Sheba Medical Center

Research Article

Keywords: Infective endocarditis, Cardiac surgery, Diabetes, Epidemiology, Mortality

Posted Date: February 25th, 2022

DOI: <https://doi.org/10.21203/rs.3.rs-1379104/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background

Type 2 diabetes mellitus (DM) is a frequent co-morbidity among patients suffering from infective endocarditis (IE). The aim of the study was to evaluate the impact of type 2 DM on the early-, intermediate- and long-term mortality of patients who underwent surgical treatment of endocarditis.

Methods

We performed an observational cohort study in the large tertiary center in Israel during 14 years. All data of patients who underwent surgical treatment of endocarditis, performed between 2006 and 2020 were extracted from the departmental database. Patients were divided into two groups: Group I (non-diabetic patients), and Group II (diabetic patients).

Results

The study population includes 420 patients. Group I (non-diabetic patients), comprise 326 patients, and Group II (diabetic patients), comprise 94 patients. Mean follow-up duration was 39.3 ± 28.1 months. Short-term, 30-day and in-hospital mortality, also intermediate-term mortality (1- and 3-year) was higher in the DM group compared with the non-DM group, but did not reach statistical significance: 11.7% vs. 7.7% ($p = 0.215$); 12.8% vs. 8.3% ($p = 0.285$); 20.2% vs. 13.2% ($p = 0.1$) and 23.4% vs. 15.6% ($p = 0.09$) respectively. Long-term, 5-year mortality was significantly higher in the DM group, compared to the non-DM group: 30.9% vs. 16.6% ($p = 0.003$). Furthermore, predictors for long-term mortality included diabetes (CI 1.056–2.785, $p = 0.029$), as demonstrated by regression analysis.

Conclusions

Diabetic patients have trend to increasing mortality at the short- and intermediate period post-surgery for IE, but this is not statistically significant. Survival of diabetic patients deteriorates after more than three years follow surgery. Diabetes is an independent predictor for long-term, 5-year mortality after surgical treatment of endocarditis, regardless of the patients age and comorbidities.

Introduction

The global burden of diabetes is rising because of increased obesity and population ageing. Infections in diabetes mellitus (DM) are relatively more common and serious. DM patients run the risk of acute metabolic decompensation during infections, and at higher risk of certain invasive infections [1]. DM is one of the most prevalent conditions in the elderly and is associated with considerable mortality, mainly from cardiovascular complications [2]. Incidence of infective endocarditis (IE), among patients with and without DM, have increased during the period 2001–2015, and significantly higher in the DM population. [3]. Also, another data suggested, that prevalence of DM among patients with native valve endocarditis increased from 22% in 2004 to 30% in 2014 [4]. In addition, recent studies have demonstrated

increased mortality in diabetic patients undergoing valve surgery for IE [4-6] compared to non-diabetic patients. The aim of the current study was to investigate the impact of diabetes mellitus on short and long-term mortality in a cohort of patients after surgery for IE.

Methods

The study protocol was approved by the Sheba Medical Center Institutional Ethics Committee (Protocol No 4257). The requirement for informed consent waived because of the retrospective nature of the study. We carried out a retrospective, observational study of prospectively collected data from consecutive patients who underwent valve surgery for IE at a large tertiary care university hospital in Israel over a 14-year period, from 01.09.2006 and 31.08.2020. Diagnosis of IE was made according to Duke criteria [7]. Patients were divided into two groups: Group I, non-diabetic patients (non-DM), and Group II, patients with mellitus type 2 (DM). DM type 2 was defined in accordance with the American Diabetes Association as: (a) hemoglobin A1C $\geq 6.5\%$; (b) fasting plasma glucose levels ≥ 126 mg/d; (c) classic symptoms of hyperglycemia or a hyperglycemic crisis, a random plasma glucose level ≥ 200 mg/dL (11.1 mmol/L) [7]; or (d) currently on pharmacologic treatment (oral antihyperglycemic drugs and/or insulin). Patients under 18 years of age and patients with type 1 diabetes mellitus were excluded. Empiric antibiotic treatment was started when needed, and specific antibiotic treatment was initiated after the results of blood cultures were available. If blood cultures were negative after 72 hours, specific serologic tests were done for culture negative endocarditis (*Q-fever*, *bartonella*, *Chlamydia*, *Brucella* *Legionella* and *Mycoplasma*).

Using de-identified patient data from our department's database, we evaluated the following variables: gender, age, systemic and pulmonary hypertension, smoking, dialysis-dependent renal failure, New York Heart Association functional class, chronic obstructive pulmonary disease (COPD), peripheral vascular disease, cerebrovascular accident (CVA)/transient ischemic attack (TIA), left ventricular function, previous cardiac surgery, affected valve, and microorganism isolated from the blood cultures or from valve tissue. Concomitant surgery which was defined as procedure, directly not connected to endocarditis, e.g. CABG or another valve replacement or repair.

Surgical intervention was performed according to the European Society of Cardiology guidelines [9]. After surgery, all patients were admitted to the intensive care unit (ICU) directly from the operating room. In the operating room and the ICU, patients from both groups received intravenous continuous infusion of regular insulin according to the Society of Thoracic Surgeons practice guideline [10]. After discharge from the ICU, patients were transferred either to a step-down unit or directly to the cardiac surgical department floor. Patients from Group I (non-DM) did not receive insulin or any other hypoglycemic medication.

Patients from Group II (DM) were restarted on their preoperative anti-glycemic regimen (per oral drugs and/or insulin) as soon as they resumed regular eating. Antibiotic's treatment was continued at least 6 weeks from the operation. We evaluated the following outcome variables: 30-day, in-hospital, 1-, 3- and 5-year mortality.

Database management and statistical analysis

At discharge, all patient data were reviewed, corrected and entered into a database. Descriptive statistics were used to summarize data, and numerical data were expressed as means (SD). Normality of the distribution of continuous data variables was analyzed using the Kolmogorov-Smirnov test. Since not all numerical data were distributed normally, Mann-Whitney U-tests were used to evaluate differences between groups. Differences between the frequencies of categorical variables were also estimated using Fisher's exact test. All outcome variables were compared between whole groups of patients.

A Cox proportional hazard model was constructed to assess the association between DM and mortality. To address the differences in baseline characteristic between the groups, subjects' propensity score is first estimated, and then the outcome is regressed on the estimated propensity score. The regression choice depends on the nature of the outcome, therefore we used logistic regression for the predicted probability, which specify the propensity score. Variables that were associated with mortality adjusted to age were included in the regression model. In addition, we included pre-specified clinically significant variables in the model. The variables included in the final model were: gender, age, DM, affected valve, previous cardiac surgery, presence of perioperative heart failure and perioperative acute kidney injury. Kaplan-Meier survival analysis was performed to compare mortality among the groups, with statistical differences tested by the log-rank test. Statistical significance was assumed when the null hypothesis could be rejected at $P < 0.05$. All P-values are the results of two-sided tests. Statistical analyses were conducted using R (version 3.4.1)

Results

The study population included 420 patients. Group I (non-diabetic patients) comprising 326 patients (77.6%), and Group II (diabetic patients) comprising 94 patients (22.4%). Mean follow-up duration was 39.3 ± 28.1 months. There were no differences regarding sex, average systolic left ventricular injection fraction, NIHA functional class and incidence of native or prosthetic valve endocarditis between groups I and II. Compared with the non-diabetic group, the diabetic group patients were older, had a higher prevalence of systemic and pulmonary hypertension, peripheral vascular disease and hyperlipidemia. Also, diabetic patients had higher EuroSCORE values and longer bypass and cross-clamp time (See Table 1). *Staphylococcus species* were more frequent pathogen in patients with DM, 52.1% vs. 41.4% ($p = 0.32$), however, not statistically significant.

Table 1
Patients demographic and perioperative data

	Group I (Not-DM patients)	Group II (DM patients)	P Value
N	326	94	
Age (years)	54.8 ± 16.1	66.4 ± 11.1	0.000
Male gender (n,%)	214 (65.6%)	61 (64.9%)	0.902
NIHA III-IV (n,%)	103 (31.6%)	37 (39.4%)	0.234
Previous cardiac surgery (n,%)	124 (38.0%)	36 (38.3%)	1.000
Native valve endocarditis (n,%)	212 (65.0%)	62 (66.0%)	0.903
EF (%)	54.6 ± 15.1	54.8 ± 15.9	0.914
Hypertension (n,%)	136 (42.0%)	76 (81.7%)	0.000
COPD (n,%)	15 (4.6%)	10 (10.8%)	0.044
Dialysis (n,%)	6 (2.1%)	5 (6.5%)	0.057
Hyperlipidemia (n,%)	100 (31.0%)	77 (81.9%)	0.000
PVD (n,%)	7 (2.2%)	12 (13.5%)	0.000
CVA/TIA (n,%)	46 (14.7%)	17 (18.5%)	0.414
Pulmonary hypertension (n,%)	40 (12.9%)	13(14.6%)	0.723
Smoking (n,%)	81 (25.3%)	35 (38.5%)	0.017
Standard EuroSCORE I	7.4 ± 3.9	10.0 ± 4.1	0.000
Logistic EuroSCORE %	16.3 ± 17.7	27.3 ± 24.4	0.000
Bypass time (min)	121.7 ± 56.1	148.8 ± 68.1	0.001
Cross-clamp (min)	86.1 ± 37.6	108.5 ± 48.3	0.000
Aortic valve involvement (n,%)	172 (52.8%)	55 (58.5%)	0.349
Mitral valve involvement (n,%)	175 (53.7%)	54 (57.4%)	0.558
Tricuspid valve involvement (n,%)	2 (0.6%)	0 (0%)	0.280
Concomitant procedure (n,%)	128 (39.3%)	54 (57.4%)	0.434

30-day, in-hospital, 1-year and 3-years mortality was higher in the DM group compared with the non-DM group, but did not reach statistical significance: 11.7% vs. 7.7%, ($p = 0.12$); 12.8% vs. 8.3% ($p = 0.285$); 20.2% vs. 13.2% ($p = 0.10$) and 23.4% vs. 15.6% ($p = 0.09$) respectively. Long-term, 5-years mortality was significantly higher in the DM group, compared to the non-DM group: 30.9% vs. 16.6% ($p = 0.003$) (See

Table 2 and Fig. 1). By Cox analysis we found that three factors impacted patients' 5-year mortality rates independently: diabetes mellitus (Group I), age and NIHA III-IV (See Table 3).

Table 2
Patient's mortality

	Group I (Non-DM) 326 patients	Group II (DM) 94 patients	<i>p</i> Value
30-days mortality	25 (7.7%)	11 (11.7%)	0.215
In-hospital mortality	27 (8.3%)	12 (12.8%)	0.285
1-year mortality	43 (13.2%)	19 (20.2%)	0.100
3-year mortality	51 (15.6%)	22 (23.4%)	0.090
5-year mortality	54 (16.6%)	29 (30.9%)	0.003

Table 3
Cox Regression for overall mortality

	B	Sig.	Exp(B)	95.0% CI for Exp(B)	
				Lower	Upper
Diabetes Mellitus	.539	.029	1.715	1.056	2.785
NIHA III-IV	.807	.001	2.241	1.408	3.568
Age (Predicted probability)	-2.213	.013	.109	.019	.632

Discussion

Gu *et al.* [11] reported, that in US population, survival of diabetic subjects was lower than that of non-diabetic subjects in all age, sex, and race groups. Also, type 2 DM is associated with reduced life expectancy at almost all ages in the Scotland [12]. Adults with type 2 DM, especially those with cardiovascular comorbidity, did not live as long than their non-diabetic people in South Korea [13]. Patients with diabetes mellitus (DM) are at increased risk of infection. Movahed *et al.* [14] studied Veterans Health Administration hospitals database and reported presence of IE in 1340 (0.5%) DM patients versus 1412 (0.3%) non-DM, control group patients (relative increase of 40%), and concluded, that patients with type II DM have significantly higher prevalence of IE. Also, Abe *et al.* [4] identified 76385 patients with native valve IE during 2004 -14 years, from the US National Inpatient Sample, of which 22284 (28%) had DM, and found, that proportion of DM among patients with IE increased from 22% in 2004 to 30% in 2014 ($p < 0.0001$). Miguel-Yanes *et al.* [3] identified 16,626 patients with IE in Spain, and found, that incidence of IE, among population with and without diabetes, have increased during the period 2001–2015 with significantly higher incidence rates in the diabetic population. However, it is not clear whether this is a true increase in the incidence of IE, or a better diagnosis.

Kourany *et al.* [15] compare outcomes between 150 diabetic and 905 non-diabetic patients with IE, from the International Collaboration on Endocarditis Merged Database. Diabetic patients underwent surgical intervention less frequently (32.0% vs. 44.9%, $p = 0.003$), and had higher overall in-hospital mortality (30.3% vs. 18.6%, $p = 0.001$). Olmos *et al.* [16] reported, that in-hospital mortality (43.5% vs. 30.0%; $p = 0.008$) were more common among patients with DM than in those without, but not described treatment modality, surgical or conservative. Farag *et al.* [17] studied 360 patients with IE, operated between 1993 and 2012, and reported, that non-survivors had higher rates of preoperative diabetes mellitus ($p = 0.005$), but not found diabetes as independent predictors of 30-day mortality. Yoshioka *et al.* [5] described 470 patients underwent valve surgery for left-sided active IE, 374 non-DM and 96 with DM. In-hospital mortality was 8% in patients without DM and 13% in patients with DM ($p = 0.187$). The overall survival rate at 1 and 5 years was 87% and 81% in patients without DM and 72% and 59% in patients with DM ($p < 0.001$). Wei *et al.* [18] studied of 866 people, who had been diagnosed with IE. They were divided into three groups: 469 patients in the normoglycaemia group, 246 patients in prediabetes group and 151 patients in diabetes group. Surgery was performed in 72.3%, 68.7% and 55.6% of cases respectively ($p = 0.001$). During median follow-up of 2.4 year, 126 (14.5%) died. Compared with those in the normoglycaemia group, higher long-term mortality was seen among people with prediabetes or diabetes (9.7% vs. 22.0% vs. 28.1%, $P < 0.001$). Lin *et al.* [19] described 412 patients with IE divided into 2 groups: group 1, patients with DM ($n = 72$) and group 2, patients without DM ($n = 340$) and concluded, that overall in-hospital mortality rate for both groups was 20.2% and was higher in group 1 than in group 2 (41.7% vs. 16.5%, $p < 0.01$).

Among the entire cohort of patients in our study, the percentage of the diabetic patients was 22% and those with DM were older. These data are comparable with other studies [5, 15, 16, 18, 19]. The possible explanation for this fact is that the long-term course of diabetes is associated with a sharp increase in the frequency of IE. Østergaard *et al.* [20] identified 300,000 patients with DM through the Danish Prescription Registry and investigated the association between the duration of DM and the incidence of IE. In patients with DM duration of 0–5 years, 5–10 years, 10–15 years, and more than 15 years, the incidence rates of IE were 0.24, 0.33, 0.58, and 0.96 cases of IE/1000 person years, respectively. Patients with DM duration 5–10 years, 10–15 years, and > 15 years were associated with a higher risk of IE compared with DM duration 0–5 years. Another study from the above mention Danish Registry [21], included 1767 patients with IE undergoing surgery, 735 patients < 60 years (24.1% female), 766 patients 60–75 years (25.8% female), and 266 patients ≥ 75 years (36.1% female). The proportions of patients undergoing surgery were 35.3, 26.9, and 9.1% for patients < 60 years, 60–75 years, and > 75 years, respectively. Mortality at 90 days was 7.5, 13.9, and 22.3% ($p < 0.001$) for three age groups. In adjusted analyses, patients 60–75 years and patients ≥ 75 years were associated with a higher mortality, HR = 1.84 (95% CI: 1.48–2.29) and HR = 2.47 (95% CI: 1.88–3.24) as compared with patients < 60 years. In addition, the diabetic patients had more co-morbidities, higher operation risk and longer operation time.

Our study carried out in a contemporary cohort of Israeli patients, who underwent surgery due to IE, demonstrates several important implications regarding the impact of type 2 DM on short-, mid- and long-term mortality. Our results are very similar to previously reported: 30-day mortality 11.7% vs. 7.7%, 1- and

5-years mortality was 86.8% vs. 79.8% and 84.4% vs. 69.1% for diabetic and non-diabetic patients respectively. Diabetic patients have trend to increasing mortality at the short- and intermediate period post-surgery for IE, but this is not statistically significant. Statistically significant difference was observed only for 5-years survival ($p = 0.003$).

There are a few limitations in our study. First, our study is retrospective in design. Second, our study was conducted in a single-center cardiac surgery department. Third, very high-risk patients are not candidate for surgery, there might be an option, that non-operated IE patients would change the survival results, Forth, we have limited microbiologic data.

Conclusions

Diabetic patients have trend to increasing mortality at the short- and intermediate period post-surgery for IE, but this is not statistically significant. Survival of diabetic patients deteriorates after more than three years follow surgery. Diabetes is an independent predictor for long-term, 5-year mortality after surgical treatment of endocarditis, regardless of the patient's age and comorbidities.

Abbreviations

DM: Diabetes mellitus;

CI: Confidence interval;

IE: Infective endocarditis;

COPD: Chronic obstructive pulmonary disease;

CVA: Cardiovascular accident;

TIA: Transient ischemic attack

CABG: Coronary artery bypass grating

ICU: Intensive Care Unit;

SD: standard deviation;

NIHA: New York Heart Association;

EuroSCORE: European System for Cardiac Operative Risk Evaluation

Declarations

Ethics approval and consent to participate

The study was conducted in accordance with the principles of the Helsinki declaration. The study protocol was approved by the Sheba Medical Center Institutional Ethics Committee (Protocol No 4257). The requirement for informed consent waived because of the retrospective nature of the study.

Consent for publication

Not applicable.

Availability of data and materials

Data, supported the findings of this study are available from corresponding author upon request.

Competing interests

The authors declare that they have no competing interests.

Funding

Not applicable.

Authors' contributions

A.K. and A.W. conceived and designed the study. J.F. contributed to the acquisition of data. Ei.R. performed the statistical analyses. A.K, A.W, Y.P-P, and L.S. interpreted of data. A.K, A.W. and J.F. drafted the manuscript. A.K, A.W, Eh.R. and L.S. contribute in writing of manuscript. All authors read and approved the final version of the manuscript.

Acknowledgements

The authors thank all the staff Cardiac Surgery ICU.

References

1. Gupta S, Koirala J, Khardori R, Khardori N. Infections in diabetes mellitus and hyperglycemia. *Infect Dis Clin North Am*. 2007 Sep;21(3):617-38, vii. doi: 10.1016/j.idc.2007.07.003. PMID: 17826615.
2. Knapp S. Diabetes and infection: is there a link?--A mini-review. *Gerontology*. 2013;59(2):99-104. doi: 10.1159/000345107. Epub 2012 Nov 24. PMID: 23182884.
3. de Miguel-Yanes JM, Jiménez-García R, Hernández-Barrera V, de Miguel-Díez J, Méndez-Bailón M, Muñoz-Rivas N, Pérez-Farinós N, López-de-Andrés A. Infective endocarditis according to type 2 diabetes mellitus status: an observational study in Spain, 2001-2015. *Cardiovasc Diabetol*. 2019 Nov 21;18(1):161. doi: 10.1186/s12933-019-0968-0. PMID: 31752887; PMCID: PMC6868776.
4. Abe T, Eyituyo HO, De Allie G, Olanipekun T, Effoe VS, Olaosebikan K, Mather P. Clinical outcomes in patients with native valve infective endocarditis and diabetes mellitus. *World J Cardiol*. 2021 Jan 26;13(1):11-20. doi: 10.4330/wjc.v13.i1.11. PMID: 33552399; PMCID: PMC7821010.

5. Yoshioka D, Toda K, Yokoyama JY, Matsuura R, Miyagawa S, Kainuma S, Sakaguchi T, Sakaki M, Nishi H, Shirakawa Y, Iwata K, Suhara H, Sakaniwa R, Fukuda H; Osaka Cardiovascular Research Group study group, Sawa Y. Diabetes mellitus adversely affects mortality and recurrence after valve surgery for infective endocarditis. *J Thorac Cardiovasc Surg*. 2018 Mar;155(3):1021-1029.e5. doi: 10.1016/j.jtcvs.2017.09.013. Epub 2017 Sep 15. PMID: 28992971.
6. Chirillo F, Bacchion F, Pedrocco A, Scotton P, De Leo A, Rocco F, Valfrè C, Olivari Z. Infective endocarditis in patients with diabetes mellitus. *J Heart Valve Dis*. 2010 May;19(3):312-20. PMID: 20583393.
7. Durack DT, Lukes AS, Bright DK. New criteria for diagnosis of infective endocarditis: utilization of specific echocardiographic findings. Duke Endocarditis Service. *Am J Med*. 1994 Mar;96(3):200-9. doi: 10.1016/0002-9343(94)90143-0. PMID: 8154507.
8. American Diabetes Association. 2. Classification and Diagnosis of Diabetes. *Diabetes Care*. 2016 Jan;39 Suppl 1:S13-22. doi: 10.2337/dc16-S005. Erratum in: *Diabetes Care*. 2016 Sep;39(9):1653. PMID: 26696675.
9. Habib G, Lancellotti P, Antunes MJ, Bongiorni MG, Casalta JP, Del Zotti F, Dulgheru R, El Khoury G, Erba PA, Iung B, Miro JM, Mulder BJ, Plonska-Gosciniak E, Price S, Roos-Hesselink J, Snygg-Martin U, Thuny F, Tornos Mas P, Vilacosta I, Zamorano JL; ESC Scientific Document Group. 2015 ESC Guidelines for the management of infective endocarditis: The Task Force for the Management of Infective Endocarditis of the European Society of Cardiology (ESC). Endorsed by: European Association for Cardio-Thoracic Surgery (EACTS), the European Association of Nuclear Medicine (EANM). *Eur Heart J*. 2015 Nov 21;36(44):3075-3128. doi: 10.1093/eurheartj/ehv319. Epub 2015 Aug 29. PMID: 26320109.
10. Lazar HL, McDonnell M, Chipkin SR, Furnary AP, Engelman RM, Sadhu AR, et al; Society of Thoracic Surgeons Blood Glucose Guideline Task Force. The Society of Thoracic Surgeons practice guideline series: Blood glucose management during adult cardiac surgery. *Ann Thorac Surg*. 2009 Feb;87(2):663-9. doi: 10.1016/j.athoracsur.2008.11.011. PMID: 19161815.
11. Gu K, Cowie CC, Harris MI. Mortality in adults with and without diabetes in a national cohort of the U.S. population, 1971-1993. *Diabetes Care*. 1998 Jul;21(7):1138-45. doi: 10.2337/diacare.21.7.1138. PMID: 9653609.
12. Walker J, Colhoun H, Livingstone S, McCrimmon R, Petrie J, Sattar N, Wild S; Scottish Diabetes Research Network Epidemiology Group. Type 2 diabetes, socioeconomic status and life expectancy in Scotland (2012-2014): a population-based observational study. *Diabetologia*. 2018 Jan;61(1):108-116. doi: 10.1007/s00125-017-4478-x. Epub 2017 Oct 26. PMID: 29075822; PMCID: PMC6448945.
13. Kang YM, Cho YK, Lee SE, Park JY, Lee WJ, Kim YJ, Jung CH. Cardiovascular diseases and life expectancy in adults with Type 2 Diabetes: A Korean National Sample Cohort Study. *J Clin Endocrinol Metab*. 2017 Sep 1;102(9):3443-3451. doi: 10.1210/jc.2017-00643. PMID: 28911137.
14. Movahed MR, Hashemzadeh M, Jamal MM. Increased prevalence of infectious endocarditis in patients with type II diabetes mellitus. *J Diabetes Complications*. 2007 Nov-Dec;21(6):403-6. doi:

- 10.1016/j.jdiacomp.2007.07.003. PMID: 17967715.
15. Kourany WM, Miro JM, Moreno A, Corey GR, Pappas PA, Abrutyn E, Hoen B, Habib G, Fowler VG Jr, Sexton DJ, Olaison L, Cabell CH; ICE MD Investigators. Influence of diabetes mellitus on the clinical manifestations and prognosis of infective endocarditis: a report from the International Collaboration on Endocarditis-Merged Database. *Scand J Infect Dis*. 2006;38(8):613-9. doi: 10.1080/00365540600617017. PMID: 16857604.
 16. Olmos C, Vilacosta I, Pozo E, Fernández C, Sarriá C, López J, Ferrera C, Maroto L, González I, Vivas D, Palacios J, San Román JA. Prognostic implications of diabetes in patients with left-sided endocarditis: findings from a large cohort study. *Medicine (Baltimore)*. 2014 Mar;93(2):114-119. doi: 10.1097/MD.0000000000000023. PMID: 24646468; PMCID: PMC4616308
 17. Farag M, Borst T, Sabashnikov A, Zerriouh M, Schmack B, Arif R, Beller CJ, Popov AF, Kallenbach K, Ruhparwar A, Dohmen PM, Szabó G, Karck M, Weymann A. Surgery for Infective Endocarditis: Outcomes and Predictors of Mortality in 360 Consecutive Patients. *Med Sci Monit*. 2017 Jul 25;23:3617-3626. doi: 10.12659/msm.902340. PMID: 28740070; PMCID: PMC5539855.
 18. Wei XB, Liu YH, Huang JL, Chen XL, Yu DQ, Tan N, Chen JY, He PC. Prediabetes and diabetes are both risk factors for adverse outcomes in infective endocarditis. *Diabet Med*. 2018 Nov;35(11):1499-1507. doi: 10.1111/dme.13761. Epub 2018 Jul 4. PMID: 29908070
 19. Lin CJ, Chua S, Chung SY, Hang CL, Tsai TH. Diabetes Mellitus: An independent risk factor of in-hospital mortality in patients with infective endocarditis in a new era of clinical practice. *Int J Environ Res Public Health*. 2019 Jun 25;16(12):2248. doi: 10.3390/ijerph16122248. PMID: 31242695; PMCID: PMC6617149.
 20. Østergaard L, Mogensen UM, Bundgaard JS, Dahl A, Wang A, Torp-Pedersen C, Gislason G, Køber L, Køber N, Dejgaard TF, Frandsen CS, Fosbøl EL. Duration and complications of diabetes mellitus and the associated risk of infective endocarditis. *Int J Cardiol*. 2019 Mar 1;278:280-284. doi:10.1016/j.ijcard.2018.09.106. Epub 2018 Oct 1. PMID: 30291010.
 21. Østergaard L, Smerup MH, Iversen K, Jensen AD, Dahl A, Chamat-Hedemand S, Bruun NE, Butt JH, Bundgaard H, Torp-Pedersen C, Køber L, Fosbøl E. Differences in mortality in patients undergoing surgery for infective endocarditis according to age and valvular surgery. *BMC Infect Dis*. 2020 Sep 25;20(1):705. doi: 10.1186/s12879-020-05422-8. PMID: 32977755; PMCID: PMC7519559.

Figures

Figure 1

5 years Survival rate by DM groups