Metagenomic pipeline for identifying co-infections among distinct SARS-CoV-2 variants of concern: Study cases from Alpha to Omicron

Jose Arturo Molina-Mora (✉ jose.molinamora@ucr.ac.cr)

Universidad de Costa Rica

Estela Cordero-Laurent

Instituto Costarricense de Investigación y Enseñanza en Nutrición y Salud (INCIENSA)

Melany Calderón-Osorno

Instituto Costarricense de Investigación y Enseñanza en Nutrición y Salud (INCIENSA)

Edgar Chacón-Ramírez

Universidad de Costa Rica

Francisco Duarte-Martínez

Instituto Costarricense de Investigación y Enseñanza en Nutrición y Salud (INCIENSA)

Research Article

Keywords: COVID-19, SARS-CoV-2, Co-infection, Omicron, Variant of concern, Delta

Posted Date: February 23rd, 2022

DOI: <https://doi.org/10.21203/rs.3.rs-1389767/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Scientific Reports on June 7th, 2022. See the published version at <https://doi.org/10.1038/s41598-022-13113-4>.

Abstract

Concomitant infection or co-infection with distinct SARS-CoV-2 genotypes have been reported as part of the epidemiological surveillance of the COVID-19 pandemic. In the context of the spread of more transmissible variants during 2021, co-infections are not only important due to the possible changes in the clinical outcome, but also the chance to generate new genotypes by recombination. However, a few approaches have developed bioinformatic pipelines to identify co-infections.

Here we present a metagenomic pipeline based on the inference of multiple fragments similar to amplicon sequence variant (ASV-like) from sequencing data and a custom SARS-CoV-2 database to identify the concomitant presence of divergent SARS-CoV-2 genomes, i.e., variants of concern (VOCs). This approach was compared to another strategy based on whole-genome (metagenome) assembly. Using single or pairs of sequencing data of COVID-19 cases with distinct SARS-CoV-2 VOCs, each approach was used to predict the VOC classes (Alpha, Beta, Gamma, Delta, Omicron or non-VOC and their combinations). The performance of each pipeline was assessed using the ground-truth or expected VOC classes. Subsequently, the ASV-like pipeline was used to analyze 1021 cases of COVID-19 from Costa Rica to investigate the possible occurrence of co-infections.

After the implementation of the two approaches, an accuracy of 96.2% was revealed for the ASV-like inference approach, which contrasts with the misclassification found (accuracy 46.2%) for the whole-genome assembly strategy. The custom SARS-CoV-2 database used for the ASV-like analysis can be updated according to the appearance of new VOCs to track co-infections with eventual new genotypes. In addition, the application of the ASV-like approach to all the 1021 sequenced samples from Costa Rica in the period October 12th - December 21th 2021 found that none corresponded to co-infections with VOCs.

In conclusion, we developed a metagenomic pipeline based on ASV-like inference for the identification of co-infection with distinct SARS-CoV-2 VOCs, in which an outstanding accuracy was achieved. Due to the epidemiological, clinical, and molecular relevance of the concomitant infection with distinct genotypes, this work represents another piece in the process of the surveillance of the COVID-19 pandemic in Costa Rica and worldwide.

Highlights

- We present a metagenomic pipeline to identify the concomitant presence of divergent SARS-CoV-2 genomes.
- The approach is based on the inference of multiple fragments similar to amplicon sequence variant (ASV-like).
- An accuracy of 96.2% was revealed for the ASV-like inference approach, which contrasts with the misclassification found for the whole-genome assembly strategy.
- A custom SARS-CoV-2 database was required for the ASV-like analysis, that can be updated according to the appearance of new genotypes.

- Analysis of 1021 sequenced samples from Costa Rica in 2021 found that none corresponded to co-infections by divergent genotypes.

Introduction

The COVID-19 pandemic, caused by the SARS-CoV-2 virus, has affected 282 million people worldwide and 570 000 people in Costa Rica by December 2021. The genomic sequencing approach is one of the hallmarks in the management of COVID-19 to follow up virus evolution and spread across the globe almost in real-time, unlike other pandemics (Gouvêa dos Santos, 2021). Thus, since the emergence of the virus, efforts have been made to map the genetic diversity of the virus and to identify genotypes with a possible selective advantage (Young et al., 2020).

The SARS-CoV-2 genotypes, which share several common mutations and are expected to have similar biological properties, can be classified as clades, PANGOLIN lineages, or variants depending on the nomenclature system. These versions of the SARS-CoV-2 virus have been reported each time faster during the last year in part due to the increased mutation rate of the virus over time (Schrörs et al., 2021). Out of the thousands of lineages that have been reported at the end of the year 2021, the World Health Organization (WHO) have recognized five of those divergent genotypes as a variant of concern (VOC), mainly due to the increased transmissibility and/or a capacity to evade inhibition by neutralizing antibodies. The divergent VOCs, namely Alpha (lineage B.1.1.7), Beta (B.1.351), Gamma (P.1), Delta (B.1.617.2), and Omicron (B.1.1.529) variants, have been initially reported in specific geographic regions, but rapidly were spread to multiple locations worldwide (Bentley et al., 2021; Schrörs et al., 2021; Team, 2021). Other genotypes, such as the variants of interest (VOI) or variants under monitoring (VUM) have been also reported, which are still under study owed to possible changes in the patterns of transmission, severity, clinical manifestations, mortality, or vaccine effectiveness (Graham et al., 2021; J. A. Molina-Mora, 2021).

Because of the amount and features of circulating variants, epidemiological surveillance of the pandemic must include the analysis of concomitant infections (co-infection) with different SARS-CoV-2 genomes (Banerjee, Mossman, & Grandvaux, 2021). Co-infection can be described as the occurrence of a re-infection when a first infection was not yet cured (Pedro et al., 2021) or the horizontal transmission of multiple genotypes (Hashim et al., 2020). Estimation of frequency and the study of effects of co-infections are relevant not only for the management of the disease at the personal (symptoms) or population (transmission) level but also the molecular surveillance of possible risky events of recombination that can be triggered (Francisco et al., 2021). However, the incidence of concomitant mixed infections with different genotypes has not been extensively reported (Hashim et al., 2020; Liu et al., 2021). Some studies have reported up to 2.6-8% of COVID-19 cases as co-infections (Liu et al., 2021; Schrörs et al., 2021; Young et al., 2020), but a more specific and confident analysis found that 0.18% of cases were concomitant infections (Zhou et al., 2021).

Regarding the bioinformatic strategies, only a few studies have been implemented analysis to detect co-infections by SARS-CoV-2 genomes (Francisco et al., 2021; Hashim et al., 2020; Liu et al., 2021; Young et al., 2020). These pipelines are based on the identification of haplotypes (sequences of each genome in the concomitant infection) using haplotype reconstruction programs (Francisco et al., 2021; Hashim et al., 2020; Liu et al., 2021; Young et al., 2020). However, haplotype identification is usually used to detect co-infections with related but distinct viruses and performs poorly for close genomes (Schirmer, Sloan, & Quince, 2014). The only specific pipeline to identify co-infections by divergent SARS-CoV-2 viruses was recently developed by (Zhou et al., 2021).

In this context and as part of the epidemiological surveillance of the pandemic in Costa Rica, we now present a new pipeline based on metagenomic analyses to detect co-infections with divergent SARS-CoV-2 viruses, specifically with VOCs. After sequencing and pre-processing, the workflow follows the inference of multiple fragments similar to amplicon sequence variant (ASV-like) from sequencing data and a taxonomy assignment with a custom database of SARS-CoV-2 sequences. Thus, this study aimed to develop a pipeline to identify co-infections with divergent SARS-CoV-2 genomes using an ASV inference approach.

Methods

General strategy

To identify cases of COVID-19 with a co-infection with two distinct SARS-CoV-2 VOCs, we implemented a strategy using genome sequencing data and two different pipelines (Fig. 1). Sequencing data of samples with different SARS-CoV-2 lineages (one or two lineages, including VOCs) were obtained (Fig. 1, A). A first strategy was the whole-genome assembly, in which a metagenomic assembler was used to build the genome sequence(s) in the sample. After the lineage assignment, genome sequences in each sample were classified into VOC classes (Alpha, Beta, Gamma, Delta, Omicron, or non-VOC) and this prediction was compared to the known or expected categories (Fig. 1, B).

In a second approach, sequencing data were used for the analysis with sequences at single-nucleotide resolution, the ASV-like calling strategy. In this case, lineage assignment was not directly done but ASVs were mapped to a custom database of SARS-CoV-2 genomes sequences. For genome sub-sequences that are shared among all the lineages, the corresponding ASV are expected to map multiple sequences, including the non-VOC genomes if they are provided first. However, ASVs carrying specific mutations of the VOC are expected to only map to the genome of the variant. Thus, for each genome sequence in the database, the mapping ASVs were counted to assign the sample into a VOC category (Fig. 1, C). The prediction of these classes was compared to the expected results to assess the performance of the pipeline.

Clinical isolates and genome sequencing

Using the sample collection of Costa Rican cases of COVID-19 in the period between March 2020 and August 2021 as part of the genomic surveillance of the SARS-CoV-2 virus, 12 samples from distinct lineages were selected (single lineages in Table 2). Genotypes included VOCs and VOIs, as well as the regional lineage B.1.1.389 circulating in Costa Rica (Table 2).

Patients had been diagnosed in INCIENSA (Instituto Costarricense de Investigación y Enseñanza en Nutrición y Salud) or different public and private clinical laboratories by real-time reverse transcription polymerase chain reaction (RT-PCR) using nasopharyngeal swab samples. The diagnosis was done using the guidelines of the Pan American Health Organization and the World Health Organization (PAHO, 2020), and the Ministry of Health of Costa Rica.

Selected samples had a CT < 25 (cycle threshold in the PCR) and genome sequencing had been done in the local sequencing service of INCIENSA. Amplicons were obtained using the protocol by (Resende et al., 2020). Sequencing libraries were prepared using the Illumina DNA Prep Kit (Illumina, San Diego, CA, USA) according to the laboratory standard operating procedure for pulsenet Nextera DNA flex library preparation (<https://www.cdc.gov/pulsenet/pathogens/wgs.html>). Paired-end sequencing was performed for each library on a MiSeq instrument using 500 cycles v2 chemistry cartridges (Illumina, San Diego, CA, USA).

Pre-processing and ground-truth genotype

FastQC v0.11.7 (Andrews, 2010) was used for the quality control of sequencing data. Trimmomatic v0.38 (Bolger, Lohse, & Usadel, 2014) was used for adapters removal and trimming of low-quality bases (Q < 30). Filtered reads were used to infer the ground-truth genotype, the *de novo* whole-genome assembly, and the ASV calling.

To identify the ground-truth (expected) genotype of the sequences, a reference-based genome assembly was implemented. BWA-MEM 0.7.5a-r405 (H. Li & Durbin, 2009) with default parameters was used to map reads to the reference genome NC_045512.2. Freebayes v1.3.1 (Garrison & Marth, 2012) (parameters -p 1 -q 20 -m 60 -min-coverage 10 -V) was implemented to call variants. Low-confidence variants were removed using VCF_filter v3.2 (https://github.com/moskalenko/vcf_filter). Annotation of variants was done using SNPeff (Cingolani et al., 2012). Genotype for each genome was allocated using the PANGOLIN lineages assigner version 3.1.17 (<https://pangolin.cog-uk.io/>). Based on the lineage, genome sequences were classified into the VOC classes (Alpha, Beta, Gamma, Delta, Omicron, or non-VOC) and the results were used as the ground-truth genotype (expected lineage or VOC class, Table 2).

Based on the SARS-CoV-2 genotypes found in the 12 samples (samples with a single lineage, Table 2), 14 new datasets were generated by combining sequencing data of two distinct cases (double lineages, Table 2). The ground-truth genotypes were inferred based on the individual genomes (expected genotypes, Table 2).

Whole-genome (metagenome) assembly

To assemble the genome of cases with a single (12 samples) or double (14 samples) genotype, a *de novo* metagenomic assembler was implemented using the filtered sequencing reads. Megahit v1.1.3 (D. Li, Liu, Luo, Sadakane, & Lam, 2015) was used due to its ability to build sequences of individual or multiple genomes (D. Li et al., 2015; J.-A. Molina-Mora, Campos-Sánchez, Rodríguez, Shi, & García, 2020). Genome assembly was evaluated based on contiguity, completeness, and correctness using the 3C criterion (J.-A. Molina-Mora et al., 2020; J. A. Molina-Mora & Garcia, 2020). Genotype for each genome was allocated using the PANGOLIN lineages assigner (<https://pangolin.cog-uk.io/>). Based on the lineage, the 26 genome sequences were classified into the VOC classes and the results were used as the prediction of this pipeline. The predicted genotypes were compared to the expected (ground-truth) genotypes (Table 2).

ASV-like inference

To call ASVs for each sample, the DADA2 package (Callahan et al., 2016) was run using the R software. The standard protocol of this software for Illumina sequencing data was implemented (<https://benjjneb.github.io/dada2/tutorial.html>), in which the only modified step was the taxonomy assignment using a custom database (see below). Briefly, sequencing data (fastq files) were filtered by quality as error rates were calculated and removed from the dereplicated reads. Possible chimeric reads were identified and removed. Finally, taxonomy was assigned using the RDP Naive Bayesian Classifier algorithm against a custom SARS-CoV-2 database, and an error-corrected table of the abundances of ASVs was obtained. In addition, because of the nature of the ASV inference in which a consensus sequence is built and results are dependent on the iteration, we completed 5 repetitions of the analysis to verify the reproducibility of the results.

Custom SARS-CoV-2 database: To assign the taxonomy to the ASVs, nine SARS-CoV-2 genome sequences were incorporated into a custom database. Sequences corresponded to VOCs (which were retrieved from <https://viralzone.expasy.org/9556>) and other non-VOC genomes circulating in Costa Rica since March 2020, which are detailed in Table 1. Because of the use of the DADA2 pipeline, the database required a format like 16S-rRNA databases for amplicon-based metagenomics (16S-rRNA). Thus, the sequence name in the fasta file required a format including the organism and subtype, i.e., ">SARS-CoV-2; VOC_class-lineage-ID" according to the data of Table 1. The SARS-CoV-2 genome database is provided as a supplementary file.

Table 1
SARS-CoV-2 sequences used to build the custom database to
classify genome sequences into VOC categories based on
ASV calling analysis.

ID	Lineage	VOC classes	Database
CRC-0381	B.1.1.519	Non-VOC	GISAID
CRC-0449	B.1.1.389	Non-VOC	GISAID
CRC-0493	B.1.525	Non-VOC	GISAID
CRC-0653	C36.3	Non-VOC	GISAID
MZ344997.1	B.1.1.7	Alpha	NCBI
MW598419.1	B.1.351	Beta	NCBI
MZ169911.1	P.1	Gamma	NCBI
MZ359841.1	B.1.617.2	Delta	NCBI
PI_ISL_6913995	B.1.1.529	Omicron	GISAID

Assessment of the pipelines performance

To assess the performance of the two pipelines to identify co-infections, the genome assembly and the ASV calling, the predictions regarding the VOC class of each approach were compared to the expected genotype. A ROC (Receiver Operating Characteristics) analysis was implemented in R software (<https://www.r-project.org/>) using the ROCR package (Sing, Sander, Beerenwinkel, & Lengauer, 2005). AUC (Area Under Curve) of the ROC curve, as well as the general accuracy of the predictions, were calculated for each pipeline.

Estimation of occurrence of co-infections in Costa Rica

Using the ASV-like approach, we run the pipeline to identify possible co-infection with VOCs in COVID-19 cases in Costa Rica. The analysis was done using all the samples which had locally sequenced (1021 cases) in the period October 12th - December 21th 2021. Samples were processed as described before.

Results

In order to identify cases of COVID-19 with a co-infection with two distinct SARS-CoV-2 VOCs, we implemented a strategy using genome sequencing data and two different pipelines (Fig. 1). First, a whole-genome analysis approach, which included the metagenome assembly, lineage assignment, and the classification into the VOC categories, resulted not suitable to identify co-infections with VOCs. With an accuracy of 46.2%, this strategy misclassified all the genome sequences for cases with two lineages, including VOCs, while only samples with a single lineage were properly identified (Table 2 and Fig. 2). The

ROC analysis found a value of $AUC = 0.500$, revealing that the performance of the VOC assignment is equivalent to a classification by chance.

Table 2

Genome classification using a whole-genome assembly strategy with sequencing data for one or two variants of the SARS-CoV-2 virus (PERF: performance of the prediction regarding the expected class)

Samples		Expected genotype		Predicted genotype		PERF
Type	ID	Expected lineage	Expected VOC class	Predicted lineage	Predicted VOC class	
Single lineage	S1	B.1.1.389	Non-VOC	B.1.1.389	Non-VOC	1
	S2	C.36.3	Non-VOC	C.36.3	Non-VOC	1
	S3	P.2	Non-VOC	P.2	Non-VOC	1
	S4	B.1.625	Non-VOC	B.1.625	Non-VOC	1
	S5	B.1.429	Non-VOC	B.1.429	Non-VOC	1
	S6	B.1.525	Non-VOC	B.1.525	Non-VOC	1
	S7	B.1.1.519	Non-VOC	B.1.1.519	Non-VOC	1
	S8	B.1.1.7	Alpha	B.1.1.7	Alpha	1
	S9	B.1.351	Beta	B.1.351	Beta	1
	S10	P.1	Gamma	P.1	Gamma	1
	S11	AY.113	Delta	AY.113	Delta	1
	S12	B.1.1.529	Omicron	B.1.1.529	Omicron	1
Double lineage	D1 (S4 + S6)	B.1.625 + B.1.525	Non-VOC (+ Non-VOC)	B.1	Non-VOC	1
	D2 (S1 + S8)	B.1.1.389 + B.1.1.7	Alpha (+ Non-VOC)	B.1.1	Non-VOC	1
	D3 (S8 + S9)	B.1.1.7 + B.1.351	Alpha + Beta	B.1	Non-VOC	1
	D4 (S8 + S10)	B.1.1.7 + P.1	Alpha + Gamma	B.1	Non-VOC	1
	D5 (S8 + S11)	AY.113 + B.1.1.7	Alpha + Delta	B.1	Non-VOC	1
	D6 (S8 + S12)	B.1.1.7 + B.1.1.529	Alpha + Omicron	B.1	Non-VOC	1
	D7 (S9 + S11)	AY.113 + B.1.351	Beta + Delta	B.1	Non-VOC	1
	D8 (S9 + S10)	P.1 + B.1.351	Beta + Gamma	B.1	Non-VOC	1
	D9 (S9 + S12)	B.1.351 + B.1.1.529	Beta + Omicron	B.1	Non-VOC	1

D10 (S11 + S10)	AY.113 + P.1	Delta + Gamma	B.1	Non-VOC	□
D11 (S10 + S12)	P.1 + B.1.1.529	Gamma + Omicron	B.1	Non-VOC	□
D12 (S11 + S2)	AY.113 + C.36.3	Delta (+ Non-VOC)	B.1.629	Non-VOC	□
D13 (S7 + S11)	AY.113 + B.1.1.519	Delta (+ Non-VOC)	B.1	Non-VOC	□
D14 (S11 + S12)	AY.113 + B.1.1.529	Delta + Omicron	B.1	Non-VOC	□

To deal with this, a second metagenomic approach was implemented using an ASV-like calling strategy. A custom database of SARS-CoV-2 genomes sequences was created to assign the taxonomy of the ASV sequences into VOCs or Non-VOC genomes. During the standardization, it was determined that the identification of a VOC was possible if at least three ASV were mapped to the VOC genome. Thus, the presence of a specific VOC was determined if the number of total mapping ASVs was ≥ 3 . This is in line with the profile of mutations for each VOC, in which a few mutations are shared and most of them are exclusive (Fig. 3).

According to Table 3, the ASV-inference pipeline was able to correctly classify all samples with two lineages but one (sample D13) into the VOC classes. In the same way, all the samples with a single genotype were correctly classified. As presented in the Supplementary file, the classification of the VOC class is consistent among the iterations for all the samples when the pipeline was run 5 times to assess reproducibility. In the case of the misclassified sample D13, the combination was done using the Delta variant and a case of the B.1.1.519 lineage. Results of the 5 iterations predicted a Beta variant in the sample.

Table 3

Genome classification using an ASV inference strategy with sequencing data for one or two variants of the SARS-CoV-2 virus (PERF: performance of the VOC prediction regarding the expected VOC class)

Samples		Expected genotype		Predicted genotype		PERF
Type	ID	Expected lineage	Expected VOC class	Score by iterations	Predicted VOC class	
Single lineage	S1	B.1.1.389	Non-VOC	5	Non-VOC	□
	S2	C.36.3	Non-VOC	5	Non-VOC	□
	S3	P.2	Non-VOC	5	Non-VOC	□
	S4	B.1.625	Non-VOC	5	Non-VOC	□
	S5	B.1.429	Non-VOC	5	Non-VOC	□
	S6	B.1.525	Non-VOC	5	Non-VOC	□
	S7	B.1.1.519	Non-VOC	5	Non-VOC	□
	S8	B.1.1.7	Alpha	5	Alpha	□
	S9	B.1.351	Beta	5	Beta	□
	S10	P.1	Gamma	5	Gamma	□
	S11	AY.113	Delta	5	Delta	□
	S12	B.1.1.529	Omicron	5	Omicron	□
Double lineage	D1 (S4 + S6)	B.1.625 + B.1.525	Non-VOC (+ Non-VOC)	5	Non-VOC (+ Non-VOC)	□
	D2 (S1 + S8)	B.1.1.389 + B.1.1.7	Alpha (+ Non-VOC)	4	Alpha (+ Non-VOC)	□
	D3 (S8 + S9)	B.1.1.7 + B.1.351	Alpha + Beta	3	Alpha + Beta	□
	D4 (S8 + S10)	B.1.1.7 + P.1	Alpha + Gamma	4	Alpha + Gamma	□
	D5 (S8 + S11)	AY.113 + B.1.1.7	Alpha + Delta	4	Alpha + Delta	□
	D6 (S8 + S12)	B.1.1.7 + B.1.1.529	Alpha + Omicron	5	Alpha + Omicron	□
	D7 (S9 + S11)	AY.113 + B.1.351	Beta + Delta	5	Beta + Delta	□
	D8 (S9 + S10)	P.1 + B.1.351	Beta + Gamma	5	Beta + Gamma	□
	D9 (S9 + S12)	B.1.351 + B.1.1.529	Beta + Omicron	5	Beta + Omicron	□

D10 (S11 + S10)	AY.113 + P.1	Delta + Gamma	5	Delta + Gamma	□
D11 (S10 + S12)	P.1 + B.1.1.529	Gamma + Omicron	5	Gamma + Omicron	□
D12 (S11 + S2)	AY.113 + C.36.3	Delta (+ Non-VOC)	5	Delta (+ Non-VOC)	□
D13 (S7 + S11)	AY.113 + B.1.1.519	Delta (+ Non-VOC)	0	Beta	□
D14 (S11 + S12)	AY.113 + B.1.1.529	Delta + Omicron	5	Delta + Omicron	□

When the profile of mutations were compared, 3 mutations (ORF1b-P314L, ORF8-S84L and S-D614G) are shared by Beta, Delta and B.1.1.519 genotypes, which made the ASV-like inference to incorrectly assign the subsequences to the Beta variant. However, this phenomenon was not a drawback for the rest of cases.

The metrics for the ASV-like calling strategy showed an accuracy value of 96.2% and AUC of 0.964 (Figure 2), indicating an outstanding performance in the identification of the lineages for cases with one or two SARS-CoV-2 variants. Based on these results, the ASV-like calling is a suitable strategy to identify co-infections with two SARS-CoV-2 VOCs.

Finally, we investigated the possible occurrence of SARS-CoV co-infections in Costa using the ASV-like approach. We found that none of 1021 samples were identified with concomitant infections with distinct VOCs in the period October 12th - December 21th 2021.

Discussion

The SARS-CoV-2 genome has rapidly evolved into multiple variants due to not only the widespread in diverse human populations but also the increase in the mutation rate during 2021 (Schrörs et al., 2021). In this context, the interaction of multiple viral sequences with each other during simultaneous infection can lead to potential differences in epidemiological behavior (Hashim et al., 2020). Thus, it is vital to reveal the frequency of co-infection events, how often it occurs in the population as well as and the exact composition of lineages (Zhou et al., 2021).

Here we presented an analysis of co-infection by divergent VOCs of SARS-CoV-2 virus, in which samples with two distinct genotypes were analyzed using a metagenomic approach by ASV inference. Similar to another work by (Zhou et al., 2021), we assumed that the existence of specific lineage-defined feature mutations of the lineages in viral quasi-species achieves the identification of co-infection events (Fig. 3). A custom SARS-CoV-2 database led to identify specific ASV belonging to VOCs, as well as non-specific ASV found in other genotypes. The metrics of the classification (VOCs classes) revealed a high performance of the pipeline with an accuracy of 96.2% and AUC of 0.964. This completely contrasted

with the metagenome assembly approach, in which the classification was suggested to be by chance (accuracy = 46.2% and AUC = 0.500). The poor performance of the metagenome assembly relies on the generation of a single consensus sequence even for cases with two distinct genomes, in which a mutation of a genotype can be overshadowed by the presence of the non-mutated nucleotide in the other sequence, creating an incorrect profile of few mutations with a erroneous VOC class assignment (Table 2).

Besides, previous to the arrival of the omicron variant by November 2021, the first version of this work was prepared and the accuracy was reported in 96.3% for the ASV-like approach and 57.7% for the whole-genome assembly (more data not shown). This update demonstrated the versatility of our approach to incorporate new genotypes to infer co-infections with distinct VOCs.

Regarding sequencing data, the high reliability of Illumina technology has been reported to keep the genomic evidence of co-infections or within-host variations (Zhou et al., 2021), which has motivated its use for co-infection studies, including this work. General approaches to identify the concomitant presence of organisms can be done using: (i) metagenomics strategies, or (ii) strategies based on the reconstruction of haplotypes by mapping. For SARS-CoV-2 co-infections, the last has been the selected method due to the availability of ready-to-use bioinformatic tools (Leviyang, Griva, Ita, & Johnson, 2017; Prabhakaran, Rey, Zagordi, Beerenwinkel, & Roth, 2014; Prosperi & Salemi, 2012). Despite this, viral haplotype reconstruction programs usually perform poorly for sequences with low divergence or rare haplotypes (Schirmer, Sloan, & Quince, 2014), which represent a possible limitation for the use among samples with simultaneous SARS-CoV-2 genomes. Because of this, assembly of single genomes and the subsequent combination into simulated co-infection data were preferred here not only for the developed pipeline but also to create the ground-truth dataset rather than the comparison to a haplotype caller.

Using analysis of haplotype reconstruction, some studies have reported events of co-infection caused by the occurrence of two distinct genotypes. In 2020, 19 cases of co-infections were identified in Iraq (Hashim et al., 2020). Up to 8% of co-infections were informed in a study from Singapore (Young et al., 2020), while at least 5% was estimated in the United Arab Emirates (Liu et al., 2021). In Brazil, a co-infection was detected with local lineages in early 2021 (Francisco et al., 2021). By September 2021, a study analyzed 30,806 raw sequence datasets, in which about 2.6% were identified as co-infections at high confidence (Schrörs et al., 2021).

To our knowledge, only one study has implemented a specific pipeline to identify co-infections by distinct SARS-CoV-2 genotypes, which used a strategy with an intra-host variant calling analysis and a hypergeometric distribution method (Zhou et al., 2021). Using sequencing data of COVID-19 cases from the United States of America, authors recognized only 53 out of 29 993 samples (0.18%) as co-infection cases. A single case was reported with three lineages, while the other 52 were identified with two genotypes. These results regarding the frequency are in line with our results of the possible occurrence of co-infections in Costa Rica. None of the 1021 analyzed cases were identified as a concomitant presence of VOCs. Due to the frequency of co-infections, which is suggested to be very low, the analyzed cases

could be not enough to identify at least a single case in this country. In addition, other factors affecting this result are the very few sequenced samples in comparison to the diagnosed cases (0.4% in Costa Rica and 0.41% in Latin America according to GISAID database), the inability to clinically differentiate cases of co-infection (see later), as well as the rapid displacement of circulating lineages by VOCs which have a higher transmissibility. However, with the co-dominance of Delta and Omicron during the transition between the years 2021 and 2022 (GISAID, 2021), the reports of co-infections could be increased in the coming months. Thereby, this work could be a useful tool to investigate the occurrence of this phenomenon.

About the biological meaning, the report of co-infections is of concern because other studies have demonstrated that this phenomenon can contribute to the recombination of RNA viruses (Banerjee et al., 2021; Gouvêa dos Santos, 2021; Zhou et al., 2021). Product of the recombination processes, the new virions may acquire different pathogenic properties (Gouvêa dos Santos, 2021) and it might impact the clinical presentation of the disease into more severe symptoms (Zhou et al., 2021).

Until now, cases with a co-infection with distinct SARS-CoV-2 genotypes have been reported with the same symptoms as other COVID-19 patients (Francisco et al., 2021; Young et al., 2020). However, a single report of co-infection, in a young and healthy female patient with severe COVID-19, suggested the concomitant infection as responsible for the clinical presentation (Pedro et al., 2021). More studies and updated statistics are required to establish the relevance of co-infections in terms of severity and mortality of COVID-19 disease (Hashim et al., 2020). Also, the detection of possible cases of co-infections is another factor to consider in the interaction between the immune system and SARS-CoV-2 mutations. It has been suggested that immunity driven by a specific SARS-CoV-2 genotype does not protect against another one but can instead lead to a more severe disease pattern (Pedro et al., 2021). However, the real immunological implications of co-infections on the cellular or humoral levels are not well known (Francisco et al., 2021).

On the other hand, in this study some considerations and limitations are needed to take into account. First, the approach using ASV inference requires a custom database that needs to be built using sequences of genomes circulating locally, i.e., local genomic surveillance is a previous step to implement the pipeline. This includes the update of VOC sequences carrying new mutations (sublineages). Second, similar to the other approaches to identify co-infections, only co-infections with divergent sequences (VOCs in this case) can be identified. A test using co-infections with VOIs and VUMs was not able to identify concomitant sequences due to the low diversity, in which a poor performance was obtained during the genotype classification. Also, *de novo* intra-host mutation cannot be identified using this approach. This does not represent a drawback for this implementation because there is a very low probability of the *de novo* appearance of mutations corresponding exactly to all the feature mutations of the VOCs. If some *de novo* mutation was equal to a feature mutation, the ASV can be discarded with the implementation of the threshold of the mapping ASVs, as we followed here. Finally, although co-infection with SARS-CoV-2 and other pathogens have been reported, such as Influenza or bacterial agents (Dadashi

et al., 2021; Musuuza et al., 2021), and this pipeline could be adapted to identify them, a metagenome assembly could be a suitable strategy rather than this approach.

Altogether, this analysis represents a new effort to track the SARS-CoV-2 genotypes circulating in Costa Rica, which are complementary to our other local studies for genomic surveillance (J. A. Molina-Mora, 2021; J. A. Molina-Mora, Cordero-Laurent, et al., 2021) as well as the identification of clinical patterns of COVID-19 patients (J. A. Molina-Mora, González, et al., 2021). Concomitant infection with distinct viral genotypes can lead to the generation of SARS-CoV-2 variants with possible new properties in terms of transmission, severity, mortality, or vaccine effectiveness. This remarks the relevance to continue with the surveillance of the dynamics of the pandemic including origin and tracking, genotyping, and clinical features of the infections worldwide, which can eventually arise new insights about co-infection events.

Conclusions

In conclusion, we developed a metagenomic pipeline based on ASV-like inference for the identification of co-infection with distinct SARS-CoV-2 VOCs, in which a 96.2% of accuracy was achieved. This performance was outstanding in comparison to the whole-genome assembly approach in which a resolution by chance was suggested with an accuracy of 46.2%. The custom SARS-CoV-2 database used for the ASV-like inference can be updated according to the appearance of new VOCs to track co-infections with eventual new genotypes. In addition, the application of the ASV-like approach to all the 1021 sequenced samples from Costa Rica in the period October 12th - December 21th 2021 found that none corresponded to co-infections with VOCs. Although a small percentage of COVID-19 cases worldwide are reported as co-infections with different SARS-CoV-2 lineages, the spread of more transmissible variants and the possibility of recombination to induce new genotypes remark the need for developing tools and pipelines to track concomitant infections with SARS-CoV-2 variants. Thus, this work represents another piece in the process of the genomic surveillance of the COVID-19 pandemic in Costa Rica and worldwide.

Declarations

Acknowledgments

We are grateful to clinicians, microbiologists, and other personnel of public (Caja Costarricense de Seguro Social CCSS) and private clinical laboratories for the samples of confirmed cases of COVID-19. We also thank Meriyeins Sibaja, Carlos Martínez and Daniel Ulate for their participation in distinct activities associated with the project.

Author contributions

J.A.M.M., E.C.L and F.D.M. participated in the conception and design of the study. E.C.L., M.C.O., and F.D.M. were involved in sample processing. J.A.M.M. implemented and standardized the bioinformatics pipelines. J.A.M.M. and E.C.M processed all data using the pipelines. J.A.M.M., E.C.L., and F.D.M.

participated in the interpretation of results. J.A.M.M. drafted the manuscript. All authors reviewed and approved the final manuscript.

Ethical approval and consent to participate

This study was approved by INCIENSA (INCIENSA-DG-of-2020-174) and the scientific committee of CIET-UCR (No. 242-2020). Samples were collected for epidemiological surveillance according to the Costa Rican regulation Law N° 8270 (May 17th, 2002), in which no additional consent was required for retrospective studies of archived and anonymized samples.

Availability of data and material

Script and the custom database used in this work are available at <https://github.com/josemolina6/sars-cov-2-co-infections>.

Declaration of Competing Interest

The authors declare that there is no conflict of interest.

Funding

This work was funded by Instituto Costarricense de Investigación y Enseñanza en Nutrición y Salud (INCIENSA) and Vicerrectoría de Investigación – Universidad de Costa Rica, with the Project “C0196 Protocolo bioinformático y de inteligencia artificial para el apoyo de la vigilancia epidemiológica basada en laboratorio del virus SARS-CoV-2 mediante la identificación de patrones genómicos y clínico-demográficos en Costa Rica (2020-2022)”.

References

- Andrews, S. (2010). FastQC A Quality Control tool for High Throughput Sequence Data. Retrieved April 10, 2018, from <https://www.bioinformatics.babraham.ac.uk/projects/fastqc/>
- Banerjee, A., Mossman, K., & Grandvaux, N. (2021). Molecular Determinants of SARS-CoV-2 Variants. *Trends in Microbiology*, 29(10), 871–873. <https://doi.org/10.1016/J.TIM.2021.07.002>
- Bentley, E. G., Kirby, A., Sharma, P., Kipar, A., Mega, D. F., Bramwell, C., ... Stewart, J. P. (2021). SARS-CoV-2 Omicron-B.1.1.529 Variant leads to less severe disease than Pango B and Delta variants strains in a mouse model of severe COVID-19. *BioRxiv*, 2021.12.26.474085. <https://doi.org/10.1101/2021.12.26.474085>
- Bolger, A. M., Lohse, M., & Usadel, B. (2014). Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics*, 30(15), 2114–2120. <https://doi.org/10.1093/bioinformatics/btu170>

- Callahan, B. J., McMurdie, P. J., Rosen, M. J., Han, A. W., Johnson, A. J. A., & Holmes, S. P. (2016). DADA2: High-resolution sample inference from Illumina amplicon data. *Nature Methods* 13:7, 13(7), 581–583. <https://doi.org/10.1038/nmeth.3869>
- Cingolani, P., Platts, A., Wang, L. L., Coon, M., Nguyen, T., Wang, L., ... Ruden, D. M. (2012). A program for annotating and predicting the effects of single nucleotide polymorphisms, SnpEff: SNPs in the genome of *Drosophila melanogaster* strain w1118; iso-2; iso-3. *Fly*, 6(2), 80–92. <https://doi.org/10.4161/fly.19695>
- Dadashi, M., Khaleghnejad, S., Abedi Elkhichi, P., Goudarzi, M., Goudarzi, H., Taghavi, A., ... Hajikhani, B. (2021). COVID-19 and Influenza Co-infection: A Systematic Review and Meta-Analysis. *Frontiers in Medicine*, 8, 681469. <https://doi.org/10.3389/FMED.2021.681469/FULL>
- Francisco, R. da S., Benites, L. F., Lamarca, A. P., de Almeida, L. G. P., Hansen, A. W., Gularte, J. S., ... Spilki, F. R. (2021). Pervasive transmission of E484K and emergence of VUI-NP13L with evidence of SARS-CoV-2 co-infection events by two different lineages in Rio Grande do Sul, Brazil. *Virus Research*, 296, 198345. <https://doi.org/10.1016/J.VIRUSRES.2021.198345>
- Garrison, E., & Marth, G. (2012). Haplotype-based variant detection from short-read sequencing. *ArXiv:1207.3907 [q-Bio.GN]*. Retrieved from <http://arxiv.org/abs/1207.3907>
- GISAID. (2021). GISAID - Clade and lineage nomenclature aids in genomic epidemiology of active hCoV-19 viruses. Retrieved November 18, 2020, from GISAID website: <https://www.gisaid.org/references/statements-clarifications/clade-and-lineage-nomenclature-aids-in-genomic-epidemiology-of-active-hcov-19-viruses/>
- Gouvêa dos Santos, W. (2021). Co-infection, re-infection and genetic evolution of SARS-CoV-2: Implications for the COVID-19 pandemic control. *Commentary Dos Santos*, 2(3), 56–61.
- Graham, M. S., Sudre, C. H., May, A., Antonelli, M., Murray, B., Varsavsky, T., ... Gunson, R. N. (2021). Changes in symptomatology, reinfection, and transmissibility associated with the SARS-CoV-2 variant B.1.1.7: an ecological study. *The Lancet Public Health*, 6(5), e335–e345. [https://doi.org/10.1016/s2468-2667\(21\)00055-4](https://doi.org/10.1016/s2468-2667(21)00055-4)
- Hashim, H. O., Mohammed, M. K., Mousa, M. J., Abdulameer, H. H., Alhassnawi, A. T. S., Hassan, S. A., & Al-Shuhaib, M. B. S. (2020). Infection with Different Strains of SARS-CoV-2 in Patients with COVID-19. *Archives of Biological Sciences*, 72(4), 575–585. <https://doi.org/10.2298/ABS201024051H>
- Leviyang, S., Griva, I., Ita, S., & Johnson, W. E. (2017). A penalized regression approach to haplotype reconstruction of viral populations arising in early HIV/SIV infection. *Bioinformatics*, 33(16), 2455. <https://doi.org/10.1093/BIOINFORMATICS/BTX187>
- Li, D., Liu, C.-M., Luo, R., Sadakane, K., & Lam, T.-W. (2015). MEGAHIT: an ultra-fast single-node solution for large and complex metagenomics assembly via succinct de Bruijn graph. *Bioinformatics*, 31(10),

1674–1676. <https://doi.org/10.1093/bioinformatics/btv033>

Li, H., & Durbin, R. (2009). Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics (Oxford, England)*, 25(14), 1754–1760. <https://doi.org/10.1093/bioinformatics/btp324>

Liu, R., Wu, P., Ogrodzki, P., Mahmoud, S., Liang, K., Liu, P., ... Zaher, W. A. (2021). Genomic epidemiology of SARS-CoV-2 in the UAE reveals novel virus mutation, patterns of co-infection and tissue specific host immune response. *Scientific Reports* 2021 11:1, 11(1), 1–14. <https://doi.org/10.1038/s41598-021-92851-3>

Molina-Mora, J.-A., Campos-Sánchez, R., Rodríguez, C., Shi, L., & García, F. (2020). High quality 3C de novo assembly and annotation of a multidrug resistant ST-111 *Pseudomonas aeruginosa* genome: Benchmark of hybrid and non-hybrid assemblers. *Scientific Reports*, 10(1), 1392. <https://doi.org/10.1038/s41598-020-58319-6>

Molina-Mora, J. A. (2021). Insights into the mutation T1117I in the spike and the lineage B.1.1.389 of SARS-CoV-2 circulating in Costa Rica. *Gene Reports*, 27, 1–24. <https://doi.org/10.1016/j.genrep.2022.101554>

Molina-Mora, J. A., Cordero-Laurent, E., Godínez, A., Calderón-Osorno, M., Brenes, H., Soto-Garita, C., ... Duarte-Martínez, F. (2021). SARS-CoV-2 genomic surveillance in Costa Rica: Evidence of a divergent population and an increased detection of a spike T1117I mutation. *Infection, Genetics and Evolution*, 92, 104872. <https://doi.org/10.1016/j.meegid.2021.104872>

Molina-Mora, J. A., & Garcia, F. (2020). The 3C criterion: Contiguity, Completeness and Correctness to assess de novo genome assemblies. *BMC Bioinformatics, Bioinformatics: From Algorithms to Applications*, 21(S20: 07), 5. <https://doi.org/10.1186/s12859-020-03838-2>

Molina-Mora, J. A., González, A., Jiménez-Morgan, S., Cordero-Laurent, E., Brenes, H., Soto-Garita, C., ... Duarte-Martínez, F. (2021). Clinical profiles at the time of diagnosis of COVID-19 in Costa Rica during the pre-vaccination period using a machine learning approach. *MedRxiv*, 2021.06.18.21259157. <https://doi.org/10.1101/2021.06.18.21259157>

Musuuza, J. S., Watson, L., Parmasad, V., Putman-Buehler, N., Christensen, L., & Safdar, N. (2021). Prevalence and outcomes of co-infection and superinfection with SARS-CoV-2 and other pathogens: A systematic review and meta-analysis. *PLOS ONE*, 16(5), e0251170. <https://doi.org/10.1371/JOURNAL.PONE.0251170>

PAHO, P. A. H. O. (2020). *Laboratory Guidelines for the Detection and Diagnosis of COVID-19 Virus Infection*. Retrieved from PAHO website: <https://www.who.int/emergencies/diseases/novel-coronavirus-2019/technical-guidance/laboratory->

- Pedro, N., Silva, C. N., Magalhães, A. C., Cavadas, B., Rocha, A. M., Moreira, A. C., ... Tavares, M. (2021). Dynamics of a Dual SARS-CoV-2 Lineage Co-Infection on a Prolonged Viral Shedding COVID-19 Case: Insights into Clinical Severity and Disease Duration. *Microorganisms* 2021, Vol. 9, Page 300, 9(2), 300. <https://doi.org/10.3390/MICROORGANISMS9020300>
- Prabhakaran, S., Rey, M., Zagordi, O., Beerenwinkel, N., & Roth, V. (2014). HIV Haplotype Inference Using a Propagating Dirichlet Process Mixture Model. *IEEE/ACM Transactions on Computational Biology and Bioinformatics*, 11(1), 182–191. <https://doi.org/10.1109/TCBB.2013.145>
- Prosperi, M. C. F., & Salemi, M. (2012). QuRe: software for viral quasispecies reconstruction from next-generation sequencing data. *Bioinformatics*, 28(1), 132–133. <https://doi.org/10.1093/BIOINFORMATICS/BTR627>
- Resende, P. C., Motta, F. C., Roy, S., Appolinario, L. R., Fabri, A., Xavier, J., ... Fundação Oswaldo Cruz. Instituto Oswaldo Cruz. Laboratório de Vírus Respiratório e do Sarampo. Rio de Janeiro, R. B. (2020). SARS-CoV-2 genomes recovered by long amplicon tiling multiplex approach using nanopore sequencing and applicable to other sequencing platforms. *BioRxiv*. <https://doi.org/10.1101/2020.04.30.069039>
- Schirmer, M., Sloan, W. T., & Quince, C. (2014). Benchmarking of viral haplotype reconstruction programmes: an overview of the capacities and limitations of currently available programmes. *Briefings in Bioinformatics*, 15(3), 431–442. <https://doi.org/10.1093/BIB/BBS081>
- Schrörs, B., Riesgo-Ferreiro, P., Sorn, P., Gudimella, R., Bukur, T., Rösler, T., ... Sahin, U. (2021). Large-scale analysis of SARS-CoV-2 spike-glycoprotein mutants demonstrates the need for continuous screening of virus isolates. *PLOS ONE*, 16(9), e0249254. <https://doi.org/10.1371/JOURNAL.PONE.0249254>
- Sing, T., Sander, O., Beerenwinkel, N., & Lengauer, T. (2005). ROCr: visualizing classifier performance in R. *Bioinformatics*, 21(20), 3940–3941. <https://doi.org/10.1093/BIOINFORMATICS/BTI623>
- Team, C. C.-19 R. (2021). SARS-CoV-2 B.1.1.529 (Omicron) Variant – United States, December 1–8, 2021. *Morbidity and Mortality Weekly Report*, 70(50), 1731. <https://doi.org/10.15585/MMWR.MM7050E1>
- Young, B. E., Fong, S. W., Chan, Y. H., Mak, T. M., Ang, L. W., Anderson, D. E., ... Ng, L. F. P. (2020). Effects of a major deletion in the SARS-CoV-2 genome on the severity of infection and the inflammatory response: an observational cohort study. *The Lancet*, 396(10251), 603–611. [https://doi.org/10.1016/S0140-6736\(20\)31757-8](https://doi.org/10.1016/S0140-6736(20)31757-8)
- Zhou, H.-Y., Cheng, Y.-X., Xu, L., Li, J.-Y., Tao, C.-Y., Ji, C.-Y., ... Wu, A. (2021). Genomic evidence for divergent co-infections of SARS-CoV-2 lineages. *BioRxiv*, 2021.09.03.458951. <https://doi.org/10.1101/2021.09.03.458951>

Figures

Figure 1

Conceptual design of the approach to identify co-infection by SARS-CoV-2 VOCs. Samples with two distinct VOCs (X and Y) of the SARS-CoV-2 virus are sequenced (A). Using a strategy for the whole-genome assembly, a single and misclassified sequence is obtained (B). In contrast, the correct identification of the two VOCs is achieved when an ASV inference is implemented, with the use of a custom database of SARS-CoV-2 for the taxonomy assignment. In this process, specific ASV are recognized for the VOCs, while shared ASVs among all the genomes are assigned to other sequences.

ROC curve

Figure 2

Performance of two distinct pipelines to identify co-infection by SARS-CoV-2 VOCs. Metrics for the whole-genome assembly, with AUC=0.5 and accuracy=46.2%, suggest no discrimination of the VOCs among samples, with a misclassification of all the cases with two variants. In contrast, the ASV-like inference was able to correctly identify VOCs in cases with one or two variants, with an outstanding performance according to the AUC=0.964 and accuracy=96.2%. FPR: False Positive Rate.

Figure 3

Comparison of the mutation profile of the SARS-CoV-2 VOCs. A few mutations are shared among the genotypes while most of them are exclusive for each genome, making possible to map and track genome subsequences to identify concomitant presence of VOCs.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Supplementaryfile.xlsx](#)