‘Surviving to Thriving’: A Meta-Ethnography of the Experiences of Healthcare Staff Caring for Persons With COVID-19

Frank Agyei

Presbyterian University College

Jonathan Bayuo (✉ jbayuo88@gmail.com)

Presbyterian University College

Prince Baffour

Komfo Anokye Teaching Hospital

Cletus Laari

Nsawam Government Hospital

Research Article

Keywords: Experiences, Healthcare professionals, Meta-ethnography

Posted Date: January 18th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-139341/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Abstract

Background: The emergence of the Coronavirus disease has heightened the experience of emotional burden among healthcare staff. To guide the development of support programmes, this review sought to aggregate and synthesise primary studies to establish a comparative understanding of the experiences of healthcare staff caring for persons with the disease.

Design: A meta-ethnography approach was used to aggregate and synthesise primary qualitative studies. Database search was undertaken from January to November 2020. A standardised tool was used to extract data from the identified primary studies. The studies were translated into each other to formulate overarching concepts/ metaphors which formed the basis of undertaking a narrative synthesis.

Results: Eight qualitative studies met the inclusion criteria. Two overarching concepts/ metaphors were formulated from the primary studies: 1) surviving to thriving in an evolving space and 2) support amid the new normal. The initial phase of entering the space of caring during the outbreak was filled with psychological chaos as healthcare staff struggled to survive within the context of an illness which was not fully understood. Gradually, healthcare staff may transition to a thriving phase characterised by resilience but still experienced heavy workload and physical/ emotional exhaustion. Fear persisted throughout their experiences. Fear of contracting the disease or infecting one's family members/ loved ones remained a key concern among healthcare staff despite infection precaution measures. Healthcare staff who contracted the disease felt isolated with additional fears of dying alone. The sources of support were varied with a strong emphasis on peer support.

Conclusions: Healthcare staff caring for persons infected with the Coronavirus disease require ongoing mental health support commensurate to their needs in the survival or thriving phase. Staff who contract the disease may require additional support to navigate through the illness and recovery. Concerted efforts are needed to strengthen support systems and build resilience among healthcare staff.

Background

The emergence of the novel Coronavirus disease (COVID-19) across the globe has led to various healthcare systems becoming overwhelmed with clinicians facing significant emotional strain and physical pressure (1, 2). Evidence from several countries suggest an increasing rate of depression, anxiety, and insomnia among healthcare workers caring for persons with COVID-19 (3–5). These emotional/ physical pressures, if left unresolved, may lead to a higher incidence of suicide and substance abuse among healthcare workers (6–9).

Caring for critically ill persons is often associated with emotional and physical exhaustion (10–13). The sudden occurrence of the COVID-19 pandemic, which healthcare systems were seemingly unprepared for alongside increasing mortality rates in some areas have contributed to the development of fear, worry and uncertainty (14). These concerns are likely to increase the burden experienced by healthcare staff creating the need for ongoing support (15). Various settings are implementing several programmes for healthcare staff but there appears to be an ever increasing need to provide ongoing evidence-based psychosocial support (16–19).

The pandemic is gradually becoming the 'new normal' implying that we may have to live with it for an unknown period (20–22). In the absence of adequate/ context-specific support programmes for our healthcare providers, their well-being may be adversely affected which can affect the overall availability of human resource and even

translate to poor patient care (23–25). So far, primary studies exploring the experiences of healthcare staff caring for patients with COVID-19 are emerging which offers some insight into their lived experiences. However, to gain a broader perspective and facilitate the design of interventions (timing and nature/ components), there is a need to establish a comparative understanding of these experiences. Thus, this review sought to identify the available primary studies, aggregate and synthesise their findings to understand the phenomenon of caring for persons diagnosed with COVID-19. The review question was “what are the experiences of healthcare professionals caring for persons with COVID-19?”

Aim

The aim of this review was to develop a comparative understanding of the experiences of healthcare staff caring for persons with COVID-19.

Methods

Review design

Noblit and Hare’s approach to meta-ethnography was utilised for this review (26). Meta-ethnography is an aggregative method of synthesis which seeks to integrate separate parts to form a whole. It involves induction and interpretation, thus resembling the primary studies it aims to synthesise (27). In simple terms, meta-ethnography is the qualitative alternative to quantitative meta-analysis (26). The product of a meta-ethnographic synthesis is the interpretation of the primary studies into one another to generate an in-depth/ new understanding of a phenomenon (28). This meta-ethnography was reported according to the eMERGe reporting guidelines (29). Additionally, the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) flowchart (30) was used to guide the process of study selection.

Search strategy/ study identification

A limited search in CINAHL, EMBASE and PubMed was initially undertaken which informed the development of a detailed search strategy. The full search sources included Cochrane Reviews Library, EMBASE, CINAHL, PubMed, OVID, Scopus, and Web of Science from December 2019 to November 2020. The following key words were used: “healthcare professionals” OR “healthcare practitioners” OR “healthcare staff” OR “healthcare workers” AND “COVID-19” OR “Coronavirus disease” OR “clinical respiratory illness” OR “clinical respiratory infection” OR “influenza-like illnesses to identify qualitative studies focusing on the phenomenon. The reference lists of identified articles were manually searched for potential studies.

Study selection and screening

Upon completing the search, all identified articles were exported into Endnote X9.2 and duplicates removed. This was followed by a selection procedure to identify primary studies for inclusion in the review. The inclusion criteria were 1) primary studies exploring the experiences of healthcare professionals caring for patients with COVID-19 2) qualitative studies irrespective of the design and 3) reported in English. Preprints, grey, non-qualitative studies, commentaries, editorials, and unpublished literature were not considered for inclusion in this review. Title screening was initially carried out to ensure that the study is relevant to the review. Abstract screening was then carried out. Full texts of the studies meeting the inclusion criteria were retrieved. These

studies proceeded to the critical appraisal stage before including them in the review. The results of the search are presented in the PRISMA flow diagram below as Fig. 1.

Quality assessment/ appraisal

Studies considered for inclusion were critically appraised using the Joanna Briggs Institute (JBI) standardized critical appraisal checklist for qualitative studies. Potential studies that received an overall appraisal as 'include' were retained in the review (see Table 1).

Table 1
Quality Appraisal

	Sun et al.,	Liu et al.,	Kackin et al.,	Karimi et al.,	Galehdar et al.,	Ardebili et al.,	Al Ghafri et al.,	Nyashanu et al.,
Is there congruity between the stated philosophical perspective and the research methodology?	Yes	Yes	Yes	Yes	Unclear	Unclear	Unclear	Yes
Is there congruity between the research methodology and the research question or objectives?	Yes	Yes	Yes	Yes	Unclear	Yes	Yes	Yes
Is there congruity between the research methodology and the methods used to collect data?	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Is there congruity between the research methodology and the representation and analysis of data?	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Is there congruity between the research methodology and the interpretation of results?	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Is there a statement locating the researcher culturally or theoretically?	Yes	Unclear	Yes	No	Unclear	Unclear	Unclear	Unclear
Is the influence of the researcher on the research, and vice- versa, addressed?	Unclear	Unclear	Yes	No	Unclear	Unclear	Unclear	Unclear

	Sun et al.,	Liu et al.,	Kackin et al.,	Karimi et al.,	Galehdar et al.,	Ardebili et al.,	Al Ghafri et al.,	Nyashanu et al.,
Are participants, and their voices, adequately represented?	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Is the research ethical according to current criteria or, for recent studies, and is there evidence of ethical approval by an appropriate body?	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Do the conclusions drawn in the research report flow from the analysis, or interpretation, of the data?	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Overall appraisal	Include	Include	Include	Include	Include	Include	Include	Include

Data Extraction And Synthesis

Data extracted from selected studies included standard information such as authors, setting, study findings and verbatim from participants reported (see Table 2). To synthesise the data, codes in the form of first and second order constructs were formulated from each primary study. These codes were organized into categories through a process of constant comparison across the studies (31). This facilitated either translation of the studies into one another (areas of agreement across the studies) or refutational synthesis (areas of disagreement across the studies) based on the emerging categories. Following the translation process, the categories were re-interpreted to formulate overarching concepts/ metaphors. These overarching concepts/ metaphors formed the basis for undertaking a narrative synthesis (31). The interpretive process was iterative with reference to the primary studies.

Table 2: Data Extraction

Author/ year/ aim	Design/ methodology	Key findings	Sample quotes	Codes	Key/ overarching concepts/ metaphors
Sun et al., (2020) To explore the psychological experiences of nurses caring for COVID-19 patients.	Descriptive phenomenology	Four themes emerged:	"After putting on protective clothing, nursing duties are awkward to carry out. Protective clothing needs to be worn for 8 hours or more without drinking water and eating food and urinating was done with adult diapers."	Initial psychological chaos; coping and thriving; support	1. Surviving to thriving in an evolving space
	Purposeful sampling approach to recruit 20 nurses caring for patients with COVID-19 in Henan, China	<p>· Significant number of negative emotions in the early stage: this theme describes participants' experiences with fatigue, discomfort and helplessness caused by the nature of caring for the patients and wearing the protective clothing. Also, there was fear of contracting the virus, anxiety caused by limited understanding of managing the infection/ presence of strangers as well as concerns about their family members</p>			
	Face to face and telephone interviews conducted		"...The moment I walked through the door of the Department of Infectious Diseases, I felt very scared. I felt much better after I got used to it. And I felt scared when I pushed the door of the negative pressure room for the first time, but I was fine the second time."		
	Data analysis by Colaizzi's 7-step method				
	Critical appraisal: Include	· Coping and self-care styles: this theme describes the psychological/ life adjustment processes to face the situation and seeking support from one's professional group.	"My method is not to think about stress, I shield it out of my life." "...I forget everything when I am busy..." "We encourage each other. It does not feel like I'm fighting alone, I'm not afraid."		2. Support amid the new normal
		· Growth	"I used to work to		

under pressure: this theme describes the personal growth experienced by participants which include the opportunity for self-reflection, gaining a deeper sense of professional identity and increased affection and feelings of gratefulness.

earn a salary, but now it feels like a responsibility."

"Maybe there was a discrimination against nurses in the society but now I am proud of my choice."

"After work I find the sky is blue and everything is beautiful."

"I never thought I could be so strong."

Positive emotions occurring simultaneously or progressively with negative emotions: the theme describes the positive emotions that emerged in the process. These included feelings of confidence in the hospital environment, happiness from multiple sources of support and calmness during the care delivery process.

"I feel that the government has strong prevention and control measures, and the epidemic will be controlled very soon. But after all, we have a large population, and it is a process."

"Patients are very cooperative with our work. Although some patients have emotions due to illness, they show great respect to us."

"My mood is much better after starting pre-job training."

"Many colleagues called me to encourage me and I felt that

			there were many people who cared about me."	Sense of responsibility to care; psychological chaos; support
Liu et al., (2020)	Descriptive phenomenology	Three themes with 10 subthemes emerged:	"We must try our best to win this battle. As health-care providers, we are at the forefront. I fight for my family, and I fight more for this society."	
To describe the experiences of nurses and doctors in the early stages of the COVID-19 outbreak	<p>Purposive and snowball sampling was employed to recruit 9 nurses and 4 physicians in Hubei Province, China</p> <p>Data collection by in-depth telephone interviews</p> <p>Haase's adaptation of Colaizzi's method was used to analyse transcripts</p> <p>Critical appraisal: Include</p>	<p>Being fully responsible for patients' wellbeing—"this is my duty": this theme describes a sense of duty expressed by participants. This included a feeling of being called to duty, caring for the affected persons, and emotionally supporting the patients.</p>	<p>"This is my duty because I am a medical worker. No matter what will happen"</p> <p>"Patients are struggling to breathe, and some can only lie in bed. They are very helpless and want care from their families."</p>	
		<p>Challenges associated with working on the COVID-19 wards: working in these wards was a completely new contexts with participants experiencing exhaustion and being overwhelmed with the workload, dealing with uncertainty and fear of contracting the virus/ infecting others, being witnesses of</p>	<p>"I was very tired. I had to lie in bed for a whole day to recover from the fatigue after work."</p> <p>"I felt very depressed on the first day in the infectious disease hospital because there was only one entrance and passage for medical staff, and it is a real isolation unit with negative pressure. I felt it was difficult to</p>	

the patients' experiences, and dealing with the healthcare provider-patient relationship amid the chaos.

breathe... This new environment brought a sense of oppression."

"I have to treat many patients who are not in my specialty. Although the country has released six editions of diagnosis and treatment guidelines [for COVID-19], there is still no effective antiviral medicine. It is an unknown disease, and everyone feels powerless."

"I recently contacted a colleague without any protection,

who was later diagnosed with COVID-19. Although my CT results did not show any abnormality, I am anxious and waiting to do the throat swab."

Many sources of social support to cope with the situation and transcendence: This theme reflects participants' sources of support including family, friends, colleagues, and the society.

"I am not overstrained because I trust our hospital.

Our hospital gives us strong logistical support, including

providing medical protective supplies, accommodations,

transportation, food, medicines, and subsidies."

"The head nurse knows we come from different

departments and infectious disease is not our specialty,

so she sent us some educational videos and materials, and we can learn after work."

"When I feel stressful, I complain to my boyfriend. He is also a nurse, and we are in the same department. We communicate with and understand each other."

Kackin et al., (2020)	Descriptive phenomenology	Three themes and ten subthemes emerged:	"Nurses I have never known or seen. They were assigned to our service unit from another one. I don't know their reactions . . . we had a dispute the other day with another Nurse . . . It feels as if working in another hospital. Different patients, a different order"	Initial sense of psychological chaos; needing ongoing psychological support; coping
To determine the experiences and psychosocial problems of nurses caring for patients diagnosed with COVID-19 in Turkey.	Purposive sampling was employed to recruit 10 nurses caring for patients diagnosed with COVID-19 in Istanbul, Turkey	· Effects of the outbreak: this theme highlights the working conditions, psychological and social effects of the outbreak. The nurses were faced with lack of equipment/ worsening work conditions, stress, feeling threatened, uncertain, depression, fear, aggression, social isolation (spending	"There is a patient lying there, you know that the patient needs you, but wearing that protective equipment, feeling his/her physical pain in	
	Data collection was conducted via questionnaires and semi-structured interviews			
	Colaizzi's method was used to			

	analyse transcripts	more time in the hospital) and stigma.	your own body, you may have to work for an hour at most once you wear the helmet. It gives you a headache. You cannot enter the isolation rooms without those garments, and those garments are extremely smothering you. Sometimes, leaving the room when we admit new patients can take 2.5-3 hours without exaggeration. When we leave, you find yourself in full of sweat “	
	Critical appraisal: Include	<ul style="list-style-type: none"> Short-term coping strategies: Participants emotions normalized (accepting the situation and thinking it is a temporal issue), refusal to dwell on the experiences, avoidance (avoiding the media/ comments about the disease), openly expressing their feelings (crying etc) and distraction (being thankful, listening to music, sports etc.) Needs: nurses required ongoing psychological support and increasing resource availability at the setting. 	<p>“I left my family alone . . . My mother suffers from high blood pressure, what happens if she becomes infected . . . there is the fear of losing her . . .”</p>	
Karimi et al., 2020	Descriptive phenomenology	Three themes and six subthemes emerged:	“Maybe I die, but I still have lots of dreams.”	Initial psychological stress; navigating an unknown disease; dealing with resource limitations
To explore the lived experiences of nurses caring for patients with COVID-19 in Iran.	<p>Purposive sampling was employed to recruit 12 nurses caring for patients diagnosed with COVID-19 in Iran</p> <p>Data collection was conducted via semi-</p>	<ul style="list-style-type: none"> Mental condition: The theme describes the psychological responses of healthcare providers which include stress, anxiety, and fear. These were related to fear of the disease, being worried 	<p>“I’m scared for my family and also for myself.”</p> <div> <p>“How horrible these days are, we’re all dying.”</p> </div>	

	structured interviews	about their families.		
	Colaizzi's method was used to analyse transcripts	· Emotional condition: the theme highlights feelings of suffering and affliction such as uncertainty about the disease process, witnessing death/ dying and separation from their own family		
	Critical appraisal: Include	· Care context: the theme highlights the turmoil and limited availability of support and equipment. Increasing work pressure, staff shortage, chaos and inexperience in handling the chaos		
Galehdar et al., 2020	Qualitative design	Eleven categories and 5 subcategories emerged:	"It is agonizing to see a person deprived of breath, his heart failing, and you can't do anything about his suffering it sometimes causes me to feel agitated and distressed and becoming really sad and confused about what I'm going to do?"	Psychological issues and navigating the death of patients
To explore nurses' experiences of psychological distress during care of patients with COVID-19.	Purposive sampling was employed to recruit 20 nurses caring for patients diagnosed with COVID-19 in Iran	· Death anxiety: the nurses experienced psychological distress witnessing the deaths of patients with feelings of helplessness when they could not do anything to alleviate the patient's symptoms; concerns regarding the high mortality rates	"I myself was caring for a patient with COVID-19, it was really painful to see a person striving to breathe to save himself "	
	Data collection was conducted via semi-structured telephone interviews			
	Conventional content analysis			

	<p>was used to analyse transcripts</p> <p>Critical appraisal: Include</p>	<ul style="list-style-type: none"> Anxiety due to the nature/ severity of the illness, rate of spread and unknown dimensions of the disease 	<p>"In my opinion, the nature of the disease is beyond what we are teaching and learning now"</p>	
<p>Ardebili et al., 2020</p> <p>To undertake an in-depth exploration of the experiences of health-care staff working during the COVID-19 crisis.</p>	<p>Qualitative design</p> <p>Purposive sampling was employed to recruit 97 healthcare professionals (pre-hospital emergency services (EMS), physicians, nurses, pharmacists, laboratory personnel, radiology technicians, hospital managers and managers in the ministry of health who work directly or indirectly with COVID-19 cases) caring for patients diagnosed with COVID-19 in Iran</p> <p>Data collection was conducted via semi-structured interviews</p> <p>Thematic analysis was used to analyse transcripts</p>	<p>Three themes and eleven subthemes emerged:</p> <ul style="list-style-type: none"> Working in the pandemic era: This was experienced as high workload and feelings of losing control over the situation, fear, anxiety, and being overwhelmed (Providing futile care) Changes in personal life and enhanced negative effects Gaining experience, normalization and adapting to the pandemic (overcoming the initial crisis, gaining experience regarding patient management, reducing referrals and increasing recoveries). Mental health issues: Experiences of loss of control, heavy workload, 	<p><i>"In the early days, our workload was very high, we had to move the wards and hospitalized corona patients in the non-infectious wards"</i></p> <p><i>"Every day a new drug is introduced, every day a new route of transmission is introduced"</i></p> <p><i>"This disease does not have a specific drug, nor can you predict with confidence who will survive and who will die. This made me feel (completely ineffectual and I felt) like I was losing control"</i></p> <p><i>"It's very difficult to wear N95 masks for twelve hours, I feel short of breath and I will definitely have problems later (Nurse)"</i></p> <p><i>They give (you) a body suit in each shift. When we wear these clothes, sweat</i></p>	<p>Initial psychological distress; facing personal changes; adjusting to the situation</p>

	Critical appraisal: Include	severe stress, the experience of a sense of futile care, fear of infection and transmission, self-isolation, and quarantine, decreased emotional relationships, fundamental changes in lifestyle, worrying about the future and the economic situation, all appeared to contribute towards the manifestation of mental health issues).	<p><i>flows from all over our bodies, we can't eat anything with these clothes, we can't drink anything too, we have to wear them for 12 hours"</i></p> <p><i>"When I was hospitalized in the ICU, I had very severe shortness of breath. When the shortness of breath was present, I thought I was dying (Nurse) I was thinking, I will die alone, without seeing my family, they will not see my body. I will not have a proper funeral"</i></p>	
<p>Al Ghafri et al., 2020</p> <p>To explore the experiences and perceptions of health care workers (HCWs) in primary health care in the management of COVID-19 with respect to medical response experiences, socio-cultural and religious reforms, psychological impressions, and lessons learned.</p>	<p>Phenomenology</p> <p>Purposive and snowball sampling was employed to recruit 40 healthcare professionals/stakeholders involved in managing patients diagnosed with COVID-19 in Oman</p> <p>Data collection was conducted via focus group discussions (6 focus group discussions conducted)</p> <p>Thematic analysis</p>	<p>Three themes emerged:</p> <ul style="list-style-type: none"> Medical response experiences: rapid restructuring of public health services, enforcing technology use and increasing burden on limited human resource available. Socio-cultural and religious reforms: having to stay away from parents and families (inability to participate in social/religious practices); empathy towards the 	<p>"we had to work for more than 12 hours continuously due to shortages of staff. This was an overburden to us"</p> <p>" "when our colleagues got infected, we all suffered physically and emotionally"</p>	Psychological concerns; adjusting the healthcare system

		vulnerable in the society		
	Critical appraisal: Include	<ul style="list-style-type: none"> Psychological issues: being at home was described as depressing, and inability to travel around were distressing; exhaustion among healthcare professionals and fear of transmitting the virus to families/ loved ones. 		
<p>Nyashanu et al., 2020</p> <p>To explore the triggers of mental health problems among frontline healthcare workers during the COVID-19 pandemic.</p>	<p>Exploratory qualitative study</p> <p>Purposive sampling was employed to recruit 40 frontline staff involved in private care homes and domiciliary care agencies in the Midlands, UK</p> <p>Data collection was conducted via semi-structured interviews</p> <p>Interpretive phenomenological analysis</p> <p>Critical appraisal: Include</p>	<p>Seven themes noted in the study:</p> <ul style="list-style-type: none"> Fear of infection and infecting others Lack of recognition Lack of guidance/ frequently changing guidelines creating doubts about operational procedures and triggering anxiety Unsafe hospital discharge Loss of professionals/ residents through death Unreliable testing and delayed results Staff shortage causing 	<p>"Unfortunately, there have been so many changes on the guidance to COVID-19. Being diabetic the government has placed responsibility on my employer to make suitable safe working arrangements which is difficult. A female learning disability nurse I am really worried with ever changing information from government on how to act during this pandemic.... Honestly, it really makes me anxious"</p> <p>"We have been using agency staff to maintain staff numbers, but we don't know where else they have been working and this brings so much anxiety."</p>	<p>Fear, anxiety, worry; lack of guidelines; limited human resources</p>

Findings

Study characteristics

Following the screening process, eight primary qualitative studies met the criteria for inclusion in this review (32-39). Although all the studies involved healthcare staff caring for patients with COVID-19, majority focused on nurses (see Table 1). The settings of the primary studies include Mainland China (37, 38), Iran (33, 34, 36), Turkey (35), Oman (32) and United Kingdom (39). Five studies utilised a phenomenological approach to uncover participants' lived experiences (32, 35-38).

Concepts/metaphors

The interpretation of the data and translation of the studies into each other led to the emergence of two overarching concepts/ metaphors: surviving to thriving in an evolving space and support amid the new normal. The relationship between the emerging concepts/ metaphors was noted to be reciprocal which facilitated the development of a line of argument to understand the phenomenon of caring for patients with COVID-19.

Surviving to thriving in an evolving space

Initial psychological/emotional chaos

The COVID-19 emerged as an infection to which healthcare staff initially had limited knowledge. Thus, being asked to work on a ward for persons with the infection created an initial sense of inner tension/ psychological chaos and an 'internal' struggle to survive in an uncertain dimension of a rapidly evolving disease (32-39). Psychological responses such as anxiety, helplessness, fear of contracting the infection and spreading to one's loved ones, and uncertainty characterised the initial survival space and trickled to the thriving phase (32-39):

"Although I volunteered to work in the Department of Infectious Diseases, I still feel very scared. After all, it is a new infectious disease and there are no specific drugs at present. I was scared to see reports of the sacrifice of medical staff in other cities." (38)

"...we are fearful of being infected. Anyone who coughs in the office causes panic. If one is infected, all medics in the unit are in danger, then the unit will be paralysed... I recently contacted a colleague without any protection, who was later diagnosed with COVID-19. Although my CT results did not show any abnormality, I am anxious and waiting to do the throat swab." (37)

"I'm not calm at all, and I do not know what's going on" (36)

The fears of some healthcare staff came to fruition as they contracted the COVID-19 disease. This led to feelings of social isolation as they received treatment and hanging in a balance as they navigated through the symptoms on their own. Within the space of contracting the disease, affected healthcare staff were faced with new fears regarding dying alone with their mortal remains not receiving the final respect required (32-36).

"When I was hospitalized in the ICU, I had very severe shortness of breath. When the shortness of breath was present, I thought I was dying (Nurse) I was thinking, I will die alone, without seeing my family, they will not see my body. I will not have a proper funeral" (33)

The initial psychological chaos experienced by the healthcare staff heightened as they witnessed varying mortality rates (33, 34, 36-38). The most challenging aspect for healthcare staff appeared to be contracting the illness themselves or witnessing the death of a colleague following a diagnosis of COVID-19 (32, 33, 39):

"When our colleagues got infected, we all suffered physically and emotionally" (32)

"It is agonizing to see a person deprived of breath, his heart failing, and you can't do anything about his suffering it sometimes causes me to feel agitated and distressed and becoming really sad and confused about what I'm going to do?" (34)

Living and functioning in a 'new body'

The survival phase was also characterised by struggling to live and function in a 'new body', that is the personal protective equipment (PPE) which appeared to be uncomfortable, yet indispensable (32, 33, 37, 38). These concerns notwithstanding, healthcare professionals felt a sense of responsibility to fight the illness, care for the persons diagnosed with COVID-19, protect themselves and their loved ones from contracting the "deadly virus" (32-39):

"After putting on protective clothing, nursing duties are awkward to carry out. Protective clothing needs to be worn for 8 hours or more without drinking water and eating food and urinating was done with adult diapers." (38)

"Wearing the whole set of PPEs is very uncomfortable. I have difficulty breathing and feel very hot and my heart rate speeds up. We keep on sweating and the clothes are soaked." (37)

"It's very difficult to wear N95 masks for twelve hours, I feel short of breath and I will definitely have problems later" (33)

"We must try our best to win this battle. As health-care providers, we are at the forefront. I fight for my family, and I fight more for this society. This is my duty because I am a medical worker. No matter what will happen" (32)

Thriving amidst chaos

As healthcare staff continued to navigate through the evolving space of care provision, received training, and identified new strategies of survival, there was a gradual move from survival to thriving which was characterised by resilience (32-38). In the thriving phase, participants did not still understand the nature of the infection fully but felt more at ease working with the affected persons. The thriving phase was also characterised by personal

and professional growth in the face of adversity with a feeling of being in a supportive environment, although resources were still limited (32-38). Further within the thriving phase, healthcare staff began to appraise the negative experiences in a positive manner as a means of coping within a context that was not fully understood (32-37, 40). Irrespective of the phase healthcare staff found themselves, they were still faced with fear, increasing workload which led to exhaustion as they navigated through patient care and their own experiences (32-39):

"My method is not to think about stress, I shield it out of my life...I forget everything when I am busy..." (32)

"In the early days, our workload was very high, we had to move the wards and hospitalized corona patients in the non-infectious wards" (33)

Support amid the 'new normal'

Support systems

The need for ongoing psychological support to help manage the 'self' was highlighted by all studies as healthcare staff navigated through survival to thriving (32-38). Support from other members of the healthcare team was considered essential as the disease was considered a common 'enemy' among staff (32-38). Beyond the confines of the hospital, some healthcare staff also received support from families and friends (37, 38). Aside managing the 'self', healthcare staff also required support in utilizing the limited resources, ongoing training to stay updated about the disease and how best to protect oneself and family (32-38):

"When I feel stressful, I complain to my boyfriend. He is also a nurse, and we are in the same department. We communicate with and understand each other." (36)

"The head nurse knows we come from different departments and infectious disease is not our specialty, so she sent us some educational videos and materials, and we can learn after work." (37)

Clinical guidelines

Rapidly changing guidelines were challenging for healthcare staff and they required more support regarding operational procedures (39):

"Unfortunately, there have been so many changes on the guidance to COVID-19. Being diabetic the government has placed responsibility on my employer to make suitable safe working arrangements which is difficult. A female learning disability nurse I am really worried with ever changing information from government on how to act during this pandemic.... Honestly, it really makes me anxious" (39).

Discussion

The review sought to gain a comparative understanding of the experiences of healthcare staff in caring for persons with the novel COVID-19 disease. The findings highlight an initial sense of psychological chaos with

healthcare staff struggling to survive as they navigated through the outbreak. Overtime, healthcare staff transitioned from survival to thriving as they continued to provide care but still experienced fear of contracting the disease and transmitting to family members/ loved ones. Some healthcare staff who contracted the disease also experienced fear of dying alone. The impact of these psychological experiences emphasise the need for early and ongoing psychosocial support as well as maintaining high standards of infection prevention and control measures to make healthcare staff feel safe. Continuing professional education on emerging trends of the disease, ensuring the availability and utilisation of safety materials, promoting team morale and providing avenues of release for healthcare professionals are also needed to support healthcare staff caring for persons with COVID-19.

The initial phase of working with persons infected with the novel virus is a critical period of transitioning to an unknown context with emotional responses heightening around a week of entering this unfamiliar space (19, 32–39). In a previous study among Korean nurses during the era of the Middle East Respiratory Syndrome (MERS), the unfamiliar space of the infection was described as a dangerous field filled with psychological and physical stressors (41). As previously noted, healthcare staff who work with critically ill persons already face several stressors. Extrapolating these stressors and experiences to the context of an unknown illness suggest the existence of a significant psychological challenge. Without professional support, healthcare staff are at a risk of several mental health issues such as depression, insomnia, anxiety, mood and eating disorders, in both short and long term (1, 42). These findings should therefore direct our attention towards early mental health intervention to identify, acknowledge and offer support commensurate to the needs of healthcare staff (19).

Further to the above, fears about contracting the COVID-19 disease and/ or infecting one's family members/ loved ones emerged as a major source of stress among healthcare staff. Even when the initial experience of psychological chaos was low with existing high standards of infection prevention strategies, fear of contracting the disease remained a significant concern among healthcare staff (43–49). Healthcare staff who contracted the disease were faced with additional fears of dying alone with feelings of being socially isolated from colleagues and family/ loved ones as they underwent treatment. The findings strengthen the need to make clinicians well-being a priority across healthcare settings and countries (50). Healthcare staff need to feel safe within the healthcare setting whilst ensuring adherence to best infection prevention and control practices (15). Avenues for healthcare staff to express their fears are needed to help them navigate through their emotions (51). Additionally, healthcare staff who contract the illness may require extra support to deal with the psychological/ emotional impact of the illness.

Despite the emergence of several stressors, some facilitators to surviving/ thriving emerged. Key among these facilitators is the support offered by peers within one's team and family. Previous studies have highlighted the significant role played by peers at the workplace as there seem to be a shared concern among these persons who are journeying together within unfamiliar territories (41, 52). In fact, lack of social support has been linked to the development of anxiety, insomnia, and depression (49, 53, 54). Peer support develops overtime and creates a sense of connectedness which may be difficult to quantify. This form of unique support requires further attention to determine ways of facilitating their development and improvement particularly in this era of journeying through a common ground.

Although the review findings offer insights into the experiences of healthcare professionals caring for persons with COVID-19, a limitation is noteworthy. Majority of the participants in the primary studies were nurses which

creates the need to engage other healthcare staff such as laboratory technicians and mortuary attendants to understand their experiences.

Conclusion

Navigating through the experiences of healthcare staff during the outbreak suggest the existence of several concerns warranting attention although some form of professional/ personal growth may be experienced. The findings imply a need for early and ongoing mental health support, support in handling fears, ensuring the availability of required equipment and identify strategies to boost team morale.

Declarations

Ethical approval: not applicable

Consent for publication: Not applicable

Availability of data and materials: The datasets used and/ or analyzed during the current study are included in this published article.

Competing interests: All authors declare no competing interests

Funding: The study received no funding

Authors' contributions: JB and FBA conceptualised the study; all authors were involved in study search, appraisal, analysis, and synthesis.

Acknowledgement: The authors wish to thank the staff of the hospital for their support in the conduct of the study.

References

1. Shen X, Zou X, Zhong X, Yan J, Li L. Psychological stress of ICU nurses in the time of COVID-19. Springer; 2020.
2. Galbraith N, Boyda D, McFeeters D, Hassan T. The mental health of doctors during the COVID-19 pandemic. *BJPsych bulletin*. 2020;1–4.
3. Lai J, Ma S, Wang Y, Cai Z, Hu J, Wei N, et al. Factors associated with mental health outcomes among health care workers exposed to coronavirus disease 2019. *JAMA network open*. 2020;3(3):e203976-e.
4. Rossi R, Socci V, Pacitti F, Di Lorenzo G, Di Marco A, Siracusano A, et al. Mental health outcomes among front and second line health workers associated with the COVID-19 pandemic in Italy. *medRxiv*. 2020.
5. Ruiz MA, Gibson C-AM. Emotional impact of the COVID-19 pandemic on US health care workers: A gathering storm. *Psychological Trauma: Theory, Research, Practice, and Policy*. 2020;12(S1):S153.
6. Montemurro N. The emotional impact of COVID-19: From medical staff to common people. *Brain, behavior, and immunity*. 2020.

7. Gold JA. Covid-19: adverse mental health outcomes for healthcare workers. British Medical Journal Publishing Group; 2020.
8. Braquehais MD, Vargas-Cáceres S, Gómez-Durán E, Nieva G, Valero S, Casas M, et al. The impact of the COVID-19 pandemic on the mental health of healthcare professionals. *QJM: An International Journal of Medicine*. 2020;113(9):613–7.
9. Cullen W, Gulati G, Kelly B. Mental health in the Covid-19 pandemic. *QJM: An International Journal of Medicine*. 2020;113(5):311–2.
10. Bayuo J, Agbenorku P. Coping strategies among nurses in the Burn Intensive Care Unit: A qualitative study. *Burns Open*. 2018;2(1):47–52.
11. Kisorio LC, Langley GC. Critically ill patients' experiences of nursing care in the intensive care unit. *Nursing in critical care*. 2019;24(6):392–8.
12. Limbu S, Kongsuwan W, Yodchai K. Lived experiences of intensive care nurses in caring for critically ill patients. *Nursing in critical care*. 2019;24(1):9–14.
13. Query LA, Olson KR, Meyer MT, Drendel AL. Minding the gap: A qualitative study of provider experience to optimize care for critically ill children in general emergency departments. *Academic Emergency Medicine*. 2019;26(7):803–13.
14. Shanafelt T, Ripp J, Trockel M. Understanding and addressing sources of anxiety among health care professionals during the COVID-19 pandemic. *Jama*. 2020;323(21):2133–4.
15. Adams JG, Walls RM. Supporting the health care workforce during the COVID-19 global epidemic. *Jama*. 2020;323(15):1439–40.
16. Dewey C, Hingle S, Goelz E, Linzer M. Supporting clinicians during the COVID-19 pandemic. *American College of Physicians*; 2020.
17. Hopman J, Allegranzi B, Mehtar S. Managing COVID-19 in low-and middle-income countries. *Jama*. 2020;323(16):1549–50.
18. Tracy DK, Tarn M, Eldridge R, Cooke J, Calder JD, Greenberg N. What should be done to support the mental health of healthcare staff treating COVID-19 patients? *The British Journal of Psychiatry*. 2020;217(4):537–9.
19. Tomlin J, Dalglish-Warburton B, Lamph G. Psychosocial support for healthcare workers during the COVID-19 pandemic. *Frontiers in Psychology*. 2020;11:1960.
20. Bowdle A, Munoz-Price LS. Preventing infection of patients and healthcare workers should be the new normal in the era of novel coronavirus epidemics. *Anesthesiology: The Journal of the American Society of Anesthesiologists*. 2020;132(6):1292–5.
21. Lee TH. Creating the new normal: The clinician response to COVID-19. *NEJM Catalyst Innovations in Care Delivery*. 2020;1(2).
22. Luker GD, Boettcher AN. Transitioning to a new normal after COVID-19: preparing to get back on track for cancer imaging. *Radiological Society of North America*; 2020.
23. Bai Y, Lin C-C, Lin C-Y, Chen J-Y, Chue C-M, Chou P. Survey of stress reactions among health care workers involved with the SARS outbreak. *Psychiatric Services*. 2004;55(9):1055–7.
24. Brooks SK, Webster RK, Smith LE, Woodland L, Wessely S, Greenberg N, et al. The psychological impact of quarantine and how to reduce it: rapid review of the evidence. *The Lancet*. 2020.

25. Marjanovic Z, Greenglass ER, Coffey S. The relevance of psychosocial variables and working conditions in predicting nurses' coping strategies during the SARS crisis: an online questionnaire survey. *International journal of nursing studies*. 2007;44(6):991–8.
26. Noblit GW, Hare RD. *Meta-ethnography: Synthesizing qualitative studies*: sage; 1988.
27. France EF, Wells M, Lang H, Williams B. Why, when and how to update a meta-ethnography qualitative synthesis. *Systematic reviews*. 2016;5(1):44.
28. France EF, Cunningham M, Ring N, Uny I, Duncan EA, Jepson RG, et al. Improving reporting of meta-ethnography: the eMERGe reporting guidance. *BMC medical research methodology*. 2019;19(1):25.
29. Higgins JP, Thomas J, Chandler J, Cumpston M, Li T, Page MJ, et al. *Cochrane handbook for systematic reviews of interventions*: John Wiley & Sons; 2019.
30. Campbell R, Pound P, Pope C, Britten N, Pill R, Morgan M, et al. Evaluating meta-ethnography: a synthesis of qualitative research on lay experiences of diabetes and diabetes care. *Social science & medicine*. 2003;56(4):671–84.
31. Al Ghafri T, Al Ajmi F, Anwar H, Al Balushi L, Al Balushi Z, Al Fahdi F, et al. The Experiences and Perceptions of Health-Care Workers During the COVID-19 Pandemic in Muscat, Oman: A Qualitative Study. *Journal of primary care & community health*. 2020;11:2150132720967514.
32. Ardebili ME, Naserbakht M, Bernstein C, Alazmani-Noodeh F, Hakimi H, Ranjbar H. Healthcare providers experience of working during the COVID-19 pandemic: A qualitative study. *American journal of infection control*. 2020.
33. Galehdar N, Kamran A, Toulabi T, Heydari H. Exploring nurses' experiences of psychological distress during care of patients with COVID-19: a qualitative study. *BMC psychiatry*. 2020;20(1):1–9.
34. Kackin O, Ciydem E, Aci OS, Kutlu FY. Experiences and psychosocial problems of nurses caring for patients diagnosed with COVID-19 in Turkey: A qualitative study. *International Journal of Social Psychiatry*. 2020:0020764020942788.
35. Karimi Z, Fereidouni Z, Behnammoghadam M, Alimohammadi N, Mousavizadeh A, Salehi T, et al. The lived experience of nurses caring for patients with COVID-19 in Iran: a phenomenological study. *Risk Management and Healthcare Policy*. 2020;13:1271.
36. Liu Q, Luo D, Haase JE, Guo Q, Wang XQ, Liu S, et al. The experiences of health-care providers during the COVID-19 crisis in China: a qualitative study. *The Lancet Global Health*. 2020.
37. Sun N, Wei L, Shi S, Jiao D, Song R, Ma L, et al. A qualitative study on the psychological experience of caregivers of COVID-19 patients. *American journal of infection control*. 2020;48(6):592–8.
38. Nyashanu M, Pfende F, Ekpenyong MS. Triggers of mental health problems among frontline healthcare workers during the COVID-19 pandemic in private care homes and domiciliary care agencies: Lived experiences of care workers in the Midlands region, UK. *Health & social care in the community*. 2020.
39. Sun N, Shi S, Jiao D, Song R, Ma L, Wang H, et al. A qualitative study on the psychological experience of caregivers of COVID-19 patients. *American Journal of Infection Control*. 2020.
40. Kim Y. Nurses' experiences of care for patients with Middle East respiratory syndrome-coronavirus in South Korea. *American journal of infection control*. 2018;46(7):781–7.
41. Pappa S, Ntella V, Giannakas T, Giannakoulis VG, Papoutsis E, Katsaounou P. Prevalence of depression, anxiety, and insomnia among healthcare workers during the COVID-19 pandemic: A systematic review and

- meta-analysis. *Brain, behavior, and immunity*. 2020.
42. Karnatovskaia LV, Johnson MM, Varga K, Highfield JA, Wolfrom BD, Philbrick KL, et al., editors. *Stress and fear: clinical implications for providers and patients (in the Time of COVID-19 and Beyond)*. Mayo Clinic Proceedings; 2020: Elsevier.
 43. Khattak SR, Saeed I, Rehman SU, Fayaz M. Impact of fear of COVID-19 pandemic on the mental health of nurses in Pakistan. *Journal of Loss and Trauma*. 2020:1–15.
 44. Labrague LJ, de Los Santos JAA. Fear of Covid-19, psychological distress, work satisfaction and turnover intention among frontline nurses. *Journal of nursing management*. 2020.
 45. Wang H, Liu Y, Hu K, Zhang M, Du M, Huang H, et al. Healthcare workers' stress when caring for COVID-19 patients: An altruistic perspective. *Nursing ethics*. 2020;27(7):1490–500.
 46. Saleem Z, Majeed MM, Rafique S, Siqqiqui Z, Ghandhi D, Tariq H, et al. COVID-19 pandemic fear and anxiety among healthcare professionals in Pakistan. 2020.
 47. Ahmed MA, Jouhar R, Ahmed N, Adnan S, Aftab M, Zafar MS, et al. Fear and practice modifications among dentists to combat Novel Coronavirus Disease (COVID-19) outbreak. *International journal of environmental research and public health*. 2020;17(8):2821.
 48. Han L, Wong FKY, She DLM, Li SY, Yang YF, Jiang MY, et al. Anxiety and depression of nurses in a north west province in china during the period of novel coronavirus pneumonia outbreak. *Journal of Nursing Scholarship*. 2020;52(5):564–73.
 49. Chirico F, Nucera G, Magnavita N. COVID-19: Protecting Healthcare Workers is a priority. *Infection Control & Hospital Epidemiology*. 2020:1-.
 50. Fofana NK, Latif F, Sarfraz S, Bashir MF, Komal B. Fear and agony of the pandemic leading to stress and mental illness: an emerging crisis in the novel coronavirus (COVID-19) outbreak. *Psychiatry Research*. 2020;291:113230.
 51. Bayuo J, Agyei Bediako F, Allotey G, Kyei Baffour P. Developing support strategies for burn care nurses through an understanding of their experiences: A meta-ethnographic study. *International journal of nursing practice*. 2019;25(2):e12685.
 52. Spoorthy MS, Pratapa SK, Mahant S. Mental health problems faced by healthcare workers due to the COVID-19 pandemic–A review. *Asian journal of psychiatry*. 2020;51:102119.
 53. Wańkowicz P, Szylińska A, Rotter I. Assessment of mental health factors among health professionals depending on their contact with COVID-19 patients. *International Journal of Environmental Research and Public Health*. 2020;17(16):5849.
 54. Wańkowicz P, Szylińska A, Rotter I. Assessment of mental health factors among health professionals depending on their contact with COVID-19 patients. *International Journal of Environmental Research and Public Health*. 2020;17(16):5849.

Figures

Figure 1

PRISMA Flowchart of Study Selection