Birth and Regulation of head Extension to Guide Manual perineal Assistance (BREGMA) study

Abdelrahman Abouelhassan (✉ abdelrahmanmohamed@med.asu.edu.eg)

Charles University Faculty of Medicine in Pilsen: Univerzita Karlova Lekarska fakulta v Plzni
<https://orcid.org/0000-0001-9366-1307>

Vladimir Kalis

Charles University Faculty of Medicine in Pilsen: Univerzita Karlova Lekarska fakulta v Plzni

Marc Schüssler

Agaplesion Allgemeines Krankenhaus Hagen gGmbH

Hassan Awwad

Ain Shams University Faculty of Medicine

Rania Hassan

Ain Shams University Faculty of Medicine

Khaled M Ismail

Charles University Faculty of Medicine in Pilsen: Univerzita Karlova Lekarska fakulta v Plzni
<https://orcid.org/0000-0001-9449-0706>

Research Article

Keywords: manual perineal protection, perineal mapping, perineal body, perineal tension

Posted Date: March 1st, 2022

DOI: <https://doi.org/10.21203/rs.3.rs-1395708/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Introduction and Hypothesis: The exact positioning of the dominant hand at the time of manual perineal protection is not clearly specified. The overarching aim of this work was to collect data on the bregma to posterior fourchette distance (BFD), as a surrogate measure for the degree of fetal head extension, and maternal perineal body length (PBL) at the time of manual perineal protection (MPP).

Methods: This was a two-center study. Term cephalic singleton nulliparous women having spontaneous vaginal delivery were considered eligible for recruitment into the study. Once crowning was diagnosed and just prior the initiation of manual perineal protection (MPP) or cutting an episiotomy, the BFD and PBL were measured using a standardized measurement protocol.

Results: A total of 100 women (50 women in each center) were recruited into the study. The overall mean BFD was 2.8 ± 0.5 cm. The overall mean PBL was 4.4 ± 0.8 cm. There were no statistically significant differences between the measurements taken in both units regarding BFD or PBL ($p = 0.81$ and 0.10 respectively). There was a weak correlation between both measurements.

Conclusion: This is the first study to generate data on the bregma to fourchette distance and the perineal body length just prior to fetal head expulsion. This information will probably be useful to inform future biomechanical computational modeling studies to optimize hands and fingers placement during manual perineal protection.

Introduction

Studies investigating perineal trauma at the time of vaginal birth found that primiparity, forceps delivery, midline episiotomy, persistent occipitoposterior position, and shoulder dystocia were significant risk factors [1]. Manual perineal protection (MPP) is an intrapartum intervention that is used to reduce the risk of severe perineal trauma and its consequences [2–4]. However, there is lack of complete understanding of how to accurately execute the technique and the intended value to be achieved by each of its components [5–9]. We believe that these, as well as the lack of objective evidence of how the technique relates to the dynamic changes occurring at the fetal head and the perineum at the time of crowning, are important limitations to the wide adoption of MPP amongst professionals.

Biomechanical analysis of a variety of MPP techniques that reduce the tension at the midpoint of the perineum identified the Finnish and Viennese maneuvers to be the most effective [4, 10–12]. In addition to the reduction in perineal tension, both techniques also aim to regulate fetal head extension using the flexed middle finger or the ulnar border of the dominant hand respectively [13–16]. This component is considered to be important for the effectiveness of MPP because as the head descends it starts to extend [17].

Stereo-photogrammetric analysis and biomechanical studies have optimized the positioning of the thumb and finger(s) of the dominant hand to minimize tension at the midpoint of the perineum [10–12,

18]. However, the exact point on the perineum where the middle finger or ulnar border of the dominant hand is supposed to be applied is not clearly specified. This lack of clarity leaves the positioning open to individual interpretation and might account for the variation in the reproducibility of the technique's effectiveness in reducing perineal trauma.

The overarching aim of this work was to collect data on the degree of fetal head extension and maternal perineal length at the time of manual perineal protection to inform a biomechanical computer model used to optimize this intervention. We proposed that the fetal bregma to the posterior fourchette distance (BFD) and the perineal body length (PBL) could be used as objective surrogate measures for these two variables respectively. Therefore, we designed this study to gather information on these two measurements in a diversified and representative group of women.

Methods

This is a prospective cohort study conducted in 2 maternity units in Hagen, Germany (unit 1) and in Cairo, Egypt (unit 2). Women having their first spontaneous vaginal birth of a singleton term baby in vertex presentation and occipitoanterior position were considered eligible to participate in the study. Suspected fetal distress or fetal anomalies, occipitoposterior position, operative vaginal delivery, and history of vaginal or perineal operations were reasons for exclusion. Women meeting the inclusion criteria were informed about the study on admission to the labor ward during the first stage of labor. Favorable ethical approvals were granted to the study by the Hospital Board of Management at Allgemeins Hospital Hagen on 02.01.2020 and the Research Ethics Committee at Ain Shams University on 05.01.2021 (Ref. FMASU MS 704/2020/2021) (supplementary information). All study participants provided a valid written informed consent.

Intrapartum care was undertaken based on the participating unit's protocol. One assessor (AA) performed all the measurements in both units. Measurements were performed while the woman was in a semi-recumbent lithotomy position with the legs flexed at the hip joints at an angle between 90° and 100° [19, 20]. Disposable sterile surgical skin marker and tape measure kits were used for the assessments (Viscot Medical, LLC; East Hanover, NJ 07936). The fixed reference points were the posterior fourchette (the posterior margin of the hymen), and the most anterior point on the anal epithelium at 12 o'clock from the center of the anal orifice [21]. PBL was defined as the distance between the posterior fourchette and the anterior margin of the anal opening [21]. We used the anterior margin of the anal opening rather than the center of the anus, as proposed by the International Continence Society, to mitigate inaccuracies resulting from the varying degrees of anal dilatation that may occur at the time of fetal head crowning.

Measurements were taken once crowning was diagnosed and just prior the initiation of MPP or cutting an episiotomy, whichever came first. The index finger was inserted through the vaginal introitus in the midline between the fetal head and perineum to palpate the fetal bregma. Using a sterile skin marker, a point was marked externally in the midline on the perineal skin corresponding to the point at which the center of the fetal bregma was palpated. Using a sterile tape measure, the distance between this marked

point on the perineum and posterior fourchette was measured (BFD). Using the same tape measure, the distance between the posterior fourchette and the anterior margin of the anal opening, (at 12 o'clock) was then measured (PBL) (Fig. 1). Additionally, we collected data on maternal age, BMI, duration of second stage of labor, birth weight, neonatal head circumference and any associated perineal trauma including episiotomy.

A convenience sample of 50 women per unit was set a priori for the study. Statistical analysis was performed using SAS 9.4 (SAS Institute Inc., Cary, NC, USA) statistical software. Difference in distribution of ranked data between groups was carried out with non-parametric ANOVA (2-sample Wilcoxon test or 2-sample Median test), while categorical variables were assessed by the χ^2 test or Fisher's exact test. The Pearson correlation and linear regression were used to assess the relation between PBL and BFD. A $p < 0.05$ was considered statistically significant.

Results

The study was conducted between January 2, 2020, and August 30, 2020, at unit 1 and between January 5, 2021, and May 30, 2021, at unit 2. A total of 100 primiparous women were recruited and contributed full data into the study in the two units. The number of patients approached and recruited to achieve the target sample is presented in Fig. 2. The demographic details and obstetrics variables are presented in Table 1. There was a statistically significant difference between the two centers regarding duration of the 2nd stage of labor ($p = 0.004$) and neonatal weight ($p = 0.02$). The overall mean BFD was 2.8 ± 0.5 cm (2.8 ± 0.6 cm in unit 1 and 2.8 ± 0.5 cm in unit 2). The overall mean PBL was 4.4 ± 0.8 cm (4.5 ± 0.8 cm in unit 1 and 4.3 ± 0.7 cm in unit 2). There were no statistically significant differences between both units regarding BFD or PBL ($p = 0.81$, and 0.10 respectively).

Table 1
The demographic data and obstetrics variables in the study group

Obstetric variable	Cohort	MEAN ± SD	MEDIAN (IQR)	p-value
Maternal age, years	overall	28.0 ± 5.9	27.5 (24.0–32.0)	0.46 *
	unit 1	28.4 ± 6.2	27.5 (24.0–33.0)	
	unit 2	27.5 ± 5.6	27.5 (24.0–31.0)	
Body mass index, kg/m2	overall	29.3 ± 4.2	29.0 (26.0–32.0)	0.63 *
	unit 1	29.4 ± 4.3	30.0 (26.0–32.4)	
	unit 2	29.2 ± 4.0	29.0 (26.0–32.0)	
Duration of the 2nd stage of labor, min	overall	82.3 ± 24.9	83.5 (65.0–98.0)	0.004 *
	unit 1	89.7 ± 27.0	92.0 (70.0–103.0)	
	unit 2	74.8 ± 20.3	75.0 (60.0–90.0)	
Birth weight, g	overall	3446 ± 403	3440 (3225–3670)	0.02 *
	unit 1	3531 ± 463	3535 (3300–3820)	
	unit 2	3361 ± 314	3345 (3150–3550)	
Neonatal head circumference, cm	overall	35.3 ± 1.5	35.0 (34.0–36.5)	0.38 *
	unit 1	35.5 ± 1.6	35.0 (34.0–37.0)	
	unit 2	35.1 ± 1.4	35.5 (34.0–36.0)	
Bregma – fourchette distance, cm	overall	2.8 ± 0.5	2.8 (2.5–3.2)	0.81 *
	unit 1	2.8 ± 0.6	2.9 (2.4–3.2)	
	unit 2	2.8 ± 0.5	2,8 (2.5–3.1)	
Perineal body length, cm	overall	4.4 ± 0.8	4.2 (3.9–4.9)	0.10 *
	unit 1	4.5 ± 0.8	4.4 (3.9–5.1)	
	unit 2	4.3 ± 0.7	4.1 (3.8–4.6)	
*) Wilcoxon Two Sample Test				

On assessing the relationship between BFD and PBL using Pearson coefficient, there was low correlation between both measurements ($r = 0.4820$) (Fig. 3).

Discussion

Summary of findings

To our knowledge, this is the first study to generate data for BFD and PBL at time of initiation of MPP. Our study demonstrated that there were no statistically significant differences between these measurements in two cohorts of women recruited in a unit in Germany and another one in Egypt despite some significant differences in participants' demographics. This observation suggests that ethnicity does not seem to affect these measurements; nonetheless, this finding may be limited by our sample size. The statistically significant difference in maternal age between the two groups may be explained by cultural and / or social differences between the two countries. There was also statistically significant difference in neonatal weight where the mean birthweight was higher in Germany compared to Egypt. Exploring the reasons for this difference is beyond the scope of our work, however, it is plausible that constitutional and / or nutritional factors have contributed to such difference.

When comparing the duration of second stage of labor between the two centers, there was a statistically significant difference between the two groups. This could be due to variations in the management protocols adopted by the two centers. Indeed, at Ain Shams University, getting women to commence active pushing as soon as full cervical dilatation is confirmed is the preferred policy to shorten the second stage of labor as much as possible to be able to manage the large number of women admitted in labor per day. Moreover, in the absence of a midwifery service in Egypt, labor is totally managed by a medical team, which might have had an impact on the observed difference.

Comparison Of Findings To Other Studies

Although our study group was not the first one to measure the PBL, to our knowledge, we are the first to measure it at the time of initiation of MPP and the first to propose and measure the BFD. Rizk et al. studied the relation between perineal length and position of the anus at the time of vaginal delivery in primigravidae. They measured PBL and the distance between the fourchette and the inferior margin of the coccyx to calculate the anal position index. Unlike our study, these measurements were obtained at the first stage of labor. In their study a short perineum and anterior displacement of the anus were associated with a traumatic vaginal delivery [22]. Interestingly, Rizk and associates' reported mean PBL measurement of 4.6 cm is comparable to the mean PBL in our study despite the difference in the stage in labor when the measurements were taken. In contrast, the mean PBL measured in our study was shorter than that reported by Meriwhether et al. where their mean reported PBL increase from 3.7 cm during the first stage to a maximum of 6.1 cm during the second stage [23]. Nonetheless, in Rizk et al and

Meriwhether et al studies PBL was defined as the distance between the posterior fourchette and the center of the anal orifice [22, 23]. However, we opted to use the anterior anal margin rather than the center of the anus to mitigate the potential inaccuracies caused by the varying degrees of anal dilatation at the later stages of labor [21]. Hence, it is plausible that, not only the difference in the stage of labor at which measurements were taken but also the reference point used for measurement could have contributed to some of the unexpected comparisons. Furthermore, Pihl and colleagues, studied the relation between the anovaginal distance (AVD) measured by transperineal ultrasound and the occurrence of external anal sphincter injury before suturing the perineal laceration. They reported that a short AVD could be a warning sign and should increase the awareness of accoucheurs of possible external sphincter injury before suturing a perineal tear [24]. Nevertheless, they measured the AVD after the vaginal birth and before suturing a perineal laceration, which does not reflect the actual perineal length at the time of crowning.

Jansova et al. investigated the role of the thumb and index finger in manual perineal protection and concluded that the optimal placement of the thumb and index finger is 2 cm anterior of the fourchette and 12 cm apart while the most effective distance and direction of movement is to approximate these digits by 1 cm on either side without changing their antero-posterior orientation to the fourchette [10–12]. We believe that the results of this study provide important information for the optimization of the placement of the flexed middle finger, in the Finnish technique, or the ulnar border of the hand, in the Viennese technique, to facilitate the delivery of the fetal head through the perineum with the least possible head circumference. Indeed, the lack of such measurements to objectively describe the degree of flexion of the fetal head at the time of instigating MPP, has been a major limitation for studies investigating this intervention.

Strengths And Limitations

We appreciate that our study has some limitations. The lack of availability of data on BFD and PBL at the time of crowning hindered our ability to perform a priori formal sample size calculation for this work. However, we were able to recruit a well-defined cohort of women from different ethnic backgrounds and were able to generate a fairly homogenous dataset for these measurements. Additionally, it could be argued that having one assessor collecting all the data could be a potential source for bias. Nevertheless, this design has negated the possibility of inter-rater variability.

Conclusion

This is the first study to generate data on the bregma to fourchette distance and the perineal body length just prior to fetal head expulsion. This information will inform future biomechanical computational modeling studies to optimize hands and fingers placement during manual perineal protection.

Declarations

Funding

This work was supported by the project no. 337 “Virtual models for optimization of medical training” within INTERREG V-A program: Cross-border cooperation between the Czech Republic and the Federal State of Germany, Bavaria, Aim European Cross-border Cooperation 2014–2020 funded jointly by the European Regional Development Fund (85% of the costs) and the state budget of the Czech Republic (5%). KI is partly funded by project No. CZ.02.1.01/0.0/0.0/16_019/0000787 “Fighting Infectious Diseases - FIND “, awarded by the Ministry of Education, Youth and Sports of the Czech Republic, financed from The European Regional Development Fund. Funders were not involved in the design, analysis, or the reporting of this work.

The funders did not have a role in the collection, analysis, and interpretation of data and in the writing of the manuscript.

Competing Interests

The authors have no relevant financial or non-financial interests to disclose.

Author Contributions

A Abouelhassan: literature search, protocol development, data collection, data analysis, prepared the first draft of manuscript.

V Kalis: conceived the idea, literature search, data analysis and interpretation, editing and revising the manuscript.

M Schüssler: data interpretation.

H Awwad: data interpretation.

R Hassan: data interpretation.

K M Ismail: protocol development, literature search, data analysis and interpretation, editing and revising the manuscript.

Ethics approval

This study was performed in line with the principles of the Declaration of Helsinki. Approvals were granted to the study by the Hospital Board of Management at Allgemeins Hospital Hagen on 02.01.2020 and the Research Ethics Committee at Ain Shams University on 05.01.2021 (Ref. FMASU MS 704/2020/2021) (supplementary information).

Consent to participate

All study participants provided a valid written informed consent.

Consent to publish

References

1. Third- and Fourth-degree Perineal Tears, Management (Green-top Guideline No. 29).
<https://www.rcog.org.uk/en/guidelines-research-services/guidelines/gtg29/>. Accessed 9 Jan 2022
2. Agliatoro M (1970) [The contribution of the Salerno School to the historical evolution of protection and suturing of the perineum in labor]. *Minerva Med* 61:5201–3
3. DEWEES WB (1889) RELAXATION AND MANAGEMENT OF THE PERINEUM DURING PARTURITION. *J Am Med Assoc* XIII: <https://doi.org/10.1001/jama.1889.02401200011001c>
4. Ismail KMK, Paschetta E, Papoutsis D, Freeman RM (2015) Perineal support and risk of obstetric anal sphincter injuries: A Delphi survey. *Acta Obstet. Gynecol. Scand.* 94
5. Kleprlikova H, Kalis V, Lucovnik M, et al (2020) Manual perineal protection: The know-how and the know-why. *Acta Obstet. Gynecol. Scand.* 99
6. Kalis V, Kalis V, Rusavy Z, et al (2020) Metrics of perineal support (MOPS) study. *BMC Pregnancy Childbirth* 20: <https://doi.org/10.1186/s12884-020-03042-3>
7. Manresa M, Kalis V, de Tayrac R, et al (2020) Hands up if you do not understand hands on. *Midwifery* 90
8. Ali-Masri H, Hassan S, Fosse E, et al (2018) Impact of electronic and blended learning programs for manual perineal support on incidence of obstetric anal sphincter injuries: A prospective interventional study. *BMC Med Educ* 18: <https://doi.org/10.1186/s12909-018-1363-3>
9. Ali HY, Vikanes Å, Anti M, et al (2017) Evaluation of an animated instructional video as a training tool for manual perineum support during vaginal delivery. *Int J Gynecol Obstet* 137: <https://doi.org/10.1002/ijgo.12115>
10. Jansova M, Kalis V, Lobovsky L, et al (2014) The role of thumb and index finger placement in manual perineal protection. *Int Urogynecol J* 25: <https://doi.org/10.1007/s00192-014-2425-7>
11. Jansova M, Kalis V, Rusavy Z, et al (2014) Modeling manual perineal protection during vaginal delivery. *Int Urogynecol J* 25: <https://doi.org/10.1007/s00192-013-2164-1>
12. Jansova M, Kalis V, Rusavy Z, et al (2017) Fetal head size and effect of manual perineal protection. *PLoS One* 12: <https://doi.org/10.1371/journal.pone.0189842>
13. Laine K, Pirhonen T, Rolland R, Pirhonen J (2008) Decreasing the incidence of anal sphincter tears during delivery. *Obstet Gynecol* 111: <https://doi.org/10.1097/AOG.0b013e31816c4402>
14. Laine K, Skjeldestad FE, Sandvik L, Staff AC (2012) Incidence of obstetric anal sphincter injuries after training to protect the perineum: Cohort study. *BMJ Open* 2: <https://doi.org/10.1136/bmjopen-2012-001649>
15. Poulsen MO, Madsen ML, Skriver-Møller AC, Overgaard C (2015) Does the Finnish intervention prevent obstetric anal sphincter injuries? A systematic review of the literature. *BMJ Open* 5
16. Kalis V, Jansova M, Rusavy Z (2017) Perineal mapping. In: *Perineal Trauma at Childbirth*

17. Cechova H, Kalis V, Havelkova L, et al (2021) Finite element modeling of maximum stress in pelvic floor structures during the head expulsion (FINESSE) study. *Int Urogynecol J* 32:.
<https://doi.org/10.1007/s00192-021-04769-z>
18. Zemčík R, Karbanova J, Kalis V, et al (2012) Stereophotogrammetry of the perineum during vaginal delivery. *Int J Gynecol Obstet* 119:.. <https://doi.org/10.1016/j.ijgo.2012.05.018>
19. Kalis V, Karbanova J, Horak M, et al (2008) The incision angle of mediolateral episiotomy before delivery and after repair. *Int J Gynecol Obstet* 103:.. <https://doi.org/10.1016/j.ijgo.2008.05.026>
20. Kalis V, Landsmanova J, Bednarova B, et al (2011) Evaluation of the incision angle of mediolateral episiotomy at 60 degrees. *Int J Gynecol Obstet* 112:.. <https://doi.org/10.1016/j.ijgo.2010.09.015>
21. Kalis V, Karbanova J, Bukacova Z, et al (2010) Anal dilation during labor. *Int J Gynecol Obstet* 109:.. <https://doi.org/10.1016/j.ijgo.2009.11.024>
22. Rizk DEE, Thomas L (2000) Relationship between the length of the perineum and position of the anus and vaginal delivery in primigravidae. *Int Urogynecol J* 11:..
<https://doi.org/10.1007/s001920050074>
23. Meriwether K V., Rogers RG, Dunivan GC, et al (2016) Perineal body stretch during labor does not predict perineal laceration, postpartum incontinence, or postpartum sexual function: a cohort study. *Int Urogynecol J* 27:.. <https://doi.org/10.1007/s00192-016-2959-y>
24. Pihl S, Uustal E, Blomberg M (2019) Anovaginal distance and obstetric anal sphincter injury: a prospective observational study. *Int Urogynecol J* 30:.. <https://doi.org/10.1007/s00192-018-3838-5>

Figures

i) Index finger was inserted through the vaginal introitus in the midline between fetal head and perineum to palpate the fetal bregma.

ii) Using a sterile skin marker, a point was marked externally in the midline on the perineal skin corresponding to the point at which the center of the fetal bregma was palpated

iii) Using a sterile tape measure, the distance between the marked points on perineum and posterior fourchette were measured.
(BFD = Distance A to B; PBL = Distance A to C)

Figure 1

Simulation of how the BFD and PBL measurements were assessed.

* Not feasible to take measurements because the participant was too distressed, labor progressed too fast or the assessor was not available at time of birth.

Figure 2

Flowchart of patient flow into the study

Figure (3): Relationship between BFP and PBL using Pearson coefficient

Figure 3

Relationship between BFP and PBL using Pearson coefficient