Precipitation Frequency in Med-CORDEX and EURO-CORDEX Ensembles from 0.44° to Convection-Permitting Resolution: Impact of Model Resolution and Convection Representation

Minh Truong Ha (✉ minh.ha@latmos.ipsl.fr)

LATMOS: Laboratoire Atmospheres Milieux Observations Spatiales <https://orcid.org/0000-0002-3755-0714>

S. Bastin

LATMOS: Laboratoire Atmospheres Milieux Observations Spatiales

P. Drobinski

LMD: Laboratoire de Meteorologie Dynamique

L. Fita

CIMA: Centro de Investigaciones del Mar y la Atmosfera

J. Polcher

LMD: Laboratoire de Meteorologie Dynamique

O. Bock

IPGP: Institut de Physique du Globe de Paris

M. Chiriaco

LATMOS: Laboratoire Atmospheres Milieux Observations Spatiales

D. Belušić

Swedish Meteorological and Hydrological Institute: Sveriges meteorologiska och hydrologiska institut

C. Caillaud

CNRM GMGEC: Centre National de Recherches Meteorologiques Groupe de Meteorologie de Grande Echelle et Climat

A. Dobler

Norwegian Meteorological Institute: Meteorologisk institutt

J. Fernandez

IFCA: Instituto de Fisica de Cantabria

K. Goergen

Jülich Research Centre Institute for Advanced Simulation: Forschungszentrum Jülich Institute for Advanced Simulation

Ø. Hodnebrog

Center for International Climate Research: CICERO

S. Kartsios

Aristotle University of Thessaloniki: Aristoteleio Panepistemio Thessalonikes

E. Katragkou

Aristotle University of Thessaloniki: Aristoteleio Panepistemio Thessalonikes

A. Lavin-Gullon

IFCA: Instituto de Fisica de Cantabria

T. Lorenz

NORCE Norwegian Research Centre AS

J. Milovac

IFCA: Instituto de Fisica de Cantabria

H.-J. Panitz

KIT: Karlsruher Institut für Technologie

S. Sobolowski

NORCE Norwegian Research Centre AS

H. Truhetz

University of Graz: Karl-Franzens-Universität Graz

K. Warrach-Sagi

University of Hohenheim: Universität Hohenheim

V. Wulfmeyer

University of Hohenheim: Universität Hohenheim

Research Article

Keywords: Precipitation triggering, convection-permitting models, convection processes, integrated water vapor

Posted Date: March 3rd, 2022

DOI: <https://doi.org/10.21203/rs.3.rs-1397006/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Abstract

Recent studies using convection-permitting (CP) climate simulations have demonstrated a step-change in the representation of heavy rainfall and rainfall characteristics (frequency-intensity) compared to coarser resolution Global and Regional Climate models. The goal of this study is to better understand what explains the weaker frequency of precipitation in the CP ensemble by assessing the triggering process of precipitation in the different ensembles of regional climate simulations available over Europe. We focus on the statistical relationship between tropospheric temperature, humidity and precipitation to understand how the frequency of precipitation over Europe and the Mediterranean is impacted by model resolution and the representation of convection (parameterized vs. explicit). We employ a multi-model data-set with three different resolutions (0.44° , 0.11° and 0.0275°) produced in the context of the MED-CORDEX, EURO-CORDEX and the CORDEX Flagship Pilot Study "Convective Phenomena over Europe and the Mediterranean" (FPSCONV). The multi-variate approach is applied to all model ensembles, and to several surface stations where the integrated water vapor (IWV) is derived from Global Positioning System (GPS) measurements. The results show that all model ensembles capture the temperature dependence of the critical value of IWV (IWV_{cv}), above which an increase in precipitation frequency occurs, but the differences between the models in terms of the value of IWV_{cv}, and the probability of its being exceeded, can be large at higher temperatures. The lower frequency of precipitation in convection-permitting simulations is not only explained by higher temperatures but also by a higher IWV_{cv} necessary to trigger precipitation at similar temperatures, and a lower probability to exceed this critical value. The spread between models in simulating IWV_{cv} and the probability of exceeding IWV_{cv} is reduced over land in the ensemble of models with explicit convection, especially at high temperatures, when the convective fraction of total precipitation becomes more important and the influence of the representation of entrainment in models thus becomes more important. Over lowlands, both model resolution and convection representation affect precipitation triggering while over mountainous areas, resolution has the highest impact due to orography-induced triggering processes. Over the sea, since lifting is produced by large-scale convergence, the probability to exceed IWV_{cv} does not depend on temperature, and the model resolution does not have a clear impact on the results.

1. Introduction

Regional climate models (RCMs) are developed to provide regional details by downscaling General Circulation Models (GCMs) output. With fine-scale information for regional features, RCMs can give some added value compared to GCMs (Giorgi, 2019). RCMs can describe more realistic near-surface wind speed over complex coastal areas (Winterfeldt and Weisse 2009), improve describing mesoscale variability of near-surface pressure and temperature (Feser and von Storch 2005), they are also able to capture well the average statistics of daily moderate precipitation on scales of a few grid boxes (Boberg et al. 2009; Kjellström et al. 2010). Despite refinements in resolution (12–50 km) and efforts to revise the dynamical and physical processes in RCMs, rainfall extremes are still poorly represented by the models even at regional scales (IPCC report AR6, 2021). Models tend to overestimate weak precipitation (Bastin et al, 2019, Stephens et al., 2010) and underestimate heavy rainfall (Lenderink and van Meijgaard, 2008).

In GCMs and RCMs, deep convection, which is a dominant source of summer precipitation, is represented by using convection parameterization schemes. However, the inherent assumptions made in such schemes have

been identified as a major source of errors and uncertainties in climate models (Hohenegger et al. 2008, Foley, 2010; Kendon et al., 2012; Ban et al., 2014). For example, RCMs with parameterized convection can produce a too early onset of daytime moist convection in the warm season (Brockhaus et al., 2008), which affects convective potential energy and leads to a simulation of too frequent light precipitation (Dai and Trenberth, 2004). Additionally, the poor representation of the interaction between subgrid-scale soil moisture and atmospheric temperature anomalies (Froidevaux et al., 2014) can exhibit a positive soil moisture-precipitation feedback and amplify perturbations of the hydrological cycle (Hohenegger et al., 2009). Recently, with the advances in high-performance computing, convection-permitting models (CPMs) with horizontal grid spacing less than ~ 4 km, which are traditionally used in weather forecasting, have begun to be applied on climate-time scales (Hohenegger et al. 2008; Kendon et al. 2017; Prein et al. 2015). At these resolutions, the parameterization of deep convection is not required (Prein et al., 2015; Weisman et al., 1997), providing hope that model uncertainties and parameter sensitivity may be reduced.

Several long-term simulations have been conducted with CPMs in the last decade over different areas, mainly Europe and the USA (e.g Ban et al, 2014, 2021; Kendon et al, 2012, 2014; Rasmussen RM et al. 2011, 2014b; Rasmussen KL et al, 2017; Keller et al, 2016; Prein et al. 2015; Leutwyler et al, 2017; Liu et al, 2017; Berthou et al, 2018; Hodnebrog et al, 2019; Fumière et al, 2020; Knist et al, 2020) and a few over other areas (e.g Kouadio et al., 2020; Berthou et al, 2019; Kendon et al 2019; Guo et al, 2019, 2020; Ahrens et al, 2020; Konduru & Takahashi, 2020; Argüeso et al., 2020). These studies emphasize the added value of CPMs in representing precipitation over orographic regions, focusing more specifically on convective organization, the diurnal cycle of convection precipitation, the spatial structure of rainfall and its duration-intensity characteristics, and the intensity of hourly precipitation extremes (see Lucas-Picher et al., 2021 for a review). However, the frequency of precipitation, which tends to be overestimated in RCMs at coarser resolution, seems to be underestimated in summer compared to available observational datasets (Ban et al., 2021). Even if these studies show promising improvements, only a few of them identify the processes underlying such improvements (e.g Rasmussen KL et al., 2017). Convection triggering processes in particular are not well investigated despite the fact that they have a strong impact on the entire precipitation distribution and more specifically on extreme precipitation.

Properly representing convection is critical to successful simulations and ensuing water resources assessments. Convection can be viewed conceptually as being driven by the buoyancy realized by parcels being lifted to their level of free convection and rising freely into the local environment. Yet, it can be argued that moisture is a key component for convection, since convective parameterization is simply a method to account for the effects of subgrid-scale saturation. From this perspective, the moisture content drives the behavior of the convective scheme, controlling the amount of convection produced in an unstable environment based upon the available moisture that can be removed from the atmosphere. Convective parameterizations attempt to represent the dependence of the statistics of moist convection on water vapor and temperature. This forms the basis of many convective parameterizations and a better knowledge of this relationship in climate models would help to improve precipitation simulation. Therefore, we analyze the triggering of precipitation based on the relationship between precipitation, integrated water vapor (IWV) and tropospheric temperature. IWV has been identified as a better proxy than saturation vapor pressure for precipitation triggering in the Tropics (Neelin et al., 2009; Sahany et al., 2012). Over Europe this relationship

was found to be consistent with studies conducted in the Tropics (Bastin et al., 2019). Comparisons of observations and some simulations in the HYMEX/Med-CORDEX programs showed deviations of IWV between models with parameterized deep convection over Europe at high temperature, giving some insights in the way this relationship could help in the investigation of model's spread.

In this study, we investigate the impact of model resolution and convection representation on the triggering of precipitation over Europe and the Mediterranean area. Model ensembles with three different resolutions (0.44° , 0.11° and 0.0275°) are used. At 0.0275° , deep convection is explicitly represented while at 0.44 and 0.11° , it is parameterized with different deep convection schemes. Simulations were performed in the frame of the EURO- (Jacob et al., 2014) and Med-CORDEX (Ruti et al., 2016) projects, and of their shared Flagship Pilot Study (FPS) on “Convective phenomena at high resolution over Europe and the Mediterranean” (Coppola et al, 2020; Ban et al., 2021).

The structure of the paper is as follows: Section 2 presents the data and methodology of this study. Section 3 presents results on the analysis of the impact of the model resolution and convection representation on precipitation triggering, and section 4 assesses the discrepancy between the different ensembles. Finally, section 5 provides a summary and conclusion.

2. Data And Methods

2.1. Model simulations

As mentioned in the introduction, we have collected those simulations from Med-CORDEX, EURO-CORDEX and FPS on convection (FPSCONV), which provide the variables required in our methodology (Section 2.3). The high resolution convection-permitting simulations (2.2 to 3 km) use explicit deep convection and cover the greater Alpine region (Fig. 1) for the period 2000–2009. These simulations are generally embedded in an outer domain covering the EURO-CORDEX area at 12 or 15 km resolution. This provides pairs of simulations suitable for the assessment of the effects of convection representation and resolution. Therefore, the analysis will be done over the FPSCONV domain, which is a common area of the Med-CORDEX, EURO-CORDEX and FPSCONV experiments, to assess: (1) the role of increased resolution and (2) the role of explicit convection over the period 2001–2008, which is the common period of the three ensembles. Additionally, three subareas (black boxes in Fig. 1) are chosen to analyze in more detail the model performance in different terrain including: (1) high orography over the mountains of the Alps ($5-13^\circ\text{E}$, $45.5-47^\circ\text{E}$), (2) the Mediterranean Sea ($4-8^\circ\text{E}$, $40.8-42.5^\circ\text{E}$) and (3) continental plains ($2-12^\circ\text{E}$, $47.5-49^\circ\text{E}$).

The ensemble at 0.44° resolution (hereafter EUR-44) includes 4 simulations of the Med-CORDEX project and 6 simulations of the EURO-CORDEX project (Table 1). For the simulations at 0.11° resolution (EUR-11) ensemble, we have collected 3 simulations of the EURO-CORDEX project and the 9 remaining members are the simulations that drive the convection-permitting models over the Alpine region (Table 2). This sub-ensemble of 9 simulations is named ALP-11 and will give fair comparisons with high resolution (0.0275°) convection-permitting simulations (ALP-3). Six simulations are using version 3.8.1 of the WRF model (Skamarock et al., 2008) with the same configuration and initial and boundary forcings. Only physics schemes differ between them (the last letter indicates a different physics configuration; Lavin-Gullon et al., 2020 gives more details on

this multi-physics ensemble). One simulation is using CNRM-ALADIN62 (Nabat et al, 2020), one HCLIM38-ALADIN (Belušić et al, 2020), and the last one is using COSMO-CLM (Rockel et al, 2008; Baldauf et al, 2011). Precipitation from all these simulations have been evaluated in Ban et al., 2021.

Table 1
List of ERA-Interim driven simulations at 0.44 degree used in this study (referred to as EUR-44).

Group Abbreviation	Group Full Name	Model	Domain
IPSL	Institut Pierre-Simon-Laplace (France)	WRF311	MED-CORDEX
UCLM	Universidad de Castilla-La Mancha (Spain)	UCLM	MED-CORDEX
CNRM	Centre National de Recherches Meteorologiques (France)	ALADIN52	MED-CORDEX
CMCC	Centro Euro-Mediterraneo sui Cambiamenti Climatici (Italy)	CCLM4-8-19	MED-CORDEX
DMI	Danish Meteorological Institute (Denmark)	HIRHAM5	EURO-CORDEX
HMS	Hungarian Meteorological Service (Hungary)	ALADIN52	EURO-CORDEX
KNMI	Royal Netherlands Meteorological Institute (Netherlands)	RACMO22E	EURO-CORDEX
MOHC	Met Office Hadley Centre (England)	HadRM3P	EURO-CORDEX
SMHI	Swedish Meteorological and Hydrological Institute (Sweden)	RCA4	EURO-CORDEX
UCAN	Universidad de Cantabria (Spain)	WRF341I	EURO-CORDEX

Table 2

List of ERA-Interim driven simulations at 0.11 degree and 0.0275 degree used in this study. The twelve 0.11° simulations are referred to as EUR-11 ensemble and the last nine simulations as ALP-11 (ALP-11 is a sub-ensemble of EUR-11). The 0.00275° simulations are referred to as ALP-3. Please note the FPSCONV WRF RCMs' ALP-3 domain is driven by a 15km European domain run, covering the same area as the EUR-11 EURO-CORDEX domain.

Group Abbreviation	Group Full Name	Model and grid spacing [km] used over EUR /Med CORDEX domain	Model and nested grid spacing [km] used over Alpine domain
DMI	Danish Meteorological Institute (Denmark)	HIRHAM5 / 12	
KNMI	Royal Netherlands Meteorological Institute (Netherlands)	RACMO22E / 12	
SMHI	Swedish Meteorological and Hydrological Institute (Sweden)	RCA4 / 12	
HCLIMcom	HARMONIE-Climate community Danish Meteorological Institute and MET Norway and Swedish Meteorological and Hydrological Institute (Sweden)	HCLIM38-ALADIN / 12	HCLIM38-AROME / 3
AUTH	Aristotle University of Thessaloniki (Greece)	WRF381BG / 15	WRF381BG / 3
CICERO	Center for International Climate Research (Norway)	WRF381BJ / 15	WRF381BJ / 3
CNRM	Centre National de Recherches Meteorologiques (France)	CNRM-ALADIN62 / 12 (using spectral nudging)	CNRM-AROME41t1. / 2.5
FZJ	Research Centre Juelich (Germany)	WRF381BB / 15	WRF381BB / 3
UCAN	Universidad de Cantabria (Spain)	WRF381BI / 15	WRF381BI / 3
UHOH	University of Hohenheim (Germany)	WRF381BD / 15	WRF381BD / 3
WEGC	Wegener Center for Climate and Global Change, Univ. of Graz (Austria)	WRF381BL / 15	WRF381BL / 3
KIT	Karlsruhe Institute of Technology (Germany)	COSMO-CLM / 25	COSMO-CLM / 3

The ALP-3 ensemble (9 simulations) has a higher resolution and a different representation of deep convection that is explicitly resolved. Two ALP-3 models have some other differences in physical parameterizations compared to their parent configurations and hence different names, as described in the corresponding references: CNRM-AROME41t1 (Caillaud et al, 2021) and HCLIM38-AROME (Belušić et al, 2020). In order to provide a comparison at the same resolution with their driving simulations, we remapped the outputs of ALP-3 to the EUR-11 grid (ALP-3remapEUR-11) using conservative remapping.

2.2. Reference data

To compare model results with observations, we used available gridded precipitation products (for an estimate of daily precipitation occurrence), and stations where we can get IWV, tropospheric temperature, and

precipitation rate.

For IWV, we selected 25 Global Positioning System (GPS) stations (Table 3) distributed throughout the domain (red dots in Fig. 1) and for which the available data coverage is at least 4 years of the evaluation period (2001–2008). The GPS dataset used in this study is based on homogeneously reprocessed GPS delay data produced by IGN. It contains 6-hourly, daily and monthly Zenith Tropospheric Delay (ZTD) and IWV estimates for more than 400 stations over France and Europe. The original ZTD data were screened and converted into IWV using surface pressure and weighted mean temperatures computed from ERA5 pressure-level data. The daily values were computed as weighted averages of 00, 06, 12, 18 UTC of the day and 00UTC of the next day. More information on the processing and post-processing procedures and options can be found in Bock, (2021), including a link to download the data.

Table 3
List of GPS stations used in this study.

Stations	Province/Country	Latitude	Longitude	Altitude (m)	Period	Number of non-missing values
AXPV	Aix-En-Provence/France	43.49121475	5.33319521	179.526398	2002–2008	1504
BSCN	Besancon/France	47.24687958	5.98938465	310.606384	2001–2008	1884
BZRG	Bolzano/Italy	46.49902344	11.3367958	279.552094	2001–2008	1391
EGLT	Egletons/France	45.40335083	2.05199409	616.033936	2001–2008	1916
GINA	France	43.67572403	5.78697968	331.485809	2001–2008	1911
GRAS	Caussols/France	43.75473785	6.92057228	1269.36096	2001–2008	2222
GRAZ	Graz/Autria	47.06713104	15.4934788	490.8508	2001–2008	2349
MARS	Marseille/France	43.27877045	5.35378599	12.4186001	2001–2008	2225
MDOR	Saint-Didier-au-Mont-d'Or/France	45.79901886	4.80895758	280.781586	2002–2008	1801
MEDI	Medicina/Italy	44.5199585	11.6468134	9.64779949	2001–2008	2250
MICH	France	43.92415619	5.71735287	577.435791	2001–2008	1929
MLVL	Marne-la-Vallee/France	48.84105682	2.58730984	115.980797	2001–2008	2075
MODA	France	45.21377563	6.71008444	1129.13391	2001–2008	1649
MTPL	France	43.63743591	3.86483932	69.8919983	2001–2008	1741
NICA	France	43.70326233	7.22726059	207.646393	2002–2008	1492
OPMT	Paris/France	48.83591843	2.33493781	78.0371018	2002–2008	1849
SJDV	Saint-Jean-des-Vignes/France	45.87908554	4.67657423	382.418915	2001–2008	1689
SMNE	Saint-Mande/France	48.84444427	2.42502046	81.7205963	2001–2008	2333

Stations	Province/Country	Latitude	Longitude	Altitude (m)	Period	Number of non-missing values
STJ9	France	48.62166214	7.68381643	188.723404	2001–2008	2010
TLMF	Toulouse/France	43.57460785	1.37509692	171.992599	2002–2008	1618
TLSE	Toulouse/France	43.56069565	1.48088932	158.102707	2001–2008	2227
TORI	Torino/Italy	45.06336594	7.66128016	262.679779	2001–2008	2157
TRYS	France	48.29424667	4.01181555	122.274399	2002–2008	1807
VFCH	Villefranche-sur-Cher/France	47.29419327	1.71967316	106.545097	2001–2008	1866
ZIMM	Zimmerwald/Switzerland	46.87709808	7.46527576	907.471191	2001–2008	2405

To complement these GPS stations, this study also uses observations collected since 2003 at the SIRTA atmospheric observatory (blue dot on Fig. 1), located 20-km South West of Paris (Haeffelin et al. 2005). This observatory has collected many observations, which are now synthesized into the so-called “SIRTA-ReOBS” dataset described in Chiriaco et al. (2018) and used in Chiriaco et al. (2014) and Bastin et al. (2018, 2019). GPS IWV and precipitation rates are available in this dataset, and the advantage of this type of station is the collocation of the measurements of all variables. For tropospheric temperature, we used the radio soundings launched twice a day a few kilometers from SIRTA.

For the other stations, tropospheric temperature is extracted from the European Centre for Medium-Range Weather Forecasts (ECMWF) reanalysis ERA5 (Hersbach et al., 2019). ERA5 is the fifth generation of the ECMWF atmospheric reanalysis that has been operational at ECMWF since 2016. This new reanalysis replaces the ERA-Interim reanalysis and covers the period from 1950 until the present (currently, the period 1950–1978 is only preliminary released). ERA5 is based on the Integrated Forecasting System (IFS) Cy41r2 which was the operational weather forecasting model in 2016, combined with the best available observational data from satellites and in situ stations around the world into a globally complete and consistent dataset.

For precipitation, we used different national gridded products of daily precipitation with a resolution below 8 km:

- EURO4M-APGD is available at a horizontal resolution of 5 km over the Alpine region from 1971–2008. This dataset is based on daily rain gauge station data and is presented in Isotta et al (2014).
- SAFRAN: The SAFRAN meteorological analysis is a mesoscale atmospheric analysis system for surface variables. It provides meteorological data over France on a 8 km grid at the hourly time step using

observed data and atmospheric simulations. Further details on the SAFRAN analysis system can be found in Quintana-Seguí et al. (2008) and Vidal et al. (2010).

- REGNIE (Regionalisierte Niederschlagshöhen) is a gridded dataset of daily precipitation with 1 km horizontal resolution provided by the hydrometeorologic department of Deutscher Wetterdienst. It covers Germany and is a multiple regression interpolation product based on geography and inverse distance weighting (Rauthe et al., 2013).

2.3. Method to analyze precipitation triggering based on the relationship between IWV and precipitation as a function of temperature

The method to establish the relationship between IWV and precipitation as a function of tropospheric temperature was introduced in detail in Bastin et al. (2019). Here we introduced some modifications to the method in order to adapt it to the new observational datasets used in this study and for the comparison between models at different resolutions. The analysis period is 2001–2008, when the maximum number of simulations and observations are available, with the exception of the SIRTa station, which covers the period 2008–2015. Using a different period can slightly affect the results, but SIRTa is still of interest because it is collocated with other measurements. An assessment of the impact of the selected period has been done in Bastin et al. (2019) and shows the match for the overlapping period. We therefore decided to keep this station so that we can have a reference from observations to compare with model outputs in the continental plains analysis subarea.

Tropospheric temperature is calculated as the average of the 2m temperature and the 850, 500, and 200hPa temperature as in Bastin et al. (2019) but, here, we split the tropospheric temperature into 1K size bins instead of 4K to discover some changes in the behavior of the model that may occur in only a small temperature range. For each temperature bin, the daily mean precipitation rates are sorted by increasing values of daily mean IWV, then the sorted IWV values are divided into different bins that contain an equal number of pairs of precipitation rates and IWV to ensure a reasonable number of days to compute the 50th quantile of precipitation. The critical IWV (IWV_{cv}) is defined by the mean value of the first IWV bin whose 50th quantile of precipitation is greater than 0.1 mm/day. Here, the 50th quantile of precipitation indicates if there are more precipitating days than non-precipitating days in each IWV bin and shows the transition between mostly non-precipitating days and mostly precipitating days. The probability of exceeding IWV_{cv} can therefore be used to assess the frequency of precipitation. For the model simulations, the precipitation frequency, IWV_{cv}, the probability that IWV exceeds IWV_{cv}, and the probability distribution function (PDF) of IWV rescaled (IWV/IWV_{cv}) are computed in the native resolution of each ensemble, and then we average the results of all grid points within the analysis domains to show the final results.

3. Occurrence Of Precipitation

Figure 2 shows the spatial distribution of precipitation frequency (when daily rainfall is greater than 0.1 mm) in winter (DJF) over the Alpine domain for simulations from EUR-44, EUR-11, ALP-11, ALP-3remapEUR-11, ALP-3 and observations. As expected, the difference of resolution between the ensembles and the observations affects this diagnostic, but it is our aim to keep the native resolution here for this comparison. The bias of precipitation frequency between ensembles and observations is shown in Table 4.

Table 4

Bias of precipitation frequency (averaged over all grid points in the different areas shown on Fig. 1) in DJF/JJA between model simulations and observation. Results indicate the MODEL-OBS value, blue shows positive bias and red shows negative bias.

Ensemble	Winter (DJF)			Summer (JJA)		
	Alpine mountains	Plains	Whole observation area	Alpine mountains	Plains	Whole observation area
EUR-44	0.11	0.18	0.13	0.10	0.13	0.11
EUR-11	0.07	0.19	0.11	0.00	0.06	0.03
ALP-11	0.06	0.18	0.09	-0.03	0.02	0.00
ALP-3remapEUR-11	0.01	0.12	0.05	-0.11	-0.08	-0.09
ALP-3	-0.03	0.11	0.03	-0.17	-0.10	-0.12

Compared to observations, almost all ensembles overestimate on average winter daily precipitation frequency, confirming the well-known problem of “too often too light precipitation” (e.g. Sun et al. 2006; Dai 2006; Brown et al. 2010). However, we note an improvement with increasing resolution: the spatial variability is fairly well captured by ALP-3 and ALP-11/EUR-11, but not by EUR-44 which smooths out the variability produced by topographic effects, for example along coastlines, or in valleys around the mountains. The winter rainfall frequency of EUR-11 and ALP-11 is indeed lower than for EUR-44 over the Mediterranean and mountains (the bias is reduced from 0.11 in EUR-44 to 0.07 in EUR-11 and 0.06 in ALP-11) but quite similar over the plains and about 0.18 for the 3 ensembles. When the convection parameterization is switched off (ALP-3 and ALP-3remapEUR-11), the occurrence is again reduced over the entire domain (bias of 0.03 and 0.05 respectively compared to 0.09 in ALP-11), also over the plains (but still overestimated compared to observations, except over the Alpine and Corsica).

Figure 3 and the second column in Table 4 displays the results for summer (JJA). With a higher resolution, EUR-11 and ALP-11 simulate fewer days of precipitation compared to EUR-44 not only over the sea and mountains as in winter but also over the plains (the bias is 0.06 with EUR-11 and 0.02 with ALP-11, while it is 0.13 with EUR-44). Without convection parameterization schemes, the frequency of summer precipitation in ALP-3 and ALP-3remapEUR-11 ensembles are lower than in EUR-11 and ALP-11 for almost all areas and they even underestimate the occurrence compared to observations (the bias is negative), as already noticed by Ban et al. (2021). In summer over the Alpine region, ALP-11 ensemble is the closest to observations in terms of precipitation occurrence, despite the fact that the resolutions of gridded datasets are closer to the resolution of the ALP-3 ensemble than to ALP-11.

To further investigate the impact of resolution in the precipitation triggering, Fig. 4 shows the precipitation frequency of all the ensembles as a function of IWV and tropospheric temperature over the 4 different selected areas represented on Fig. 1: the whole FPSCONV domain, the Alpine mountains range, the Mediterranean Sea

and the continental plains. When surface stations are available in the area, green solid lines indicate the averaged value of IWV as a function of temperature for each station included in the considered area.

For all areas, at low temperature, the range of IWV values for which precipitation occurs is quite narrow (about 6 kg m^{-2}), and a good agreement between the different ensembles and observations is found (green lines in figures represent observed values from stations located within each area). When temperature increases, the range of IWV values becomes larger (greater than 20 kg m^{-2} when the whole range is considered). Values of IWV for rainy occurrence increase at a rate close to Clausius-Clapeyron (hereafter C-C, lower than C-C for the Alpine domain) until a certain value of temperature (T_{break}) at which the curve angles away from C-C, as already noticed and discussed in Bastin et al., (2019). Physically, T_{break} corresponds to the temperature over which the fraction of convective precipitation over total precipitation becomes close to 1 and precipitation triggering is thus yielded by conditional instability, which occurs at lower relative humidity rate than saturation.

When considering the whole domain (Fig. 4a, b), we can see that this threshold of IWV value is simulated by the 4 ensembles and visible in the observations over the different stations included in the studied area. The green line with the lowest values of IWV corresponds to the Caussols station which is located close to the coastline and surrounded by complex terrain (see Fig. 1 and Table 3), where triggering of precipitation is favored by the frequent occurrence of sea breeze which intensifies the upslope winds (Bastin et al., 2006), explaining the occurrence of precipitation for lower values of IWV. This line is however still within the model spread.

The model ensembles clearly simulate T_{break} over the Mediterranean and continental plains. The value of the simulated T_{break} over the Mediterranean seems a bit weaker than over the plains (less than 266 K for the sea versus around 268K in average for the land). Over the Alpine domain, model ensembles do not simulate this limit, while it is visible in the observations (but only from two stations). Also, the values of IWV for which rain occurs are generally lower over the Alpine domain where triggering processes are favored by the complex terrain and because the depth of the atmospheric column is reduced. Generally, the differences between ensembles are not obvious; we can only see higher IWV values for EUR-11 ensemble at low temperatures over the Alpine domain compared to EUR-44 (Fig. 4c).

Entrainment of dry versus moist environmental air into the convective plume conditions the onset of convection (e.g Wulfmeyer et al., 2016, Kuo et al., 2017; Schiro et al., 2016). Since parameterization physics treat entrainment differently, we expect to detect differences in the convective transition statistics (see below) between ensembles and among models more than in the temperature break after which convective precipitation dominates. That is why we need to go a step further in the triggering process.

In the next section, we consider IWV_{cv} over which the median of precipitation occurrence starts to increase (i.e when the probability that it rains becomes higher than 50%). We also examine the distribution of values around this critical value to better assess the triggering process in the different ensembles and over the different areas. As demonstrated by Kuo et al., (2017), over the tropics (continental and ocean), the enhancement of precipitation with increased IWV only exists when entrainment is strong enough, and the value of IWV_{cv} increases with stronger entrainment until a certain value when IWV_{cv} ceases to be sensitive to entrainment of environmental air.

4. Influence Of Resolution And Convection Representation On The Triggering Of Precipitation

4.1 Precipitation enhancement

Figure 5 compares the IWVcv from different simulations as a function of tropospheric temperature over: (a) the FPSCONV domain, (b) Mediterranean, (c) Alpine mountains, and (d) continental plains. The total distribution of IWVcv values for all grid points of evaluation domain and all models is represented by a boxplot, and the mean value of IWVcv for each simulation of the different model ensembles is represented as a filled-color dot. Each color corresponds to a different ensemble: EUR-44 simulations are in blue, EUR-11 simulations in green, ALP-11 in pink, ALP-3remapEUR-11 in brown and ALP-3 simulations in red. The diamonds indicate the standard deviation for each ensemble. It has been multiplied by 10 for a better visualization. The black stars show IWVcv from the observations at the stations which are included in the considered domain. Note the similarity with the range of values from ALP-11 and EUR-11, although ALP-11 ensemble has fewer members (e.g. Figure 5a), which suggests that the three different models of the ALP-11 ensemble (with 9 simulations in total) cover an acceptable representation of model variability in this study.

a) Whole FPSCONV domain

For the whole FPSCONV domain, Fig. 5 clearly shows the increase of IWVcv with tropospheric temperature as well as the increase of the spread between models, the interquartile range of the total distribution being 2.8 kg.m⁻² for the lower temperature bin, and being around 7.5 kg.m⁻² for the higher temperature bin. As in Fig. 4, IWVcv does not increase any further beyond a certain temperature (above about 265 K), and we can see that Tbreak over the sea is lower than over the land. Concerning observations, most of the IWVcv values for the observed stations are contained within the model boxplot range. For lower temperatures, most mean IWVcv values from EUR-11 simulations are equal or higher to those from EUR-44 simulations (difference of the ensemble mean is about 0.5 kg.m⁻²) and the spread between models (represented by the standard deviation as diamonds in the figure) is similar. At higher temperatures, the difference between IWVcv ensemble mean values increases, and the standard deviation becomes very large in EUR-44 ensemble. Standard deviation of EUR-11 simulation is lower than EUR-44 simulation by about 1.9 kg.m⁻² in the 265 K temperature bin. However, it is important to note that we are considering several extra independent models in EUR-44 (UCLM, RACMO22E, HadRM3P) with respect to EUR-11. This extra model diversity can influence the value of the standard deviation. For the other ensembles, mean values of IWVcv for ALP-3 simulations are higher than for ALP-11 in all temperature bins, suggesting that higher resolution increases the value of IWVcv. This difference becomes larger at high temperature, when convective processes become predominant, and the convection parameterization schemes become more important. Since ALP-3 has both higher resolution and a different representation of convective processes, it is suspected that it is not only the resolution which affects the IWVcv value. Indeed, ALP-3remapEUR-11 has very similar values to ALP-3 and higher than ALP-11, although the resolution of the ALP-3remapEUR-11 is similar to that of the ALP-11. The fact that the standard deviation is not much different between ALP-3 and ALP-3remapEUR-11 simulations and lower than in ALP-11 also confirms that activation of convection parameterization schemes contributes to increased spread between models because of the inherent assumptions in convection parameterization schemes.

When considering the probability of exceeding IWV_{cv} (Fig. 6), the models' deficiencies in their ability to represent the distribution of humidity is evident. Indeed, the mean probability decreases when temperature increases but the range is quite large for both low and high temperatures, which reveals the difficulty for models to correctly simulate the distribution of critical humidity throughout the column at all temperatures, not only when convection is active and when surface processes are important. Most of the observations show lower probability to exceed the threshold than the median of boxplot at low temperature and the mean of each model at all temperature bins. It has already been noticed that models generally overestimate low values of humidity (Bastin et al., 2019; Ning et al., 2013) and do not perfectly simulate the phasing of the diurnal cycle (Ning et al., 2013) and the threshold of IWV values observed at high temperature (Bastin et al., 2019). However, some differences between ensembles exist: In EUR-44 ensemble, some models do not really capture IWV_{cv} well because they too frequently exceed IWV_{cv} threshold. At low temperature, EUR-11 simulates probability lower than EUR-44 but the spread is not much different showing a similar degree of agreement between models of each ensemble. At high temperature bin (265K), the probability in EUR-11 is not lower than EUR-44 but the spread reduces more significantly showing a larger similarity among models response at higher resolution. When convection parameterization schemes are switched off, ALP-3 and ALP-3remapEUR-11 simulate probability to exceed IWV_{cv} lower than ALP-11 and the standard deviations are lower, too. At high temperature bins, we cannot see a big difference between ALP-3 and ALP-3remapEUR-11, which means that the impact of convection representation is likely more important in these bins. However, the probability to exceed the critical value is both linked to the value of IWV_{cv} itself, and to the moisture content. Surface humidity can strongly help to spend more time close to the critical value but does not affect the critical value itself.

To better understand the role of resolution versus the role of convection parameterization, the results over the three different areas are analyzed: over the sea, neither resolution nor convection representation should matter a lot, since the triggering process of precipitation is mainly linked to large scale convergence. Over Alpine topography, resolution should be predominant and over the plains both are expected to matter.

b) Mediterranean Sea

What we obtain over the Mediterranean (Fig. 5c and 6c) confirms what we expect: for similar temperature, IWV_{cv} over the sea is higher than over land because of less convective processes (Lucas et al., 1994), so cloud formation needs more humidity in the lower layer to overcome the saturation deficiency of the lower free troposphere, which increases when temperature increases.

The impact of model resolution on the values of IWV_{cv} is not clear. EUR-11 and ALP-11 indeed simulate lower values of IWV_{cv} than both EUR-44 and ALP-3, while ALP-3 values are within the range of EUR-44. But the spread between models is reduced when convection parameterization is switched off.

Figure 6c shows that the probability to exceed the IWV_{cv} does not depend on temperature, which is consistent with the fact that convection is generally weak over the sea, the main process of lifting being the large scale convergence of moisture, which does not depend on temperature. And as for IWV_{cv}, it does not show a sensitivity to resolution while switching off convection parameterization decreases the probability to exceed

the critical value (probably because the critical value is larger) and decreases the standard deviation between models.

c) Alpine domain

Over the Alpine mountains, IWV_{cv} increases at a lower rate than over other areas but does not stop increasing. The absence of a T_{break} value can be explained by the fact that topographic lifting is strong enough to ensure that saturation is reached independently of temperature. With increased model resolution, the mean value of IWV_{cv} in EUR-11 is higher than in EUR-44 for all tropospheric temperature bins, while the spread becomes smaller at higher temperatures. This suggests that resolution significantly impacts the value of IWV_{cv} over mountains compared to other areas. This is because dynamical processes linked to orography play an important role in triggering convection. When considering ALP-11, ALP-3remapEUR-11 and ALP-3, we can detect some differences between the three ensembles with, once again, higher values of IWV_{cv} for ALP-3 than for ALP-11. At higher temperatures, when comparing the mean value of IWV_{cv}, there is a much bigger difference between ALP-11 and ALP-3remapEUR-11 than between ALP-3 and ALP-3remapEUR-11. However, the fact that we detect differences between ALP-3 and ALP-3remapEUR-11 confirms that over mountains, data resolution impacts the diagnostics and that we should be careful to use gridded datasets that have consistent resolution and use representative stations to evaluate models (Lundquist et al., 2020). However, we cannot exclude the impact of convection representation on IWV_{cv} in the simulations beside increasing model resolutions.

Higher resolutions result in lower probabilities and standard deviations in almost all temperature bins. With convection permitting simulations, the probability to exceed IWV_{cv} is lower and differences between ALP-3remapEUR-11 and ALP-11 are larger than between ALP-3 and ALP-11. However, we still cannot really conclude on the impact of switching off convection schemes compared to resolution at higher temperature because we are aware that ALP-3remapEUR-11 simulation is not equivalent to performing a simulation with explicit convection at 0.11° resolution.

d) Plains

Over the plains, when increasing model resolutions, the value of IWV_{cv} in EUR-11 is not really increased compared to EUR-44 but the spread is reduced significantly between models. When considering ALP-11, ALP-3remapEUR-11 and ALP-3, we can see a bigger impact of convection representation, with very similar results for ALP-3remapEUR-11 and ALP-3, and significant differences with EUR-11 and ALP-11.

At lower temperatures, increasing resolution does not reduce the probability to exceed IWV_{cv} but the spread between models decreases a little in EUR-11 compared to EUR-44. At higher temperature bins, both the impact of resolution and the impact of convection representation become more effective.

To conclude this section, when considering the whole domain, if we only increase model resolution, the values of IWV_{cv} are not higher for every model but it can help to reduce the uncertainty between different models. But with convection-permitting simulations, the value of IWV_{cv} is increased everywhere and the spread is lower in all temperature bins, especially in higher temperature bins. As shown in Fig. 3, this impacts the occurrence of precipitation, which becomes too infrequent in summer compared to available observational datasets.

However, high resolution simulations can be more credible than gridded datasets over topography as suggested by Lundquist et al. (2020). Resolution does not impact the results substantially over the sea but is more important over mountains where the processes of convective triggering are linked to orography. The more significant effects of higher resolution on precipitation triggering over the Alps are consistent with previous studies by Knist et al. (2020) and Fosser et al. (2015). We cannot exclude the effect of explicitly resolved convective processes over the Alpine mountains. Some other studies show that it is both the reduced grid spacing and explicit convection schemes in CPMs that improve the simulation of precipitation over the Alps (Bauer et al., 2011; Adinolfi et al., 2021; Ban et al., 2014; Giorgi et al., 2016; Hohenegger et al., 2008; Keller et al., 2016; Langhans et al., 2013; Leutwyler et al., 2017; Lind et al., 2016; Lüthi et al., 2019; Reder et al., 2020). An increase in the resolution may even play a role in improving the timing of the diurnal cycle and intensity of convective precipitation (Prein, et al., 2013) or precipitation triggering in winter (Piazza et al., 2019) over the Eastern Alpine region. Although the deactivation of convection parameterization does not show a clear effect on IWV_{cv} over mountainous areas except at higher temperatures, it should be an important argument regarding temperature evolution over the Mediterranean. According to the results obtained over the tropics by several studies (Kuo et al., 2017), entrainment may explain part of the differences obtained here but it is difficult to extract the information from ensemble experiments.

4.2 Distribution of occurrence of IWV

Several studies (e.g. Neelin et al, 2009; Sahany et al, 2014) suggest that the shape of the distribution of IWV near and above IWV_{cv} may be a robust indicator of transition to deep convection over the tropical oceans and can be useful in integrating properties of various statistical measures of precipitation. Figures 7 and 8 show the number of occurrences of IWV rescaled by the corresponding critical values for each of the temperature bins, considering only the precipitating days for the period from 2001 to 2008. Figure 7 is for the whole domain, for the model ensembles ALP-11 and ALP3remapEUR11 and for observations (all stations are considered in order to generate sufficient sample size). Figure 8 presents the results for each box separately only for the two above-mentioned model ensembles because of insufficient sampling when observations are restricted to few stations. The occurrence for each area is calculated by averaging all grid points in that box.

In general, and consistent with previous studies (e.g. Neelin et al, 2009; Sahany et al, 2014), the occurrence of IWV for precipitating days features a peak around IWV_{cv} and models capture the shape of distribution fairly well compared to observations. The shapes of the distributions look quite similar between the parametrized and explicit convection ensemble with very few differences between temperature bins below IWV_{cv} - except above the Mediterranean Sea - and larger differences between temperature bins after IWV_{cv}, with a more abrupt decrease of occurrence at high temperature than at low temperature. However, we can note some differences between the two ensembles: over the high mountain and the plains, maximum of IWV counts for precipitation days occurs when IWV exceeds IWV_{cv} in ALP-11 simulations, while without convection parameterization schemes, it is the highest around IWV_{cv}. For both ensembles, after the highest frequency reached, IWV occurrence decreases at a higher rate for higher temperature bins but the decrease is sharper in ALP3remapEUR11 than in ALP-11 simulations. The fact that the maximum of IWV count for precipitation days occurs at higher rescaled IWV values and that the rate of decrease after the maximum is weaker in ALP-11 ensemble suggests weaker dissipative effects when convection is parametrized. With convection schemes in

the ALP-11 ensemble, the models potentially increase occurrence of convective precipitation that occurs primarily above criticality.

Over the Mediterranean the distributions peak at IWV_{cv} for both ensembles and all temperature bins, but they are much sharper than over the other domains. Important differences appear between temperature bins, with flatter distribution at higher temperature bins in both ensembles but much more pronounced in ALP3remapEUR11 ensemble. This flattening is the result of strong saturation deficit (Seager et al., 2015) at higher temperatures without triggering effects over the sea, so we do not have enough precipitating days to plot the distribution for the highest temperature bins in ALP3remapEUR11 ensemble. More specifically, over the sea, more precipitation occurs at lower IWV_{cv}, and the asymmetry of the distribution below and above IWV_{cv} is the opposite to the one exhibited over mountains or plains. Over the sea, precipitation triggering is determined by large-scale dynamics operating on longer time scales, so the frequency of occurrence is higher below IWV_{cv} and is not constrained by loss of water vapor by precipitation processes. In contrast, convective precipitation processes are determined by short time and small space scales over the land, at higher temperatures when the critical value and the associated precipitation pickup are a property of the convection itself, depending only on the tropospheric water vapor and temperature. So IWV_{cv} plays an important role for precipitation triggering and a larger fraction of precipitation occurs around IWV_{cv}.

5. Concluding Remarks

This work uses different model ensembles from EURO, Med-CORDEX and CORDEX FPS –« Convective phenomena at high resolution over Europe and the Mediterranean » to investigate the impact of model resolution and convection representation (parameterized versus explicit) in simulating precipitation frequency over Europe and the Mediterranean. We investigate the differences between the model ensembles by focusing on triggering processes. In this study, precipitation is considered in the relationship with tropospheric temperature and IWV, which has been shown to be a better proxy of deep convection onset than saturation over tropical oceans (e.g Neelin et al., 2009; Sahany et al., 2011) as well as over Europe (Bastin et al., 2019).

The results indicate that with higher resolution, models simulate fewer precipitating days in both summer and winter. Precipitation frequency is reduced in CPMs without convection parameterization schemes, and even becomes lower than the observations (other simulations overestimate precipitation frequency) in summer. This result has been noticed in previous studies (e.g Ban et al, 2021; Berthou et al, 2018) but here we show that the less frequent occurrence of precipitation is not only explained by higher temperatures in CPMs (e.g Sangelantoni et al., submitted) but also by a higher critical value of IWV (IWV_{cv}) that must be reached to trigger precipitation at similar temperatures, and a lower probability of exceeding this critical value. Moreover, in CPM simulations, it is shown that the occurrence of precipitation is more peaked around the critical value, indicating stronger dissipative effects than when convection is parametrized. This reduces the occurrence of rainfall and this effect will likely be more pronounced when we will consider hourly time scales in the next study. Finally, the spread between models in simulating IWV_{cv} and its exceedance probability is reduced in the ensemble of models with explicit convection over land, especially at higher temperatures, when fraction of convective precipitation to total precipitation becomes more important. This is likely because inherent assumptions in convection parameterization schemes (different expressions of entrainment and convective updrafts) alter the statistics for the transition to deep convection (Kuo et al., 2017; Tompkins, 2001).

Considering the Alpine mountains, model resolution has a larger impact than over other areas in simulating IWV_{cv} because dynamical processes linked to orography play an important role in triggering convection. However, we cannot completely exclude the influence of convection parameterization schemes. Over the plains, both model resolution and convection representation affect the precipitation triggering, especially at higher temperatures. Some studies have shown that in CPMs, land surface/atmosphere interactions are modified compared to RCMs with drier soils and weaker latent heat fluxes in most models (Sangelantoni et al., submitted). This explains the higher temperatures as well as the lack of moisture in the boundary layer making the deficit to saturation more difficult to sustain. The moistening of the lower free troposphere is then more difficult and conditional instability that yields the onset of convection occurs at higher values of IWV. In contrast to the land, over the sea, precipitation triggering depends on large scale convergence which is the main process allowing lifting of the particles, but the saturation deficit increases with temperature and becomes much more difficult to break through in the higher temperature bins. This effect is increased in the ensemble with explicit convection, while surface humidity is not a limiting factor. Additional information about vertical structure of humidity and temperature would help in the understanding of these differences, in particular to differentiate between the impacts of the boundary layer and lower free-troposphere.

In this study, the impact of convection representation (explicit versus parameterized) cannot be assessed fully because of the lack of simulations without convective schemes at 0.11° resolution. For the first time and very recently, an ensemble of climate simulations using explicit convection at 0.11° has been performed and the analysis of these simulations will help to disentangle what comes from the resolution and what comes from the convection representation.

Finally, the daily time scale is considered because of the availability of data in EURO- and Med-CORDEX ensembles. This is not the best time scale for addressing the rapid processes in convective events, which often occur over much shorter periods (Schiro et al, 2016). The next step will be to assess the processes of triggering at hourly time scales and to use an ensemble of climate projections to investigate the evolution of these convective transition processes and of precipitation distribution in a warming climate.

Declarations

Acknowledgments:

All authors gratefully acknowledge the WCRP-CORDEX-FPS on Convective phenomena at high resolution over Europe and the Mediterranean (FPSCONVALP- 3) and the research data exchange infrastructure and services provided by the Jülich Supercomputing Centre, Germany, as part of the Helmholtz Data Federation initiative. To process the data, this study benefited from the IPSL mesocenter ESPRI facility which is supported by CNRS, UPMC, Labex L-IPSL, CNES and EcolePolytechnique, and received funding from the HORIZON 2020 EUCP (European Climate Prediction System) project (<https://www.eucp-project.eu>, grant agreement No. 776613). IPSL, CNRM and SMHI also acknowledge funding from the HORIZON 2020 EUCP. IPSL's simulation was granted access to the HPC resources of IDRIS under the allocations of the project 0227 " Fonctionnement de la Mediterranee : circulation et ecosystemes" and HPC resources of TGCC under the allocation A0090106877 made by GENCI. The GPS and SIRTAReOBS datasets are maintained by the French national center for Atmospheric data and services AERIS and Research Infrastructure ACTRIS-FR. J.M. acknowledges the support

of the Spanish Government through the Agencia Estatal de Investigación (project PID2019-111481RB-I00 and Unidad de excelencia María de Maeztu" MDM-2017-0765). Ø.H. has received support from the project GREAT, funded by the Research Council of Norway (grant no. 275589), and acknowledge computing resources from Notur (NN9188K). AUTH simulations were supported by computational time granted from the National Infrastructures for Research and Technology S.A. (GRNET S.A.) in the National HPC facility - ARIS - under project pr009020_thin. UCAN simulations have been carried out on the Altamira Supercomputer at the Instituto de Física de Cantabria (IFCA-CSIC), member of the Spanish Supercomputing Network. JF acknowledges support from project CORDyS (PID2020-116595RB-I00) funded by MCIN/AEI/10.13039/501100011033. K.G. acknowledges computing time granted by the John von Neumann Institute for Computing (NIC) and through JARA-HPC on the supercomputer JURECA at Jülich Supercomputing Centre (JSC).

Author contributions

The analysis has been performed by Ha T-M, with contribution from Bastin S. and Drobinski P. and datasets from all the other authors. All authors have read, given feedbacks to improve the paper and agreed to the published version of the manuscript.

Data Availability

Output from EURO-CORDEX simulations are available at <https://esgf-node.ipsl.upmc.fr/search/cordex-ipsi>; Output from Med-CORDEX simulations are available at <https://www.medcordex.eu/search/index.php>; FPSCONV simulations during the current study are not publicly available but are available from the co-authors on reasonable request; ERA5 tropospheric temperature is available at <https://www.ecmwf.int/en/forecasts/datasets/reanalysis-datasets/era5>; SIRTAReOBS, GNSS IWV and gridded precipitation are provided by co-authors.

References

1. Adinolfi, M., Raffa, M., Reder, A., & Mercogliano, P. (2021). Evaluation and expected changes of summer precipitation at convection permitting scale with COSMO-CLM over alpine space. *Atmosphere*, 12, 54. <https://doi.org/10.3390/atmos1201005>
2. Ahrens, B., Meier, T., & Brisson, E. (2020). Diurnal cycle of precipitation in the Himalayan foothills—Observations and model results. In A. Dimri, B. Bookhagen, M. Stoffel, & T. Yasunari (Eds.), *Himalayan weather and climate and their impact on the environment*. Springer. https://doi.org/10.1007/978-3-030-29684-1_5
3. Argüeso, D., Romero, R., & Homar, V. (2020). Precipitation features of the maritime continent in parameterized and explicit convection models. *Journal of Climate*, 33, 2449–2466. <https://doi.org/10.1175/JCLI-D-19-0416.1>
4. Baldauf M, Seifert A, Förstner J, Majewski D, Raschendorfer M, Reinhardt T (2011) Operational convection-scale numerical weather prediction with the COSMO model: description and sensitivities. *Mon Weather Rev* 139:3887–3905
5. Ban, N., Caillaud, C., Coppola, E. et al. The first multi-model ensemble of regional climate simulations at kilometer-scale resolution, part I: evaluation of precipitation. *ClimDyn* 57, 275–302 (2021).

<https://doi.org/10.1007/s00382-021-05708-w>

6. Ban N, Schmidli J, Schär C (2014) Evaluation of the convection-resolving regional climate modeling approach in decade-long simulations. *J Geophys Res* 119(13):7889–7907.
<https://doi.org/10.1002/2014JD021478>
7. Bastin, S., Chiriaco, M., and Drobinski, P. (2018): Control of radiation and evaporation on temperature variability in a WRF regional climate simulation: comparison with colocated long-term ground based observations near Paris, *Clim. Dynam.*, 51, 985–1003, <https://doi.org/10.1007/s00382-016-2974-1>
8. Bastin S., Drobinski P. (2006): Sea breeze induced mass transport over complex terrain in southeastern France: a case study. *Q J R Meteorol Soc* 132:405–423
9. Bastin, S., Drobinski, P., Chiriaco, M., Bock, O., Roehrig, R., Gallardo, C., Conte, D., Domínguez Alonso, M., Li, L., Lionello, P., and Parracho, A. C. (2019): Impact of humidity biases on light precipitation occurrence: observations versus simulations, *Atmos. Chem. Phys.*, 19, 1471–1490, <https://doi.org/10.5194/acp-19-1471-2019>
10. Bauer, H.-S., T. Weusthoff, M. Dorninger, V. Wulfmeyer, T. Schwitalla, T. Gorgas, M. Arpagaus, and K. Warrach-Sagi, 2011: Predictive skill of a subset of the D-PHASE multi-model ensemble in the COPS region. *Q. J. Roy. Meteor. Soc.* 137(S1), 287-305. DOI:10.1002/qj.715
11. Belušić D, de Vries H, Dobler A, Landgren O, Lind P, Lindstedt D, Pedersen RA, Sánchez-Perrino JC, Toivonen E, van Ulft B, Wang F, Andrae U, Batrak Y, Kjellström E, Lenderink G, Nikulin G, Pietikäinen JP, Rodríguez-Camino E, Samuelsson P, van Meijgaard E, Wu M (2020) HCLIM38: A flexible regional climate model applicable for different climate zones from coarse to convection permitting scales. *Geosci Model Dev* 13:1311–1333. <https://doi.org/10.5194/gmd-13-1311-2020>
12. Berthou S, Kendon EJ, Chan SC, Ban N, Leutwyler D, Schär C, Fosser G (2018) Pan-European climate at convection-permitting scale: a model intercomparison study. *ClimDyn*. <https://doi.org/10.1007/s00382-018-4114-6>
13. Berthou, S., Rowell, D.P., Kendon, E.J. et al. Improved climatological precipitation characteristics over West Africa at convection-permitting scales. *ClimDyn* 53, 1991–2011 (2019). <https://doi.org/10.1007/s00382-019-04759-4>
14. Boberg F, Berg P, Thejll P, Gutowski WJ, Christensen JH (2009) Improved confidence in climate change projections of precipitation evaluated using daily statistics from the PRUDENCE ensemble. *ClimDyn* 32:1097–11
15. Bock, O. (2021). GNSS Integrated Water Vapour data over France and Europe, for EURO- and Med-CORDEX. [Data set]. AERIS. <https://doi.org/10.25326/294>
16. Brockhaus, P., D. Lüthi, and C. Schär (2008), Aspects of the diurnal cycle in a regional climate model, *Meteorol. Z.*, 17, 433–443
17. Brown, J.R., Jakob, C. and Haynes, J.M. 2010. An evaluation of rainfall frequency and intensity over the Australian region in a global climate model. *J. Clim.*, 23, 6504–25
18. Caillaud, C., Somot, S., Alias, A., Bernard-Bouissières, I., Fumière, Q., Laurantin, O., Seity, Y., Ducrocq, V. (2021) Modelling Mediterranean heavy precipitation events at climate scale: an object-oriented evaluation of the CNRM-AROME convection-permitting regional climate model. *ClimDyn*, 56, 1717–1752

19. Chiriaco, M., Bastin, S., Yiou, P., Haefelin, M., Dupont, J.-C., Klenov, L., and Stéfanon, M.: European heat-wave in July 2006: observations and modelling showing how local processes amplify conducive large-scale conditions, *Geophys. Res. Lett.*, 41, 5644–5652, 2014
20. Chiriaco, M., Dupont, J.-C., Bastin, S., Badosa, J., Lopez, J., Haefelin, M., Chepfer, H., and Guzman, R. (2018): ReOBS: a new approach to synthesize long-term multi-variable dataset and application to the SIRTAsupersite, *Earth Syst. Sci. Data*, 10, 919–940, <https://doi.org/10.5194/essd-10-919-2018>
21. Coppola E, Sobolowski S, Pichelli E, Rafaele F, Ahrens B, Anders I, Ban N, Bastin S, Belda M, Belusic D et al (2020) A first-of-its-kind multi-model convection permitting ensemble for investigating convective phenomena over Europe and the Mediterranean. *ClimDyn*. <https://doi.org/10.1007/s00382-018-4521-8>
22. Dai, A., 2006: Precipitation characteristics in eighteen coupled climate models. *J. Climate*, 19, 4605–4630, doi:10.1175/JCLI3884.1
23. Dai, A., and K. E. Trenberth (2004), The diurnal cycle and its depiction in the community climate system model, *J. Clim.*, 17, 930–951
24. Dickman, R., A. Vespignani, and S. Zapperi, 1998: Self-organized criticality as an absorbing-state phase transition. *Phys. Rev. E*, 57, 5095–5105
25. Feser, F., and H. von Storch, 2005: A spatial two-dimensional discrete filter for limited area model evaluation purposes. *Mon. Wea. Rev.*, 133, 1774–1786
26. Foley, A. M. (2010). Uncertainty in regional climate modelling: A review. *Progress in Physical Geography*, 34, 647–670. <https://doi.org/10.1177/0309133310375654>
27. Fosser, G., Khodayar, S., & Berg, P. (2015). Benefit of convection permitting climate model simulations in the representation of convective precipitation. *Climate Dynamics*, 44(1–2), 45–60. <https://doi.org/10.1007/s00382-014-2242-1>
28. Froidevaux, P., L. Schlemmer, J. Schmidli, W. Langhans, and C. Schär (2014), Influence of the background wind on the local soil moisture-precipitation feedback, *J. Atmos. Sci.*, 71, 782–799
29. Fumière Q, Déqué M, Nuissier O, Somot S, Alias A, Caillaud C, Laurantin O, Seity Y (2020) Extreme rainfall in Mediterranean France during the fall: added value of the CNRM-AROME convection-permitting regional climate model. *ClimDyn*. <https://doi.org/10.1007/s00382-019-04898-8>
30. Giorgi F (2019) Thirty years of regional climate modeling: Where are we and where are we going next? *J Geophys Res Atmos* 124(11):5696–5723. <https://doi.org/10.1029/2018JD030094>. <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2018JD030094>
31. Giorgi, F., Torma, C., Coppola, E., Ban, N., Schär, C., & Somot, S. (2016). Enhanced summer convective rainfall at Alpine high elevations in response to climate warming. *Nature Geoscience*, 9, 584–589. <https://doi.org/10.1038/ngeo2761>
32. Guo, Z., Fang, J., Sun, X., Tang, J., Yang, Y., & Tang, J. (2020). Decadal long convection-permitting regional climate simulations over eastern China: Evaluation of diurnal cycle of precipitation. *Climate Dynamics*, 54, 1329–1349. <https://doi.org/10.1007/s00382-019-05061-z>
33. Guo, Z., Fang, J., Sun, X., Yang, Y., & Tang, J. (2019). Sensitivity of summer precipitation simulation to microphysics parameterization over eastern China: Convection-permitting regional climate simulation.

34. Haefelin, M., Barthès, L., Bock, O., Boitel, C., Bony, S., Bouniol, D., Chepfer, H., Chiriaco, M., Cuesta, J., Delanoë, J., Drobinski, P., Dufresne, J.-L., Flamant, C., Grall, M., Hodzic, A., Hourdin, F., Lapouge, F., Lemaître, Y., Mathieu, A., Morille, Y., Naud, C., Noël, V., O'Hirok, W., Pelon, J., Pietras, C., Protat, A., Romand, B., Scialom, G., and Vautard, R. (2005): SIRTa, a ground-based atmospheric observatory for cloud and aerosol research, *Ann. Geophys.*, 23, 253–275, <https://doi.org/10.5194/angeo-23-253-2005>
35. Hersbach, H.; Bell, B.; Berrisford, P.; Horányi, A.; Sabater, J.M.; Nicolas, J.; Radu, R.; Schepers, D.; Simmons, A.; Soci, C.; et al. Global Reanalysis: Goodbye ERA-Interim, Hello ERA5. ECMWF Newsl. 2019. Available online: <https://www.ecmwf.int/en/newsletter/159/meteorology/global-reanalysis-goodbye-era-interim-hello-era5>
36. Hohenegger, C., P. Brockhaus, and C. Schar, 2008: Towards climate simulations at cloud-resolving scales. *Meteor. Z.*, 17, 383–394
37. Hohenegger, C., P. Brockhaus, C. Bretherton, and C. Schär (2009), The soil moisture-precipitation feedback in simulations with explicit and parameterized convection, *J. Clim.*, 22, 5003–5020
38. Hodnebrog, Ø., Marelle, L., Alterskjær, K., Wood, R. R., Ludwig, R., Fischer, E. M., Richardson, T. B., Forster, P. M., Sillmann, J., and Myhre, G.: Intensification of summer precipitation with shorter time-scales in Europe, *Environmental Research Letters*, 14, 12, 124050, [10.1088/1748-9326/ab549c](https://doi.org/10.1088/1748-9326/ab549c), 2019
39. IPCC, 2021: Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change [Masson-Delmotte, V., P. Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T.K. Maycock, T. Waterfield, O. Yelekçi, R. Yu, and B. Zhou (eds.)]. Cambridge University Press. In Press.
40. Isotta F, Frei C, Weilguni V, Tadić MP, Lasségués P, Rudolf B, Pavan V, Cacciamani C, Antolini G, Ratto SM, Munari M, Micheletti S, Bonati V, Lussana C, Ronchi C, Panettieri E, Marigo G, Vertačnik G (2014) The climate of daily precipitation in the Alps: development and analysis of a high-resolution grid dataset from pan-Alpine rain-gauge data. *Int J Climatol* 34(5):1657–1675. <https://doi.org/10.1002/joc.3794>
41. Jacob D, Petersen J, Eggert B, Alias A, Christensen OB, Bouwer LM, Braun A, Colette A, Déqué M, Georgievski G, Georgopoulou E, Gobiet A, Menut L, Nikulin G, Haensler A, Hempelmann N, Jones C, Keuler K, Kovats S, Kröner N, Kotlarski S, Kriegsman A, Martin E, van Meijgaard E, Moseley C, Pfeifer S, Preuschmann S, Radermacher C, Radtke K, Rechid D, Rounsevell M, Samuelsson P, Somot S, Soussana JF, Teichmann C, Valentini R, Vautard R, Weber B, Yiou P (2014) Euro-cordex: new high-resolution climate change projections for european impact research. *Reg Environ Change* 14(2):563–578. <https://doi.org/10.1007/s10113-013-0499-2>
42. Keller, M., Fuhrer, O., Schmidli, J., Stengel, M., Stöckli, R., & Schär, C. (2016). Evaluation of convection-resolving models using satellite data: The diurnal cycle of summer convection over the Alps. *Meteorologische Zeitschrift*, 25, 165–179. <https://doi.org/10.1127/metz/2015/0715>
43. Kendon EJ, Ban N, Roberts NM, Fowler HJ, Roberts MJ, Chan SC, Evans JP, Fosse G, Wilkinson JM (2017) Do convection-permitting regional climate models improve projections of future precipitation change? *Bull Am Meteor Soc* 98(1):79–93. <https://doi.org/10.1175/BAMS-D-15-0004.1>

44. Kendon EJ, Roberts NM, Senior CA, Roberts MJ (2012) Realism of rainfall in a very high-resolution regional climate model. *J Clim* 25(17):5791–5806. <https://doi.org/10.1175/JCLI-D-11-00562.1>
45. Kendon EJ, Roberts NM, Fowler HJ, Roberts MJ, Chan SC, Senior CA (2014) Heavier summer downpours with climate change revealed by weather forecast resolution model. *Nat Clim Change* 4:570– 576. <https://doi.org/10.1038/nclimate2258>
46. Kendon, E.J., Stratton, R.A., Tucker, S. et al. Enhanced future changes in wet and dry extremes over Africa at convection-permitting scale. *Nat Commun* 10, 1794 (2019). <https://doi.org/10.1038/s41467-019-09776-9>
47. Kjellström E, Boberg F, Castro M, Christensen JH, Nikulin G, Sánchez E (2010) Daily and monthly temperature and precipitation statistics as performance indicators for regional climate models. *Clim Res* 44:135–150. <https://doi.org/10.3354/cr00932>
48. Knist, S., Goergen, K., & Simmer, C. (2020). Effects of land surface inhomogeneity on convection-permitting WRF simulations over Central Europe. *Meteorology and Atmospheric Physics*, 132, 53–69. <https://doi.org/10.1007/s00703-019-00671-y>
49. Konduru, R. T., & Takahashi, H. G. (2020). Effects of convection representation and model resolution on diurnal precipitation cycle over the Indian monsoon region: Toward a convection-permitting regional climate simulation. *Journal of Geophysical Research: Atmospheres*, 125, e2019JD032150. <https://doi.org/10.1029/2019JD032150>
50. Kouadio, K., Bastin, S., Konare, A. and Ajayi, V.O. (2020) Does convection-permitting simulate better rainfall distribution and extreme over Guinean coast and surroundings? *Clim. Dyn.* 55, pp153-174. doi :10.1007/s00382-018-4308-y. (insu-01821831)
51. Kuo Y.-H., Neelin D. J., Mechoso R. C., 2017: Tropical Convective Transition Statistics and causality in the water-vapor-precipitation relation. doi:10.1175/JAS-D-16-0182.1
52. Langhans, W., Schmidli, J., Fuhrer, O., Bieri, S., & Schär, C. (2013). Long-term simulations of thermally driven flows and orographic convection at convection-parameterizing and cloud-resolving resolutions. *Journal of Applied Meteorology and Climatology*, 52, 1490–1510. <https://doi.org/10.1175/JAMC-D-12-0167.1>
53. Lavin-Gullon A., Fernandez J., Bastin S., Cardoso R. M., Fita L., Giannaros T., Goergen K., Gutierrez J. M., Kartsios S., Katragkou E., Lorenz T., Milovac J., Sobolowski S., Soares P.M., Warrach-Sagi K., 2020: Internal variability vs multi-physics uncertainty in a regional climate model. *Int. Jour. of Clim.* Doi : 10.1002/joc.6717. insu-02885105
54. Lenderink G and van Meijgaard E 2008 Increase in hourlyprecipitation extremes beyond expectations from temperaturechanges*Nat. Geosci.*1511–4
55. Leutwyler D, Lüthi D, Ban N, Fuhrer O, Schär C (2017) Evaluation of the convection-resolving climate modeling approach on continental scales. *J Geophys Res Atmos.* <https://doi.org/10.1002/2016JD026013>
56. Lind, P, Lindstedt, D., Kjellström, E., & Jones, C. (2016). Spatial and temporal characteristics of summer precipitation over central Europe in a suite of high-resolution climate models. *Journal of Climate*, 29, 3501–3518. <https://doi.org/10.1175/JCLI-D-15-0463.1>

57. Liu C, Ikeda K, Rasmussen R, Barlage M, Newman AJ, Prein AF, Chen F, Chen L, Clark M, Dai A, Dudhia J, Eidhammer T, Gochis D, Gutmann E, Kurkute S, Li Y, Thompson G, Yates D (2017) Continental-scale convection-permitting modeling of the current and future climate of North America. *ClimDyn* 49(1):71–95. <https://doi.org/10.1007/s00382-016-3327-9>
58. Lucas, C., Zipser, E. J., & Lemone, M. A. (1994). Vertical velocity in oceanic convection off tropical Australia. *Journal of the Atmospheric Sciences*, 51, 3183–3193. [https://doi.org/10.1175/1520-0450\(2001\)040<1075:ACASFI>2.0.CO;2](https://doi.org/10.1175/1520-0450(2001)040<1075:ACASFI>2.0.CO;2)
59. Lucas-Picher P, Argüeso D., Brisson E., Trambly Y., Berg P, Lemonsu A, Kotlarski S., Caillaud C.: Convection-permitting modeling with regional climate models: Latest developments and next steps. *Wiley Interdisciplinary Reviews Climate Change*. <https://doi.org/10.1002/wcc.731>
60. Lundquist J, Hughes M, Gutmann E, Kapnick S (2020) Our skill in modeling mountain rain and snow is bypassing the skill of our observational networks. *Bull Am Meteorol Soc* 100(12):2473– 2490. <https://doi.org/10.1175/BAMS-D-19-0001.1>
61. Lüthi, S., Ban, N., Kotlarski, S., Steger, C. R., Jonas, T., & Schär, C. (2019). Projections of alpine snow cover in a high-resolution climate simulation. *Atmosphere*, 10(8), 463. <https://doi.org/10.3390/atmos1008046>
62. Nabat, P., Somot, S., Cassou, C., Mallet, M., Michou, M., Bouniol, D., Decharme, B., Drugé, T., Roehrig, R., and Saint-Martin, D. (2020) Modulation of radiative aerosols effects by atmospheric circulation over the Euro-Mediterranean region, *Atmos. Chem. Phys.*, 20, 8315–8349, <https://doi.org/10.5194/acp-20-8315-2020>
63. Neelin, J. D., O. Peters, and K. Hales, 2009: The transition to strong convection. *J. Atmos. Sci.*, 66, 2367–2384, doi:10.1175/ 2009JAS2962.1
64. Ning T, Elgered G, Willén U, Johansson JM (2013) Evaluation of the atmospheric water vapor content in a regional climate model using ground-based GPS measurements. *J Geophys Res Atmos* 118(2):329–339
65. Piazza, M., Prein, A. F., Truhetz, H., & Csaki, A. (2019). On the sensitivity of precipitation in convection-permitting climate simulations in the Eastern Alpine region. *Meteorologische Zeitschrift*, 28(4), 323–346. <https://doi.org/10.1127/metz/2019/0941>
66. Prein AF, Langhans W, Fosser G, Ferrone A, Ban N, Goergen K, Keller M, Tölle M, Gutjahr O, Feser F, Brisson E, Kollet S, Schmidli J, van Lipzig NPM, Leung R (2015) A review on regional convection-permitting climate modeling: demonstrations, prospects, and challenges. *Rev Geophys* 53(2):323–361. <https://doi.org/10.1002/2014RG000475>
67. Prein, A. F., Gobiet, A., Suklitsch, M., Truhetz, H., Awan, N. K., Keuler, K., & Georgievski, G. (2013). Added value of convection permitting seasonal simulations. *Climate Dynamics*, 41, 2655–2677. <https://doi.org/10.1007/s00382-013-1744-6>
68. Quintana Seguí, P., P. Le Moigne, Y. Durand, E. Martin, F. Habets, M. Baillon, L. Franchisteguy, S. Morel, and J. Noilhan (2008), Analysis of near-surface atmospheric variables: Validation of the SAFRAN analysis over France, *J. Appl. Meteorol. Climatol.*, 47(1), 92– 107
69. Rasmussen KL, Prein AF, Rasmussen RM, Ikeda K, Liu C (2017) Changes in the convective population and thermodynamic environments in convection-permitting regional climate simulations over the United States. *ClimDyn* 1–26
70. Rasmussen RM et al (2011) High-resolution coupled climate runoff simulations of seasonal snowfall over Colorado: a process study of current and warmer climate. *J Clim* 24:3015–3048

71. Rasmussen RM et al (2014b) Climate change impacts on the water balance of the Colorado headwaters: high-resolution regional climate model simulations. *J Hydrometeorol* 15:1091–1116
72. Rauthe, M., Steiner, H., Riediger, U., Mazurkiewicz, A. and Gratzki, A. (2013) A central European precipitation climatol-ogy? Part I: generation and validation of a high-resolution gridded daily data set (HYRAS). *Meteorologische Zeitschrift*, 22, 235–256. <https://doi.org/10.1127/0941-2948/2013/0436>
73. Reder, A., Raffa, M., Montesarchio, M., & Mercogliano, P. (2020). Performance evaluation of regional climate model simulations at different spatial and temporal scales over the complex orography area of the Alpine region. *Natural Hazards*, 102, 151–177. <https://doi.org/10.1007/s11069-020-03916-x>
74. Rockel B, Will A, Hense A (2008) The regional climate model COSMO-CLM (CCLM). *Meteorol Z* 17:347–348. <https://doi.org/10.1127/0941-2948/2008/0309>
75. Ruti PM, Somot S, Giorgi F, Dubois C, Flaounas E, Obermann A, Dell'Aquila A, Pisacane G, Harzallah A, Lombardi E, Ahrens B, Akhtar N, Alias A, Arsouze T, Aznar R, Bastin S, Bartholy J, Béranger K, Beuvier J, Bouffies-Cloch S, Brauch J, Cabos W, Calmanti S, Calvet JC, Carillo A, Conte D, Coppola E, Djurdjevic V, Drobinski P, Elizalde-Arellano A, Gaertner M, Galn P, Gallardo C, Gualdi S, Goncalves M, Jorba O, Jord G, L'Heveder B, Lebeaupin-Brossier C, Li L, Liguori G, Lionello P, Macis D, Nabat P, nol B, Raikovic B, Ramage K, Sevault F, Sannino G, Struglia MV, Sanna A, Torma C, Vervatis V (2016) Med-cordex initiative for mediterranean climate studies. *Bull Am Meteorol Soc* 97(7):1187–1208. <https://doi.org/10.1175/BAMS-D-14-00176.1>
76. Sahany, S., J. D. Neelin, K. Hales, and R. B. Neale, 2012: Temperature–moisture dependence of the deep convective transition as a constraint on entrainment in climate models. *J. Atmos. Sci.*, 69, 1340–1358, doi:10.1175/JAS-D-11-0164.1
77. Sahany, S., J. D. Neelin, K. Hales, and R. B. Neale, 2014: Deep convective transition characteristics in the NCAR CCSM and changes under global warming. *J. Climate*, 27, 9214–9232, doi:10.1175/JCLI-D-13-00747.1
78. Schiro, K. A., Neelin, J. D., Adams, D. K., and Lintner, B. R. (2016): Deep Convection and Column Water Vapor over Tropical Land versus Tropical Ocean: A Comparison between the Amazon and the Tropical Western Pacific, *J. Atmos. Sci.*, 73, 4043-4063, <https://doi.org/10.1175/JAS-D-16-0119.1>
79. Seager, R., A. Hooks, A. P. Williams, B. Cook, J. Nakamura, and N. Henderson (2015), Climatology, variability, and trends in the U.S. Vapor pressure deficit, an important fire-related meteorological quantity, *J. Appl. Meteorol. Climatol.*, 54(6), 1121–1141
80. Skamarock WC (2008) A description of the advanced research WRF Version 3. NCAR Technical Notes NCAR/TN-4751STR. <https://opensky.ucar.edu/islandora/object/technotes%3A500/datastream/PDF/view>
81. Stephens, G. L., and et al., 2010: Dreary state of precipitation in global models. *J. Geophys. Res.*, 115, D24211, <https://doi.org/10.1029/2010JD014532>
82. Sun, Y., S. Solomon, A. Dai, and R. W. Portmann, 2006: How often does it rain? *J. Climate*, 19, 916–934, doi:10.1175/JCLI3672.1
83. Tang, C., and P. Bak, 1988: Critical exponents and scaling relations for self-organized critical phenomena. *Phys. Rev. Lett.*, 60, 2347–2350, doi:10.1103/PhysRevLett.60.2347
84. Tompkins, A. M., 2001: Organization of tropical convection in low vertical wind shears: The role of water vapor. *J. Atmos. Sci.*, 58, 529–545

85. Vidal J.-P., Martin E., Franchistéguy L., Baillon M., Soubeyroux J.-M. A 50-year high-resolution atmospheric reanalysis over France with the Safran system. *Int. J. Climatol.*, 30 (2010), pp. 1627-1644, 10.1002/joc.2003
86. Weisman, M. L., Skamarock, W. C., & Klemp, J. B. (1997). The resolution dependence of explicitly modeled convective systems. *Monthly Weather Review*, 125, 527–548. [https://doi.org/10.1175/1520-0493\(1997\)125<0527:TRDOEM>2.0.CO;2](https://doi.org/10.1175/1520-0493(1997)125<0527:TRDOEM>2.0.CO;2)
87. Winterfeldt, J., and R. Weisse, 2009: Assessment of value added for surface marine wind speed obtained from two regional climate models. *Mon. Wea. Rev.*, 137, 2955–2965
88. Wulfmeyer, V., Muppa, S. K., Behrendt, A., Hammann, E., Spath, F., Sorbjan, Z., Turner, D. D., and Hardesty, R. H.: Determination of Convective Boundary Layer Entrainment Fluxes, Dissipation Rates, and the Molecular Destruction of Variances: Theoretical Description and a Strategy for Its Confirmation with a Novel Li-dar System Synergy, *J. Atmos. Sci.*, 73, 667–692, 2016

Figures

Figure 1

Evaluation domain with the GPS locations (red points) and SIRTAs stations (blue points) used for comparison with simulations. The three black boxes indicate subareas (mountains of the Alps, the Mediterranean Sea, and continental plains) analyzed in more detail in the study.

Figure 2

Precipitation frequency (>0.1 mm/day) in DJF for the different model ensembles and observations: a) EUR-44; b) EUR-11; c) ALP-11; d) ALP-3 remap on EUR-11; e) ALP-3; f) Observation.

Figure 3

Same as Fig.2 but for JJA.

Figure 4

Precipitation frequency (%) as a function of IWV and tropospheric temperature over (a, b) the FPSCONV domain, (c, d) Alpine mountains, (e, f) Mediterranean, and (g, h) plains (see boxes in Fig.1). Ensembles are compared 2 by 2: upper row is for EUR-44 and EUR-11 and lower row is for ALP-11 and ALP-3. Green solid lines show the IWV as a function of tropospheric temperature when it rains from the different stations included in each box. The orange line shows the C-C theoretical saturation humidity as a function of temperature.

Figure 5

IWVcv (kg m^{-2}) from all model's simulations (boxplot), EUR-44 (blue), EUR-11 (green), ALP-11 (pink), ALP-3remapEUR-11 (brown) and ALP-3 (red) over a) FPSCONV domain, b) Mediterranean, c) Alpine mountain and d) plains. The boxplot denotes the 1%, 25%, 50%, 75%, 99% probability of IWVcv to get this value by considering all grid points and all models used in this study; Black stars represent the observations. See more details in the text.

Figure 6

Same as Fig.5 but for probability to exceed IWV_{cv} (%), the diamonds only show the standard deviation for each ensemble and have not been multiplied by 10.

Figure 7

Number of occurrence of IWV rescaled by the corresponding onset threshold values IWV_{cv} for precipitating days between ALP-11 (a), ALP3remapEUR-11 (b) and observation (c) over the FPSCONV domain.

Figure 8

Same as Fig. 7 but for ALP-11 (left), ALP3remapEUR-11 (right) over the Alpine mountain (a, b), Mediterranean (c, d) and plains (e, f), respectively.