Association between egg consumption and arterial stiffness: a longitudinal study

Naiwen Ji

The Pennsylvania State University

Zhe Huang

Kailuan General Hospital

Xinyuan Zhang

The Pennsylvania State University

Yuanyuan Sun

Peking Union Medical College

Shumao Ye

Tufts University

Shuohua Chen

Kailuan General Hospital

Katherine Tucker

University of Massachusetts Lowell

Shouling Wu

Kailuan General Hospital

Xiang Gao (✉ xxg14@psu.edu)

the Pennsylvania State University, University Park <https://orcid.org/0000-0003-2617-6509>

Research

Keywords: Egg consumption, arterial stiffness, cardiovascular disease, longitudinal study

Posted Date: January 8th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-140036/v1>

License: (cc) (i) This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Version of Record: A version of this preprint was published at Nutrition Journal on July 13th, 2021. See the published version at <https://doi.org/10.1186/s12937-021-00720-6>.

Abstract

Background

Inconsistent associations between egg consumption and cardiovascular disease (CVD) risk have been observed in previous studies. This study aims to longitudinally investigate the association between egg consumption and altered risk of arterial stiffness, a major pre-clinical pathogenic change of CVD, which was assessed by brachial-ankle pulse wave velocity (baPWV).

Methods

A total of 7,315 Chinese participants from the Kailuan Study, free of CVD and cancer were included in this study. Egg consumption was assessed by a semi-quantitative validated food frequency questionnaire in 2014. baPWV was repeatedly measured at baseline and during follow-up (mean follow-up: 3.41 years). General linear regression was used to calculate means of baPWV change rate across different egg consumption groups, adjusting for age, sex, baseline baPWV, healthy eating index, total energy, social-economic status, blood pressure, obesity, smoking, lipid profiles, and fasting glucose concentrations.

Results

Compared to the annual baPWV change rate in participants with 0-1.9 eggs/wk (adjusted mean: 35.9 ± 11.2 cm/s/y), those consuming 3-3.9 eggs/wk (adjusted mean: 0.2 ± 11.4 cm/s/y) had the lowest increase in baPWV during follow-up (P-difference = 0.002). Individuals with low (0-1.9 eggs/wk) vs. high (5+ eggs/wk) egg intake showed similar changes in baPWV.

Conclusions

In this large-scale longitudinal analysis, we did not find a significant difference in arterial stiffness, as assessed by baPWV level, between low and high egg consumption groups. However, moderate egg consumption (3-3.9 eggs/wk) appeared to have beneficial effects on arterial stiffness.

Background

Eggs are a type of nutrient-dense food, an excellent source of protein, and a good source of lutein/zeaxanthin, vitamins and minerals. (1) Eggs are also one of the major sources of dietary cholesterol in the human diet. (2) A meta-analysis with 55 studies reported that high dietary cholesterol intake was associated with elevated serum cholesterol. (3) However, inconsistent associations between egg consumption and CVD risk have been observed in previous studies, and no clear association between egg consumption and cardiovascular disease (CVD) risk has been established. Several recent systematic reviews and meta-analyses showed no significant association between egg consumption and CVD mortality (4, 5) or altered status of CVD risk factors. (6, 7) In contrast, a recent pooled analysis of prospective study

reported that each 0.5 egg/wk was associated with 6% higher risk of CVD and 8% higher risk of all-cause mortality.(8)

With the purpose of better understanding the relation between egg consumption and CVD, it is important to understand whether egg intake is associated with altered risk of major pre-clinical pathogenic progressions of CVD. Arterial stiffness is recognized as an important consequence of aging that has been shown to provoke deleterious vascular phenotypes in diseases such as diabetes, atherosclerosis and renal diseases among others.(9, 10) Pulse wave velocity (PWV) is currently considered the gold standard in the evaluation of arterial stiffness. (11) PWV assesses the velocity of the blood pressure wave as it travels a known distance between two anatomic sites within the arterial system, and indicates the elasticity and other properties of the artery. (12) Arterial stiffness, as assessed by Brachial-ankle pulse wave velocity (baPWV), has been shown to be associated with subsequent CVD risk.(13) We, thus, conducted analysis with data from a large-scale longitudinal study, to examine whether egg intake was associated with change in PWV during 3 years of follow-up, in 7,315 adults without CVD.

Methods

Study population

The data utilized are from the Kailuan Study, a population-based prospective cohort launched in 2006 in Tangshan, China (The Chinese Clinical Trial Registry #: ChiCTR-TNRC-1101489). Participants recruited in the study were followed via biennial face-to-face physical examination, anthropometry, laboratory assessment and annual comprehensive surveillance of medical records and death certificates, as detailed previously. (10,14,15)

BaPWV measurement was introduced into the testing routine in 2010 for a sub-cohort of the Kailuan study, as described previously. (10,16) Dietary data were collected in 2014 (baseline for the current analysis), and 8,638 participants had completed dietary information and two baPWV assessments at and after the baseline. We further excluded 336 participants with CVD or cancer and 987 participants with energy intake out of range (for men, < 800 kcal/d or >4000 kcal/d; for women <500 kcal/d or >3500 kcal/d; Figure 1), leaving a total of 7,315 participants in the current analysis. The primary outcome variable was longitudinal change in baPWV (mean follow-up: 3.41 years; interquartile range:1.99-5.13 years). In the secondary analysis for incident arterial stiffness, 3,865 participants without arterial stiffness (baPWV<1,400 cm/s) at baseline were included. The study was performed according to the guidelines of the Helsinki Declaration and was approved by the Ethics Committee of the Kailuan General Hospital.

Assessment of egg consumption and other dietary components

In 2014, a 40-item (33 food items and 7 condiments) semi-quantitative food frequency questionnaire was used to collect possible food items participants consumed, which was validated in Chinese population previously. (17) The food frequency questionnaire was administered along with other assessments during the face-to-face interview. (14) For each food item, participants were asked to provide information regarding the frequency (never, daily, weekly, or monthly) and quantity (liangs, 1 liang =50g) of their consumption.

Specifically, a question regarding egg consumption was included in this food frequency questionnaire: “Based on your recent 1-year diet, do you usually eat egg including chicken egg, duck egg, etc.?” Participants answered this question by selecting following options, “very often, __times/day”, “often, __times/week”, “sometimes, __times/month”, “never” and participants were then asked to provide how many eggs they consumed each time, on average. Based on self-reported egg consumption, we grouped the participants into 5 categories (0-1.9, 2-2.9, 3-3.9, 4-4.9, or ≥ 5 eggs/wk).

Overall diet quality, a potential confounder that might be associated with both egg consumption and arterial stiffness status, was evaluated via a modified Healthy Eating Index 2015 (mHEI-2015) score. The original HEI-2015, based on the 2015-2020 Dietary Guidelines for Americans, (18) includes 13 components: total fruits, whole fruits, total vegetables, greens and beans, whole grains, dairy, total protein foods, seafood and plant proteins, fatty acids, refined grains, sodium, added sugars and saturated fats. (18) Because intakes of greens and beans were not specifically assessed, and fatty acids, added sugar or saturated fat information was not available in Chinese food composition table, these items were excluded from calculation. As a consequence, mHEI-2015 score was based on the remaining 9 components, calculated proportionately according to intake between the minimum and maximum standards. (18) Score ranges from 0(worst) to 65(best). A higher mHEI-2015 score indicates a better diet quality. The Food Pattern Equivalents Database was used to convert grams to cups or ounces during calculation of the HEI-2015 score. (19)

Measurement of arterial stiffness

Arterial stiffness status was assessed at baseline and follow-up (mean 3.41y) using baPWV, which has been used in Asian populations (10,20) and has previously been described.(10,16) Briefly, participants were asked to refrain from smoking and to rest for at least five minutes before the measurement of baPWV. Cuffs were placed on both brachial area of the arm and the ankle to assess the pulse transit time which defined as the travel time of the pulse wave from brachia to ankle. In our study, baPWV was measured with a BP-203RPE networked arteriosclerosis detection device (OMRON Healthcare (China) Co., LTD.). The measurement was repeated twice at each visit to ensure accuracy, and the baPWV value was read by trained health professionals. Therefore, in this study, arterial stiffness was noted using a cut-off point of baPWV ≥ 1400 cm/s. (13,22)

Assessment of covariates

Baseline information was assessed with a validated questionnaire, administered in a face-to-face interview. The questionnaire included basic information (e.g. age, sex, marriage status, education level, smoking status and occupation). Physical activity was assessed with the International Physical Activity Questionnaire (IPAQ) short form, which assesses the duration and frequency of sitting, walking, moderate-intensity activities and vigorous-intensity activities, and has been validated in Chinese population.(23,24)

Height, weight, and blood pressure were measured by trained health professionals, as detailed previously. (10) Body Mass Index was calculated as weight, in kilograms, divided by height, in meters, squared. Blood samples were collected the morning after a minimum of 8 hours of fasting. Fasting blood glucose (FBG),

high-density lipoprotein cholesterol (HDL-c) and low-density lipoprotein cholesterol (LDL-c) were measured in the central lab, Kailuan General Hospital, as previously described. (10)

Statistical analysis

All statistical analyses were performed in SAS version 9.4 (SAS Institute Inc., USA), and all tests were two-sided. In the primary analysis, we compared the rate of longitudinal change in baPWV across egg consumption groups. Change in baPWV was calculated as the difference of baPWV measurements (cm/s) between two visits divided by the time interval (years). General linear model regression was used to calculate adjusted means of baPWV change rate in each group. Model 1 adjusted for age, sex, baseline baPWV, total energy intake and HEI score; model 2 further adjusted for physical activity, marriage, employment, education level, alcohol use, smoking status and heart rate; and model 3 further adjusted for systolic blood pressure, FBG, LDL-c, HDL-c.

We further tested interactions between egg consumption and age, sex, BMI and mHEI score in the fully adjusted model. When significant, subgroup analyses were conducted.

In the secondary analyses, we examined egg consumption and risk of developing arterial stiffness during the follow-up among participants who were free of arterial stiffness at baseline. Cox proportional hazard models were used to calculate hazard ratios (HRs) and 95% confidence intervals (CIs), after adjusting for aforementioned covariates.

Results

Compared to participants with low egg intake (< 2 eggs/ wk), participants with higher egg intake were more likely to have higher mHEI score and higher degree of education, but lower physical activity (Table 1). However, there were no significant differences of baseline characteristics crossing every groups.

Compared to the participants consuming 0-1.9 eggs/wk, participants consuming 3-3.9 eggs/wk had lowest increment in baPWV during follow-up (Table 2). In the final model, the difference in baPWV change rate between this group (adjusted mean: 0.2 ± 11.4 cm/s per year) and the group with consumption of 0-1.9 eggs/wk (35.9 ± 11.2 cm/s per year) was significant (P-difference= 0.002). Similar results were observed when restricting to participants without arterial stiffness at baseline (Table 2). Further adjustment for other blood pressure indices, such as diastolic blood pressure, mean arterial pressure, and pulse pressure, did not materially change observed results (supplementary table 1). There was no significant association between egg consumption and risk of developing arterial stiffness (Table 3). None of the interactions between egg consumption and age, sex, BMI and HEI score, in relation to baPWV change rate were significant (p for interaction > 0.2 for all).

Discussion

In this longitudinal community-based study, individuals with low (<2 eggs/wk) vs. high (5+ eggs /wk) egg intake showed similar changes in baPWV and risk of developing arterial stiffness over a mean 3.41 y follow-

up. Interestingly, those with egg consumption of 3-3.9/wk experienced the slowest aging-related increment in baPWV during the follow-up, among all participants, independent of age, sex, overall diet quality and health status. In this study, we showed an average lower increase in baPWV of 35.7 cm/s per year between the moderate and lowest egg consumption groups, which amounted to ~122 cm/s over 3.41 years of follow-up. The difference in accumulated change may be associated with ~ 8% higher risk of CVD, based on a recent meta-analysis. (25)

To the best of our knowledge, this study is the first to examine the association between egg consumption and arterial stiffness in a community-based population without CVD. The non-significant difference in baPWV status between two extreme egg consumption categories is consistent with previous systematic reviews and meta-analyses regarding egg intake and mortality, CVD risk, and CVD biomarkers. A recent meta-analysis reported a non-significant difference, for high (7+/wk) vs low (<1/wk) egg consumption, for death due to ischemic heart disease (pooled HR=0.97, 95%CI: 0.90, 1.05), ischemic stroke (pooled HR=0.91; 95% CI 0.82, 1.01) and hemorrhagic stroke (pooled HR=0.88; 95%CI: 0.68, 1.14). (5) Similar results were reported in another meta-analysis of egg consumption and total and cause-specific mortality. (4) A systematic review concluded that “high-quality intervention studies have found nonsignificant effects of increasing the consumption of eggs on risk markers for CVD and type 2 diabetes (T2D) in healthy subjects and subjects with T2D” (26). Another two recent meta-analyses of randomized controlled trials also reported non-differential effects of egg consumption (>4/wk) on inflammation (7), blood pressure (6), and lipid profiles (6). In contrast, a recent pooled study based on 6 US cohorts reported that higher egg intake was associated with risk of incident CVD and all-cause mortality.(8) However, there was no significant difference when comparing individuals who consumed <1 eggs/d with those with no egg consumption. For example, the adjusted HR for CVD was 1.06 (95%CI: 0.98, 1.14) for <3.5 eggs/wk, and 1.05 (95% CI: 0.92, 1.20) for 3.5-6.9 eggs/wk, relative to 0 eggs/wk. (8)

We found that participants with modest egg consumption (3-3.9/wk) had significantly lower age-related increment in baPWV across all egg consumption groups. Of note, when the binary variable of arterial stiffness was used as the outcome, a similar trend was observed, but it became non-significant. Several previous studies have reported low risk of CVD and mortality being associated with modest egg consumption. For example, in a recent prospective study including 146,011 individuals from 21 countries, those who consumed 3-4.9 eggs/wk had the lowest risk of mortality and CVD across all egg consumption groups (27). The adjusted HR, comparing 3-4.9 eggs/wk with <1 egg/wk, was 0.89 (95%CI: 0.84, 0.96) for composite outcome (mortality and CVD), and 0.88 (95%CI: 0.80, 0.96) for major CVD. In contrast, the difference in risk of mortality and/or CVD was not significant for 5+eggs /wk vs <1 egg/wk (27). In a prospective study in Sweden, individual consuming 3-6 eggs/wk had the lowest risk for heart failure, myocardial infarction, and hemorrhagic stroke, but not ischemic stroke, across all egg consumption groups, although the difference was not significant (28). A similar non-significant U-shaped trend was also observed for the association between eggs and total mortality (4,5,25, 27).

Arterial stiffness is the stiffening of the blood vessel wall. When the blood vessel loses its elasticity, its ability to control the blood flow is compromised. The blood vessel needs more force to push blood through, which increases the pulse pressure and may introduce damage to the blood vessels. (9) Eggs, as a type of

nutrient-dense food, are an accessible source of L-arginine, vitamin D, vitamin K, choline, and xanthophyll carotenoids, specifically lutein and zeaxanthin (2), which have been shown to be protective against arterial stiffness. The amino acid, L-arginine, is a precursor of nitric oxide. Nitric oxide inhibits the expression of adhesion and further improve endothelial function. (30,31) Xanthophyll carotenoids, lutein and zeaxanthin, and choline have been shown to decrease inflammation. (32–34) However, choline is also a source for trimethylamine-N-oxide, which has been shown to be associated with high risk of atherosclerosis and other cardiovascular events. (35,36) On the other hand, eggs are a major source of dietary cholesterol and saturated fat. (2) A previous meta-analysis showed that high egg consumption was associated with elevated serum cholesterol, (3) a potential risk factor for arterial stiffness. (37,38) High saturated fat intake has also been associated with high aortic pulse wave velocity. (39) Given both positive and negative nutritional components in eggs, it is not surprising that our result, and others, appears to be non-linear, with modest, but not low, egg consumption associated with the desired health outcome. However, further studies are needed to confirm this association.

However, given the lack of consistency in findings, more research is needed. As a majority of existing studies were conducted in the western countries, more investigations at various geographic areas are necessary. This will be beneficial for future meta-analyses.

This study has some limitations. Egg consumption was assessed with a single self-reported question, which may introduce measurement error (misclassification) or recall bias. Undifferentiated misclassification could bias toward the null. In this study, total energy intake was likely to be underestimated due to the small number of food items in the food frequency questionnaire, which might miss some important foods. Thus, total energy intake was adjusted as a covariate in all multiple regression models in order to reduce the influence of this systematic error. Because this study is a preliminary study, the follow-up (mean 3.41 years) is relatively short, which is another limitation. Because the majority of participants were in the 4-4.9 eggs/wk group, the sample size in the extreme categories was small, which may limit the statistical power to detect small-to-moderate effects. The study population is localized, and the observed results may not generalize to other ethnic groups with different eating habits and social economic backgrounds. Only 2406 women were included in the analysis, which is also one of the major limitations.

In conclusion, in this large-scale longitudinal analysis, we did not find a significant difference in arterial stiffness, as assessed by baPWV level, between low and high egg consumption groups (0-1.9/wk vs. ≥ 5 /wk). However, moderate egg consumption (3-3.9 eggs/wk) appeared to have beneficial effects on arterial stiffness. Further studies, investigating association between egg consumption and arterial stiffness with longer follow-up years and more women participants, are warranted to replicate our findings.

Abbreviations

CVD

cardiovascular diseases;

PWV

pulse wave velocity;

baPWV
brachial-ankle pulse wave velocity;
mHEI-2015
modified Healthy Eating Index 2015;
IPAQ
International Physical Activity Questionnaire;
FBG
fasting blood glucose;
HDL-c
high-density lipoprotein cholesterol;
LDL-c
low-density lipoprotein cholesterol;
HRs
hazard ratios;
CIs
confidence intervals;
T2D
type 2 diabetes;
DBP
diastolic blood pressure;
MAP
mean arterial pressure;
PP
pulse pressure.

Declarations

Ethics approval and consent to participate

The data utilized are from the Kailuan Study, a population-based prospective cohort launched in 2006 in Tangshan, China (The Chinese Clinical Trial Registry #: ChiCTR-TNRC-1101489).

Consent for publication

Not applicable

Availability of data and materials

The datasets used and analyzed during the current study are available from the corresponding author on reasonable request.

Competing interests

The authors declare that they have no competing interests.

Funding

The Penn State University and the start-up grant from the college of health and human development and the department of nutritional sciences, Penn State University.

Authors' contributions

NJ and ZH interpreted the analyzed data regarding the egg consumption and baPWV and were major contributors in writing the manuscript. XZ conducted the statistical analysis. All authors read and approved the final manuscript.

Acknowledgements

Not applicable

References

- [1] Dietary Guidelines for Americans 2015-2020. U.S. Department of Health and Human Services and U.S. Department of Agriculture, <https://health.gov/dietaryguidelines/2015/>; 2015 [accessed 17 August 2020].
- [2] US Department of Agriculture. Agricultural Research Service, Nutrient Data Laboratory. USDA National Nutrient Database for Standard Reference, Release 28; 2015. <https://ndb.nal.usda.gov/ndb/>; 2015 [accessed 17 August 2020].
- [3] Vincent MJ, Allen B, Palacios OM, Haber LT, Maki KC. Meta-regression analysis of the effects of dietary cholesterol intake on LDL and HDL cholesterol. *Am J Clin Nutr.* 2019;109:7-16. <https://doi.org/10.1093/ajcn/nqy273>.
- [4] Mazidi M, Katsiki N, Mikhailidis DP, Pencina MJ, Banach M. Egg Consumption and Risk of Total and Cause-Specific Mortality: An Individual-Based Cohort Study and Pooling Prospective Studies on Behalf of the Lipid and Blood Pressure Meta-Analysis Collaboration (LBPMC) Group. *J Am Coll Nutr.* 2019;38:552-63. <https://doi.org/10.1080/07315724.2018.1534620>.
- [5] Xu L, Lam TH, Jiang CQ, et al. Egg consumption and the risk of cardiovascular disease and all-cause mortality: Guangzhou Biobank Cohort Study and meta-analyses. *Eur J Nutr.* 2019;58:785-96. <https://doi.org/10.1007/s00394-018-1692-3>.
- [6] Wang MX, Wong CH, Kim JE. Impact of whole egg intake on blood pressure, lipids and lipoproteins in middle-aged and older population: A systematic review and meta-analysis of randomized controlled trials. *Nutr Metab Cardiovasc Dis.* 2019;29:653-64. <https://doi.org/10.1016/j.numecd.2019.04.004>.
- [7] Hezaveh ZS, Sikaroudi MK, Vafa M, Clayton S, Soltani S, Soltani S. Effect of egg consumption on inflammatory markers: a systematic review and meta-analysis of randomized controlled clinical trials. *J Sci Food Agric.* 2019;99:6663-70. <https://doi.org/10.1002/jsfa.9903>.

- [8] Zhong VW, Van Horn L, Cornelis MC, et al. Associations of Dietary Cholesterol or Egg Consumption With Incident Cardiovascular Disease and Mortality. *JAMA*. 2019;321:1081-95. <https://doi.org/10.1001/jama.2019.1572>.
- [9] Shirwany NA, Zou MH. Arterial stiffness: a brief review. *Acta Pharmacol Sin*. 2010;31:1267-76. <https://doi.org/10.1038/aps.2010.123>.
- [10] Wu S, Jin C, Li S, et al. Aging, Arterial Stiffness, and Blood Pressure Association in Chinese Adults. *Hypertension*. 2019;73:893-9. <https://doi.org/10.1161/HYPERTENSIONAHA.118.12396>.
- [11] Mancia G, Fagard R, Narkiewicz K, et al. 2013 ESH/ESC Guidelines for the management of arterial hypertension: the Task Force for the management of arterial hypertension of the European Society of Hypertension (ESH) and of the European Society of Cardiology (ESC). *J Hypertens*. 2013;31:1281-357. <https://doi.org/10.1097/01.hjh.0000431740.32696.cc>.
- [12] Miyatani M, Masani K, Oh PI, Miyachi M, Popovic MR, Craven BC. Pulse wave velocity for assessment of arterial stiffness among people with spinal cord injury: a pilot study. *J Spinal Cord Med*. 2009;32:72-8. <https://doi.org/10.1080/10790268.2009.11760755>.
- [13] Yamashina A, Tomiyama H, Arai T, et al. Brachial-ankle pulse wave velocity as a marker of atherosclerotic vascular damage and cardiovascular risk. *Hypertens Res*. 2003;26:615-22. <https://doi.org/10.1291/hypres.26.615>.
- [14] Wu S, An S, Li W, et al. Association of Trajectory of Cardiovascular Health Score and Incident Cardiovascular Disease. *JAMA Netw Open*. 2019;2:e194758. <https://doi.org/10.1001/jamanetworkopen.2019.4758>.
- [15] Jin C, Chen S, Vaidya A, et al. Longitudinal Change in Fasting Blood Glucose and Myocardial Infarction Risk in a Population Without Diabetes. *Diabetes Care*. 2017;40:1565-72. <https://doi.org/10.2337/dc17-0610>.
- [16] Zheng X, Jin C, Liu Y, et al. Arterial Stiffness as a Predictor of Clinical Hypertension. *J Clin Hypertens (Greenwich)*. 2015;17:582-91. <https://doi.org/10.1111/jch.12556>.
- [17] Li Y, Song J, Pan H, Yao M, Hu X, Ma G. Validity of Food Frequency Questionnaire to investigate the dietary energy and nutrients intake. *Ying Yang Xue Bao* 2006; 28:143-7
- [18] Krebs-Smith SM, Pannucci TE, Subar AF, et al. Update of the Healthy Eating Index: HEI-2015. *J Acad Nutr Diet*. 2018;118:1591-602. <https://doi.org/10.1016/j.jand.2018.05.021>.
- [19] Bowman S, Clemens J, Friday J, Schroeder N, Shimizu M, LaCombs R, et al. Food Patterns Equivalents Intakes by Americans: What We Eat in America, NHANES 2003-2004 and 2015-2016, <http://www.ars.usda.gov/nea/bhnrc/fsrg/>; 2018 [accessed 17 August 2020].

- [20] Chen S, Li W, Jin C, et al. Resting Heart Rate Trajectory Pattern Predicts Arterial Stiffness in a Community-Based Chinese Cohort. *Arterioscler Thromb Vasc Biol*. 2017;37:359-64. <https://doi.org/10.1161/ATVBAHA.116.308674>.
- [21] Yamashina A, Tomiyama H, Arai T, et al. Brachial-ankle pulse wave velocity as a marker of atherosclerotic vascular damage and cardiovascular risk. *Hypertens Res*. 2003;26:615-22. <https://doi.org/10.1291/hypres.26.615>.
- [22] The International Physical Activity Questionnaire, <http://www.ipaq.ki.se/>; 2015 [accessed 17 August 2020].
- [23] The IPAQ Group. Guidelines for Data Processing and Analysis of the International Physical Activity Questionnaire (IPAQ) – Short and Long Forms Contents. <http://www.ipaq.ki.se/>; 2015
- [24] Deng HB, Macfarlane DJ, Thomas GN, et al. Reliability and validity of the IPAQ-Chinese: the Guangzhou Biobank Cohort study. *Med Sci Sports Exerc*. 2008;40:303-7. <https://doi.org/10.1249/mss.0b013e31815b0db5>.
- [25] Ohkuma T, Ninomiya T, Tomiyama H, et al. Brachial-Ankle Pulse Wave Velocity and the Risk Prediction of Cardiovascular Disease: An Individual Participant Data Meta-Analysis. *Hypertension*. 2017;69:1045-52. <https://doi.org/10.1161/HYPERTENSIONAHA.117.09097>.
- [26] Geiker NRW, Larsen ML, Dyerberg J, Stender S, Astrup A. Egg consumption, cardiovascular diseases and type 2 diabetes. *Eur J Clin Nutr*. 2018;72:44-56. <https://doi.org/10.1038/ejcn.2017.153>.
- [27] Virtanen JK, Mursu J, Virtanen HE, et al. Associations of egg and cholesterol intakes with carotid intima-media thickness and risk of incident coronary artery disease according to apolipoprotein E phenotype in men: the Kuopio Ischaemic Heart Disease Risk Factor Study. *Am J Clin Nutr*. 2016;103:895-901. <https://doi.org/10.3945/ajcn.115.122317>.
- [28] Larsson SC, Åkesson A, Wolk A. Egg consumption and risk of heart failure, myocardial infarction, and stroke: results from 2 prospective cohorts. *Am J Clin Nutr*. 2015;102:1007-13. <https://doi.org/10.3945/ajcn.115.119263>.
- [29] Schwingshackl L, Schwedhelm C, Hoffmann G, et al. Food groups and risk of all-cause mortality: a systematic review and meta-analysis of prospective studies. *Am J Clin Nutr*. 2017;105:1462-73. <https://doi.org/10.3945/ajcn.117.153148>.
- [30] Siasos G, Tousoulis D, Antoniadou C, Stefanadi E, Stefanadis C. L-Arginine, the substrate for NO synthesis: an alternative treatment for premature atherosclerosis?. *Int J Cardiol*. 2007;116:300-8. <https://doi.org/10.1016/j.ijcard.2006.04.062>.
- [31] Tousoulis D, Böger RH, Antoniadou C, Siasos G, Stefanadi E, Stefanadis C. Mechanisms of disease: L-arginine in coronary atherosclerosis—a clinical perspective. *Nat Clin Pract Cardiovasc Med*. 2007;4:274-83. <https://doi.org/10.1038/ncpcardio0878>.

- [32] Mayer O Jr, Seidlerová J, Wohlfahrt P, et al. Synergistic effect of low K and D vitamin status on arterial stiffness in a general population. *J Nutr Biochem*. 2017;46:83-9. <https://doi.org/10.1016/j.jnutbio.2017.04.010>.
- [33] Kim JE, Leite JO, DeOgburn R, Smyth JA, Clark RM, Fernandez ML. A lutein-enriched diet prevents cholesterol accumulation and decreases oxidized LDL and inflammatory cytokines in the aorta of guinea pigs. *J Nutr*. 2011;141:1458-63. <https://doi.org/10.3945/jn.111.141630>.
- [34] Detopoulou P, Panagiotakos DB, Antonopoulou S, Pitsavos C, Stefanadis C. Dietary choline and betaine intakes in relation to concentrations of inflammatory markers in healthy adults: the ATTICA study. *Am J Clin Nutr*. 2008;87:424-30. <https://doi.org/10.1093/ajcn/87.2.424>.
- [35] Miller CA, Corbin KD, da Costa KA, et al. Effect of egg ingestion on trimethylamine-N-oxide production in humans: a randomized, controlled, dose-response study. *Am J Clin Nutr*. 2014;100:778-86. <https://doi.org/10.3945/ajcn.114.087692>.
- [36] Tang WH, Wang Z, Levison BS, et al. Intestinal microbial metabolism of phosphatidylcholine and cardiovascular risk. *N Engl J Med*. 2013;368:1575-84. <https://doi.org/10.1056/nejmoa1109400>.
- [37] R. O. Alvim, J. Mourão, G. L. Magalhães et al. Non-HDL cholesterol is a good predictor of the risk of increased arterial stiffness in postmenopausal women in an urban Brazilian population. *Clinics (Sao Paulo)*. 2017;72:106-10. [http://dx.doi.org/10.6061/clinics/2017\(02\)07](http://dx.doi.org/10.6061/clinics/2017(02)07)
- [38] Zhao W, Gong W, Wu N, et al. Association of lipid profiles and the ratios with arterial stiffness in middle-aged and elderly Chinese. *Lipids Health Dis*. 2014;13:37. <https://doi.org/10.1186/1476-511X-13-37>.
- [39] Livingstone KM, Givens DI, Cockcroft JR, Pickering JE, Lovegrove JA. Is fatty acid intake a predictor of arterial stiffness and blood pressure in men? Evidence from the Caerphilly Prospective Study. *Nutr Metab Cardiovasc Dis*. 2013;23:1079-85. <https://doi.org/10.1016/j.numecd.2012.12.002>.

Tables

Table 1. Baseline characteristics based on egg consumption among 7315 Chinese adults

	Egg consumption				
	0-1.9/wk	2-2.9/wk	3-3.9/wk	4-4.9/wk	≥5/wk
	n=161	n=138	n=158	n=3012	n=396
Age, y	46.0±8.8	44.2±7.7	46.2±9.9	45.2±9.1	43.9±8.4
Men, %	78.3	67.4	58.9	52.6	73.7
Modified Healthy Eating Index	29.6±6.24	32.4±6.64	34.4±5.47	34.4±6.19	32.8±7.68
Total energy intake, kcal	1560±522	1724±569	1909±493	1638±447	1910±605
Body Mass Index, kg/m ²	25.3±3.3	25.5±3.1	24.6±3.3	24.5±3.4	24.9±2.9
Physical activity					
Low, %	51.9	56.1	50.9	87.9	61.8
Moderate, %	34.9	40.6	40.6	10.8	32.9
High, %	13.3	3.4	8.6	1.3	5.3
Systolic blood pressure, mmHg	134±15.9	136±18.2	136±17.3	130±18.5	135±15.9
Fasting blood glucose, mmol/L	5.85±1.43	5.94±1.54	6.00±1.72	5.66±1.70	5.95±1.64
Low-density lipoprotein cholesterol, mmol/L	2.97±0.80	2.99±0.78	3.05±0.86	2.75±0.81	2.96±0.75
High-density lipoprotein cholesterol, mmol/L	1.37±0.99	1.40±0.47	1.40±0.36	1.35±0.36	1.35±0.60
Married, %	95	100	98.1	95.4	95.7
Work type, manual, %	84.7	80.3	88.2	79.9	81.9
Alcohol consumption, %	40.1	36.2	30.7	55.3	30.4
College education or above, %	8.97	7.94	9.62	18.4	9.57
Current smoker, %	46.9	39.0	39.7	36.5	47.6
Baseline baPWV, cm/s	1426±17	1417±19	1391±18	1432±13	1436±15
Follow-up baPWV, cm/s	1483±20	1477±22	1436±21	1491±15	1499±18

Abbreviation: baPWV, brachial-ankle pulse wave velocity.

Table 2. Adjusted means (95% confidence intervals) of baPWV change rate (cm/s/year) by egg consumption group

	Egg consumption					
	0-1.9/wk	2-2.9/wk	3-3.9/wk	4-4.9/wk	≥5/wk	p for trend
n	325	266	316	5611	797	
Model 1	33.0±8.1	21.3±8.8	2.1±8.1*	14.4±2.0	23.4±5.4	0.38
Model 2	38.9±10.9	28.8±11.7	7.5±11.1*	18.9±7.7*	30.1±9.5	0.39
Model 3	35.9±11.2	21.7±12.0	0.2±11.4*	18.6±8.1	25.5±9.8	0.57
Excluding participants with arterial stiffness at baseline ^a	42.1±12.4	28.3±13.1	10.5±12.6*	24.0±8.7	32.0±10.6	0.80

Model 1 adjusted for age, sex, baseline baPWV, total energy intake and modified healthy eating index.

Model 2 adjusted for age, sex, baseline baPWV, total energy intake, modified healthy eating index, physical activity, marriage, employment, education level, alcohol consumption and smoking status, and heart rate.

Model 3 adjusted for age, sex, baseline baPWV, total energy intake, modified healthy eating index, physical activity, marriage, employment, education level, alcohol consumption, smoking status, heart rate, systolic blood pressure, fasting blood glucose, low-density lipoprotein cholesterol, and high-density lipoprotein cholesterol.

Compare to egg consumption 0-1.9/wk, * p<0.05

^a Based on the model 3

Table 3. Hazard ratios and 95% confidence intervals for incident arterial stiffness, according to baseline egg consumption.

	Egg consumption					
	0-1.9/wk	2-2.9/wk	3-3.9/wk	4-4.9/wk	≥5/wk	p for trend
Case/total	47/161	41/138	43/158	828/3012	116/396	
Model 1	1	1.03(0.67,1.57)	0.99(0.64,1.51)	1.09(0.81,1.48)	0.97(0.69,1.38)	0.87
Model 2	1	1.07(0.70,1.64)	0.96(0.62,1.49)	1.07(0.78,1.48)	0.99(0.70,1.42)	0.94
Model 3	1	1.07(0.70,1.64)	0.89(0.57,1.39)	1.15(0.83,1.60)	0.99(0.69,1.42)	0.71

Model 1 adjusted for age, sex, total energy intake and modified healthy eating index.

Model 2 adjusted for age, sex, total energy intake, modified healthy eating index, physical activity, marriage,

employment, education level, alcohol consumption, smoking status and heart rate.

Model 3 adjusted for age, sex, total energy intake, modified healthy eating index, physical activity, marriage, employment, education level, alcohol consumption, smoking status, heart rate, systolic blood pressure, fasting blood glucose, low-density lipoprotein cholesterol, and high-density lipoprotein cholesterol.

Figures

Figure 1

Flow chart of study participants selection.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Additionalfile1.docx](#)
- [STROBEchecklistcohort.docx](#)