Comparative Study of Phytochemical Profiles and Morphological Properties of Some Damask Rose From Iran

Mohammad Omid (✉ m.omidighale@ut.ac.ir)

University of Tehran

Azizollah Khandan-Mirkohi

University of Tehran

Mohsen Kafi

University of Tehran

Zabihollah Zamani

University of Tehran

Omid Rasouli

Tarbiat Modares University

Arezo Shaghaghi

University of Urmia

Mahnaz Kiani

Texas A&M University

Research Article

Keywords: Damask rose, Essential oil, GC-mass spectrometry, Perfume, Volatile compounds

Posted Date: March 7th, 2022

DOI: <https://doi.org/10.21203/rs.3.rs-1401571/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Abstract

Background: *Rosa damascena* is an aromatic rose species, is cultivated for its essential oil, and has wide application in perfume, cosmetic, pharmaceutical, and food industries in the world. This experiment was conducted to evaluate essential oil and morphological variations of 26 Damask rose genotypes. For this purpose, the effect of harvest time i.e. early morning or evening, and sampling type i.e. fresh or dried petals was evaluated on oil content. In addition, the composition of essential oil of the genotypes was determined using Gas chromatography–mass spectrometry (GC-MS).

Results: Results showed that early morning was the preferable time for flower collection based on oil content quantification. Furthermore, the oil yield of fresh petals was higher than the dried petals. Twenty-five volatile compounds were found in the extracted oils. β -damascenone, a key marker for the quality of rose oil, was found in 22 genotypes and was more than 1.5% concentration in G3, G6, and G11 genotypes. The highest components of the oil in Damask rose genotypes were nonadecane (42.51%), β -citronellol (40.82%), n-heneicosane (34.69%), geraniol (27.76%), and n-tricosane (14.2%). A wide variation in flower characteristics, such as petal color (from white to nearly red) and petal numbers from about 25 to 95, were also recorded. The G2, G5, and G15 genotypes originated from Isfahan, Fars, and Kerman respectively were selected based on petal number, flower weight, and essential oil content in fresh and dried petals.

Conclusions: Results suggest that the morphological and biochemical diversity of Damask rose genotypes can be used effectively to characterize genetic diversity between different genotypes and to select special traits in the breeding program

Background

Damask rose (*Rosa damascena* Mill.) is one of the supreme fragrance species in the Rosaceae family that is derived from *Rosa gallica* and *Rosa moschata* [1]. This species is cultivated for its ornamental value and also for essential oil extraction in most parts of the northern hemisphere [2, 3]. Iran has been introduced as the genetic diversity center and the origin of Damask roses [4–10]. Nowadays, this species is cultivated extensively in Bulgaria, Iran, Turkey, France, Italy, Morocco, the USA, and India [11]. The global production of rose oil is about 4.5 tones per year [12]. The global rose oil market was valued 278.7 million USD in 2018 [13]. Products of Damask rose, including essential oil, rose water, rose concrete, dried petals, dried flower buds, and rose absolute, are used in perfume, cosmetic, pharmaceutical, and food industries. Several pharmacological attributes, such as antibacterial, antioxidant, and anti-HIV effects have been found in rose oil [14–16].

Rosa damascena is cultivated in widespread ecological conditions, but specific climatic conditions are needed to produce high-quality essential oil. The quality of essential oil and flower yield of *R. damascena* is mainly affected by geographical origin and climate conditions, time and stage of flower harvesting, method of extraction [5], and agriculture practices [17–21]. *Rosa damascena* grows wild in some parts of Iran and has been propagated vegetatively and cultivated for a long time [6]. Thus, various cultivars of Damask rose have been selected during the long cultivation history, and it has also been crossed naturally with local species [22]. The phenotypic homogeneity caused by continuous vegetative reproduction and environmental effects makes it possible for mass production of this crop to produce rose oil [5, 9, 23].

The most important compounds of rose oil are β -citronellol, nonadecane, geraniol, eugenol, heneicosane, and phenols such as eugenol [24, 25]. In addition, some factors such as the concentration of ethanol used for extraction, storage period, and production conditions of flowers, also would affect the amounts of key compounds in rose oil [5]. Several studies have been conducted on the chemical composition of essential oil in various populations of Damask rose by GC/MS through different extraction methods [3, 9, 15, 26–28], and also on genetic and morphological diversity of *Rosa damascena* [6, 10, 29–32]. The effect of micro-climate on Damask rose cultivation and the oil composition has been also reported [32, 33].

As we have access to the wide genetic diversity of *Rosa damascena* in Iran, it will be valuable to characterize their specific morphology and biochemical characters in more detail. Thus, the present study was carried out to determine the variations in the flower yield and morphological and chemical compositions of 26 different Iranian Damask rose genotypes by using gas chromatography-mass spectrometry (GC-MS).

Materials And Methods

Plant materials and collection site

Damask rose genotypes that were selected for this study consisted of 26 genotypes. These genotypes were previously collected from several parts of Iran (Table 1) and established in the research station of the Horticultural sciences department, the University of Tehran, Karaj, Iran (latitude 35° 77' N, longitude 50° 93' E and altitude 1251 m), based on a randomized block design with three replications in 2004 [8]. All samples were the same because of their yearly pruning. The average plant high was 165 cm and plant diameter was 123 cm. The current experiment was carried out during 2016–2018.

Table 1
Damask rose genotypes assessed in this study, their province of origin, collection site, and petal color

Genotype No.	Province of origin	Collecting site	Longitude (°)	Latitude (°)	Altitude	Petal color
G1	East Azerbaijan	Kashan, collection of Taghtiran Company	51.05	34.02	1814	Pink
G2	Isfahan	Kashan, collection of Taghtiran Company	51.05	34.02	1814	Dark pink
G3	Tehran	Kashan, collection of Taghtiran Company	51.05	34.02	1814	Pink
G4	Isfahan	Kashan, collection of Taghtiran Company	51.05	34.02	1814	Dark pink
G5	Fars	Kashan, collection of Taghtiran Company	51.05	34.02	1814	White
G6	Fars	Darab, Lyzangan	54.98	28.66	2018	Pale pink
G7	Fars	Darab, Rostagh	55.06	28.44	1314	Pink
G8	Fars	Darab, Ghale Biaban	54.87	28.52	1339	Pink
G9	Fars	Darab, Lyzangan	54.99	28.67	2070	Pale pink
G10	Fars	Maimand, Sahra sefid	52.79	28.83	1480	White
G11	Fars	Maimand, Kang	52.83	28.87	1649	Pink
G12	Fars	Maimand	52.76	28.86	1548	Pink
G13	Kerman	Bardsir	56.58	29.90	2070	Pale pink
G14	Kerman	Bardsir	56.61	29.87	2095	Pink
G15	Kerman	Mahan	57.24	30.12	1823	Pink
G16	East Azerbaijan	Osco	46.13	37.92	1567	Pink
G17	East Azerbaijan	Tabriz	46.43	38.01	1673	Pink
G18	East Azerbaijan	Osco	46.11	37.89	1575	Pink
G19	East Azerbaijan	Osco	46.18	37.90	1685	Dark pink
G20	East Azerbaijan	Ahar	47.04	38.44	1387	Pink
G21	Isfahan	Kashan	51.47	33.94	979	Pink
G22	Isfahan	Kashan	51.53	33.94	974	Pink
G23	Isfahan	Kashan	51.61	33.93	946	Pale pink
G24	Razavi Khorasan	Mashhad	59.46	36.61	1122	Pink
G25	Razavi Khorasan	Mashhad	59.43	36.38	1092	Pink
G26	East Azerbaijan	Tabriz	46.40	37.98	1783	White

Evaluation of plant vegetative and flower characteristics

The morphological characters, such as plant height, crown diameter, number of main stems in each plant, number of flowers in main stems, angle of the secondary branches, internode length, and thorn density (one to five from high to low), were determined in twelve years old plants. Additionally, the length of stipule, peduncle, receptacle, and flower bud's length and diameter prior to the opening stage were measured. Petal colors were determined visually and also measured with a colorimeter (Minolta CR-400 Chroma meter, Konica Minolta Sensing, Inc., Osaka, Japan) using the following parameters: L* (lightness), a* (redness), and b* (yellowness). Color parameters were obtained through reflectance values and Chroma calculated by the following formula [34].

$$\text{Chroma} = \sqrt{a^2 + b^2}$$

Isolation and content of essential oil

For the extraction of essential oil, fresh flowers from each accession were randomly collected (20 fresh flowers for each accession). Flowers were harvested both in the morning and evening. For dried samples, the collected flowers were spread on wire shelves and kept in the shade for two weeks at room temperature [35]. A total of 200 g of fresh petals and 55 g of dry petals (equivalent to 200 g fresh petals) subjected to hydrodistillation using 400 ml distilled water in a Clevenger for 3 hours with three replications [64]. The essential oil was measured directly in the extraction burette and the oil content (v/w) in flower was expressed as percentage on a fresh weight basis of essential oil per 200 g of fresh petals. The extracted oils were transferred into vials and stored at 4°C in the dark until used.

Gas chromatography (GC-FID) and (GC-MS)

GC-MS analysis of the oil samples was performed on a Thermo–UFM (Ultra-Fast model, Italy) gas chromatograph equipped with a P5 (non-polar) capillary column (10 m × 0.1 mm), which employed helium (0.5 ml/min) as the carrier gas to split injection at 1:100. The oven temperature was set at 60°C for 30 min, FID detector temperature was programmed at 285°C at the rate of 80°C/min and injector temperature was 280°C. The relative amounts of individual components were calculated based on the GC peak areas by using a normalization method regarding response factor. The essential oil constituents were identified following an injection of n-alkanes (C₈–C₂₄) under the same conditions and confirmed according to Wiley 275-L library and literature [36–38]. The compounds were identified using commercial mass spectral libraries (NIST 05, Wiley 7th Mass spectra register) [37].

Statistical analysis

For the evaluation of morphological characteristics of vegetative and flower parts, the experiment was arranged in a randomized complete block design (RCBD) with three replications. Mean values were compared at 95% ($p \leq 0.05$) and 99% ($p \leq 0.01$) confidence intervals using the LSD test by Minitab ver.16. [39].

Results

Oil content in fresh and dry petals

In the majority of selected Damask rose genotypes, petals harvested in the morning time for dried and fresh petals had higher oil content in comparison with samples collected in the evening time with the exception of G9, G11 and G23 genotypes that opposite trend was found (Fig. 1). The highest oil content in fresh petals was found in G21 for morning and evening harvest time 0.14 and 0.15 (v/w%) respectively. Additionally, in G18, G19, G20, G24, and G25 genotypes the total volume of essential oil content in both harvesting times was similar (Fig. 1). In dried petals, the time of harvesting also affected the oil content. However, the oil content in dried petals generally was lower than the fresh petals. Although in most of the genotypes, the higher oil content was recorded in the morning time harvested dried petals, three genotypes (G1, G2, and G4) had the same content of essential oil in both dried morning and evening times harvested petals. On the other hand, due to later flower blooming in G22 (0.05 v/w%) and G24 (0.04 v/w%) genotypes, the oil content in dried petals harvested in the evening time was more than the morning time collected samples (Fig. 2).

Morphological traits

There were clear differences in morphological characteristics between selected Damask rose genotypes (Table 2). The correlation matrix among morphological traits of *R. damascena* showed that the plant height was significantly ($P = 0.01$) and positively correlated with the branch angle ($r = 0.21$), number of stems per plant ($r = 0.57$), flower bud length ($r = 0.46$) and flower peduncle length ($r = 0.45$) (Table 3). Thorn density was positively correlated with the number of petals ($P = 0.01$). Moreover, a negative correlation ($r = -0.39$) was found between thorn density and branch angle. A significant ($P = 0.01$) negative ($r = -0.56$) correlation was found between flower peduncle length and thorn density. Thorn density had a negative relation with all traits, except with the bud diameter (Table 3). The flower peduncle length was positively correlated with most traits evaluated in this study. Different petal colors, from white to dark purple, were observed in the selected Damask rose genotypes (Table 1). The measurements of color parameters gave different values of L*, a*, and b*. The results showed highly significant differences among genotypes for all color traits (Table 4). Chroma values also were different between genotypes.

Table 2

Measured morphological characters (mean $\pm$ SE) of 26 Damask rose genotypes

Genotype No	Plant height (cm)	Plant crown diameter(cm)	No of main stems per plant	No of nodes in the branch	Internode length (mm)	Secondary branches angle	Thorns density	Leaf stipule length (mm)	Leaf stipule width (mm)	Peduncle length (mm)	Receptacle length (mm)
G1	111.8 $\pm$ 5.5	69.4 $\pm$ 5.13	14.2 $\pm$ 2.0	10.5 $\pm$ 1.7	23.9 $\pm$ 1.2	73.1 $\pm$ 5.5	4.0 $\pm$ 00	15.9 $\pm$ 1.0	6.9 $\pm$ 0.4	14.8 $\pm$ 1.1	10.2 $\pm$ 1.3
G2	160.5 $\pm$ 4.7	123.9 $\pm$ 8.7	19.0 $\pm$ 1.8	27.2 $\pm$ 3.3	34.0 $\pm$ 3.9	83.3 $\pm$ 1.6	4.0 $\pm$ 00	16.8 $\pm$ 1.4	20.4 $\pm$ 1.9	22.9 $\pm$ 1.8	7.7 $\pm$ 0.3
G3	111.2 $\pm$ 3.5	104.0 $\pm$ 3.7	14.2 $\pm$ 0.8	19.5 $\pm$ 1.7	29.3 $\pm$ 2.7	81.1 $\pm$ 2.3	4.0 $\pm$ 00	19.0 $\pm$ 1.2	8.4 $\pm$ 0.8	18.8 $\pm$ 1.2	14.3 $\pm$ 1.0
G4	185.2 $\pm$ 3.7	139.2 $\pm$ 4.2	16.4 $\pm$ 0.9	34.1 $\pm$ 3.2	32.8 $\pm$ 3.1	85.5 $\pm$ 1.3	2.0 $\pm$ 00	18.5 $\pm$ 0.7	10.1 $\pm$ 0.6	26.7 $\pm$ 1.5	10.6 $\pm$ 0.4
G5	129.4 $\pm$ 4.5	107.8 $\pm$ 4.8	16.9 $\pm$ 1.0	23.8 $\pm$ 1.4	29.0 $\pm$ 2.4	66.7 $\pm$ 1.6	5.0 $\pm$ 00	20.0 $\pm$ 1.0	15.3 $\pm$ 0.9	22.2 $\pm$ 2.0	6.4 $\pm$ 0.2
G6	174.1 $\pm$ 6.1	152.9 $\pm$ 5.6	18.3 $\pm$ 1.7	27.2 $\pm$ 2.0	27.0 $\pm$ 2.3	78.3 $\pm$ 2.8	3.1 $\pm$ 0.2	14.6 $\pm$ 1.0	7.6 $\pm$ 0.5	28.9 $\pm$ 2.5	8.9 $\pm$ 0.4
G7	177.1 $\pm$ 2.8	135.8 $\pm$ 2.3	17.2 $\pm$ 1.1	32.1 $\pm$ 5.7	36.5 $\pm$ 4.0	83.3 $\pm$ 1.6	1.0 $\pm$ 00	26.6 $\pm$ 1.9	8.0 $\pm$ 0.5	32.2 $\pm$ 2.7	10.6 $\pm$ 0.4
G8	194.4 $\pm$ 3.3	148.0 $\pm$ 3.3	15.3 $\pm$ 0.6	46.4 $\pm$ 4.1	37.2 $\pm$ 2.6	86.1 $\pm$ 1.4	1.0 $\pm$ 00	25.2 $\pm$ 1.1	10.6 $\pm$ 0.8	29.5 $\pm$ 1.8	10.8 $\pm$ 0.2
G9	168.3 $\pm$ 8.9	133.5 $\pm$ 4.3	15.0 $\pm$ 0.7	41.9 $\pm$ 6.4	36.3 $\pm$ 2.6	79.4 $\pm$ 2.4	5.0 $\pm$ 00	26.2 $\pm$ 1.7	11.4 $\pm$ 0.7	24.1 $\pm$ 0.9	10.2 $\pm$ 0.7
G10	200.6 $\pm$ 4.8	113.0 $\pm$ 3.6	22.2 $\pm$ 1.5	35.9 $\pm$ 2.6	32.8 $\pm$ 3.2	82.2 $\pm$ 2.2	2.0 $\pm$ 00	22.0 $\pm$ 1.2	9.4 $\pm$ 0.9	30.0 $\pm$ 1.7	9.8 $\pm$ 0.4
G11	116.5 $\pm$ 2.3	104.0 $\pm$ 4.8	11.4 $\pm$ 0.9	16.9 $\pm$ 2.4	33.2 $\pm$ 3.1	86.6 $\pm$ 1.1	1.0 $\pm$ 00	24.1 $\pm$ 1.2	10.6 $\pm$ 0.4	29.7 $\pm$ 1.5	10.8 $\pm$ 0.2
G12	205.4 $\pm$ 3.0	153.9 $\pm$ 4.4	22.1 $\pm$ 1.3	38.2 $\pm$ 4.5	35.0 $\pm$ 1.7	78.3 $\pm$ 2.0	1.0 $\pm$ 00	17.0 $\pm$ 0.7	8.1 $\pm$ 0.4	32.6 $\pm$ 1.7	10.1 $\pm$ 0.5
G13	213.8 $\pm$ 5.5	150.0 $\pm$ 2.8	35.3 $\pm$ 1.1	32.0 $\pm$ 2.1	33.7 $\pm$ 3.6	75.0 $\pm$ 2.0	2.3 $\pm$ 0.1	19.5 $\pm$ 0.4	7.5 $\pm$ 0.3	29.5 $\pm$ 2.6	9.4 $\pm$ 0.4
G14	182.5 $\pm$ 3.5	114.0 $\pm$ 3.1	19.8 $\pm$ 1.4	31.4 $\pm$ 2.8	30.2 $\pm$ 2.4	80.0 $\pm$ 2.3	1.0 $\pm$ 00	20.5 $\pm$ 1.6	9.4 $\pm$ 0.7	30.2 $\pm$ 1.9	9.2 $\pm$ 0.4
G15	201.3 $\pm$ 4.3	160.0 $\pm$ 3.3	28.1 $\pm$ 0.9	33.0 $\pm$ 2.0	28.1 $\pm$ 1.9	77.8 $\pm$ 2.6	2.3 $\pm$ 0.1	21.5 $\pm$ 1.2	7.5 $\pm$ 0.8	35.7 $\pm$ 2.0	9.2 $\pm$ 0.5
G16	147.5 $\pm$ 3.9	118.3 $\pm$ 6.3	17.5 $\pm$ 0.9	16.2 $\pm$ 1.6	28.9 $\pm$ 2.8	79.4 $\pm$ 1.7	4.0 $\pm$ 00	18.9 $\pm$ 0.9	11.3 $\pm$ 0.8	20.6 $\pm$ 2.5	6.9 $\pm$ 0.2
G17	122.7 $\pm$ 5.1	101.2 $\pm$ 7.4	10.1 $\pm$ 1.2	11.4 $\pm$ 1.2	29.0 $\pm$ 2.8	84.4 $\pm$ 1.7	4.5 $\pm$ 0.1	17.7 $\pm$ 0.6	11.9 $\pm$ 0.4	20.4 $\pm$ 1.1	7.1 $\pm$ 0.4
G18	139.5 $\pm$ 4.0	117.8 $\pm$ 4.0	21.2 $\pm$ 0.9	20.4 $\pm$ 2.0	30.7 $\pm$ 2.3	66.1 $\pm$ 0.1	4.0 $\pm$ 00	19.4 $\pm$ 1.8	11.0 $\pm$ 0.6	21.0 $\pm$ 1.2	7.9 $\pm$ 0.5
G19	143.8 $\pm$ 1.9	146.2 $\pm$ 6.7	16.7 $\pm$ 0.6	23.4 $\pm$ 3.5	30.2 $\pm$ 3.2	66.7 $\pm$ 1.6	4.0 $\pm$ 00	21.5 $\pm$ 1.2	11.0 $\pm$ 0.6	18.7 $\pm$ 1.5	7.3 $\pm$ 0.0
G20	115.7 $\pm$ 4.8	121.9 $\pm$ 3.9	14.1 $\pm$ 1.0	21.9 $\pm$ 2.8	28.7 $\pm$ 3.3	62.8 $\pm$ 1.7	4.0 $\pm$ 00	20.2 $\pm$ 1.2	12.4 $\pm$ 1.4	22.2 $\pm$ 2.3	8.0 $\pm$ 0.3
G21	167.8 $\pm$ 3.6	125.9 $\pm$ 6.8	14.5 $\pm$ 1.2	28.1 $\pm$ 3.1	32.7 $\pm$ 3.4	80.0 $\pm$ 1.1	1.0 $\pm$ 00	47.2 $\pm$ 2.0	10.4 $\pm$ 0.6	32.4 $\pm$ 1.5	10.7 $\pm$ 0.2
G22	200.4 $\pm$ 3.5	143.9 $\pm$ 3.0	20.4 $\pm$ 1.5	34.4 $\pm$ 3.5	33.0 $\pm$ 3.3	86.1 $\pm$ 1.4	1.0 $\pm$ 00	23.2 $\pm$ 1.3	10.6 $\pm$ 0.7	29.2 $\pm$ 1.8	9.8 $\pm$ 0.4
G23	186.2 $\pm$ 4.8	129.7 $\pm$ 5.4	22.0 $\pm$ 0.8	40.7 $\pm$ 4.7	36.3 $\pm$ 2.5	82.2 $\pm$ 1.8	1.0 $\pm$ 00	24.0 $\pm$ 1.8	9.6 $\pm$ 0.7	35.9 $\pm$ 2.8	11.6 $\pm$ 0.4
G24	192.4 $\pm$ 8.3	136.1 $\pm$ 8.3	24.1 $\pm$ 1.5	28.5 $\pm$ 2.9	35.0 $\pm$ 0.3	80.5 $\pm$ 2.4	4.0 $\pm$ 00	20.5 $\pm$ 1.3	11.3 $\pm$ 1.2	26.5 $\pm$ 1.3	7.5 $\pm$ 0.2
G25	154.1 $\pm$ 3.4	134.5 $\pm$ 5.4	13.0 $\pm$ 0.5	13.7 $\pm$ 2.7	35.5 $\pm$ 3.8	83.3 $\pm$ 1.1	1.0 $\pm$ 00	25.1 $\pm$ 0.9	17.5 $\pm$ 1.7	35.9 $\pm$ 2.4	8.6 $\pm$ 0.6
G26	158.1 $\pm$ 3.4	140.8 $\pm$ 3.8	21.1 $\pm$ 1.7	25.0 $\pm$ 2.8	27.2 $\pm$ 2.5	75.0 $\pm$ 2.0	5.0 $\pm$ 00	17.6 $\pm$ 1.3	7.8 $\pm$ 0.6	26.9 $\pm$ 1.5	8.2 $\pm$ 0.3

Table 3
Correlation coefficients between the main morphological characters in *R. damascena* genotypes

Parameters	Plant height	Plant crown diameter	No of main stems per plant	No of nodes in the branch	Internode length	Branches angle	Thorns density	Leaf stipule length	Leaf stipule Width	Peduncle length	Receptacle length	Flower bud length	Flow bud diam
Plant height	1												
Plant crown diameter	0.55**	1											
No of main stems per plant	0.57**	0.33**	1										
No of nodes in the branch	0.51**	0.36**	0.23**	1									
Internode length	0.17**	0.12	0.05	0.16**	1								
Branches angle	0.21**	0.09	-0.08	0.015	0.17**	1							
Thorns density	-0.47**	-0.23**	-0.09	-0.29**	-0.19**	-0.39**	1						
Leaf stipule length	0.07	0.05	-0.08	0.07	0.12	0.07	-0.01	1					
Leaf stipule Width	-0.05	0.04	-0.04	-0.07	0.04	-0.04	0.07	-0.05	1				
Peduncle length	0.45**	0.36**	0.18**	0.28**	0.07	0.24**	-0.56**	0.08	0.05	1			
Receptacle length	0.07	-0.01	-0.04	0.17	0.06	0.23**	-0.36**	0.06	-0.16	0.18**	1		
Flower bud length	0.46**	0.27**	0.11	0.21**	0.21**	0.27**	-0.56**	0.06	-0.04	0.36**	0.27**	1	
Flower bud diameter	-0.09	-0.01	-0.01	0.00	-0.10	-0.02	0.27**	-0.09	0.06	-0.11	0.05	-0.04	1
No of petals	-0.27**	-0.16	-0.10	-0.20**	-0.03	-0.11	0.39**	-0.10	0.17**	-0.31**	-0.29**	-0.35**	0.07

Table 4
Color indices of petals of 26 Damask rose landraces (mean $\pm$ SE)

Genotype	L*	a*	b*	Chroma
G1	61.4 $\pm$ 4.4	31.59 $\pm$ 3.6	-0.4 $\pm$ 0.4	38.6 $\pm$ 1.0
G2	37.3 $\pm$ 0.7	52.41 $\pm$ 2.1	8.8 $\pm$ 2.4	41.9 $\pm$ 4.1
G3	65.0 $\pm$ 3.2	32.5 $\pm$ 5.6	-5.0 $\pm$ 2.1	39.9 $\pm$ 0.7
G4	49.0 $\pm$ 1.7	47.8 $\pm$ 5.0	-1.8 $\pm$ 0.7	53.9 $\pm$ 2.1
G5	68.2 $\pm$ 1.2	23.4 $\pm$ 6.5	24.7 $\pm$ 3.8	41.3 $\pm$ 3.0
G6	67.3 $\pm$ 2.9	28.8 $\pm$ 3.1	0.9 $\pm$ 0.4	35.8 $\pm$ 2.0
G7	70.8 $\pm$ 1.4	26.5 $\pm$ 1.9	-0.9 $\pm$ 0.5	32.5 $\pm$ 3.1
G8	71.6 $\pm$ 1.4	28.1 $\pm$ 2.4	0.0 $\pm$ 0.6	34.1 $\pm$ 5.1
G9	74.8 $\pm$ 1.1	22.0 $\pm$ 3.4	22.3 $\pm$ 1.5	38.2 $\pm$ 2.9
G10	76.4 $\pm$ 3.3	27.5 $\pm$ 5.3	-3.1 $\pm$ 0.9	43.1 $\pm$ 2.5
G11	64.3 $\pm$ 1.0	36.3 $\pm$ 0.6	-3.6 $\pm$ 0.3	42.5 $\pm$ 3.0
G12	67.0 $\pm$ 1.9	25.9 $\pm$ 2.2	4.5 $\pm$ 1.6	31.3 $\pm$ 0.6
G13	59.5 $\pm$ 3.1	29.4 $\pm$ 2.8	-2.5 $\pm$ 0.5	36.5 $\pm$ 1.0
G14	55.9 $\pm$ 5.1	42.1 $\pm$ 2.3	-1.6 $\pm$ 1.7	49.1 $\pm$ 2.0
G15	60.6 $\pm$ 3.3	37.3 $\pm$ 4.5	-0.9 $\pm$ 3.0	44.3 $\pm$ 3.2
G16	41.7 $\pm$ 1.7	53.0 $\pm$ 1.8	-8.8 $\pm$ 0.8	60.7 $\pm$ 4.0
G17	63.2 $\pm$ 2.9	34.4 $\pm$ 4.0	-2.5 $\pm$ 0.5	41.5 $\pm$ 0.4
G18	43.6 $\pm$ 5.1	37.0 $\pm$ 2.9	1.6 $\pm$ 0.6	43.0 $\pm$ 7.9
G19	39.0 $\pm$ 2.4	35.0 $\pm$ 2.7	3.3 $\pm$ 1.8	42.2 $\pm$ 6.3
G20	55.8 $\pm$ 4.9	35.7 $\pm$ 7.6	3.7 $\pm$ 3.4	41.9 $\pm$ 4.5
G21	60.2 $\pm$ 0.2	25.7 $\pm$ 5.0	2.0 $\pm$ 0.4	32.8 $\pm$ 1.9
G22	66.2 $\pm$ 3.9	29.9 $\pm$ 6.5	-0.6 $\pm$ 1.6	36.9 $\pm$ 0.2
G23	65.7 $\pm$ 1.5	24.3 $\pm$ 3.0	-2.2 $\pm$ 0.8	41.3 $\pm$ 3.7
G24	66.3 $\pm$ 2.9	32.1 $\pm$ 3.4	0.3 $\pm$ 2.7	39.1 $\pm$ 1.4
G25	65.4 $\pm$ 3.1	33.1 $\pm$ 3.1	1.4 $\pm$ 1.4	40.1 $\pm$ 2.1
G26	79.0 $\pm$ 0.1	26.2 $\pm$ 0.4	19.3 $\pm$ 0.7	33.4 $\pm$ 1.8

Essential oil components

Significant differences were found between chromatographic characteristics of the genotypes, indicating differences in their chemical compositions. In total, twenty-five compounds were identified in extracted oils of fresh petals of selected 26 Iranian genotypes (Table 5). The principal components of the essential oils were n-heneicosane, citronellol, and nonadecane in all genotypes. Results show that geraniol is the highest component in Damask rose oil, except in G14 and G18. The highest concentration of geraniol was found in G5 (27.76%), G9 (27.33%), and G2 (27.27%) respectively. Geraniol has been reported to be one of the main essential oil components in Damask rose [5]. According to GC-MS results, the highest nonadecane contents (42.51%, 35.06%, 30.91%, and 30.26%) were found in essential oils of G12, G14, G11, and G19, respectively. Several studies indicated that heneicosane, heptadecane, nonadecane, and eicosane were abundant hydrocarbons in rose oil [32]. Based on the results, G14 (34.69%), G23 (30.92%), and G21 (30.82%) had the highest content of heneicosane. In the current study, damascone (Z)- α and β -damascenone had been determined in most of the genotypes, but β -damascone was less abundant. The highest concentrations of damascone (Z)- α (2.88%), β -damascone (0.96%), and β -damascenone (1.76%) were found in G20, G3, and G3, respectively (Table 5). Damascone (Z)- α , β -damascone, and β -damascenone are the trace components and quality markers for Damask rose oil, playing an important organoleptic role in rose oil [40]. The highest quantity of β -citronellol (40%) was recorded in G26 which is one of the most abundant acyclic terpenes in rose oil.

Table 5
Quantitative data for the twenty-five components of the essential oil content (%) from 26 Damask roses genotypes, determined by GC-MS

No	Compound	Rt ^a	LRI ^b	RI ^c	G1	G2	G3	G4	G5	G6	G7	G8	G9	G10	G11
1	Phenyl ethyl alcohol	11.75	1106	1111	-	0.334	-	-	0.309	1.541	-	0.302	0.400	0.240	-
2	Dihydro linalool	12.82	1131	1145	0.556	1.451	-	1.057	0.826	41.189	0.388	0.831	1.271	0.716	-
3	β-Citronellol	16.80	1223	1225	34.660	29.268	14.911	37.574	21.812	15.608	27.728	31.830	30.389	30.77	10
4	Neral	17.32	1235	1237	2.281	10.258	0	3.893	7.347	6.221	2.239	3.791	10.832	3.914	0
5	Geraniol	17.95	1249	1250	6.182	27.270	2.931	12.182	27.768	7.986	6.935	10.863	27.335	10.364	2.1
6	Linalyl acetate	18.16	1254	1257	0.220	-	-	-	0.470	-	-	-	0.291	0.277	-
7	Geranial	18.62	1264	1267	0.202	0.877	-	0.332	0.363	-	-	-	0.728	0.268	-
8	Dihydro citronellol acetate	21.04	1319	1320	0.232	0.454	-	0.278	0.408	-	-	-	0.360	0.242	-
9	Damascone (Z)-α	22.64	1355	1358	0.368	1.847	-	0.538	2.226	1.666	0.487	0.394	1.281	0.466	-
10	Damascenone (E)-β	23.87	1383	1384	0.762	-	1.766	0.526	0.308	1.761	0.704	0.812	0.274	0.768	1.7
11	Damascone (E)-β	25.17	1413	1414	-	0.528	0.967	-	-	-	-	-	0.253	0.299	0.1
12	Dodecen-1-ol (2E)	27.55	1469	1471	-	0.717	-	-	-	-	-	0.736	2.099	0.199	0.1
13	n Heptadecane	36.74	1700	1700	1.979	0.435	1.315	2.311	1.609	-	1.271	0.583	0.529	1.923	1.0
14	z z-Farnesol	36.68	1698	1698	0.638	1.026	0.503	-	-	-	-	0.584	1.367	0.432	-
15	(e e)-Farnesyl acetate	42.01	1845	1846	0.845	4.002	-	1.138	2.249	-	1.173	1.399	3.756	1.295	0.1
16	n-Octadecane	40.40	1800	1800	-	0.360	0.567	0.262	-	-	-	-	0.429	-	0.4
17	n-Hexadecanol	43.03	1874	1875	2.918	0.317	4.184	5.202	3.815	1.174	2.634	0.605	0.343	3.308	3.4
18	Nonadecane	43.92	1900	1903	27.229	5.580	28.927	19.81	14.812	5.455	23.574	10.710	4.347	23.396	30
19	n-Eicosane	47.33	2000	2004	2.587	1.362	4.330	1.850	1.542	-	3.260	2.646	1.242	2.488	4.8
20	n-Heneicosane	50.47	2100	2100	13.860	7.729	21.574	9.429	9.522	4.784	21.243	19.543	6.749	13.208	27
21	n-Docosane	53.54	2200	2195	0.286	0.405	0.724	-	0.224	9.335	0.578	0.749	0.342	0.310	0.8
22	n-Tricosane	56.48	2300	2303	3.236	4.874	6.506	2.480	3.295	2.635	6.039	11.177	3.905	3.908	7.0
23	e-e Farnesol	38.30	1742	1743	0.363	0.522	0.439	-	-	-	0.359	0.524	0.861	-	0.1
24	Phytol	45.37	1942	1943	-	-	-	-	-	-	-	0.311	-	-	3.7
25	Trans rose oxide	12.45	1122	1125	-	-	0.605	-	-	-	-	-	-	-	0.1
Total					99.413	99.625	90.256	98.873	98.912	99.360	98.615	98.400	99.393	98.801	97

^aRt: Retention time (min)

^bLRI: RI from literature (Adams, 2017).

^cRI: Experimentally determined retention index (RI)

Table 5
(continue). Quantitative data for the twenty-five components of the essential oil content (%) from 26 Damask rose genotypes, determined

No	Compound	Rt ^a	LRI ^b	RI ^c	G14	G15	G16	G17	G18	G19	G20	G21	G22	G23	G24
1	Phenyl ethyl alcohol	11.75	1106	1111	0	0.206	0.224	-	-	0.103	0.187	-	0.165	0.409	0.1
2	Dihydro linalool	12.82	1131	1145	2.559	10.647	3.064	4.434	23.869	0.553	0.989	-	0.723	0.484	0.9
3	β-Citronellol	16.80	1223	1225	6.155	29.638	15.961	18.993	14.855	33.163	19.397	11.808	29.668	14.940	12
4	Neral	17.32	1235	1237	-	3.675	5.110	0.734	-	0.414	3.549	2.867	3.018	0.308	1.7
5	Geraniol	17.95	1249	1250	-	9.394	12.120	5.707	-	2.211	13.797	6.415	19.244	4.208	10
6	Linalyl acetate	18.16	1254	1257	-	0.692	2.762	-	-	-	2.377	-	0.152	-	0.7
7	geranial	18.62	1264	1267	-	0.219	0.285	-	-	-	-	-	-	-	-
8	Dihydro citronellol acetate	21.04	1319	1320	-	-	0.320	-	-	0.221	0.471	-	0.180	-	0.7
9	Damascone (Z)-α	22.64	1355	1358	-	0.314	2.081	-	1.768	0.233	2.880	-	0.571	-	-
10	Damascenone (E)-β	23.87	1383	1384	-	0.594	0.247	0.885	-	0.785	0.199	0.415	0.775	0.979	0.3
11	Damascone (E)-β	25.17	1413	1414	-	0.592	1.270	0.418	-	0.170	-	-	0.166	0.519	-
12	Dodecen-1-ol (2E)	27.55	1469	1471	-	-	-	0.423	1.396	0.195	-	-	0.144	-	0.7
13	n Heptadecane	36.74	1700	1700	0.854	0.569	1.459	0.76	1.300	2.532	0.644	1.174	1.296	0.545	2.7
14	z z-Farnesol	36.68	1698	1698	-	-	1.749	3.017	-	-	0.364	-	-	0.707	0.7
15	(e e)-Farnesyl acetate	42.01	1845	1846	-	-	2.834	0.870	-	0.224	2.279	0.603	1.534	0.336	1.3
16	n-Octadecane	40.40	1800	1800	-	-	-	-	2.214	-	-	-	-	-	-
17	n-Hexadecanol	43.03	1874	1875	2.386	0.497	1.386	1.119	9.166	3.829	1.270	2.033	2.302	1.979	1.4
18	Nonadecane	43.92	1900	1903	35.061	9.54	19.044	25.131	12.882	30.264	14.43	29.524	18.540	22.789	20
19	n-Eicosane	47.33	2000	2004	5.811	1.797	1.9154	3.975	1.821	2.919	2.323	4.193	2.212	4.469	3.6
20	n-Heneicosane	50.47	2100	2100	34.695	13.389	16.763	24.094	11.685	15.751	18.080	30.824	13.986	30.925	20
21	n-Docosane	53.54	2200	2195	1.065	0.607	0.590	0.704	0	0.346	0.824	0.718	0.330	1.056	0.4
22	n-Tricosane	56.48	2300	2303	10.017	8.290	10.170	6.019	5.288	4.215	14.205	8.920	3.575	11.113	4.7
23	e-e Farnesol	38.30	1742	1743	0	8.282	0	0		0	0	0	0	0	0.7
24	Phytol	45.37	1942	1943		0	0	1.059		0	0	0	0	3.375	0
25	Trans rose oxide	12.45	1122	1125	0	0.208	0	0	0	0.187	0	0	0.100	0.439	0
Total					98.605	98.660	99.362	98.348	86.255	98.322	98.281	99.498	98.690	99.584	79
^a Rt: Retention time (min)															
^b LRI: RI from literature (Adams, 2017).															
^c RI: Experimentally determined															

In the current study, neral was present in all genotypes, except in G3, G11, G14, and G18. A major concentration of neral was in G9 (10.83%) and G2 (10.25%). Geranial was found in G1, G2, G4, G5, G9, G10, G15, and G16 genotypes at low levels (Table 5). Neral and geranial are citral isomers, which have been found in Damask rose essential oil [28]. Farnesol, natural sesquiterpene alcohol in essential oils, was found to have the potential for alleviating massive inflammation, oxidative stress, and lung injury [41, 42]. Farnesol has been widely used in cosmetics, pharmaceuticals, industrial materials, and as a material for carotenoid

and tocopherol [43]. Farnesol is a sesquiterpene trans and exists in some Damask genotypes, a higher amount of it was found in G17 (3.01%) and the highest *e-e* Farnesol was observed in G15 (8.28%).

Rose oxide is an insignificant component of rose oil [44]. In this study, the rose oxide has been found in low concentrations in G3, G11, G15, G19, G22, G23, and G26 (Table 5). Phenethyl alcohol is an enjoyable floral perfume belonging to aromatic alcohols, and one of the main components of rose hydrosols which mainly is used in perfumery [2]. However, this compound was detected at low levels only in some of the selected Damask rose genotypes including G6 (1.54%), G23 (0.40%), and G2 (0.33%). Phytol is one of the major components of plant-derived essential oils which has been recognized for its wide range of pharmacological effects on the nervous system, including anxiolytic, antidepressant, and antimicrobial [45–47]. Several recent studies have suggested that some phytol-derivatives (phytanol, phytanyl amine, and phytanyl mannose) by induction of the expression of a range of chemokines and cytokines effects target tumor cells [48, 49]. Other hydrocarbon-like ingredients, *n*-docosane and *n*-tricosane, were also identified in Damask roses essential oil. Quantities of *n*-tricosane were much more than that of *n*-docosane in all genotypes. In the present study, G6 (9.33%) and G20 (14.20%) genotypes showed the highest contents of *n*-tricosane and *n*-docosane, respectively.

Discussion

Several studies have been conducted to date on the genetic diversity of *R. damascena* in Iran, which has shown a high diversity and genetic variation of this species [6, 8, 50]. In this study, *R. damascena* genotypes showed a remarkable diversity in petal color from dark pink (G3/Tehran genotype) to pale pink (G9/Fars genotype) and white (G2/ Isfahan and G26/ East Azarbayjan genotypes). However, the majority of them were pink or pinkish (Fig. 3; Table 4). Some anthocyanins such as pelargonidin and cyanidin in the petal cells are responsible for the color of rose flowers [51]. Petals of industrial oil-bearing damask roses grown in the world are typically pink, while wild roses usually have pink or white flowers [52]. Karami et al [33] reported a positive relationship between essential oil content and anthocyanin concentration in Damask rose.

The number of petals is a very important indicator of the total essential oil. Significant negative correlations between thorn density and morphological characters, excluding bud diameter, were observed. Additionally, there was a significant positive correlation (0.39**) between the number of petals and thorn density (Table 3). Therefore, it is possible to select genotypes with a higher flower weight and number of flowers in attempts to improve the flower yield and essential oil content [7, 32].

According to the results (Fig. 1, Fig. 2), harvesting time had a major effect on essential oil content, and the morning time harvested flowers had a higher essential oil content. Also, there is no positive relation between oil content and petal number. This is consistent with some reports that the oil content of the damask rose flowers depended on the time of harvesting, and the petals harvested in the morning time had a higher oil content [5,23]. Results of the current study also showed that the essential oil content was influenced by harvesting time in the majority of 26 genotypes of the Damask roses, confirming that morning time was the best time to harvest petals, which is similar to some other reports [53–56].

Large differences were observed between 26 selected Damask rose genotypes for the content of essential oils (Table 5) which was in agreement with the results of researches where reported high variations in the volatile compounds of Damask rose oil [11, 18, 32]. It has been reported that the quantity and composition of essential oil ingredients are significantly influenced by the genotype and agronomic conditions, as well as plant and flower developmental stage and harvesting time [57–59]. Overall, the content of monoterpenes (citronellol, nerol, and *E*-geraniol), sesquiterpenes, and aliphatic hydrocarbons was high (Table 5). Furthermore, *e*-geraniol, one of the major rose-oil components, was high in all 26 selected Damask rose genotypes. The percentage of four major hydrocarbons (heptadecane, nonadecane, eicosane, and heneicosane) were also high in the extracted essential oils (Table 5). Similar to other reports, this study revealed high variations between *Rosa damascena* genotypes regarding oil content and components, morphological diversity, and petal color [9, 60–63].

Conclusions

Results from this study revealed that Damask rose genotypes in Iran have significant diversity in morphological characters, oil content, and also composition. The harvesting time of Damask rose flowers significantly affected the essential oil yield, and for most genotypes harvesting recommended to be performed in the morning, but for higher oil content of G2 and G5 genotypes, evening harvesting time might be recommended. The varied deviations in petal colors, petal numbers, and essential oil content in genotypes were observed in this experiment. Thus, the existence of these characters and a good chemical variation shown in profiling reveals that the studied collection of Damask rose is a good source for the selection of the industrial oil-bearing damask rose cultivars and those that could be used as an ornamental plant in the landscape because of its unique scented flowers. Compared with the other genotypes, G5 and G21 had the highest essential oil content. 25 volatile compounds were identified in the essential oil of Damask rose genotypes. The highest concentration of geraniol, β -citronellol, nonadecane, and β -damascenone were found in G12 (42.51%), G26 (40.82%), G5 (27.76%), and G3 (1.76%) genotypes, respectively. It has been found that the most abundant compounds are of several main classes including alcohols (citronellol, geraniol, nerol) and hydrocarbons (heptadecane, nonadecane, eicosane, and heneicosane). As a final conclusion, results revealed that the morphological and biochemical diversity of Damask rose genotypes can be used effectively to characterize genetic diversity between different genotypes and to select special traits in the breeding program.

Declarations

Acknowledgements

Not applicable

Authors' contributions

MO, AKH, and MK conceived and designed the study; MO and ASH contributed to literature research; MO performed the experiments and collected the results; OR and MO analyzed and interpreted the data; MO and MK were major contributors in writing the manuscript; ZZ, AKH and MK guided all aspects of the research project and revised the manuscript; All authors read and approved the final manuscript.

Funding

This work was supported by Department of Horticulture Science, College of Agriculture and Natural Resources, University of Tehran, Karaj 31587, Iran

Availability of data and materials

All data generated or analysed during this study are included in this published article.

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

References

1. Iwata H, Kato T, Ohno S. Triparental origin of Damask roses. *Gene*. 2000; 259(2): 53-59. [https://doi.org/10.1016/S0378-1119\(00\)00487-X](https://doi.org/10.1016/S0378-1119(00)00487-X)
2. Moein M, Zarshenas MM, Delnavaz S. Chemical composition analysis of rose water samples from Iran. *Pharma Biol*. 2014; 52(10): 1358-1361. <https://doi.org/10.3109/13880209.2014.885062>
3. Moein M, Etemadfar H, Zarshenas MM. Investigation of different Damask rose (*Rosa damascena* Mill.) oil samples from traditional markets in Fars (Iran); focusing on the extraction method. *Trends in Pharma Sci*. 2016; 2(1): 51-58.
4. Rusanov K, Kovacheva N, Vosman B, Zhang L, Rajapakse S, Atanassov A, Atanassov I. Microsatellite analysis of *Rosa damascena* Mill. genotypes reveal genetic similarity between genotypes used for rose oil production and old Damask rose varieties. *Theo and Applied Genetics*. 2005; 111(4): 804-809. <https://doi.org/10.1007/s00122-005-2066-9>
5. Baydar H, Baydar NG. The effects of harvest date, fermentation duration and Tween 20 treatment on essential oil content and composition of industrial oil rose (*Rosa damascena* Mill.). *Ind.Crops Prod*. 2005; 21(2): 251-255. [https://doi.org/10.1016/S0926-6690\(04\)00056-1](https://doi.org/10.1016/S0926-6690(04)00056-1)
6. Babaei A, Tabaei-Aghdai SR, Khosh-Khui M, Omidbaigi R, Naghavi MR, Esselink GD, Smulders MJ. Microsatellite analysis of Damask rose (*Rosa damascena* Mill.) genotypes from various regions in Iran reveals multiple genotypes. *BMC Plant Biol*. 2007; 7(1): 1-6. <https://doi.org/10.1186/1471-2229-7-12>
7. Tabaei-Aghdai SR, Babaei A, Khosh-Khui M, Jaimand K, Rezaee MB, Assareh MH, Naghavi MR. Morphological and oil content variations amongst Damask rose (*Rosa damascena* Mill.) landraces from different regions of Iran. *Sci Horti*. 2007; 113(1): 44-48. <https://doi.org/10.1016/j.scienta.2007.01.010>
8. Kiani M, Zamani Z, Khalighi A, Fatahi R, Byrne DH. Wide genetic diversity of *Rosa damascena* Mill. germplasm in Iran as revealed by RAPD analysis. *Sci Horti*. 2008; 115(4): 386-392. <https://doi.org/10.1016/j.scienta.2007.10.013>
9. Kazaz S, Erbas S, Baydar H, Dilmacunal T, Koyuncu MA. Cold storage of oil rose (*Rosa damascena* Mill.) flowers. *Sci Horti*. 2010; 126(2): 284-290. <https://doi.org/10.1016/j.scienta.2010.06.018>
10. Nasri F, Fadakar A, Yousefi B, Zahedi B. Evaluation of genetic diversity of some Damask rose (*Rosa damascena* Mill.) genotypes of Kurdistan province using morphological traits. *J of Ornamental Plants*. 2016; 6(4): 237-243.
11. Sharma S, Kumar, R. Effect of temperature and storage duration of flowers on essential oil content and composition of Damask rose (*Rosa damascena* Mill.) under western Himalayas. *J. of Applied Res on Med and Aroma Plants*. 2016; 3(1): 10-17. <https://doi.org/10.1016/j.jarmap.2015.10.001>
12. Kovacheva N, Rusanov K, Atanassov I. Industrial cultivation of oil bearing rose and rose oil production in Bulgaria during 21st century, directions and challenges. *Biotech & Biotech Equip*. 2010;24(2): 1793-1798. <https://doi.org/10.2478/V10133-010-0032-4>
13. Global "Rose Essential Oil Market" 2019 Industry Research Report. Market research report. 2019; p.80.
14. Kaul K, Karthigeyan S, Dhyani D, Kaur N, Sharma RK, Ahuja PS. Morphological and molecular analyses of *Rosa damascena* × *R. bourboniana* interspecific hybrids. *Sci Horti*. 2009; 122(2): 258-263. <https://doi.org/10.1016/j.scienta.2009.05.027>
15. Boskabady MH, Shafei MN, Saberi Z, Amini S. Pharmacological effects of *Rosa damascena*. *I. J. B.M. Sci*. 2011; 14(4): 295-307.
16. Gorji-Chakespari A, Nikbakht AM, Sefidkon F, Ghasemi-Varnamkhasti M, Valero EL. Classification of essential oil composition in Damask Rose Mill. genotypes using an electronic nose. *J. of Applied Res on Medi and Aroma Plants*. 2017; 4: 27-34. <https://doi.org/10.1016/j.jarmap.2016.07.004>
17. Dobrev A, Kovacheva N. Daily dynamics of the essential oils of *Rosa damascena* Mill. and *Rosa alba* L. *Agri Sci and Tech*. 2010; 2(2): 71-74.
18. Rusanov K, Kovacheva N, Rusanova M, Atanassov I. Traditional *Rosa damascena* flower harvesting practices evaluated through GC/MS metabolite profiling of flower volatiles. *Food Chem*. 2011; 129(4): 1851-1859. <https://doi.org/10.1016/j.foodchem.2011.05.132>

19. Pal PK, Singh RD. Understanding crop-ecology and agronomy of *Rosa damascena* Mill. for higher productivity. *Aus J of Crop Sci.* 2013;7(2): 196-205.
20. Koksai N, Aslançan H, Sadighzadeh S, Kafkas E. Chemical investigation on *Rosa damascena* Mill. volatiles; Effects of storage and drying conditions. *Acta Scie Polonorum: Hortorum Cultus.* 2015; 14(1): 105-114.
21. Krupčík J, Gorovenko R, Špánik I, Sandra P, Armstrong DW. Enantioselective comprehensive two-dimensional gas chromatography. A route to elucidate the authenticity and origin of *Rosa damascena* Miller essential oils. *Journal of Separation Scie.* 2015; 38(19): 3397-3403. <https://doi.org/10.1002/jssc.201500744>
22. Widrlechner MP. History and utilization of *Rosa damascena*. *Economic Botany.* 1981;35(1): 42-58. <https://doi.org/10.1007/BF02859214>
23. Rusanov K, Kovacheva N, Stefanova K, Atanasov A, Atanasov I. *Rosa damascena*—genetic resources and capacity building for molecular breeding. *Biotech & Biotech Equipt.* 2009; 23(4): 1436-1439. <https://doi.org/10.2478/V10133-009-0009-3>
24. Erbas S, Baydar H. Variation in scent compounds of oil-bearing rose (*Rosa damascena* Mill.) produced by headspace solid phase microextraction, hydrodistillation and solvent extraction. *Rec of Natural Prod.* 2016; 10(5): 555-565.
25. Saint-Lary L, Roy C, Paris JP, Martin JF, Thomas OP, Fernandez X. Metabolomics as a tool for the authentication of rose extracts used in flavor and fragrance area. *Metabolomics.* 2016; 12(3): 49, 1-11. <https://doi.org/10.1007/s11306-016-0963-3>
26. Aydinli M, Tutaş M. Production of rose absolute from rose concrete. *Flavor and Fragr.* 2003; 18(1): 26-31. <https://doi.org/10.1002/ffj.1138>
27. Abdel-Hameed ESS, Bazaid SA, Hagag HA. Chemical characterization of *Rosa damascena* Miller var. Trigintipetala Dieck essential oil and its in vitro genotoxic and cytotoxic properties. *J. of Esse Oil Res.* 2016; 28(2): 121-129. <https://doi.org/10.1080/10412905.2015.1099120>
28. Nedeltcheva-Antonova D, Stoicheva P, Antonov L. Chemical profiling of Bulgarian rose absolute (*Rosa damascena* Mill.) using gas chromatography–mass spectrometry and trimethylsilyl derivatives. *Ind. Crops Prod.* 2017; 108: 36-43. <https://doi.org/10.1016/j.indcrop.2017.06.007>
29. Kiani M, Zamani Z, Khalighi A, Fatahi R, Kiani M. Collection and evaluation of morphological diversity of Damask rose genotypes of Iran. *Iranian J of Horti Sci.* 2011; 41: 223-233.
30. Yousefi B. Screening of *Rosa damascena* Mill. landraces for flower yield and essential oil content in cold climates. *Folia Horticulturae.* 2016; 28(1): 31-40. <https://doi.org/10.1515/fhort-2016-0005>
31. Mitrofanova I, Grebennikova O, Brailko V, Paliy A, Marko N, Lesnikova-Sedoshenko N, Mitrofanova O. Physiological and biochemical features of some cultivars in essential oil rose (*Rosa damascena* Mill.) growing in situ and in vitro. *Int J of PharmTech Res.* 2016; 9(7): 226-232.
32. Baydar H, Erbas S, Kaza, S. Variations in floral characteristics and scent composition and the breeding potential in seed-derived oil-bearing roses (*Rosa damascena* Mill.). *Turkish J. of Agri and Forestry.* 2016; 40(4): 560-569. <https://doi.org/10.3906/tar-1512-57>
33. Karami A, Khosh-Khui M, Salehi H, Saharkhiz MJ, Rowshan V. Headspace analysis of floral scent from two distinct genotypes of Iranian Damask rose (*Rosa damascena* Mill.). *J. of Esse Oil Bear Plant.* 2013; 16(4): 489-498. <https://doi.org/10.1080/0972060X.2013.813266>
34. McGuire RG. Reporting of objective color measurements. *Hort Sci.* 1992; 27(12): 1254-1255. <https://doi.org/10.21273/HORTSCI.27.12.1254>
35. Khan MA, Rehman SU. Extraction and analysis of essential oil of *Rosa* species. *Int. J. Agric. Biol.* 2005; 7:973-974.
36. Shibamoto T. Retention Indices in Essential Oil Analysis (Vol. 259). Huethig Verlag, New York. 1987; p. 259-274.
37. McLafferty FW, Stauffer DB. The Wiley NBS Registry of Mass Spectral Data; Wiley. New York. 1989; p. 518.
38. Adams RP. Identification of Essential Oils by Ion Trap Mass Spectroscopy. Academic press. San Diego, California. 2012; p. 302.
39. Meyer, R., Krueger, D. (2001). Minitab guide to statistics. Prentice Hall PTR.80 p
40. Ohloff G, Demole E. Importance of the odoriferous principle of Bulgarian rose oil in flavour and fragrance chemistry. *Journal of Chroma.* 1987; 406: 181-183. [https://doi.org/10.1016/S0021-9673\(00\)94029-9](https://doi.org/10.1016/S0021-9673(00)94029-9)
41. Derengowski LS, De-Souza-Silva C, Braz SV, Mello-De-Sousa TM, Bão SN, Kyaw CM, Silva-Pereira I. Antimicrobial effect of farnesol, a *Candida albicans* quorum sensing molecule, on *Paracoccidioides brasiliensis* growth and morphogenesis. *Annals of Clinical Microbiology and Antimicrobials.* 2009; 8(1): 13. <https://doi.org/10.1186/1476-0711-8-13>
42. Hammer KA, Carson CF. Antibacterial and antifungal activities of essential oils. *Lipids and essential oils as antimicrobial agents.* 2011; 255-306.
43. Clarke S. Chapter 3-Families of compounds that occur in essential oils. *Essential Chemistry for Aromatherapy.* 2008; 41-77.
44. Leffingwell JC, Leffingwell D. GRAS flavor chemicals-detection thresholds. *Perfumer & Flavorist.* 1991;16(1): 1-19.
45. Murbach Teles Andrade BF, Nunes Barbosa L, da Silva Probst I, Fernandes Júnior A. Antimicrobial activity of essential oils. *Journal of Essential Oil Research.* 2014; 26(1): 34-40. <https://doi.org/10.1080/10412905.2013.860409>
46. Pereira JC, Mario LRJ. Phytol a Natural Diterpenoid with Pharmacological Applications on Central Nervous System: A Review. Recent patents on biotechnology. 2014; 8(3): 194-205.
47. Gutbrod K, Romer J, Dörmann P. Phytol metabolism in plants. *Progress in lipid research.* 2019; 74: 1-17. <https://doi.org/10.1016/j.plipres.2019.01.002>
48. Achoui Y, Chowdhury RR, Fitch RW, Ghosh SK. Molecular signatures of phytol-derived immunostimulants in the context of chemokine–cytokine microenvironment and enhanced immune response. *Cellular immunology.* 2011; 271(2):227-238. <https://doi.org/10.1016/j.cellimm.2011.07.001>
49. Chowdhury RR, Fitch RW, Ghosh SK. Efficacy of phytol-derived diterpenoid immunoadjuvants over alum in shaping the murine host's immune response to *Staphylococcus aureus*. *Vaccine.* 2013;31(8):1178-1186. <https://doi.org/10.1016/j.vaccine.2012.12.069>
50. Kiani M, Zamani Z, Khalighi A, Fatahi R, Byrne DH. Microsatellite analysis of Iranian Damask rose (*Rosa damascena* Mill.) germplasm. *Plant Breeding.* 2010; 129(5): 551-557. <https://doi.org/10.1111/j.1439-0523.2009.01708.x>
51. Schmitzer V, Veberic R, Osterc G, Stampar F. Color and phenolic content changes during flower development in groundcover rose. *A Soci Horti Scie.* 2010; 135(3): 195-202. <https://doi.org/10.21273/JASHS.135.3.195>

52. Tanaka Y, Sasaki N, Ohmiya A. Biosynthesis of plant pigments: Anthocyanin's, betalains and carotenoids. *Plant J.* 2008 54(4): 733-749. <https://doi.org/10.1111/j.1365-313X.2008.03447.x>
53. Ackermann IE, Banthorpe DV, Fordham WD, Kinder JP, Poots I. β -Glucosides of aroma components from petals of *Rosa* species: Assay, occurrence, and biosynthetic implications. *Plant Physiol.* 1989; 134(5): 567-572. [https://doi.org/10.1016/S0176-1617\(89\)80148-8](https://doi.org/10.1016/S0176-1617(89)80148-8)
54. Winterhalter P, Sefton MA, Williams PJ. Two-dimensional GC-DCCC analysis of the glycoconjugates of monoterpenes, norisoprenoids, and shikimate-derived metabolites from Riesling wine. *J. of Agri and Food Chem.* 1990; 38(4): 1041-1048. <https://doi.org/10.1021/jf00094a028>
55. Cherchi G, Deidda D, Gioannis BD, Marongiu B, Pompei R, Porcedda S. Extraction of *Santolina insularis* essential oil by supercritical carbon dioxide: Influence of some process parameters and biological activity. *Flavour and Fragr J.* 2001; 16(1): 35-43. [https://doi.org/10.1002/1099-1026\(200101/02\)16:1%3C35::AID-FFJ942%3E3.0.CO;2-Y](https://doi.org/10.1002/1099-1026(200101/02)16:1%3C35::AID-FFJ942%3E3.0.CO;2-Y)
56. Jamoussi B, Romdhane M, Abderraba A, Hassine BB, Gadri AE. Effect of harvest time on the yield and composition of Tunisian myrtle oils. *Flavour and Fragr J.* 2005; 20(3): 274-277. <https://doi.org/10.1002/ffj.1453>
57. Marotti M, Piccaglia R, Giovanelli E, Deans SG, Eaglesham E. Effects of planting time and mineral fertilization on peppermint (*Mentha x piperita* L.) essential oil composition and its biological activity. *Flavour and Fragr J.* 1994; 9(3): 125-129. <https://doi.org/10.1002/ffj.2730090307>
58. Croteau R, Gershenzon J. Genetic control of monoterpene biosynthesis in mints (*Mentha*: Lamiaceae). In: *Genetic Engineering of Plant Secondary Metabolism*. 1994. pp: 193-229. https://doi.org/10.1007/978-1-4615-2544-8_8
59. Kothari SK, Singh UB. The effect of row spacing and nitrogen fertilization on scotch spearmint (*Mentha gracilis* Sole). *J of Esse Oil Rese.* 1995; 7(3): 287-297. <https://doi.org/10.1080/10412905.1995.9698521>
60. Farooq A, Khan MA, Ali A, Riaz A. Diversity of morphology and oil content of *Rosa damascena* landraces and related *Rosa* species from Pakistan. *Pakistan Journal of Agri Sci.* 2011; 48(3):177-183.
61. Baydar H, Schulz H, Krüger H, Erbas S, Kineci S. Influences of fermentation time, hydro-distillation time and fractions on essential oil composition of Damask Rose (*Rosa damascena* Mill.). *J. of Essential Oil Bearing Plants.* 2008; 11(3): 224-232. <https://doi.org/10.1080/0972060X.2008.10643624>
62. Kumar R, Sharma S, Kaundal M, Sood S, Agnihotri VK. Variation in essential oil content and composition of Damask Rose (*Rosa damascena* Mill) flowers by salt application under mid hills of the western Himalayas. *J of Esse Oil Bearing Plants.* 2016; 19(2): 297-306. <https://doi.org/10.1080/0972060X.2016.1153985>
63. Mohamadi M, Mostafavi A, Shamspur T. Effect of storage on essential oil content and composition of *Rosa damascena* Mill. petals under different conditions. *J of Essen Oil Bearing Plants.* 2011; 14(4), 430-441. <https://doi.org/10.1080/0972060X.2011.10643598>
64. European Pharmacopoeia. European directorate for the quality of medicines and healthcare, Council of Europe. 8th edn. 2014. Vol. 1.

Figures

Figure 1

Essential oil content of fresh petals of *Rosa damascena* harvested in the morning and evening.

Figure 2
Essential oil content of dry petals of *Rosa damascena* harvested at morning and evening

Figure 3

Full blooming flowers of some Rosa damascena genotypes in the present study