

Assessment of Mutations on RBD in the Spike Protein of SARS-CoV-2 Alpha, Delta and Omicron Variants

Clauber Henrique Souza Costa

Federal University of Para

Camila Auad Beltrão Freitas

Federal University of Para

Cláudio Nahum Alves

Federal University of Para

Jerônimo Lameira (✉ lameira@ufpa.br)

Federal University of Para

Research Article

Keywords:

Posted Date: March 17th, 2022

DOI: <https://doi.org/10.21203/rs.3.rs-1401835/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

The severe acute respiratory syndrome (SARS) coronavirus 2 (CoV-2) variant Omicron spread more rapid than the other variants of SARS-CoV-2 virus. Mutations on the spike (S) protein receptor-binding domain (RBD) are critical for the antibody resistance and infectivity of the SARS-CoV-2 variants. In this study, we have used accelerated molecular dynamics (aMD) simulations and free energy calculations to present a systematic analysis of the affinity and conformational dynamics along with the interactions that drive the binding between Spike protein RBD and ACE2 receptor. We evaluate the impacts of the key mutation that occur in the RBDs Omicron and other variants in the binding with the human ACE2 receptor. The results shows that S protein Omicron have stronger binding to the ACE2 than other variants. The evaluation of the decomposition energy per residue shows the mutations N440K, T478K, Q493R and Q498R observed in Spike protein of SARS-CoV-2 provided a stabilization effect for the interaction between the SARS-CoV-2 RBD and ACE2. Overall, the results demonstrate that faster spreading of SARS-CoV-2 omicron may be correlated with binding affinity of S protein RBD to ACE2 and mutations of uncharged residues to positively charged residues such as Lys and Arg in key positions in the RBD.

Introduction

First reported in the city of Wuhan, China ^{1,2}, *Coronavirus disease* (COVID-19) named by World Health Organization (WHO) was declared a global pandemic on March 2020 ³. COVID-19 is caused by the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2). ^{4,5} ^{1,2} The spread of SARS-CoV-2 have cost millions of lives and caused many implications for health, society and the economy ^{6,7}. In January 2022, the WHO reported over 304 million confirmed cases of COVID-19 and over 5.4 million fatalities have been reported since the beginning of the outbreak ⁸. Vaccines are effective for reducing the number deaths by COVID-19 ⁹⁻¹¹. On the other hand, variants may cause impact on the virus recognition by antibody-mediated vaccines ¹²⁻¹⁴.

Different mutations have been reported in the gene encoding the S protein of SARS-CoV-2 ^{15,16}, and recently, the world have faced rapid increase in COVID-19 mediated by new variants ¹⁷. The last variant detected was named Omicron (B.1.1.529) ¹⁸, identified in numerous countries in November 2021, first reported in South African with a large number of mutations, including K417N, S477N, T478K, E484A, and N501Y, which are also found in other variants ¹⁹⁻²¹ and evidences suggests there may be an increased risk of reinfection involving this variant ^{22,23} due to improve viral escape or binding affinity to angiotensin-converting enzyme 2 (ACE2) ²⁴⁻²⁶.

A recent study reported that 45 point mutations was identified and found that the Omicron spike protein sequence was subjected to stronger positive selection than that of any reported SARS-CoV-2 variants ²⁷. Additionally, These mutations and deletions in the S-protein sequence can alter the structure, affecting its stability and function, further exacerbating SARS-CoV-2 infectivity ^{16,28}. However, N501Y mutation is a key contact residue in the RBD, enhancing virus binding affinity to ACE2 ²⁹⁻³¹ making the virus more

contagious and the deletions H69/V70 is required for increase optimal infectivity of Alpha variant, that drives by higher levels of spike incorporation into virions³².

Coronaviruses use spike glycoprotein, with S1 subunit and S2 subunit in each spike monomer, anchored in the virion envelope to bind to their cellular receptors^{33,34} and mediates the recognition of the host-cell receptors and facilitates the cell attachment (S1 subunit) and the cell membrane fusion (S2 subunit) during the viral infection³⁵. The receptor-binding domain (RBD) located in the S1 region (318 to 510 sequence region) performs strongly binds to the peptidase domain of ACE2^{36,37}, leading to a critical virus-receptor interaction and reflects viral host range, tropism and infectivity³⁸. the RBD of S1 undergoes conformational changes that transiently conceal or reveal the determinants of receptor binding^{24,39}

The spike (S) protein of SARS-CoV-2 consists in an extracellular N-terminus, a transmembrane (TM) domain and a intracellular C-terminal segment⁴⁰. S protein has a total length of 1273 amino acids³⁵ and molecular weight of 180–200 kDa⁴¹. It has a signal peptide (1–13) at the N-terminus, followed by S1 subunit (14–685) and the S2 subunit (686–1273)⁴². The structure of the RBD allows for ways to alter its genetic material, developing variants by the changes in spike protein amino acids and as viruses replicate¹⁶, copying errors of itself, resulting in mutations that arise in their genomes generating several strains of SARS-CoV-2^{17,43} that differ in transmission, infectivity and severity of the disease⁴⁴.

ACE2 primary physiological role is in the maturation of angiotensin (Ang)⁴⁵, a peptide hormone that controls vasoconstriction and blood pressure, is a type I membrane protein expressed in lungs, heart, kidneys, and intestine^{25,46}, thereat, decreased expression of ACE2 is associated with cardiovascular diseases⁴⁷. The structural features of RBD increase its binding affinity to the ACE2 receptor and it is a significant step for SARS-CoV-2 to enter into target cells^{33,48}. Computer modelling studies of the interaction between the SARS-CoV-2 RBD and ACE2 were able to identify the residues involved in this interaction and elucidate how the structural change benefits receptor recognition and virus entry into the host cell, that occurs by proteolytic processing of the spike protein to promote cell-virus fusion⁴⁹. Therefore, atomic details may clarify the importance and significance of investigating the changes in residues that facilitate efficient cross-species infection and human-to-human transmission³⁴. Whereas the essential evolution and consequent mutation of SARS-CoV-2 takes place remotely from the RBD in the spike protein, such evolution may facilitate the conformational change in specific residues, punctually interfering with the infection process that occurs after the virus binds to ACE2⁵⁰.

Recently, Warshel and co-workers studied the mechanism of the binding affinity changes for mutations at different spike protein domains of SARS-CoV-2 using coarse-grained potential surface⁵¹. More recently, Chen and co-workers used machine learning model to analyze how the RBD mutations on the Omicron variant may affect the viral infectivity and efficacy of existing vaccines and antibody drugs⁵². In this study, we have used all-atom accelerated molecular dynamics simulations^{53,54} to explore the impacts of the substitutions that occur in the Spike RBD of Alpha, Delta and Omicron variants in the binding with the

human ACE2 receptor. In order to address the question whether variant infectivity and spreading is related to its binding to the receptor.

Methods

SARS-CoV-2 Spike protein (S protein) is a class I fusion homotrimer glycoprotein that is composed a total length 1273 residues⁵⁵ and the binding between the virus and the host cell is mediated by the interaction of the protein S receptor binding domain (RBD, located in the S1 domain) with the angiotensin converting enzyme receptor 2 (ACE2). Here, for the sake of simplicity, S protein RBD from SARS-CoV-2 was renamed as RBD_x, where x represents the identification of each SARS-CoV-2 variant. The initial systems were built considering the coordinates of the RBD complex and the ACE2 (PDB code 6M0J)³³. The protonation states of the protein residues were defined through the propKa program at pH 7⁵⁶. The amino acids were treated with the ff14SB force field⁵⁷ using TLeap module included in AMBER 16⁵⁸. Each system was solvated using TIP3P water⁵⁹ model in a cubic box with 10.0 Å of the amino acid at the end for all Cartesian directions. Then, each system was neutralized using Na⁺ as contra-ions.

We used four minimization steps with 10.000 cycles for each step, applying minimization first to water, contra-ions and protein, in the last step the minimization was applied to all atoms in the system in order to decrease energy, adjust interactions and decrease contacts with conjugate gradient and steepest descent. The systems were heated linearly from 0 to 300 K (temp1 = 0.0; temp0 = 300.0) to avoid excessive and sudden fluctuations of the solute in a time of 5ns in NVT ensemble employing Langevin dynamics as thermostat (collision frequency of 2 ps) had been used to guarantee a system equilibrium. The SHAKE algorithm⁶⁰ was employed to constraints all bonds involving hydrogen atoms.

First, we have performed 10 ns of Classical molecular dynamics (cMD) simulations to calculate the average dihedral and total potentials energies to be taken as reference for the accelerated molecular dynamics (aMD) simulations. Then 200 ns of aMD simulations was carried out for each system: RBD_{WT}-ACE2, RBD_{alpha}-ACE2, RBD_{Omicron}-ACE2 and RBD_{Delta}-ACE2 complex in NPT ensemble. The aMD technique is used to enhance sampling in the protein's conformational space, artificially reducing the energy barriers that separate different states of a given system^{53,54,61-64}. Additionally, we used the Bio3D package⁶⁵ to perform the principal component analysis (PCA). The PCs were obtained from the diagonalization of the covariance matrix obtained from the Cartesian coordinates of the superposed Ca atoms of complex structure. To avoid an underestimate of the atomic displacement, an iterated superposition procedure was applied before the PCA, where residues displaying the largest positional differences were excluded at each round until only the invariant 'core' residues remained⁶⁶⁻⁶⁹.

Protein-protein binding free energy

In this study, we also evaluated the binding energy differences between the complexes and then the decomposition energy was added to assess the energy contribution of each amino acid during the

binding of RBD to ACE2. The binding free energy for the each RBD-ACE2 complex was obtained using:

$$\Delta G_{\text{bind}} = G_{\text{RBD-ACE2}} - G_{\text{RBD}} - G_{\text{ACE2}} \quad (1)$$

Here, $G_{\text{RBD-ACE2}}$ represent the average over the snapshots of a single trajectory of the MD RBD-ACE2 complex, G_{RBD} and G_{ACE2} corresponds to the free energy of RBD and ACE2 protein, respectively. The binding free energy was obtained using MMGBSA method^{70,71} implemented in AMBER 16⁵⁸.

In order to calculate free energy with MMGBSA (Eq. 2) 5000 frames were taken from the 10 ns of MD production using:⁷²⁻⁷⁴

$$\Delta G_{\text{bind,MMGBSA}} = \Delta E_{\text{MM}} + \Delta G_{\text{sol}} - T\Delta S \quad (2)$$

Where, ΔE_{MM} is total gas phase energy (sum of $\Delta E_{\text{internal}}$, $\Delta E_{\text{electrostatic}}$, and ΔE_{vdw}); ΔG_{sol} is sum of polar (ΔG_{GB}) and non-polar (ΔG_{SA}) contributions to solvation. It is important to note that the entropy contribution was not included in the calculations due to the difficulty of accurately calculating entropy for a large protein-protein complex.⁷⁵ It is also worth to note that the frames were taken from the most stable structure observed in PCA graphics.

Results And Discussion

Analysis of Molecular Dynamics of RBD-ACE2 complex

All-atom aMD simulations allowed to explore the conformations of protein-protein complex over time for each system: RBD_{WT}-ACE2, RBD_{alpha}-ACE2, RBD_{Omicron}-ACE2 and RBD_{Delta}-ACE2 complex. Figure 1 shows the RMSD during 200 ns of aMD for each system with respect to the reference structure of the equilibrium step. RBD_{WT}-ACE2, RBD_{alpha}-ACE2 and RBD_{Delta}-ACE2 complexes were within fluctuation in a range of 1 to 3 Å (Figure 1), while the RBD_{Omicron}-ACE2 complex the present the different variation during simulation in a range of 1 to 4 Å (Figure 1). Therefore, the structural equilibrium was reached for all system (Figure 1).

In order to obtain insight into flexibility of each residue in protein-protein complex, we have analyzed the Root-Mean-Square Fluctuations (RMSF) taken into consideration the fluctuations of the backbone atoms. In the RMSF analysis (Figure 2) ACE2 shows the greatest fluctuation in regions 123 to 178 (in magenta), 395 to 425 (in red) and in the region of residues 248 to 368 (in yellow), that moves to interact with the viral RBD. the RBD_{alpha} residues show less fluctuation compared to the WT and its last variants (Delta and Omicron).

In this study, we also explore the flexible region in protein-protein complex. through essential dynamics analysis. The PCA graphs, were obtained using the combinations of PC1vsPC2, PC2vsPC3 and PC3vsPC1 (Figure S2), in which the clusters demonstrate two possible states for all systems in PC1vsPC2. The color scales represent the trajectory time of the MD, separating the beginning of the structures in the initial time of the final structures of the MD, however, the Alpha variant already has a greater number of clusters, where each time interval is separated into small clusters.

For Omicron system the structures are visibly separated into blue structures and red structures (see SI, Figure S2), indicating that the initial structures differ from the final ones, leading to variations in the aMD structures (Figure 3). The PCA analysis showed that the RBD_{WT} and the RBD_{omicron} variant present greater conformational fluctuations, however, the RBD_{alpha} variant stands out for its greater stability. In PC1 there are not many movements in RBD and ACE2 (Figure 3). The main movement of RBD_{WT} and RBD_{omicron} is similar because they have a greater number of movements. The Spike protein, via RBD, when it binds, causes changes in ACE2, as shown in Figure 3. The other conformational changes are shown in PC2 and PC3 in Figure S3 for all systems.

Binding Free Energy MMGBSA and Decomposition by Residue

To assess the affinity of the virus for the human receptor and a possible potential risk of immune evasion by the variants, we calculated the free energy using MM/GBSA (ΔG_{bind} (MMGBSA)) based on the points of greatest stability of the aMD trajectory (see Table 1). The RBD_{omicron} shows the highest binding affinity to ACE2, reflecting the infectivity process, but its conformational fluctuations is similar to the other variants. RBD_{omicron} present an adaptive and non-aggressive process when compared to the RBD_{alpha} (with free energy of binding equal to -62.7836 kcal/mol), which demonstrated the lower free energy than RBD_{WT} (-59.7205 kcal/mol). Based on the higher conformational stability of the Alpha variant the high risk is evident and demonstrates a worrying risk of immune evasion due to its degrees of affinity with ACE2.

The RBD_{Delta} has a higher binding affinity with the human receptor compared to the RBD_{WT} (-66.1357 kcal/mol), which demonstrates the great concern of infections based on this variant. The high risk of infectivity is pointed out as greater among the variants because they have a more favorable ΔG_{bind} in comparison to RBD_{WT}. Therefore, the risk of evolution and emergence of new variants may represent a major health concern due to the degree of affinity that evolves the greater affinity for the human receptor.

Table 1: Binding free energy for native systems (SARS-CoV-2) and variants (Alpha/Delta/Omicron).

Energy (kcal/mol)	WT	Alpha	Delta	Omicron
ΔE_{vdw}	-95.6(0.18)	-107.3(0.21)	-103.4(0.16)	-96.4(0.16)
ΔE_{ele}	-625.8(0.94)	-608.5(0.91)	-955.1(1.05)	-1381.7(1.24)
ΔE_{GB}	675.0(0.87)	667.5(0.87)	1006.3(1.01)	1416.2(1.15)
ΔE_{surf}	-13.4(0.02)	-14.5(0.02)	-13.9(0.02)	-13.5(0.02)
ΔG_{gas}	-721.3(0.96)	-715.8(0.91)	-1058.5(1.09)	-1478.1(1.24)
ΔG_{sol}	661.6(0.86)	653.0(0.86)	992.4(0.99)	1402.7(1.15)
$\Delta G_{\text{bind (MMGBSA)}}$	-59.7(0.28)	-62.8(0.23)	-66.1(0.21)	-75.4(0.23)

The effect of mutations can be investigated through the free energy calculations that track the influence of changes in certain positions⁷⁶. The results of the energy of decomposition by residue for RBD_{WT}-ACE2, RBD_{Alpha}-ACE2, RBD_{Omicron}-ACE2 and RBD_{Delta}-ACE2 complex demonstrate that the RBD is the region that has more energy variations, attractive and repulsive, when evaluated the electrostatic contributions (see Figure 4, Figure S4, Figure S5 and Figure S6). The evaluation of the decomposition energy per residue shows the mutations N440K, T478K, Q493R and Q498R observed in RBD_{Omicron} provide favorable interaction between RBD_{Omicron} and ACE2. Curiously, all these mutations include positively charged residues Lys or Arg (see Table 2). For example, K478 in RBD_{Omicron} present a stabilization effect (-85.8 kcal/mol), while T478 in RBD_{WT} has a destabilization effect (0.7 kcal/mol), see Table 2. Additionally, Table S2 shows the hydrogen bonds in the protein-protein interaction for the SARS-Cov-2, Alpha, Delta and Omicron system.

The N501Y mutation in the RBD_{Alpha} has a very similar contribution to the native RBD system, indicating that this mutation does not cause such apparent changes in the energetic contributions, therefore the main feature that contributes to the better binding of RBD_{Alpha} to ACE2, compared to the RBD_{WT}, it is its conformational stability that differs from other spikes. The alterations in the Delta variant cause a highly attractive energy, in which the residue L352R had an energetic contribution of -90,524 kcal/mol and T478K equal to -82,654 kcal/mol (see table 2), indicating that there is a great improvement in the binding with the receptor. The mutations present in RBD_{Omicron} demonstrate that during the gain in the energetic contribution of the residues.

Some mutations present in RBD_{Omicron} (N440K, T478K, Q493R, Q498R) demonstrate that substitutions for positively charged residues guide an improvement in the contribution to the interaction with ACE2 (Figure S7). T478K is located in a more solvent-oriented region, allowing interaction with ACE2, due to the increase in the side chain Figure S7a. As well, the Q493R substitution allows favorable interaction with

negatively charged residues of ACE2 such as Asp38 and Glu35, improving the binding with the receptor and increasing the affinity of the spike protein (Figure S7b). The N440K in the omicron is located in the region most focused on the solvent, increasing the contribution of this region with the medium (Figure S7c), whereas the Q498R substitution improves the protein-protein interaction since this contribution is 24 times greater in relation to the WT, demonstrating that these substitutions are essential for improving interaction with ACE2 (Figure S7d).

Table 2: Decomposition energies per residue in kcal/mol for the main mutation positions of RBD WT, Alpha, Delta and Omicron.

SARS-CoV-2		Alpha		Delta		Omicron	
G339	0.7		0.8		0.8	G339D	68.4
S371	0.8		0.4		1.1	S371L	0.8
S373	1.0		0.6		0.9	S373P	0.6
S375	-0.3		-0.3		-0.1	S375F	-0.4
K417	-121.2		-131.5		-112.4	K417N	-2.3
N440	-0.4		-0.3		-0.2	N440K	-98.6
G446	0.3		-0.2		-0.2	G446S	1.2
L452	-0.7		-0.6	L452R	-90.5	L452	-1.4
S477	-1.1		-1.7		-1.4	S477N	-0.6
T478	0.7		-2.4	T478K	-82.6	T478K	-85.8
E484	88.2		94.4		93.8	E484A	0.1
Q493	-8.7		-11.6		-8.8	Q493R	-163.7
G496	-3.6		-3.1		-4.9	G496S	-6.3
Q498	-6.7		-2.1		-7.4	Q498R	-161.0
N501	-8.6	N501Y	-8.1		-10.1	N501Y	-2.2
Y505	-7.4		-5.5		-8.0	Y505H	-1.4

Conclusion

In this study, we evaluated the effect of residues mutation on structural and energetics of Spike protein RBD from SARS-CoV-2 variants in complex with the human ACE2 receptor. All-atoms accelerated Molecular Dynamics simulations and PCA analysis shows that that the RBD_{Omicron}-ACE2 complex present similar fluctuation in comparison to S protein from Native, Delta and Alpha variants. The binding

affinity of each RBD_x to ACE2 was obtained using MM-GBSA methods. The results shows that the trend in the calculated binding free energies correlates well with virus infectivity of each variant. The mutation in RBD_{Omicron} increase the affinity of Spike protein for ACE2 and may explain Omicron's high transmissibility in comparison with other SARS-CoV-2 variants. The stabilization effect RBD_{Omicron}-ACE2 complex is achieved manly due the substitution of uncharged residues by positively charged residues: Lys and Arg in key positions. Overall, our results may explain at molecular level the effect of key mutations in the Spike protein for virus infectivity.

Declarations

Data availability

All necessary files to conduct this work (.pdb and .parm7) can be found attached as the Supporting Information. The AMBER18 suite of programs and the Amber ff14SB force field were used to carry out the MD simulations and can found at <https://ambermd.org/>.

Acknowledgements

Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) and Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) for their financial support. We also thank the access of the computational resources of the Supercomputer Santos Dumont (SDumont) provided by the Laboratório de Computação Científica (LNCC), Apollo 2000 and CABANO – UFPA.

Author information

These authors contributed equally: Clauber Henrique Souza da Costa, Camila Auad Beltrão de Freitas.

References

1. Wang, D. *et al.* Clinical Characteristics of 138 Hospitalized Patients With 2019 Novel Coronavirus–Infected Pneumonia in Wuhan, China. *JAMA* **323**, 1061–1069 (2020).
2. Huang, C. *et al.* Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* **395**, (2020).
3. Who, W. H. O. Coronavirus disease (COVID-19) outbreak (World Health Organization, 2020). 2020 (2020).
4. Wu, F. *et al.* A new coronavirus associated with human respiratory disease in China. *Nature* **579**, 265–269 (2020).
5. Zhu, N. *et al.* A Novel Coronavirus from Patients with Pneumonia in China, 2019. *N. Engl. J. Med.* **382**, 727–733 (2020).
6. Osterrieder, A. *et al.* Economic and social impacts of COVID-19 and public health measures: results from an anonymous online survey in Thailand, Malaysia, the UK, Italy and Slovenia. *BMJ Open* **11**,

- e046863 (2021).
7. Clemente-Suárez, V. J. *et al.* The Impact of the COVID-19 Pandemic on Social, Health, and Economy. *Sustainability* **13**, (2021).
 8. WHO. COVID-19 weekly epidemiological update. *World Heal. Organ.* 1–23 (2021).
 9. Fiolet, T., Kherabi, Y., MacDonald, C.-J., Ghosn, J. & Peiffer-Smadja, N. Comparing COVID-19 vaccines for their characteristics, efficacy and effectiveness against SARS-CoV-2 and variants of concern: a narrative review. *Clin. Microbiol. Infect. Off. Publ. Eur. Soc. Clin. Microbiol. Infect. Dis.* (2021). doi:10.1016/j.cmi.2021.10.005
 10. Alencar, C. H. *et al.* High Effectiveness of SARS-CoV-2 Vaccines in Reducing COVID-19-Related Deaths in over 75-Year-Olds, Ceará State, Brazil. *Trop. Med. Infect. Dis.* **6**, (2021).
 11. Gupta, S. *et al.* Vaccinations Against COVID-19 May Have Averted Up To 140,000 Deaths In The United States. *Health Aff. (Millwood)*. **40**, 1465–1472 (2021).
 12. Buchan, S. A. *et al.* Effectiveness of COVID-19 vaccines against Omicron or Delta infection. *medRxiv* 2021.12.30.21268565 (2022). doi:10.1101/2021.12.30.21268565
 13. Eyre, D. W. *et al.* Effect of Covid-19 Vaccination on Transmission of Alpha and Delta Variants. *N. Engl. J. Med.* (2022). doi:10.1056/NEJMoa2116597
 14. Lopez Bernal, J. *et al.* Effectiveness of Covid-19 vaccines against the B. 1.617. 2 (Delta) variant. *N Engl J Med* 585–594 (2021).
 15. Dawood, A. A. Mutated COVID-19 may foretell a great risk for mankind in the future. *New microbes new Infect.* **35**, 100673 (2020).
 16. Korber, B. *et al.* Tracking Changes in SARS-CoV-2 Spike: Evidence that D614G Increases Infectivity of the COVID-19 Virus. *Cell* **182**, 812-827.e19 (2020).
 17. Villoutreix, B. O., Calvez, V., Marcelin, A.-G. & Khatib, A.-M. In Silico Investigation of the New UK (B.1.1.7) and South African (501Y.V2) SARS-CoV-2 Variants with a Focus at the ACE2-Spike RBD Interface. *Int. J. Mol. Sci.* **22**, (2021).
 18. Who, W. H. O. Classification of Omicron. 11–12 (2021).
 19. Hodcroft, E. B. CoVariants: SARS-CoV-2 Mutations and Variants of Interest. (2021).
 20. Hadfield, J. *et al.* Nextstrain: real-time tracking of pathogen evolution. *Bioinformatics* **34**, 4121–4123 (2018).
 21. Wang, L. & Cheng, G. Sequence analysis of the emerging SARS-CoV-2 variant Omicron in South Africa. *J. Med. Virol.* **n/a**, (2021).
 22. Who, W. H. O. Update on Omicron. *World Health Organization* 1–5 (2021).
 23. Pulliam, J. R. C. *et al.* Increased risk of SARS-CoV-2 reinfection associated with emergence of the Omicron variant in South Africa. *medRxiv* 2021.11.11.21266068 (2021). doi:10.1101/2021.11.11.21266068
 24. Lim, H. *et al.* Hot spot profiles of SARS-CoV-2 and human ACE2 receptor protein protein interaction obtained by density functional tight binding fragment molecular orbital method. *Sci. Rep.* **10**, 16862

- (2020).
25. Zhang, H. *et al.* The digestive system is a potential route of 2019-nCov infection: a bioinformatics analysis based on single-cell transcriptomes. *bioRxiv* 2020.01.30.927806 (2020). doi:10.1101/2020.01.30.927806
 26. Wang, P. *et al.* Increased resistance of SARS-CoV-2 variant P. 1 to antibody neutralization. *Cell Host Microbe* **29**, 747–751 (2021).
 27. Wei, C. *et al.* Evidence for a mouse origin of the SARS-CoV-2 Omicron variant. *J. Genet. Genomics* (2021). doi:https://doi.org/10.1016/j.jgg.2021.12.003
 28. Berger, I. & Schaffitzel, C. The SARS-CoV-2 spike protein: balancing stability and infectivity. *Cell Res.* **30**, 1059–1060 (2020).
 29. Starr, T. N. *et al.* Deep Mutational Scanning of SARS-CoV-2 Receptor Binding Domain Reveals Constraints on Folding and ACE2 Binding. *Cell* **182**, 1295-1310.e20 (2020).
 30. Tian, F. *et al.* N501Y mutation of spike protein in SARS-CoV-2 strengthens its binding to receptor ACE2. *Elife* **10**, e69091 (2021).
 31. Luan, B., Wang, H. & Huynh, T. Molecular Mechanism of the N501Y Mutation for Enhanced Binding between SARS-CoV-2's Spike Protein and Human ACE2 Receptor. *bioRxiv* 2021.01.04.425316 (2021). doi:10.1101/2021.01.04.425316
 32. Meng, B. *et al.* Recurrent emergence of SARS-CoV-2 spike deletion H69/V70 and its role in the Alpha variant B.1.1.7. *Cell Rep.* **35**, 109292 (2021).
 33. Lan, J. *et al.* Structure of the SARS-CoV-2 spike receptor-binding domain bound to the ACE2 receptor. *Nature* **581**, 215–220 (2020).
 34. Li, F., Li, W., Farzan, M. & Harrison, S. C. Structure of SARS coronavirus spike receptor-binding domain complexed with receptor. *Science* **309**, 1864–1868 (2005).
 35. Huang, Y., Yang, C., Xu, X., Xu, W. & Liu, S. Structural and functional properties of SARS-CoV-2 spike protein: potential antiviral drug development for COVID-19. *Acta Pharmacol. Sin.* **41**, 1141–1149 (2020).
 36. Xiao, X., Chakraborti, S., Dimitrov, A. S., Gramatikoff, K. & Dimitrov, D. S. The SARS-CoV S glycoprotein: expression and functional characterization. *Biochem. Biophys. Res. Commun.* **312**, 1159–1164 (2003).
 37. Wong, S. K., Li, W., Moore, M. J., Choe, H. & Farzan, M. A 193-amino acid fragment of the SARS coronavirus S protein efficiently binds angiotensin-converting enzyme 2. *J. Biol. Chem.* **279**, 3197–3201 (2004).
 38. Junxian, O. *et al.* V367F Mutation in SARS-CoV-2 Spike RBD Emerging during the Early Transmission Phase Enhances Viral Infectivity through Increased Human ACE2 Receptor Binding Affinity. *J. Virol.* **95**, e00617-21 (2021).
 39. Daniel, W. *et al.* Cryo-EM structure of the 2019-nCoV spike in the prefusion conformation. *Science* (80-.). **367**, 1260–1263 (2020).

40. Bosch, B. J., van der Zee, R., de Haan, C. A. M. & Rottier, P. J. M. The coronavirus spike protein is a class I virus fusion protein: structural and functional characterization of the fusion core complex. *J. Virol.* **77**, 8801–8811 (2003).
41. Hoffmann, M. *et al.* SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is Blocked by a Clinically Proven Protease Inhibitor. *Cell* **181**, 271-280.e8 (2020).
42. Huang, Y., Yang, C., Xu, X., Xu, W. & Liu, S. Structural and functional properties of SARS-CoV-2 spike protein: potential antiviral drug development for COVID-19. *Acta Pharmacol. Sin.* **41**, 1141–1149 (2020).
43. Harvey, W. T. *et al.* SARS-CoV-2 variants, spike mutations and immune escape. *Nat. Rev. Microbiol.* **19**, 409–424 (2021).
44. Harvey, W. T. *et al.* SARS-CoV-2 variants, spike mutations and immune escape. *Nat. Rev. Microbiol.* **19**, 409–424 (2021).
45. Donoghue, M. *et al.* A novel angiotensin-converting enzyme–related carboxypeptidase (ACE2) converts angiotensin I to angiotensin 1-9. *Circ. Res.* **87**, e1–e9 (2000).
46. Zhao, Y. *et al.* Single-cell RNA expression profiling of ACE2, the receptor of SARS-CoV-2. *bioRxiv* 2020.01.26.919985 (2020). doi:10.1101/2020.01.26.919985
47. Guo, J., Huang, Z., Lin, L. & Lv, J. Coronavirus Disease 2019 (COVID-19) and Cardiovascular Disease: A Viewpoint on the Potential Influence of Angiotensin-Converting Enzyme Inhibitors/Angiotensin Receptor Blockers on Onset and Severity of Severe Acute Respiratory Syndrome Coronavirus 2 Infection. *J. Am. Heart Assoc.* **9**, e016219 (2020).
48. Shang, J. *et al.* Structural basis of receptor recognition by SARS-CoV-2. *Nature* **581**, 221–224 (2020).
49. Yushun, W. *et al.* Receptor Recognition by the Novel Coronavirus from Wuhan: an Analysis Based on Decade-Long Structural Studies of SARS Coronavirus. *J. Virol.* **94**, e00127-20 (2021).
50. Bai, C. & Warshel, A. Critical Differences between the Binding Features of the Spike Proteins of SARS-CoV-2 and SARS-CoV. *J. Phys. Chem. B* **124**, 5907–5912 (2020).
51. Bai, C. *et al.* Predicting Mutational Effects on Receptor Binding of the Spike Protein of SARS-CoV - 2 Variants. (2021). doi:10.1021/jacs.1c07965
52. Chen, J., Wang, R., Gilby, N. B. & Wei, G. Omicron Variant (B . 1 . 1 . 529): Infectivity , Vaccine Breakthrough , and Antibody Resistance. (2021). doi:10.1021/acs.jcim.1c01451
53. Hamelberg, D., Mongan, J. & McCammon, J. A. Accelerated molecular dynamics: a promising and efficient simulation method for biomolecules. *J. Chem. Phys.* **120**, 11919–11929 (2004).
54. Kukol, A. *Molecular modeling of proteins: Second edition. Molecular Modeling of Proteins: Second Edition* **1215**, (2014).
55. Watanabe, Y., Allen, J. D., Wrapp, D., McLellan, J. S. & Crispin, M. Site-specific glycan analysis of the SARS-CoV-2 spike. *Science* **369**, 330–333 (2020).
56. Søndergaard, C. R., Olsson, M. H. M., Rostkowski, M. & Jensen, J. H. Improved Treatment of Ligands and Coupling Effects in Empirical Calculation and Rationalization of pKa Values. *J. Chem. Theory*

- Comput.* **7**, 2284–2295 (2011).
57. Maier, J. A. *et al.* ff14SB: Improving the Accuracy of Protein Side Chain and Backbone Parameters from ff99SB. *J. Chem. Theory Comput.* **11**, 3696–3713 (2015).
58. D.A. Case, R.M. Betz, D.S. Cerutti, T.E. Cheatham, III, T.A. Darden, R.E. Duke, T.J. Giese, H. Gohlke, A.W. Goetz, N. Homeyer, S. Izadi, P. Janowski, J. Kaus, A. Kovalenko, T.S. Lee, S. LeGrand, P. Li, C. Lin, T. Luchko, R. Luo, B. Madej, D. Mermelstein, L. X. and P. A. K. AMBER 2016. *Univ. California, San Fr.* (2016).
59. Jorgensen, W. L., Chandrasekhar, J., Madura, J. D., Impey, R. W. & Klein, M. L. Comparison of simple potential functions for simulating liquid water. *J. Chem. Phys.* **79**, 926–935 (1983).
60. Kräutler, V., van Gunsteren, W. F. & Hünenberger, P. H. A fast SHAKE algorithm to solve distance constraint equations for small molecules in molecular dynamics simulations. *J. Comput. Chem.* **22**, 501–508 (2001).
61. Patrick, R. *et al.* Using Accelerated Molecular Dynamics Simulation to elucidate the effects of the T198F mutation on the molecular flexibility of the West Nile virus envelope protein. 1–6 (2020). doi:10.1038/s41598-020-66344-8
62. Markwick, P. R. L. & McCammon, J. A. Studying functional dynamics in bio-molecules using accelerated molecular dynamics. *Phys. Chem. Chem. Phys.* **13**, 20053–20065 (2011).
63. Roe, D. R., Bergonzo, C. & Cheatham, T. E. Evaluation of Enhanced Sampling Provided by Accelerated Molecular Dynamics with Hamiltonian Replica Exchange Methods. *J. Phys. Chem. B* **118**, 3543–3552 (2014).
64. Li, C. *et al.* Conformational Changes of Glutamine 5'-Phosphoribosylpyrophosphate Amidotransferase for Two Substrates Analogue Binding: Insight from Conventional Molecular Dynamics and Accelerated Molecular Dynamics Simulations. *Front. Chem.* **9**, 51 (2021).
65. Grant, B. J., Rodrigues, A. P. C. C., ElSawy, K. M., McCammon, J. A. & Caves, L. S. D. D. Bio3d: An R package for the comparative analysis of protein structures. *Bioinformatics* **22**, 2695–2696 (2006).
66. da Costa, C. H. S. *et al.* Assessment of the PETase conformational changes induced by poly(ethylene terephthalate) binding. *Proteins Struct. Funct. Bioinforma.* **n/a**, (2021).
67. Costa, C. H. S. *et al.* Computational study of conformational changes in human 3-hydroxy-3-methylglutaryl coenzyme reductase induced by substrate binding. *J. Biomol. Struct. Dyn.* **37**, 4374–4383 (2019).
68. Costa, C. H. S. da *et al.* Unraveling the conformational dynamics of glycerol 3-phosphate dehydrogenase, a nicotinamide adenine dinucleotide-dependent enzyme of *Leishmania mexicana*. *J. Biomol. Struct. Dyn.* 1–12 (2020). doi:10.1080/07391102.2020.1742206
69. Grosso, M., Kalstein, A., Parisi, G., Roitberg, A. E. & Fernandez-Alberti, S. On the analysis and comparison of conformer-specific essential dynamics upon ligand binding to a protein. *J. Chem. Phys.* **142**, 245101 (2015).
70. Srinivasan, J., Cheatham, T. E., Cieplak, P., Kollman, P. A. & Case, D. A. Continuum Solvent Studies of the Stability of DNA, RNA, and Phosphoramidate–DNA Helices. *J. Am. Chem. Soc.* **120**, 9401–9409

(1998).

71. Kollman, P. a *et al.* Calculating structures and free energies of complex molecules: combining molecular mechanics and continuum models. *Acc. Chem. Res.* **33**, 889–97 (2000).
72. Lill, M. A. & Thompson, J. J. Solvent interaction energy calculations on molecular dynamics trajectories: increasing the efficiency using systematic frame selection. *J. Chem. Inf. Model.* **51**, 2680–2689 (2011).
73. Case, D. A. *et al.* The Amber biomolecular simulation programs. *J. Comput. Chem.* **26**, 1668–1688 (2005).
74. Cui, Q. *et al.* Molecular Dynamics–Solvated Interaction Energy Studies of Protein–Protein Interactions: The MP1–p14 Scaffolding Complex. *J. Mol. Biol.* **379**, 787–802 (2008).
75. Yang, Y., Liu, H. & Yao, X. Understanding the molecular basis of MK2-p38 α signaling complex assembly: insights into protein-protein interaction by molecular dynamics and free energy studies. *Mol. Biosyst.* **8**, 2106–2118 (2012).
76. Bai, C. *et al.* Predicting Mutational Effects on Receptor Binding of the Spike Protein of SARS-CoV-2 Variants. *J. Am. Chem. Soc.* **143**, 17646–17654 (2021).

Figures

Figure 1

RMSD for RBD_{WT}-ACE2, RBD_{alpha}-ACE2, RBD_{Omicron}-ACE2 and RBD_{Delta}-ACE2 complexes.

Figure 2

three-dimensional structure of ACE2 and RBD with RMSF regions for SARS-CoV-2, Alpha, Delta and Omicron systems.

Figure 3

Movements described for the first principal component (PC1) for each structure of ACE2 and RBD. A) moving in PC1 to the native RBD complex (SARS-CoV-2) and ACE2 receptor. B) changes in PC1 to RBD_{Alpha} and ACE2. C) Change moving of PC1 to RBD_{Delta} and ACE2. D) moving from PC1 to the complex between RBD_{Omicron} and ACE2. In turquoise, the initial structure of the movement, in dark magenta, the final structure and in gray, the intermediate structures of the movement. The conformational dynamics were obtained from 200 ns of aMD simulations.

Figure 4

a) three-dimensional structure of the RBD_{WT} and ACE2 complex with the electrostatic energy regions. B) Decomposition energy per residue for the RBD_{WT} system connected to ACE2. The label in orange is from the ACE2 region and in purple is from RBD.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SISARS100322.docx](#)