

Nurses' Knowledge and Attitudes Towards Palliative Care and Death: A Learning Intervention

Yanping Hao

College of Nursing of Guangzhou Medical University

Lixuan Zhan

Department of Neurology of the Second Affiliated Hospital of Guangzhou Medical University

Meiling Huang

Nursing Department of the Third Affiliated Hospital of Guangzhou Medical University

Xianying Cui

Nursing Department of Chinese Hospital in Yuexiu District

Ying Zhou

College of Nursing of Guangzhou Medical University

En Xu (✉ enxu@163.net)

The Second Affiliated Hospital of Guangzhou Medical University

Research article

Keywords: Palliative care, nursing, Internet-based intervention, knowledge, attitude

Posted Date: January 12th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-141816/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published on March 25th, 2021. See the published version at <https://doi.org/10.1186/s12904-021-00738-x>.

Abstract

Background

In many countries, nurses are ill-prepared to provide care to patients with terminal illnesses. Limited education and training affect their ability to deliver proper palliative care. Only a few studies have explored appropriate and effective training methods of palliative care in China. Therefore, we aimed to provide evidence for a palliative care training system by appraising the effects of a mixed-method intervention on participants' knowledge of palliative care and attitudes towards dying patients and death.

Methods

An e-learning intervention approach was adopted for 97 nurses from oncology departments across five hospitals, using a mobile terminal combined with a virtual forum and face-to-face interactions. We conducted a pre- and post-training evaluation through the Palliative Care Quiz of Nursing (PCQN), Frommelt Attitude Toward Care of the Dying Scale Form B (FATCOD-B), and Death Attitude Profile-Revised (DAP-R).

Results

After a three-week intervention, there was a significant increase in the PCQN and FATCOD-B scores as compared to the baseline. For PCQN, the total score increased from 10.3 ± 1.9 to 11.1 ± 2.2 ($p = .011$) and the score for management of pain and other symptoms increased from 7.7 ± 1.7 to 8.4 ± 1.7 ($p = .003$). FATCOD-B scores increased noticeably from 104.0 ± 7.6 to 106.4 ± 8.9 ($p = .016$). The DAP-R scores showed no obvious difference between pre- and post-intervention results.

Conclusions

The mixed-method intervention was effective in improving participants' knowledge and attitudes about palliative care. The implementation of training nurses at appropriate intervals during both education and professional life are required, especially for the improvement in participants' attitudes towards death. Therefore, palliative care training in China should receive more attention.

Background

Global population ageing and the rising prominence of chronic non-communicable diseases have resulted in an increased demand for palliative care nursing.[1] With the proportion of patients with incurable diseases increasing each day,[2] the need for high-quality palliative care has also consequently increased challenges for healthcare educators.[3, 4] The implementation of palliative care in China is widely restricted by limited availability of services, shortage of skilled staff, and the lack of appropriate training systems.[5, 6] Many patients suffering from life-threatening illnesses are struggling with poor health conditions due to unmet care needs towards the end of their lives.[5, 7] Providing professionally

trained palliative care staff is a potential solution for improving the quality of life of patients facing life-threatening illnesses.

The palliative care competence of nurses is a strong contributor to improving the outcomes of high-quality care.[8] However, Chinese nurses face barriers in their clinical competencies as they do not receive adequate palliative care education in nursing schools,[9, 10] and associated professional training in this field at their workplaces.[11, 12] Accordingly, educational training is a crucial element for enhancing nurses' competence.[13] To eliminate this educational gap, establishment of a professional training system in palliative care for nurses in China is imperative.

Several key factors are associated with the competence of nurses in palliative care, including their knowledge and attitudes towards palliative care and death.[14–16] Previous studies indicated that educational training for healthcare professionals had a positive influence on their confidence, knowledge of palliative care, and attitudes towards caring for dying patients and death.[17–20]

Traditional face-to-face training is increasingly being replaced by online education. Electronic learning (e-learning) is a feasible and acceptable approach towards the delivery of high-quality education. The effectiveness of online training in palliative care has mainly been demonstrated through computer-assisted learning.[21, 22] Few studies have reported on these educational programmes, particularly by mobile terminals. Because the location of learning is no longer restricted to a fixed place, it is easy to take the course at any given location.

A professional training system of palliative care is urgently needed in China and evaluating the effectiveness of this system will assemble reliable evidence and quality scientific studies to move forward. In this study, we aimed to provide evidence for this future system by investigating the training needs of palliative care nurses and customising an educational intervention according to their feedback. We implemented the intervention based on e-learning via a mobile terminal, combined with a virtual discussion forum and face-to-face interactions, which was followed by evaluating the training effect on nurses' knowledge of palliative care and their attitudes towards caring for dying patients and death.

Methods

Study design

This study employed a non-randomised, single-arm design to explore the effect of an intervention programme, for nurses, on the knowledge of palliative care and attitudes towards caring for dying patients and death. Questionnaires were distributed among participating nurses from five hospitals.

Settings and participants

From September to December 2018, we conducted the study across five hospitals: The Second Affiliated Hospital of Guangzhou Medical University, The Third Affiliated Hospital of Guangzhou Medical University, Community Health Service Centre in Hongshan Street, People' Hospital in Liwan District, and Hospital of

Traditional Chinese Medicine in Yuexiu District, Guangzhou, China. One hundred and twenty-five nurses from the oncology departments of these hospitals were screened for eligibility and 109 of them were enrolled. All participants were registered nurses, aged 18 to 60 years. The sample size was estimated by

$$N = \left[\frac{(t_{\alpha} + t_{\beta})s}{\delta} \right]^2 \approx 96, \alpha = 0.05, \beta = 0.1, \delta = 0.3S.[23]$$

Measurements

The demographic characteristics collected from the participants were: gender, age, education level, received school education in palliative care, work experience, training experience in palliative care before the intervention, and training needs in palliative care (Table 1). The major outcomes of the study were measured by the Palliative Care Quiz of Nursing (PCQN),[24] the Frommelt Attitude Toward Care of the Dying Scale Form B (FATCOD-B),[25] and the Death Attitude Profile-Revised (DAP-R).[26] All scales had confirmed reliability and validity.[27-30]

TABLE 1 Demographic characteristics of participating nurses

Variable	n	%
Gender		
female	94	96.9
Age (years)		
18- 30	72	74.2
31- 40	18	18.6
41- 50	7	7.2
Education		
associate' s degree (3 years)	12	12.4
bachelor's degree	37	38.1
master's degree or above	48	49.5
Working experience (years)		
≤2	16	16.5
3- 5	49	50.5
6- 9	13	13.4
≥10	19	19.6
Training needs on palliative care		
no	3	3.1
yes	94	96.9
Training experience in palliative care		
no	58	59.8
yes	39	40.2

The PCQN covers three categories in end-of-life care: (i) philosophy and principles of palliative care (4 items), (ii) management of pain and other symptoms (13 items), and (iii) psychosocial aspects of care (3 items). Total scores ranged from 0 to 20, with higher scores indicating better knowledge.

The FATCOD-B comprised 30 items. The scores ranged from 30 to 150, with higher scores indicating a more positive attitude towards care for the dying.

The DAP-R, a 32-item scale comprised five subscales: fear of death (7 items), death avoidance (5 items), neutral acceptance (5 items), escape acceptance (5 items), and approach acceptance (10 items). Each

subscale, with scores ranging from 1 to 7, represented one of the five attitudes towards death and a higher score indicated a greater tendency to share the corresponding attitude towards death.

Procedure

At the beginning of the research, the participants were provided a thorough explanation of the study and notified that the participation was voluntary, and they could withdraw from the study at any time. All questionnaires were anonymous, and the participants were informed about data confidentiality. The self-administered, written questionnaire was delivered to each participant to complete at two time-points: pre-intervention and post-intervention. The first phase of the survey involved participants' demographic characteristics, training requirements for palliative care, and assessment of their knowledge of palliative care and attitudes towards caring for dying patients and death. The training course, as an intervention, was customised according to the information collected from participants regarding training requirements and implemented over a three-week period. Participants were then given the PCQN, FATCOD-B, and DAP-R questionnaires for the second phase of the survey immediately upon completion of the intervention. Finally, baseline data were compared with post-intervention data to evaluate the effects of training on nurses' knowledge of palliative care and their attitudes towards caring for the dying patients and death. The Ethics Committee of Guangzhou Medical University (L180920) approved the study protocol.

Intervention

The training content of the educational intervention was designed based on the data collected on participants' training requirements in palliative care. It consisted of three modules: life and death education, bereavement, and specialised nursing in palliative care. The courses in these modules were video-recorded and conducted by a professor who specialised in humanity science. Participants completed 45 video segments in the three modules. They could access the educational material with ease, free of charge, and could repeatedly play each video segment via the We-chat application on their mobile phones or tablets. They could complete the learning tasks in their own time and environment via Wi-Fi or a 4G mobile communication network.

One researcher acted as the administrator and principal investigator of the study, designing the learning progress plan (course segments of Module 1 were asked to be completed in two weeks and Modules 2 and 3 to be completed in one week) and following the status of learning through both virtual and face-to-face interactions. For virtual interactions, the researcher established a forum via We-chat ensuring that all participants joined through their mobile phones. During this interaction, all topics related to the training programme could be discussed by the participants by sending instant messages during their learning experience.

For face-to-face interactions, the researcher held weekly meetings with participants in a hospital conference room. The researcher reviewed the status of their learning and helped participants understand the importance of the course. The participants could freely share their feedback with each other. The training lasted for three weeks (21 ± 1 day) under the supervision of head nurses in participating

hospitals who had been instructed on the training plan prior to the commencement of the study. Additional details about the course segments of each module are shown in Table 2.

TABLE 2 Intervention programme modules

Module	Video segment
Module 1: Life & Death	1st week
I Life	1 Origin of life 15'14"
	2 Where does the human being come from 12'54"
	3 Thinking about an individual life 9'35"
	4 Living in the moment 9'9"
I Death	5 Diversity of lives 3'34"
	6 Meaning of life 12'8"
	7 Nature of death 10'42"
	8 Death is someone else's business 8'2"
	9 Being towards death 11'19"
I Death with Dignity	10 Fear of death 11'5"
	11 Surviving beyond death 12'33"
	12 Meaning of death 11'57"
	2nd week
I Dying	13 What is dignity 9'57"
	14 Could death be achieved with dignity 10'30"
	15 Understanding of death with dignity 9'8"
	16 Necessity of death with dignity 8'29"
	17 Influencing factors of death with dignity 8'45"
I Death Education	18 Enlightenment of death with dignity 3'29"
	19 What is dying 14'11"
	20 What is dying experience 8'33"
	21 Kubler Roth's view of dying experience 12'10"
	22 Moody's research on dying experience 7'17"
	23 Research status on dying experience 9'17"
	24 Cases of dying experience 18'27"
	25 Non-soul theory about dying experience 3'46"
	26 Death culture and education of life and death 30 '24"

	27 What is death education 8'29"
	28 Aim of death education 9'33"
	29 Theme of death education 9'16"
	30 Development process of death education 5'22"
	31 Practice of death education 6'22"
Module 2: Palliative Care Nursing	3rd week
	32 Concept of palliative care 8'32"
	33 Philosophy of palliative care 6'29"
	34 Development process of palliative care 7'59"
	35 Target population of palliative care 10'14"
	36 Importance of palliative care 6'50"
	37 Theoretical basis of palliative care 7'40"
	38 Pain relief and palliative care 7'52"
	39 Meaning of palliative care 4'44"
Module 3: Counselling for Bereavement & Grief	40 What is bereavement and grief counselling 10'30"
	41 Necessity of counselling for bereavement and grief 5'29"
	42 Aim of counselling for bereavement and grief 6'45"
	43 Practice of counselling for bereavement and grief 6'34"
	44 Principle of counselling for bereavement and grief 10'38"
	45 Contents and methods of counselling for bereavement and grief 9'19"

Statistical analysis

SPSS 25.0 (SPSS, Inc., Chicago, IL, USA) was used for statistical analyses in this study. For all statistical procedures, $p < .05$ was deemed significant. Categorical variables were expressed as numbers and percentages. The effect of the educational programme was determined by comparing the PCQN, FATCOD-B, and DAP-R post-intervention scores with the pre-intervention scores. Differences were calculated and analysed with paired-samples *t*-tests.

Results

Initially, 125 candidates were eligible and 16 nurses were excluded for their personal further education. Then 109 candidates representing five hospitals attended the educational intervention programme and took the survey promptly after completing their training. Ninety-seven participants gave an adequate number of responses for the analyses to be conducted (89%). Participant flow is shown in Fig. 1.

Participants' characteristics

Participants' characteristics are shown in Table 2. Most of the participants were women, aged 18 to 30 years. Most participants had five or fewer years of experience. Approximately two-thirds of the participants had not previously received any palliative care training. Almost all reported that they required palliative care training.

Main outcomes

After participation in the educational programme, significant changes in PCQN and FATCOD-B scores were noted. The total score of PCQN changed significantly from pre-intervention (10.3 ± 1.9) to post-intervention (11.1 ± 2.2 ; $p = .011$). Management of pain and other symptoms in PCQN increased markedly from 7.7 ± 1.7 to 8.4 ± 1.7 ($p = .003$). The FATCOD-B pre-intervention score of 104.0 ± 7.6 increased to a post-intervention score of 106.4 ± 8.9 ($p = .016$). Through a self-matching test, the DAP-R score showed no obvious difference between pre- and post-intervention ($p > .05$; Fig. 2).

Discussion

Main findings

This study identified that the knowledge of palliative care of participants could be improved significantly by an educational intervention, which is also consistent with prior findings.[18, 19] It was found that the nurses' attitudes towards caring for dying patients also significantly improved after the intervention programme. This is also in line with earlier studies that examined educational interventions for palliative care nurses.[18, 29]

However, our study could not demonstrate any significant changes in participants' pre- and post-intervention attitudes towards death. Some studies reported a significant effect of interventions tackling this issue in other countries.[31, 32] The DAP-R subscale mean scores for fear of death and approaching acceptance significantly increased at the end of training in a prior study.[32] The duration of the intervention in our study may not have been sufficient to change nurses' attitudes towards death. Additionally, cultural values, ethical standards, and religion may differ in China compared to other countries, where studies have shown changes in nurses' attitudes towards death after interventions.[33, 34] Talking about death is considered a taboo in China, like several other Asian cultures.[30, 34] In different Asian cities, nurses' attitudes towards death may also be influenced by the social, cultural, and organisational circumstances of practice.[33] We recommend that further studies on palliative care education be conducted in China.

Strengths and limitations of the study

To reduce sample bias, participants were recruited from the oncology departments of five hospitals, and not just from a single institution.[31, 36, 37] This ensured sampling diversity and representativeness, as well as provided sufficient data for our study. All participants were engaged in caring for dying patients as part of their daily routine work.

Our study applied a mobile-based e-learning approach instead of the web-based method used in other studies,[38] which ensured participants learned with more flexibility and portability without time and location restrictions. Moreover, sending instant messages through the forum enabled real-time communication. Weekly face-to-face Q&A interactions were established to help resolve any learning issues, which made communication more direct and efficient. The integration of multiple methods adopted into the educational programme allowed learning to become more united, convenient, and economical.

There were several limitations to this study, including a relatively small sample size. To verify the effects of the intervention programme, a control group needed to be established.[31] It is difficult to establish a control group across multiple hospitals and ensure the baselines of the control group match those of the experimental one. Further, this study only examined the immediate effects of the educational intervention. Ideally, follow-up studies with the same participants could determine its continued effects on nurses' knowledge and attitudes towards palliative care and care delivery.[17] Additionally, the duration of the intervention was relatively short. The effectiveness of intervention in participants' attitudes towards death may require highly integrated design courses, when it comes to the training contents and training models. And the implementation of training to nurses at appropriate intervals during both education and professional life are requisite.

What this study adds

It is worth noting that the strong demand of participants for palliative care training reflected the current deficit of existing professional training in caring for dying patients. Therefore, a curriculum provision of palliative care education in medical colleges and implementation of workplace training are strongly recommended as quickly as possible, in China. Undoubtedly, this initiative has the potential to guide the creation of a more professional educational training system to eliminate the gap between the training requirements and the provision of palliative care education.

This study showed that a combined-method educational strategy was effective and feasible. It considered how to deploy the available resources to organise and choose a suitable training method that was no longer restricted by factors such as time, space, and finances. It also indicated that a pre-survey based on the training needs of participants could ensure that the training courses are effectively designed and delivered to meet the expectations. The educational strategy developed in this study may be worth developing and disseminating in future initiatives.

Moreover, this study recommended that unified and professional-access criteria be established to form an authorised and recognised qualification system in palliative care in China. In other words, the competence of palliative care staff could be specified and professional standards of staff training could be clarified, which would give access to more palliative care staff to attain the required qualification in this domain.

Conclusions

In summary, our findings provided convincing evidence that mobile terminal-based e-learning combined with virtual and face-to-face interaction was an effective approach to educational training. We confirmed that educational training could exert a positive impact on trainee nurses, enhancing their knowledge of palliative care and improving their attitudes towards caring for the dying patients. This study provided scientific and promising evidence for future education and training systems in palliative care.

Abbreviations

PCQN

Palliative Care Quiz of Nursing

FATCOD-B

Frommelt Attitude Toward Care of the Dying Scale Form B

DAP-R

Death Attitude Profile-Revised

Declarations

ETHICS APPROVAL AND CONSENT TO PARTICIPATE

Ethical approval was sought from the Ethics Committee of Guangzhou Medical University (L180920). Participants provided written informed consent.

CONSENT FOR PUBLICATION

Not applicable.

AVAILABILITY OF DATA AND MATERIALS

All data relevant to the study are included in the article. No additional data available.

COMPETING INTERESTS

The authors declare that they have no competing interests.

FUNDING

The authors disclosed receipt of the following financial support for the research, authorship, and publication of this article: The 13th Five-year-planned Project of Philosophy and Social Sciences Development of Guangzhou, Guangdong Province, China [grant no. 2018GZYB133] and the National Social Science Fund of China [grant no. 18BRK034]. The funders played no role in the design of the study, or in the collection, analysis and interpretation of study findings. There is no conflict of interest related to funding to declare.

AUTHORS' CONTRIBUTIONS

YPH, EX, and YZ contributed to the study conception and design, data acquisition, data analysis and interpretation, and drafting. LXZ, MLH, and XYC, contributed to the study implementation, data acquisition and interpretation. All authors have agreed on the final version.

ACKNOWLEDGEMENTS

The authors thank all the hospitals and nurses for participating in this study. We thank Professor Junrong Liu, Guangdong Research Society of Medical Social Work, and Dr. Yunling Wang, Shandong University, for providing us with the training courses used on our platform. We would like to thank Editage (www.editage.com) for English language editing. We also thank Min Zou, Qingxia Meng, and Hailin Zhu for permitting the use of the FATCOD-B, PCQN, and DAP-R, respectively. Finally, we thank Professor Jinxiang Ma and Zhuoyuan Guo, Guangzhou Medical University, for their support with data analysis.

References

- [1] World Health Organization. Global Health Observatory, Geneva: World Health Organization. who.int/gho/database/en/ (2018). Accessed 21 June 2018.
- [2] Bray F, Ferlay J, Soerjomataram I, et al. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin.* 2018; 68(6):394–424. DOI:10.3322/caac.21492.
- [3] Hall S, Petkova H, Tsouros AD, et al. Palliative care for older people: better practices. Copenhagen: World Health Organization; 2011.
- [4] The Hospice and Palliative Nurse Association. Ensuring high quality in palliative care. *J Hosp Palliat Nurs.* 2018;20(6):E10–E12. DOI:10.1097/NJH.0000000000000520.
- [5] Economist Intelligence Unit. The 2015 Quality of Death Index ranking palliative care across the world. <https://perspectives.eiu.com/healthcare/2015-quality-death-index> (2015). Accessed 19 March 2020.
- [6] Lu Y, Gu Y, Yu W. Hospice and palliative care in China: development and challenges. *Asia Pac J Oncol Nur.* 2018;5(1):26–32. DOI:10.4103/apjon.apjon_72_17.

- [7] Sepulveda C, Marlin A, Yoshida T, et al. Palliative care: the World Health Organization's global perspective. *J Pain Symptom Manage*. 2002;24(2):91–96. DOI: 10.1016/s0885-3924(02)00440-2.
- [8] Shen Y, Nilmanat K, Promnoi C. Palliative care nursing competence of Chinese oncology nurses and its related factors. *J Hosp Palliat Nurs*. 2019;21(5):404–411. DOI: 10.1097/NJH.0000000000000581.
- [9] Jiang X, Liao Z, Hao J, et al. Palliative care education in China: insight into one medical university. *J Pain Symptom Manage*. 2011;41(4):796–800. DOI: 10.1016/j.jpainsymman.2010.07.012.
- [10] Ye G, Mao J, Hu J, et al. Palliative care medical student education: a cross-sectional medical school survey in mainland China. *BMJ Support Palliat Care*. 2019;0:1–4. DOI:10.1136/bmjspcare-2019-002044.
- [11] Xiang YR, Ning XH. Recognition of palliative care in Chinese clinicians: how they feel and what they know. *Chin Med Sci J*. 2018;33(4):221–227. DOI:10.24920/003509.
- [12] Ge N, Qu X, Ning XH, et al. Needs of continuing education on hospice and palliative care in China: a questionnaire-based survey. *Zhongguo Yi Xue Ke Xue Yuan Xue Bao*. 2018;40(3):390-394. DOI:10.3881/j.issn.1000-503X.2018.03.016.
- [13] Selman L, Robinson V, Klass L, et al. Improving confidence and competence of healthcare professionals in end-of-life care: an evaluation of the 'Transforming End of Life Care' course at an acute hospital trust. *BMJ Support Palliat Care*. 2016;6(2):231–236. DOI:10.1136/bmjspcare-2015-000879.
- [14] Nguyen LT, Yates P, Osborne Y. Palliative care knowledge, attitudes and perceived self-competence of nurses working in Vietnam. *Int J Palliat Nurs*. 2014;20(9):448–456. DOI:10.12968/ijpn.2014.20.9.448.
- [15] Ayed A, Sayej S, Harazneh L, et al. The nurses' knowledge and attitudes towards the palliative care. *J Educ Pract*. 2015;6(4):91–99.
- [16] Abu-Saad Huijjer H, Abboud S, Dimassi H. Palliative care in Lebanon: knowledge, attitudes and practices. *Int J Palliat Nurs*. 2009;15(7):346–353. DOI:10.12968/ijpn.2009.15.7.43425.
- [17] Reed E, Todd J, Lawton S, et al. A multi-professional educational intervention to improve and sustain respondents' confidence to deliver palliative care: A mixed-methods study. *Palliat Med*. 2018;32(2):571–580. DOI:10.1177/0269216317709973.
- [18] Loftus LA, Thompson E. An evaluation of a palliative care course for generic nurses. *Int J Palliat Nurs*. 2002;8(7):354–360. DOI:10.12968/ijpn.2002.8.7.10677.
- [19] Kim HS, Kim BH, Yu SJ, et al. The effect of an end-of-life nursing education consortium course on nurses' knowledge of hospice and palliative care in Korea. *J Hosp Palliat Nurs*. 2011;13(4):222–229. DOI:10.1097/NJH.0b013e318210fdec.

- [20] Henoeh I, Melin-Johansson C, Bergh I, et al. Undergraduate nursing students' attitudes and preparedness toward caring for dying persons - a longitudinal study. *Nurse Educ Pract*. 2017;26:12–20. DOI:10.1016/j.nepr.2017.06.007.
- [21] Ferrell BR, Dahlin C, Campbell ML, et al. End-of-life Nursing Education Consortium (ELNEC) training program: improving palliative care in critical care. *Crit Care Nurs Q*. 2007;30(3):206–212. DOI:10.1097/01.CNQ.0000278920.37068. e9.
- [22] Petersen CL, Callahan MF, McCarthy DO, et al. An online educational program improves pediatric oncology nurses' knowledge, attitudes, and spiritual care competence. *J Pediatr Oncol Nurs*. 2017;34(2):130–139. DOI:10.1177/1043454216646542.
- [23] Y Lu, JQ Fang, & L Tian, et al. *Advanced Medical Statistics*. 2nd Edition. Singapore: World Scientific Publishing Co. Pte. Ltd; 2015. <https://doi.org/10.1142/9080>
- [24] Ross MM, McDonald B, McGuinness J. The palliative care quiz for nursing (PCQN): the development of an instrument to measure nurses' knowledge of palliative care. *J Adv Nurs*. 1996;23(1):126–137. DOI:10.1111/j.1365-2648.1996.tb03106.x.
- [25] Frommelt KH. The effects of death education on nurses' attitudes toward caring for terminally ill persons and their families. *Am J Hosp Palliat Care*. 1991; 8(5): 37–43. DOI:10.1177/10499091-9100800509.
- [26] Wong PTP, Reker GT, Gesser AG. *The Death Attitude Profile revised: a multidimensional measure of attitudes towards death*. New York: Taylor and Francis; 1994.
- [27] Zou, M. Investigation and Analysis on Status Quo of Grasping Knowledge about Palliative Nursing of Nurses in Shanghai. *Chinese Journal of Practical Nursing*, 2015; 31 (18), 1386-1390.
- [28] Wang LP, Li YJ, Yan WZ, et al. Development and psychometric testing Chinese version of the Frommelt Attitude Toward Care of the Dying Scale, Form B in nurses and nursing Students. *J Cancer Educ*. 2016;31(1):123–130. DOI:10.1007/s13187-015-0810-7.
- [29] Meng ZX, Li YL, Zhi CX. Correlation between nurses' attitudes towards dying patients and their personal attitudes towards death. *J Nurs Sci*. 2014;29:51–53.
- [30] Zhu HL. *DAP-R Chinese edition and application in middle-aged and the older Christian believers*. Tian Jin Medical University. Tianjin: China; 2011.
- [31] Hamayoshi M, Goto S, Matsuoka C, et al. Effects of an advance care planning educational programme intervention on the end-of-life care attitudes of multidisciplinary practitioners at an acute hospital: a pre- and post-study. *Palliat Med*. 2019;33(9):1158–1165. DOI:10.1177/02692163-19860707.

- [32] Goris S, Tasci S, Ozkan B, et al. Effect of terminal patient care training on the nurses' attitudes toward death in an oncology hospital in Turkey. *J Cancer Educ.* 2017;32(1):65–71. DOI:10.1007/s13187-015-0929-6.
- [33] Chan MF, Lou FL, Arthur DG. A survey comparing the attitudes toward perinatal bereavement care of nurses from three Asian cities. *Eval Health Prof.* 2010;33(4):514–33. DOI:10.1177/0163278710381092.
- [34] Gama G, Barbosa F, Vieira M. Factors influencing nurses' attitudes toward death. *Int J Palliat Nurs.* 2012;18(6):267–273. DOI:10.12968/ijpn.2012.18.6.267.
- [35] Lin CP, Cheng SY, Chen PJ. Advance care planning for older people with cancer and its implications in Asia: highlighting the mental capacity and relational autonomy. *Geriatrics (Basel).* 2018;3(3):43. DOI:10.3390/geriatrics3030043.
- [36] Downing J, Batuli M, Kivumbi G, et al. A palliative care link nurse programme in Mulago Hospital, Uganda: an evaluation using mixed methods. *BMC Palliat Care.* 2016;15:40. DOI:10.1186/s12904-016-0115-6.
- [37] Machira G, Kariuki H, Martindale L. Impact of an educational pain management programme on nurses pain knowledge and attitudes in Kenya. *Int J Palliat Nurs.* 2013;19(7):341-346. DOI:10.12968/ijpn.2013.19.7.341.
- [38] Sinclair PM, Kable A, Levett-Jones T, et al. The effectiveness of Internet-based e-learning on clinician behaviour and patient outcomes: a systematic review. *Int J Nurs Stud.* 2016;57:70–81. DOI:10.1016/j.ijnurstu.2016.01.011.

Notes: References 27, 29, and 30 addressed the Chinese versions of the scales used in this study: PCQN, FATCOD-B, DAP-R.

Figures




Figure 1

Participants' flow (N = 97)




Figure 2

PCQN, FATCOD-B, and DAP-R scores pre- and post-intervention (N = 97) A, Total PCQN scores ranged from 0 to 20, with higher scores indicating greater knowledge. Scores in philosophy and principles of palliative care ranged from 0 to 4, scores in management of pain and other symptoms ranged between 0 and 13, and scores in psychosocial aspects of care ranged from 0 to 3. PCQN: Palliative Care Quiz of Nursing. B, Scores on the scale ranged from 0 to 150 with higher scores indicating a favourable attitude

towards care for the dying. FATCOD-B: Frommelt Attitudes Towards Care of the Dying Scale Form B. C, Scores on each subscale, ranging from 1 to 7, represented the five attitudes towards death. A higher score indicated a greater tendency to share the corresponding attitude towards death. DAP-R: Death Attitude Profile-Revised. Data were presented as estimated means and deviations. The height of the error bar in Figure 2 is equal to the mean value of the variable plus the standard deviation. Paired t-tests were used for analysis of the difference between pre- and post-intervention scores. * $p \leq .05$ for differences between pre- and post-intervention.