Quantitative assessment of German Holstein dairy cattle colostrum and impact of thermal treatment on quality of colostrum viscosity and immunoglobulins

Abdulwahed Ahmed Hassan (✉ abdulwahed-ahmed.hassan@vetmed.uni-giessen.de)

Justus-Liebig-Universität Giessen Fachbereich Veterinärmedizin <https://orcid.org/0000-0001-9088-2905>

Sebastian Ganz

Justus Liebig Universität Giessen

Florian Schneider

Justus Liebig Universität Giessen

Axel Wehrend

Justus Liebig Universität Giessen

Izhar U.H. Khan

Agriculture and Agri-Food Canada

Klaus Failing

Justus Liebig Universität Giessen

Michael Bülte

Justus Liebig Universität Giessen

Amir Abdulmawjood

Tierärztliche Hochschule Hannover

Research note

Keywords: Holstein dairy cattle colostrum; color; fat; dynamic viscosity; colostrometer; digital %Brix refractometer; thermal treatment

Posted Date: February 17th, 2020

DOI: <https://doi.org/10.21203/rs.2.23694/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at BMC Research Notes on March 30th, 2020. See the published version at <https://doi.org/10.1186/s13104-020-05019-z>.

Abstract

Objective: This study aimed to determine the color, fat, viscosity, IgG concentration, %Brix and refractive index of fresh postpartum colostrum of German Holstein dairy cattle and assess the impact of different thermal treatments on the visual and dynamic viscosity, in association to IgG concentration, of colostrum that can be used for pasteurization process.

Results: Of the total 40 fresh postpartum colostrum, the color of colostrum (white-pale yellow to yellow and dark-yellowish), fat ($1.4 - 8.2 \text{ } 100 \text{ g}^{-1}$), IgG ($4 - 116 \text{ mg mL}^{-1}$), %Brix (8.5 - 35.4%), refractive index ($1.3454 - 1.3905 \text{ nD}$), visual (ranging from watery to liquid and thick) and dynamic ($4.9 - 219 \text{ cp}$) viscosity, were recorded. Statistical analysis between visual and dynamic viscosity of fresh colostrum showed significant correlation coefficients ($r_s = 634$). Moreover, a significant correlation between viscosity and three IgG concentrations was observed. Heat-treated colostrum showed dynamic viscosity ranged from 25 to 3066 cP, where dynamic viscosity of colostrum before- and after heat-treatment showed no significant correlation. Treated colostrum containing IgG concentration $\leq 80 \text{ mg mL}^{-1}$ at $60^\circ\text{C}/60 \text{ min}$ and $\leq 68 \text{ mg mL}^{-1}$ at $63.5^\circ\text{C}/30 \text{ min}$ showed no significant change in the viscosity and can successfully be applied for pasteurization of first postpartum colostrum.

Introduction

Colostrum delivers a vital source of immunoglobulins (Igs), non-specific immune factors and other essential nutrients for passive transfer of immunity to newborn calves (1–3). After parturition, newborn calves need to ingest 3–4 liters of high-quality colostrum containing $\geq 50 \text{ mg mL}^{-1}$ of IgG within the first 6 h after birth to protect them from pathogens until their own immune system is developed (1, 4). Therefore, estimating the quality of first-milking colostrum on farms by measuring the specific gravity or IgG concentration prior to feeding calves is a useful management tool to improve calf health (5–7). The present study was designed to evaluate color, fat, and IgG (mg mL^{-1} , %Brix and refractive index) as well as visual and dynamic viscosity of first postpartum colostrum of German Holstein dairy cattle. For establishing an optimal heat process that can minimally impact on viscosity and immunoglobulins, different thermal treatments (temperature and time) were investigated in this study.

Methodology

Physico-chemical analysis

Initially, 16 German Holstein dairy cattle farms, located in Hessen, Germany, were selected for collection of 40 colostrum samples immediately after parturition, aged between three and seven years. All samples, collected between March 2017 to Oct. 2018, were visually analyzed for color and viscosity. Dynamic viscosity was measured for native colostrum program (1 rpm at $30^\circ\text{C}/60 \text{ sec.}$) using a viscometer DV3T

touchscreen rheometer (AMETEK Brookfield, Germany), and two programs (1.0 and 0.1 rpm at 30 °C/60 sec) used for samples treated at 63.5 °C/30 min. The fat content was determined using the Gerber method (8).

Immunoglobulins

Colostrum IgG (mg mL⁻¹) was measured by a colostrometer (9, 10), with an optimum 37 °C temperature (Pfizer Animal Health GmbH, Germany). A digital handheld refractometer DR201-95 (A. Krüss Optronic, Germany) was used to determine the %Brix and refractive index (nD) (11). The Brix refractometer scale of ≥ 20% (50 mg mL⁻¹) IgG corresponds to a good as compared to ≤ 20% Brix (≤ 50 mg mL⁻¹) IgG a low quality colostrum.

Thermal treatments

Three thermal treatments (60 °C/60 min, 63.5 °C/30 min and 72.0 °C/15 sec) were performed in a water bath to determine the cut-point temperature that may impact on the viscosity in relation to the IgG concentration (12, 13). The water bath was calibrated and adjusted to ± 0.5 °C with an additional digital thermometer (Testo 112 Type NTC, Germany) and thermometer type UT330A (Reichelt Elektronik, Germany).

Statistical analysis

The statistical data analysis was performed using BMDP v8.1.0 and StatXact v9.0.0 statistical software packages (14, 15). The values are described as arithmetic mean, standard error of mean (± SEM) and standard deviation (mean ± SD), P < 0.05 deemed as significant. For analyzing the relationship between fat content, visual and dynamic viscosity and IgG (mg mL⁻¹, %Brix and nD) concentration, Pearson correlation coefficient (r) and rank correlation coefficient analyses were performed for ordinal-scaled data in accordance with Spearman's rank-order correlation coefficient (r_s) and logarithmic methods.

Results

Color and fat content

The color of 37 samples ranged from white-pale yellow to yellow and dark-yellowish. The other three samples collected from healthy dairy cattle showed a slightly pinkish due to hemorrhage per diapedesis. The fat concentration of colostrum samples ranged from 1.4 to 8.8 100 g⁻¹ (mean 5.4 100 g⁻¹) (Table 1). There was no significant correlation observed between the fat and color gradation as well as between fat and IgG (mg mL⁻¹, %Brix and nD) concentrations (Table 2).

Table 1:

Analysis of fat and IgG (concentration, %Brix, refractive index (nD)), and dynamic viscosity of fresh and heat-treated dairy cattle colostrum samples.

Parameter	Mean (average)	$\pm$ SD	$\pm$ SEM*	Minimum	Maximum
Fat (100 g ⁻¹)	5.41	2.03	0.32	1.4	8.8
IgG (mg mL ⁻¹)	57.65	32.71	5.17	4.0	116.0
%Brix	20.32	6.12	0.96	8.5	35.4
nD	1.3640	0.010	0.001	1.3454	1.3905
Dynamic viscosity (cP)†	34.55	41.27	6.52	4.9	219.0
Dynamic viscosity (cP)††	658.87	911.09	145.89	5.2	3.066

* SEM: standard error of mean; †: first postpartum colostrum; ††: heat-treated colostrum (63.5°C for 30 min)

Table 2

Parameter	Mean (average)	$\pm$ SD	$\pm$ SEM*	Minimum	Maximum
Fat (100 g ⁻¹)	5.41	2.03	0.32	1.4	8.8
IgG (mg mL ⁻¹)	57.65	32.71	5.17	4.0	116.0
%Brix	20.32	6.12	0.96	8.5	35.4
nD	1.3640	0.010	0.001	1.3454	1.3905
Dynamic viscosity (cP)†	34.55	41.27	6.52	4.9	219.0
Dynamic viscosity (cP)††	658.87	911.09	145.89	5.2	3.066
* SEM: standard error of mean; †: first postpartum colostrum; ††: heat-treated colostrum (63.5 °C for 30 min)					

Immunoglobulins

The IgG (mg mL⁻¹) concentration of fresh colostrum samples using a colostrometer ranged from 4.0 to 116 mg mL⁻¹. Based on the cut-point (≤ 50 mg mL⁻¹) concentration, 15 (37.5%) and 25 (62.5%) samples were classified as low and high quality colostrum, respectively. Similarly, the %Brix of IgG ranged from 8.5 to 35.4%Brix, where based on cut-point ($\leq 20\%$ Brix) value, 17 (42.5%) and 23 (57.5%) samples were classified as low and high quality colostrum, respectively. Furthermore, the refractive index of IgG values ranged from 1.3454 to 1.3905 nD with a similar percentage (42.5 and 57.5%) of samples, based on the cut-point (≤ 1.3596) value, were classified as low and high quality colostrum (Table 1). The relationship congruency between IgG mg mL⁻¹ and %Brix values revealed that 38 (95%) samples were in agreement with the colostrometer and refractometer methods. The statistical relationship revealed significant coefficient correlation between IgG mg mL⁻¹ and %Brix ($r = 0.894$), IgG and nD ($r = 0.887$) and %Brix and nD ($r = 0.991$) (Table 2), while linear regression correlations are illustrated in Fig. 1 (Panels A-C).

Visual and dynamic viscosity

The visual viscosity of first fresh colostrum was categorized as watery ($n = 14$; 35%), liquidy ($n = 17$; 42.5%) and thick ($n = 9$; 22.5%) (Table S1). The 17 samples containing high-fat ($7.0\text{--}8.8\text{ }100\text{ g}^{-1}$) were classified into: watery ($n = 9$; 53%), liquidy ($n = 6$; 35%) and thick ($n = 2$; 12%). Although the statistical analyses revealed no significant correlation between fat and visual viscosity, a significant correlation between visual viscosity and IgG (mg mL^{-1} , %Brix and nD) concentration was recorded (Table 2; Fig. S1: Panels A-C).

The dynamic viscosity of first colostrum ranged from 4.9 to 219 centipoise (cP) (Table 1), where similar to visual viscosity, significant correlation was found between dynamic viscosity and IgG (mg mL^{-1} , %Brix and IgG nD) concentration but not with the fat content (Table 2; Fig. S1: Panels D-F). Statistical analysis results, using Spearman's rank-order correlation method, showed significant correlation coefficients ($r_s = 0.634$; $p < 0.001$) between visual and dynamic viscosity for all colostrum samples (Table 2; Fig. S1: Panel G).

Analysis of thermally treated colostrum

After treating colostrum at $60\text{ }^{\circ}\text{C}/60\text{ min}$, all 14 watery and 15 liquid samples, the viscosity was changed to liquid, while the two liquid and nine thick samples showed slight to moderate coagulation, where $\leq 80\text{ mg mL}^{-1}$ pasteurization cut-point IgG concentration was recorded. The viscosity of watery colostrum samples treated at $63.5\text{ }^{\circ}\text{C}/30\text{ min}$ changed to liquid, while the liquid and thick samples formed slight to moderate coagulation with $\leq 68\text{ mg mL}^{-1}$ cut-point IgG concentration. However, 17 samples treated at $72.0\text{ }^{\circ}\text{C}/15\text{ sec}$ showed moderate coagulation (cut point: $\leq 45\text{ mg mL}^{-1}$) as compared to the remaining 23 samples that changed into solid due to excessive coagulation (Table S1).

After heat-treatment of all colostrum samples at $63.5\text{ }^{\circ}\text{C}/30\text{ min}$, only 28 samples, measured at program 1.0 rpm at $30\text{ }^{\circ}\text{C}/60\text{ sec}$, showed dynamic viscosity ranging from 25 to 310 cP. The remaining 12 samples, tested at program 0.1 rpm at $30\text{ }^{\circ}\text{C}/60\text{ sec}$, showed dynamic viscosity ranging from 1.407 to 3066 cP (Table 1). There was no significant correlation observed between dynamic viscosity of fresh and heat-treated ($63.5\text{ }^{\circ}\text{C}/30\text{ min}$) samples (Table 2).

Discussion

To our knowledge, this is the first detailed study in Germany describing vital physico-chemical properties of first fresh postpartum colostrum samples collected from healthy German Holstein dairy cattle. Furthermore, this study evaluated the relationship between the impact of heat-treatment on the colostrum viscosity and threshold of IgG concentrations. Generally, physical properties of colostrum such as color gradation and visual viscosity provides an initial impression of the quality status of colostrum. In this study, color was assessed visually that ranged from white-pale yellow to yellow and dark-yellow, gradation of color and degree of viscosity (watery, liquid and thick) showed a significant ($p \leq 0.05$)

relationship. Gross et al. (16) reported that colostrum has a wide range of color spectra ranging from pale-white to dark-brown/red compared to the dairy cow milk color. The color gradation of colostrum increased progressively from pale to dark with more fat, protein and IgG as well as other factors including dietary composition also considered as contributory factors (16–18). However, a correlation between color and viscosity was not previously reported. Furthermore, the fat content varied ($1.4\text{--}8.8\ 100\ \text{g}^{-1}$; mean $5.4\ 100\ \text{g}^{-1}$) in contrast to Quigley et al. (19) where a higher fat content ($9.2\text{--}31.6\ 100\ \text{g}^{-1}$; mean: $23.6\ 100\ \text{g}^{-1}$) was reported. On the other hand, Kehoe et al. (20) measured the average colostrum fat $6.7\ 100\ \text{g}^{-1}$ as compared to $3.6\ 100\ \text{g}^{-1}$ (21). In contrary to the previous study (16), the present study results showed no significant relationship between color gradation and fat as well as between fat and various degrees of viscosity. This weak correlation with these parameters can possibly be due to the color variation as compared to the viscosity and fat concentration.

For assessing IgG concentration, previous studies have recommended on-farm tools to provide precise and reproducible results (6, 11, 19, 22–26). Based on these studies, we used colostrometer and refractometer as alternative indirect, rapid and accurate tools to assess the quality of German Holstein dairy cattle colostrum by estimating the IgG concentration. Both methods showed a high degree ($\geq 95\%$) of similarity in the classification of colostrum IgG concentration where strong correlation coefficients (0.894 and 0.887) with %Brix and nD were observed. Interestingly, these results are higher than previously reported results (23, 25) where the correlation coefficients of IgG and %Brix were relatively low (0.71 and 0.75). However, our results were similar to Morrill et al. (26) with a correlation coefficient of 0.86. The colostrum quality, in the present study, revealed disparate individual IgG concentrations leading to a high variation ($4\text{ to }116\ \text{mg mL}^{-1}$; mean $58\ \text{mg mL}^{-1}$) but lower ($69.9\ \text{mg mL}^{-1}$) than the previously reported IgG concentration (19). The IgG concentration data obtained from colostrometer (mg mL^{-1}) and refractometer (%Brix and nD), in our study, revealed that 37.5%, 42.5% and 42.5% samples did not show congruence to the recommended IgG cut-points ($\geq 50\ \text{mg mL}^{-1}$; $\geq 20\%$ Brix and $\geq 1.3596\ \text{nD}$). On the other hand, $72.9\ \text{mg mL}^{-1}$ IgG mean concentration was determined in Jersey dairy cattle colostrum ranging from $12.8\text{ to }154.3\ \text{mg mL}^{-1}$, and 32.8% of samples had $< 50\ \text{mg mL}^{-1}$ IgG concentration with a mean (21.24%) %Brix of fresh colostrum, %Brix values ranging from 10.5 to 28.6% with recommended breed-specific $\geq 18\%$ cut-point (26). In congruence with our %Brix results, conventional (62.5%) and organic (56.1%) Danish dairy cattle colostrum samples had equal or exceeded 22%Brix cut-point, with significant variation ranging from 8.3 to 35.1% (7). Our cut-point level of good quality colostrum was determined at $\geq 20\%$ Brix corresponded to $\geq 50\ \text{mg mL}^{-1}$ IgG concentration using colostrometer. Chigerwe et al. (11) and Biemann et al. (25) suggested 22%Brix as an optimal cut-point level as compared to the recommended levels (18%, 21% and 23%) for Jersey dairy cattle colostrum (6, 13, 23–27).

Heat-treatment of colostrum, either in experimental conditions or commercial batch pasteurization system, was previously investigated to determine the efficiency of pasteurization on viability of microorganisms to reduce calf exposure to bacterial pathogens, change in viscosity (degree of coagulation) and degradation of IgG (5, 12, 27). The present study indicates that pasteurizing colostrum

at 60 °C/60 min containing $\leq 80 \text{ mg mL}^{-1}$ IgG concentration have a minimal impact on the viscosity, whilst pasteurizing colostrum at 63.5 °C/30 min containing $\leq 68 \text{ mg mL}^{-1}$ IgG concentration have a moderate impact on the viscosity. Our study results correspond to the previous study where pasteurizing colostrum at 63.5 °C/30 min using a commercial batch pasteurizer produced a mildly thick coagulation viscosity compared to 72 °C/15 sec where heat-treatment caused a solid form of colostrum especially in samples containing IgG concentration $> 50 \text{ mg mL}^{-1}$ (5). Similar to our findings, colostrum treated at various temperatures (57, 60 and 63 °C) and time (30, 60 and 90 min) did not affect viscosity; however, the cut-point of pasteurization, in our study, at different heat-treatment maintained the viscosity, IgG concentration and high quality of colostrum (12).

Conclusion

Analysis of quantitative IgG concentration of first colostrum using colostrometer or refractometer method is useful prior to initiating on-farm pasteurization step using commercial colostrum pasteurizers. Pasteurization of first colostrum containing IgG ≤ 80 and $\leq 68 \text{ mg mL}^{-1}$ at 60 °C/60 min and 63.5 °C/30 min did not significantly impact on visual and dynamic viscosity and quality of colostrum. Therefore, these thermal treatments can successfully be applied commercially for pasteurizing colostrum. Moreover, the treatments, subsequently, allow to adjust the volume of colostrum, without altering properties, for feeding and successful transfer of passive immunity to the calves.

Limitation

In this study, IgG-ELISA test on colostrum samples before and after different heat-treatments and their correlation were not performed due to limited sources of funding.

Abbreviations

Igs:immunoglobulins; IgG:immunoglobulin G; nD:refractive index; cP:centipoise; (r):Pearson correlation coefficient; (r_s):Spearman's rank-order correlation; LHL:Landesbetrieb Hessisches Landeslabor.

Declarations

- **Ethics approval and consent to participate**

A written approval and consent from the Klinik für Geburtshilfe, Gynäkologie und Andrologie der Groß- und Kleintiere mit Tierärztlicher Ambulanz, Justus-Liebig-University Gießen was obtained to use the animals in this study. This study was approved by the animal ethic committee Regierungspräsidium Gießen (V54-19 c2015 h01). Also, the same institute received the consent from farm owners to collect samples from dairy cattle for this project.

- **Funding**

This study is a part of the project: “Validierung und Etablierung eines neuartigen Pasteurisierungsverfahrens für bovine Kolostralmilch zur Reduzierung von *Mycobacterium avium* subsp. *paratuberculosis* (MAP) – Minimierungsstrategie für Milchviehbestände” This study was funded, under Project #: 819 589, by the Federal Ministry of Food, Agriculture and Consumer Protection (Bundesministerium für Ernährung, Landwirtschaft und Verbraucherschutz) via the special federal funds (Zweckvermögen des Bundes) at the Landwirtschaftlichen Rentenbank (LR).

- **Availability of data and material**

The dataset used in the study can be provided by the corresponding author upon request

- **Consent for publication**

Not applicable.

- **Competing Interest**

The authors declare that they have no competing interests to report

- **Authors contributions**

MB, AW, AAH and SG, proposed and designed the experiment. AAH, SG and FS collected and analyzed samples. AAH, MB, AW, IK and AA processed and analyzed data. KF performed statistical data analysis. AW, MB, IK and AA drafted and edited the manuscript. All authors reviewed and approved the final manuscript.

- **Acknowledgments**

We would like to thank farm owners for their participation and untiring help and support to conduct this study. Also, we would like to thank Dr. Manuela Fischer-Zorn from (LHL), Gießen, Germany for her assistance with laboratory analysis.

References

1. Larson BK, Heary HK, Devery JE. Immunoglobulin production and transport by the mammary gland. J Dairy Sci. 1980; 63:665–71.
2. McGrath BA, Patrick FF, McSweeney PLH, Kelly AL. Composition and properties of bovine colostrum: a review. Dairy Sci. & Technol. 2016;96:133–
3. Ganz S, Bülte M, Gajewski Z, Wehrend A. Inhaltsstoffe des bovinen Kolostrums – Eine Übersicht. Tierarztl Prax Ausg G. 2018;46:178–89.
4. Fischer AJ, Song Y, He Z, Haines DM, Guan LL, Steele MA. Effect of delaying colostrum feeding on passive transfer and intestinal bacterial colonization in neonatal male Holstein calves. 2018;101:3099-109.

5. Godden SM, Smith S, Feirtag JM, Green LR, [Wells SJ](#), [Fetrow JP](#). Effect of on-farm commercial batch pasteurization of colostrum on colostrum and serum immunoglobulin concentrations in dairy calves. *J Dairy Sci.* 2003;86:1503–12.
6. Bartier AL, Windeyer MC, Doepel L. [Evaluation of on-farm tools for colostrum quality measurement](#). *J Dairy Sci.* 2015;98:1878–
7. Mejer, T. Bovine colostrum and factors impacting colostrum quality in conventional and organic dairy herds. Master thesis. Aarhus University. Denmark. 2015. pp. 1–
8. ISO 488:2008. Milk - Determination of fat content - Gerber butyrometers.
9. Fleenor, WA, Stott GH. Hydrometer test for estimation of immunoglobulin concentration in bovine colostrum. *J Dairy Sci.* 1980;63:973–77.
10. Pritchelt LC, Gay CC, Hancock DD, Besser TE. Evaluation of the hydrometer for testing immunoglobulin G1 concentrations in Holstein colostrum. *J Dairy Sci.* 1994;77:1761–67.
11. Chigerwe M, Tyler JW, Schultz LG, Middleton JR, Spain JN, Dill JS, Steevens BJ. Comparison of four methods to assess colostral IgG concentration in dairy cows. *J Am Vet Med Assoc.* 2008;233:761–66.
12. Elizondo-Salazar J, Jayarao B, Heinrichs A. Effect of heat treatment of bovine colostrum on bacterial counts, viscosity, and immunoglobulin G concentration. *J Dairy Sci.* 2010;93: 961–67.
13. Elsohaby I, McClure JT, Dow N, Keefe GP. Effect of heat-treatment on accuracy of infrared spectroscopy and digital and optical Brix refractometers for measuring immunoglobulin G concentration in bovine colostrum. *J Vet Intern Med.* 2018;32:491–96.
14. Dixon WJ (chief editor). *BMDP Statistical Software Manual*. University of California Press, Berkeley, Los Angeles, London. Volume 1 and 2. 1993.
15. Cytel Studio StatXact Vers. 9.0.0. Statistical Software for Exact Nonparametric Inference, User Manual. 2010. CYTEL Inc., Cambridge, MA 02139, U.S.A.
16. [Gross JJ](#), [Kessler EC](#), [Bruckmaier RM](#). Colour measurement of colostrum for estimation of colostral IgG and colostrum composition in dairy cows. *J Dairy Res.* 2014;81:440-4
17. Madsen BD, Rasmussen MD, Nielsen MO, Wiking L, Larsen LB. Physical properties of mammary secretions in relation to chemical changes during transition from colostrum to milk. *J Dairy Res.* 2004;71:263–72.
18. Solah VA, Staines V, Honda S, Limley HA. Measurement of milk color and composition: effect of dietary intervention on western Australian Holstein-Friesian cow's milk quality. *J Food Sci.* 2007;72 S560–S566.
19. Quigley, JD III, Martin KR, Dowlen HH, Wallis LB, Lamar K. Immunoglobulin concentration, specific gravity, and nitrogen fractions of colostrum from Jersey cattle. *J Dairy Sci.* 1994;77:264–69.
20. Kehoe SI, Jayarao BM, Heinrichs AJ. A survey of bovine colostrum composition and colostrum management practices on Pennsylvania dairy farms. *J Dairy Sci.* 2007;90:4108–16.

21. Tsioulpas A, Grandison AD, Lewis MJ. Changes in physical properties of bovine milk from the colostrum period to early lactation. *J Dairy Sci.* 2007;90:5012–17.
22. Elsohaby I, McClure JT, Cameron M, Heider LC, Keefe G P. Rapid assessment of bovine colostrum quality: How reliable are transmission infrared spectroscopy and digital and optical refractometers?. *J Dairy Sci.* 2017;100:427–35.
23. Løkke MM, Engelbrecht R, Wiking L. Covariance structures of fat and protein influence the estimation of IgG in bovine colostrum. *J Dairy Res.* 2016;83:58–
24. Biemann V, Gillan J, Perkins NR, Skidmore AL, Godden S, Leslie KE. An evaluation of Brix refractometry instruments for measurement of colostrum quality in dairy cattle. *J. Dairy Sci.* 2010;93:3713–
25. Quigley JD, Lago A, Chapman C, Erickson P, Polo J. Evaluation of the Brix refractometer to estimate immunoglobulin G concentration in bovine colostrum. *J Dairy Sci.* 2013;96:1148–55.
26. Morrill KM, Robertson KE, Spring MM, Robinson AL, Tyler HD. Validating a refractometer to evaluate immunoglobulin G concentration in Jersey colostrum and the effect of multiple freeze-thaw cycles on evaluating colostrum quality. *J Dairy Sci.* 2015;98:595–
27. Green L, Godden S, Feirtag J. Effect of batch and high temperature-short time pasteurization on immunoglobulin G concentrations in colostrum. *J Dairy Sci.* 2003;86:246.

Supplementary Files Legend

Supplementary Figure S1:

Statistical correlation analysis of dairy cattle colostrum samples, Panels A, B and C: Visual viscosity with IgG (mg mL^{-1} , %Brix and nD) concentrations. Panels D, E and F: Dynamic viscosity with IgG (mg mL^{-1} , % Brix and nD) concentrations. Panel G: statistical correlation analysis between visual and dynamic viscosity.

Supplementary Table S1:

Categorization of IgG concentrations (mg mL^{-1}) and visual viscosity of fresh first postpartum and heat-treated colostrum samples.

Figures

Figure 1:

Statistical correlation analysis of fresh dairy cattle colostrum samples between IgG mg mL⁻¹ and %Brix (Panel A); IgG mg mL⁻¹ and nD (Panel B); and %Brix and nD (Panel C).

1

Figure 1

Statistical correlation analysis of fresh dairy cattle colostrum samples between IgG mg mL⁻¹ and %Brix (Panel A); IgG mg mL⁻¹ and nD (Panel B); and %Brix and nD (Panel C).

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [TableS1.docx](#)
- [Fig.S1.docx](#)