

Chronic hypoxia leads to cognitive impairment by promoting HIF-2 alpha -mediated alpha-synuclein hyperphosphorylation

Gaifen Li

Capital Medical University

Jia Liu

Capital Medical University <https://orcid.org/0000-0001-6711-3841>

Mengyuan Guo

Capital Medical University

Yakun Gu

Capital Medical University

Yuying Guan

Capital Medical University

Qianqian Shao

Capital Medical University

Wei Ma

Capital Medical University

Xunming Ji (✉ jixm@ccmu.edu.cn)

<https://orcid.org/0000-0002-0527-2852>

Research article

Keywords: chronic hypoxia, alpha-synuclein, cognitive impairment, hypoxia-inducible factor, ceramide

Posted Date: March 25th, 2022

DOI: <https://doi.org/10.21203/rs.3.rs-1427101/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background: Cognitive impairment is involved in various central nervous system diseases. Hypoxia is a critical common factor that induces cognitive impairment, but its specific molecular mechanism remains unclear. Understanding the mechanism of cognitive impairment induced by hypoxia will provide potential molecular targets for the intervention of cognitive disorders.

Methods: We used hypoxic chambers to establish a chronic hypoxic model in mice. Cognitive impairment, hippocampal neurodegeneration, and alpha-synuclein (α -syn) pathology were assessed in these mice. The α -syn knockout mice or mice administered a competing peptide targeting serine 129 (ser129) were used to investigate the role and molecular mechanism of α -syn pathology in chronic hypoxia-induced cognitive impairment. Finally, we investigated the protective role of intermittent hypoxia preconditioning in chronic hypoxia-induced α -syn pathology and cognitive impairment.

Results: Chronic hypoxia resulted in α -syn hyperphosphorylation at ser129 and protein aggregation, hippocampal neurodegeneration, and irreversible cognitive deficits. The latter could be reversed by α -syn knockdown or an administered short peptide competing at α -syn ser129. These results suggest that α -syn hyperphosphorylation at ser129 mediates hippocampal degeneration and cognitive impairment following chronic hypoxia. Furthermore, we found that chronic hypoxia enhanced ceramide catabolism by inducing hypoxia-inducible factor (HIF)-2 α and HIF-2 α -dependent transcriptional activation of alkaline ceramidase 2 (Acer2). Thus, the enzymatic activity of protein phosphatase 2A (PP2A), a specific phosphatase for α -syn, is inhibited, leading to sustained induction of α -syn hyperphosphorylation and aggregation. Finally, we found that enhancing hypoxia tolerance through intermittent hypoxic preconditioning protected against subsequent chronic hypoxia-induced hippocampal neurodegeneration and cognitive impairment by preventing α -syn pathology.

Conclusion: These results proved the critical role of α -syn pathology in hypoxia-afforded cognitive impairment and revealed a novel mechanism underlying hypoxic-specific induction of cognitive impairment. The findings bear implications in developing novel therapeutic interventions for hypoxia-related cognitive disorders.

Background

Cognitive impairment is a common feature of many central nervous system diseases, such as Alzheimer's disease [1], Parkinson's disease (PD) [2], dementia of Lewy body (DLB) [3], and vascular dementia [4]. Searching for common factors and molecular mechanisms is a potential strategy for intervention. Hypoxia is an important stressor that induces cell damage, especially in the central nervous system (CNS), which is highly susceptible [5]. Neurons are the most abundant cell population in the CNS and are highly susceptible to hypoxia. Their damage induces the occurrence and development of various CNS diseases [6]. Chronic hypoxia, including environmental hypoxia and tissue hypoxia, inducing neuronal injury is common [7, 8]. For example, environmental hypoxia caused by altitude exposure can

cause irreversible impairment in cognitive ability and memory ability [9]. With aging, the brain's ability to use oxygen gradually decreases, which leads to the development of a range of neurological diseases, such as neurodegenerative diseases [10]. However, the molecular mechanism of neuronal injury induced by hypoxia remains unclear, limiting the development of interventions for many neurological diseases.

α -Synuclein (α -syn) is a protein that is highly expressed in neurons. Although it has a powerful physiological function, more studies have focused on its pathological role, mainly because it is involved in the pathogenesis of several neurodegenerative diseases, such as Parkinson's disease (PD) and dementia of Lewy body (DLB) [11, 12]. Abnormal changes in posttranslational modifications, especially phosphorylation at serine 129 (ser129) (p- α -syn), are critical for the transition of α -syn from a physiological to pathological function [13]. The proportion of p- α -syn in the normal state is very low, and an abnormal increase in p- α -syn induces the formation of pathological aggregates that contribute to neurotoxicity [14]. Previous studies have shown that hypoxia could promote α -syn phosphorylation and pathological aggregation [15], but its molecular mechanism is still unclear. Further exploration of this mechanism will be of great significance for understanding or intervening in hypoxia-related diseases and α -syn-related diseases.

In this study, we found that chronic hypoxia induced irreversible cognitive impairment in mice by promoting α -syn phosphorylation at ser129 and abnormal aggregate formation in the hippocampus. Both α -syn knockout mice and mice administered a short peptide competing at α -syn ser129 were able to resist chronic hypoxia-induced cognitive impairment. Such results suggested that α -syn phosphorylation played an important role in chronic hypoxia-induced cognitive impairment. Our further study showed that chronic hypoxia-induced α -syn pathology promoted the hydrolytic metabolism of ceramide and decreased protein phosphatase 2A activity by activating the hypoxia-inducible factor (HIF)-2 α /alkaline ceramidase (Acer)2 signaling pathway. In conclusion, our study has revealed a novel mechanism of cognitive deficits induced by chronic hypoxia, highlighted the role of α -syn pathology in this process, and provided a potential new target for the intervention of hypoxia-related cognitive disorders.

Methods

1. Animals

Adult male C57BL mice were purchased from SPF Biotechnology (Beijing, China). α -Syn knockout mice were provided by Professor Yang Hui of Capital Medical University. All animals were housed at room temperature under a 12/12 h light/dark cycle and had free access to food and water. All animal experiments were approved by the Animal Care and Use Committee of the Institute of Animal Management, Capital Medical University, and conducted in accordance with ethical requirements.

2. Hypoxic treatment

All mice were administered hypoxic treatment in a closed hypoxic chamber (China Innovation Instrument Co., Ltd, Ningbo, Zhejiang, China), which accurately set the desired hypoxic concentration and pattern. For

chronic hypoxia, mice were treated continuously with 13% O₂ for 1, 3, 7, and 14 days. The hypoxic chamber was opened briefly for food and water additions every 3 days. Intermittent hypoxic mice were treated with 10 cycles of 5-min 13% O₂ (hypoxia) and 5-min 21% O₂ (normoxia) per day for 14 days, followed by an additional continuous hypoxic treatment. Hypoxic-reoxygenated mice were fed normoxic for 28 days after receiving chronic hypoxia.

3. The siP-Syn peptide administration

The polypeptide we used was based on a previous study[16]. The sequence was 5-FITC-(Acp)-YGRKKRRQRRRYEMPSEEGYQD. The peptide was formed by fusing the HIV Tat (transcriptional counter-activator) signal (YGRKKRRQRRR) with a competing peptide spanning amino acids 125 to 135 (YEMPSEEGYQD), the phosphorylation site (ser129) in α -syn (siP-Syn).

The siP-Syn peptide (Chinapeptides, Shanghai, China) was injected via the tail vein every 3 days for 6 weeks at a dose of 10 mg/kg. The mice were treated in a hypoxia-inducing environment and tested for behavior two weeks after injection. Polypeptides were stored at -20°C and protected from light. Repeated freezing and thawing were prevented.

4. Behavioral tests

The cognitive function of the mice in each group was evaluated by novel object recognition tests and Morris water maze tests. For the novel object recognition tests, the experimental device was a rectangular box with a length, width, and height of 40 cm. It had no lid, and the camera was placed on the top. The experiment was divided into three stages: adaptation stage, familiarity stage, and experimental stage. Adaptation stage: Each mouse was put into the experimental device and left to explore freely for 5 min to adapt. Familiarity stage: Two identical objects A (old objects) were placed in the experimental device, and the mice were allowed to explore freely for 5 min. Experimental stage: One object A was removed from the experimental device, and a new object B was put into it (the color and shape were different from A). The mice were put into the experimental device and left to move freely for 5 min. The time that the mice explored the old and new objects were recorded. Discrimination index = (new object recognition time - object recognition time)/(new object recognition time + object recognition time).

The water maze device was a circular device with a diameter of 120 cm. It was divided into four quadrants and had a small platform. Each quadrant had a different color and marker shape for spatial memory recognition by the mice. Water was added to the pool until it was 1–2 cm higher than the platform. The water temperature was about 25°C. Training stage: The mice were placed in the pool at the same position in one of the four quadrants, and the time spent finding the platform was recorded. If the mice could not find the platform within 60 seconds, the time spent finding the platform was recorded as 60 seconds. The mice were then guided to swim to the platform and allowed to stay there for 5 seconds. The training was performed once a day for 5 days in total. Test stage: On day 6, the platform was removed and the mice were placed in the quadrant farthest from the platform. The time at which the mice

first found the location of the original platform and the number of times of crossing within 60 seconds were recorded.

5. Nissl staining

Brain tissue from the mice were cut into 10 μm sections on a frozen slicer and pasted on a slide. The sections were then fixed in 70% ethanol and successively dehydrated in 100% ethanol, 90% ethanol, 80% ethanol, and 70% ethanol for 2 minutes each. After the xylene became transparent, the sections were placed in 1% tar purple (Solarbio, G1430) and incubated for 30 min. The sections were cleaned with distilled water and separated with 70% alcohol for several minutes. The sections were dehydrated with 70% ethanol, 80% ethanol, and 95% ethanol for 2 min each, dehydrated with 100% ethanol, and sealed with neutral gum.

6. Separation of soluble and insoluble components

Single mouse hippocampal tissue was isolated, and 300 μL mixed protein lysis solution with RIPA (Applygen, C1053), a protease inhibitor (Merck, 539134), and a phosphatase inhibitor (Merck, 524628) (=100:1:2) were added. After ultrasonic crushing, ice cracking was performed for 30 min. The preparation was centrifuged at 4°C at 12,000 rpm for 15 min. The supernatant contained the RIPA-soluble protein components. The precipitate at the bottom of the tube was the insoluble fraction. Urea (8 M) was added to the precipitate to dissolve the RIPA-insoluble protein components.

7. Separation of nuclear components

Nuclear separation was performed using a nuclear separation kit (Applygen, P1202) according to the instructions. Briefly, the mouse hippocampus was put into a pre-cooled glass homogenizer, and 500 μL of Buffer1 reagent was added. After grinding and mixing, the homogenate was placed on ice for 10 min for full lysis. The homogenate was transferred to a new pre-labeled centrifuge tube and centrifuged at 800 $\times g$ for 5 min at 4°C. After centrifugation, the nuclear crude extract was at the bottom of the tube. Then, 500 μL Buffer2 was added to the bottom of the tube containing the crude nuclear extract. The crude nuclear extract was suspended and mixed by shock. Following centrifugation at 4,000 $\times g$ at 4°C for 5 min, the supernatant was discarded. Then, 500 μL Buffer2 was added again after suspension and centrifuged again to obtain relatively pure nuclei. Finally, 50 μL suspension buffer was used to re-suspend and obtain complete and unbroken nuclei.

8. Western blots

Mouse hippocampus protein lysates were resolved by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) and subsequently immunoblotted onto polyvinylidene difluoride (PVDF) membranes. Membranes were blocked with 5% nonfat milk at room temperature for 1 h. After TBST washing (three times, 5 min per wash), the membranes were incubated with the indicated primary antibodies at 4°C overnight with shaking. The primary antibodies included β -actin (Sigma-Aldrich, A5316), lamin B1 (HUABIO Antibodies, ET1606-27), HIF-1 α (Abcam, ab228649), HIF-2 α (Novus, NB100-122), Acer2

(PA5-39016), PP2A (Invitrogen, MA5-32920), p-PP2A (Invitrogen, MA5-32158), α -syn (BD, 8249518), and p- α -syn (Abcam, ab51253). After incubation, the membranes were washed three times and then incubated at room temperature for 1 h with secondary antibodies, including IRDye 680RD goat anti-mouse IgG (H+L) (Licor, 926-68070), IRDye 680RD goat anti-rabbit IgG (H+L) (Licor, 926-68071), IRDye 800CW goat anti-mouse IgG (H+L) (Licor, 926-32210), IRDye 800CW goat anti-rabbit IgG (H+L) (Licor, 926-32211). Membranes were scanned using a detection system (Odyssey, USA), and band intensities were normalized to β -actin. Statistical analyses were performed using ImageJ and GraphPad software.

9. Immunofluorescence

Mice brain sections were boiled in citric acid buffer (pH = 6.0) for 10 min for antigen repair. Sections were placed in 0.01 M PBS and washed 3 times for 10 min each. Sections were placed in 1% PBST and incubated at room temperature for 30 min. Sections were then placed in 5% BSA and sealed for 2 h at room temperature. The sections were then incubated overnight at 4°C in primary antibody: p- α -syn (WAKO, 015-25191) or MAP2 (Abcam, ab32454). The sections were washed in 0.01 M PBS 3 times for 10 min each. Sections were placed in fluorescent secondary antibody and incubated at room temperature for 2 h, protected from light. The secondary antibodies were goat anti-mouse IgG (H+L) highly cross-adsorbed secondary antibody, Alexa Fluor 488 (Invitrogen, A11029), and goat anti-rabbit IgG (H+L) highly cross-adsorbed secondary antibody, Alexa Fluor 594 (Invitrogen, A11037). DAPI was used for nuclei staining before sections were mounted (Sigma, D9542). A fluorescence microscope was used to observe and photograph the images.

10. Enzyme activity detection

A PP2A enzyme activity detection kit (Enzyme-Linked Biotechnology, JK56324), CK enzyme activity detection kit (Enzyme-Linked Biotechnology, JK68421), and PLK enzyme activity detection kit (Enzyme-Linked Biotechnology, JK89575) were used according to the instructions to detect enzyme activities. Briefly, a standard hole, blank hole, and sample hole were set. Then, 50 μ L standard was added to the standard well, 50 μ L sample dilution buffer to the blank well, and 10 μ L sample with 40 μ L dilution buffer to the sample well. Then, 100 μ L horseradish peroxidase (HRP)-labeled antibody was added to each well. The wells were sealed and incubated at 37°C for 60 min. The plate was washed 5 times before 50 μ L substrates A and B were added. The plate was incubated at 37°C for 15 min away from light. Stop solution (50 μ L) was added, and the OD value of each well was measured at 450 nm within 15 min. A standard curve with $R^2 > 0.99$ was drawn to prove the existence of a linear relationship. Finally, the concentrations and activities of PP2A, CK, and PLK were calculated.

11. Targeted metabolomics

The mouse hippocampus homogenate was melted at 4°C for 30–60 minutes. Then, 100 μ L homogenate was transferred to a labeled 1.5 mL centrifuge tube, 300 μ L methanol added, and the contents mixed on a vortex mixer for 1 h. The precipitate was collected by centrifugation at 12,000 rpm for 10 min. Then, 95 μ L supernatant was transferred to a labeled 1.5-mL microcentrifuge tube, and 5 μ L of an internal

standard of Sph/Cer was added. After mixing well and centrifugation at 12,000 rpm for 10 min, the supernatant was transferred to a 250 μ L vial insert for analysis. The ACQUITY UPLC HSS T3 2.1 \times 100 mm, 1.8 μ m column (Waters, Milford, MA, USA) was used as the stationary phase. It was suited for lipid retention. The mobile phase consisted of solvent A (0.1% formic acid-acetonitrile/water (3:2 v/v)) and solvent B (0.1% formic acid-isopropanol/acetonitrile (9:1 v/v)) with a gradient elution (0–1.5 min, 68–68% A; 1.5–15.5 min, 68–15% A; 15.5–15.6 min, 15–3% A; 15.6–18.0 min, 3–3% A; 18.0–18.1 min, 3–68% A; 18.1–20 min, 68–68% A). The flow rate of the mobile phase was 0.26 mL/min. The column temperature was maintained at 55°C. The injection volume was 4 μ L. A Waters TQ-S mass spectrometer equipped with an ESI probe was employed. The positive ESI source temperature was 150°C, the capillary voltage was 3.0 kV, the cone voltage was 20 V, the desolvation temperature was 500°C, and the desolvation gas flow was 1,000 L/h. The LC-MS system was controlled by Masslynx software. Data were collected and processed with the same software.

12. RT-PCR

An RNeasy kit (Qiagen, 74104) was used to extract total RNA from mice hippocampal tissue, and then the Transcriptor High Fidelity cDNA synthesis kit (Roche, 5081963001) was used to reverse transcribe the RNA into cDNA. All operations were according to the instructions. The following primers were used: mouse β -actin, forward: 5'-GGCTGTATTCCTCCATCG-3', reverse: 5'-CCAGTTGGTAACAATGCCATGT-3'; mouse Acer2, forward: 5'-TGGTGGCAGGACAACACTACTATC-3', reverse 5'-AAGCATCTGACCCAGGAAACTC-3'.

13. Primary neuron culture and processing

A newborn mouse was disinfected and its head was cut off immediately. The cerebral cortex was removed and the hippocampus was separated under a stereomicroscope. The tissue was cut to less than 1 mm³ in DMEM medium (Gibco, C11995500B7). A 1:40 volume ratio of papain solution was added with subsequent digestion at 37°C for 30 min. The papain was removed, DMEM culture medium was added, and the tissue was blown to make a cell suspension. The cell suspension was passed through a 40 μ m filter (Falcon, 352340). The cells were diluted to a certain density and then moved to polylysine (Sigma, P1399)-coated plates for culturing at 37°C in a 5% CO₂ incubator. After 4–8 h, DMEM was replaced with Neurobasal medium (Gibco, 21103049) containing 50 \times B27 (Gibco, 17504044), 100 \times glutamine (Invitrogen, 25030081), and 100 \times penicillin-streptomycin (Thermo, 15140122). Half of the medium was changed every 3 days after cell adhesion.

14. Cell viability detection

Cell viability was detected by the MTT assay kit (Sigma, M2128). MTT (5 mg) was placed in a 5 mL EP tube, and PBS buffer (1 mL) added to dissolve the MTT. After blending, the stock solution was kept at -20°C and protected from light. After the experimental treatments, the cell supernatant in the 96-well plate were discarded, 10 μ L MTT solution and 90 μ L culture solution were added to the pre-designed cell wells. The cells were incubated at 37°C with 5% CO₂ for 4 hours. The supernatant was discarded and 100 μ L

DMSO was added and mixed gently for 10 min while avoiding light. The OD value at 490 nm was detected by an enzyme marker.

15. Cytotoxicity detection

Cytotoxicity was detected by the LDH assay (Roche, 4744926001). The powder was dissolved in ddH₂O and mixed thoroughly to make the catalytic solution. Then, 250 μ L of the catalytic solution was added to the staining solution (11.25 mL) and mixed thoroughly. Then, 100 μ L cell supernatant of each group was added to the new 96-well plate. The LDH reaction solution (100 μ L) was added with subsequent incubation at room temperature away from light for 30 min. After the incubation, 50 μ L stop solution was added to each well and gently mixed for 10 min. The OD value of each well was measured at 490 nm by a microplate reader.

16. Cell death detection

PI/Hoechst detection was used to detect the cell death rate. Hoechst labels all cells as blue fluorescence, and PI labels only dead cells as red fluorescence. Therefore, the ratio of red to blue can be used to calculate the cell death rate. After the cells were treated, the original medium was discarded, and the cells were rinsed three times with PBS. The PI and Hoechst mixture was added into the cell culture well and incubated at 37°C for 10 min under dark conditions. The cells were removed from the incubator, the mixture of PI and Hoechst was discarded, and the cells were rinsed three times with PBS. Confocal microscopy was used for observation and imaging.

17. Statistical analysis

Excel and GraphPad Prism 9.0 software were used for data preservation, recording, statistics, and analyses. Image data were analyzed by ImageJ and other software. All results were analyzed using the *t*-test, one-way ANOVA, and two-way ANOVA as appropriate. Data are expressed as the mean \pm standard error (mean \pm SEM), with $P \leq 0.05$ as a significant difference.

Results

1. Chronic hypoxia causes cognitive impairment in mice by promoting hippocampal neurodegeneration.

Adult C57BL mice were continuously treated with 13% O₂ for 0–14 days to construct a mouse hypoxia model. This hypoxic concentration is approximately equal to that at Lhasa on the Tibetan Plateau, the highest plateau city[17]. Mice at 0, 1, 3, 7, and 14 days of hypoxia (Con, H1d, H3d, H7d, H14d) underwent behavioral tests and postmortem histological tests to assess the neurological damage (Figure 1A). New object recognition tests and the Morris water maze tests were used to assess the novel object exploration and spatial memory abilities. The results of these two behavioral tests showed that H1d and H3d mice did not experience cognitive decline, while H7d and H14d mice showed a significant weakening in novel object recognition and impaired spatial memory abilities (Figure 1B–D). Mouse hippocampal neurons

stained by Nissl staining showed that the hippocampal neurons of mice in the Con, H1d, and H3d groups had an orderly arrangement but a disordered arrangement after prolonged hypoxia, especially in the H14d group. These results are consistent with the cognitive decline observed in mice with chronic hypoxia (Figure 1E). Fluoro-Jade C (FJC) staining showed neurodegeneration in the hippocampal CA1 and DG regions in mice in the H7d and H14d groups compared with the Con group (Figure 1F). Therefore, these results suggested that chronic and persistent hypoxia induced cognitive impairment in mice by inducing degeneration of hippocampal neurons.

2. Chronic hypoxia promotes α -syn phosphorylation at ser129 and abnormal aggregation in the mouse hippocampus.

Hyperphosphorylation and abnormal aggregation of α -syn are closely related to neuronal degeneration. To investigate whether abnormal changes in α -syn were involved in neuronal injury induced by chronic hypoxia, we detected the levels of α -syn and p- α -syn in the hippocampi of mice in each group. The α -syn and p- α -syn levels in the Radio Immunoprecipitation Assay (RIPA)-soluble fraction from the hippocampi of mice increased significantly with hypoxic treatment for one day compared with the Con group (Figure 2A, C, D). In addition, α -syn and p- α -syn levels increased in the insoluble fraction of RIPA following hypoxic treatment for 7 days, suggesting that hypoxia promoted the phosphorylation of α -syn and long-term hypoxia promoted α -syn to form abnormal aggregates in the insoluble fraction (Figure 2B, E, F).

Immunofluorescence staining was carried out to further observe whether the increased p- α -syn induced by hypoxia was aggregated. The results showed almost no significant p- α -syn staining in the hippocampal tissue of mice in the Con group. However, a large amount of p- α -syn staining aggregated in the hippocampal tissue of mice in the H14d group (Figure 2G–H).

Primary neurons from mouse hippocampi were cultured for 7 days and then treated with 1% O₂ for 0–6 hours (Con, H1h, H2h, H4h, H6h). We tested cell viability and cytotoxicity of the neurons in each group by 3-(4,5)-dimethylthiazoliazol-2-yl-5-carboxymethyl imidazole hydrochloride (MTT) and lactate dehydrogenase (LDH) assays. The results showed no significant decrease in cell viability within 2 h of hypoxic treatment but decreased cell viability after 4 h of hypoxia. We chose 4 h of hypoxia for subsequent experimental detection (Supplemental Figure 1A, B). We used Propidium Iodide (PI)/Hoechst double staining to detect cell mortality. The results also showed that hypoxia for 4 h significantly increased the mortality of the primary neurons (Supplemental Figure 1C, D). Similar to in vivo experiments, we also found that hypoxia promoted α -syn phosphorylation and increased its level in primary neurons (Supplemental Figure 2). Immunofluorescence staining of α -syn and p- α -syn showed that aggregates formed in the cytoplasm and axons of neurons after hypoxic treatment. In particular, when p- α -syn formed aggregates, the neuron marker Map2 disappeared, indicating that the neurons were dead (Supplemental Figure 3). These results suggested that chronic hypoxia resulted in an increase in p- α -syn phosphorylation and abnormal aggregation.

3. The α -syn knockout mice or mice administered a competing peptide targeting ser129 resist chronic hypoxia-induced cognitive impairment.

The above studies showed that chronic hypoxia induced cognitive impairment in mice and was accompanied by abnormal changes in α -syn. To investigate whether abnormal changes in α -syn were a critical factor in chronic hypoxia-induced nerve injury, we used α -syn knockout mice. Heterozygous α -syn knockout mice received the same chronic hypoxia-inducing treatment. New object recognition tests and Morris water maze tests were used to detect the cognitive behavior of the mice (Figure 3A). The α -syn levels in the hippocampi of heterozygous α -syn knockout mice were significantly lower than in wild-type mice (Figure 3B, J). The behavioral test results showed that α -syn knockdown significantly alleviated the chronic hypoxia-induced decline in exploration ability to novel objects and spatial memory impairment (Figure 3C–E). These results suggested that α -syn played a critical role in neuronal injury induced by chronic hypoxia. western blot analysis also showed that chronic hypoxia did not induce an excessive increase in α -syn and p- α -syn in the hippocampi of heterozygous α -syn knockout mice compared with wild-type mice (Figure 3F–H). Immunofluorescence staining also showed that chronic hypoxic treatment in heterozygous α -syn knockout mice did not result in significant p- α -syn aggregate formation (Figure 3I). Similarly, in primary neurons, we also found that knockdown of A-SYN by siRNA lentivirus mitigated hypoxia-induced neuronal death (Supplemental Figure 4). These data suggest that abnormal alterations in α -syn are important in chronic hypoxia-induced nerve injury.

Hyperphosphorylation of α -syn at ser129 is an important step in the transformation of α -syn both from soluble monomers to insoluble aggregates and from a non-toxic to a toxic form. Our previous experiments have confirmed that chronic hypoxia promotes increased levels of α -syn phosphorylation. Therefore, we further explored whether preventing α -syn phosphorylation would resist chronic hypoxia-induced nerve damage. We used a previously reported small-molecule peptide that competently inhibits α -syn hyperphosphorylation at ser129[16] and administered it to mice via the tail vein (Figure 4A). Western blots showed that p- α -syn levels were significantly reduced in the peptide-treated mice (siP-Syn) (Figure 4B, J). The siP-Syn mice were also subjected to chronic hypoxia as in the previous experiments. The behavioral tests showed that, unlike wild-type mice, siP-Syn mice were able to resist cognitive impairment induced by chronic hypoxia (Figure 4C–E). This occurred mainly due to the resistance of siP-Syn mice to the hyperphosphorylation and abnormal aggregation of α -syn in the hippocampus induced by chronic hypoxia (Figure 4F–I). These results suggested that chronic hypoxia induced abnormal aggregation of α -syn by promoting its hyperphosphorylation, which was a key factor in inducing nerve damage and cognitive impairment in mice.

4. Chronic hypoxia induces a vicious cycle of decreased PP2A activity and increased p- α -syn.

We have found that hypoxia promoted α -syn phosphorylation, but the molecular mechanism remained unclear. Phosphorylation of α -syn is regulated by kinases and phosphatases. Polo-like kinase (PLK) and creatine kinase (CK) are the main kinases promoting α -syn phosphorylation[18], while PP2A is the main phosphatase promoting α -syn dephosphorylation[19]. To explore how hypoxia causes α -syn hyperphosphorylation, we tested the enzyme activities of these enzymes. The results showed that the activity of kinases CK and PLK in the hippocampal tissue of mice in the hypoxic group increased compared with the control group. However, there was no statistical significance. The activity of

phosphatase PP2A in the hypoxia group decreased significantly (Figure 5A–C). We further detected the level of PP2A and its phosphorylation at tyrosine 307 (p-PP2A) by western blotting, which characterized its inactivity. The results showed that the p-PP2A/PP2A ratio increased significantly after hypoxia, indicating that the PP2A activity decreased (Figure 5D, G). Interestingly, we found that α -syn knockout mice or siP-Syn mice were resistant to a chronic hypoxia-induced decline in PP2A activity, showing no increase in the p-PP2A/PP2A ratio after hypoxia (Figure 5E, F, H, I).

Similarly, in primary neurons, we also found that hypoxic treatment resulted in decreased activity in PP2A in neurons. We treated cells with both iHAP1[20], a PP2A agonist, and LB-100[21], an inhibitor of PP2A, and showed that iHAP1 mediated the activation of PP2A mitigated hypoxia-induced neuronal death (Supplemental Figure 5).

These results suggested that chronic hypoxia promotes hyperphosphorylation of α -syn through inhibition of PP2A activity and further inhibition of PP2A activity through p- α -syn, creating a vicious cycle.

5. Chronic hypoxia promotes ceramide hydrolytic metabolism by activating the HIF-2 α /Acer2 signaling pathway, and thereby downregulates PP2A activity.

The above study found that hypoxia caused α -syn hyperphosphorylation by inhibiting PP2A activity. Therefore, we further explored the underlying mechanism. Ceramides are essential for maintaining PP2A activity in vivo[22]. For this reason, we tested different species of ceramides by using targeted metabolomics. The results showed that hypoxia reduced the levels of almost all ceramide species, and this was significantly correlated with the duration of hypoxia (Figure 6A–E). Recently, it was reported that the ceramide hydrolase alkaline ceramidase (Acer)2 is a novel downstream target molecule of hypoxia-inducible factor (HIF)-2 α [23]. For this reason, we investigated whether hypoxia caused a reduction in ceramide levels through this pathway. HIF-1 α , HIF-2 α , and Acer2 levels were detected by western blots. The results showed that HIF-1 α increased temporarily in the H1d group and then decreased in groups with longer hypoxia (Figure 6F–G). In contrast, HIF-2 α , which is more indicative of chronic hypoxia, increased steadily with prolonged hypoxia time (Figure 6F, H). Similarly, Acer2, downstream of HIF-2 α , also showed a sustained and stable increase after hypoxia at both protein and mRNA levels (Figure 6F, I). To further confirm that the increased Acer2 level after hypoxia was caused by transcriptional activation, we detected the expression of Acer2 by RT-PCR and obtained similar results. That is, chronic hypoxia steadily promoted the expression of Acer2 (Figure 6J). These results suggested that chronic hypoxia promoted the expression of Acer2 by activating HIF-2 α . Increased Acer2 promoted the hydrolytic metabolism of ceramides. It may be an important reason for the inhibition of PP2A activity and the hyperphosphorylation of α -syn.

6. Chronic hypoxia-induced α -syn pathology and cognitive impairment in mice remain irreversible after reoxygenation.

We have found that chronic hypoxia promoted the abnormal pathology of α -syn through an HIF-2 α -dependent pathway, which is an important link in inducing neuronal injury and cognitive impairment in

mice. However, it is not known whether α -syn pathology and cognitive impairment induced by chronic hypoxia are reversible or irreversible. Therefore, the hypoxic mice were reoxygenated for a long time to observe the recovery of the mice (Figure 7A). The results showed that the mice that experienced hypoxia for 14 days still had obvious cognitive behavioral impairment after reoxygenation for 28 days, suggesting that the neurological damage induced by chronic hypoxia was irreversible (Figure 7B–F). By western blot analysis, we still found a large amount of p- α -syn in the hippocampal tissues of reoxygenated mice, suggesting that α -syn pathology induced by chronic hypoxia was also irreversible for an extended period of time (Figure 7G–I). These results suggested that α -syn pathology and cognitive impairment induced by chronic hypoxia were irreversible over an extended reoxygenation period in mice.

7. Intermittent hypoxia preconditioning alleviates chronic hypoxia-induced cognitive impairment in mice by inhibiting α -syn phosphorylation and aggregation.

The above study found that chronic persistent hypoxia induced irreversible cognitive impairment in mice. Compared with persistent hypoxia, intermittent hypoxia may be a more moderate pattern of hypoxia. Previous studies have proposed that intermittent hypoxia preconditioning (IHPC) could increase the body's tolerance to subsequent more severe injury[24]. Therefore, we investigated whether IHPC could resist the α -syn pathology and cognitive impairment in mice induced by chronic persistent hypoxia. Our IHPC treatment pattern was as follows: mice were treated with 10 cycles of intermittent hypoxia per day. Each cycle of 13% O₂ hypoxic treatment for 5 min was followed by 21% O₂ normoxia for 5 min. Mice were treated with IHPC for a total of 14 days, followed by continuous treatment for another 14 days in 13% O₂ hypoxic conditions (Figure 8A). Novel object recognition tests and the Morris water maze tests showed that IHPC significantly alleviated cognitive impairment induced by persistent hypoxia in mice (Figure 8B–G). We detected α -syn and p- α -syn in the hippocampal tissues of mice by western blots and immunofluorescence staining. The results showed that IHPC could significantly alleviate α -syn pathology induced by chronic hypoxia (Figure 8H–K). These results suggested that IHPC alleviated chronic hypoxia-induced cognitive impairment and α -syn pathology in mice.

Discussion

In this study, we found that chronic hypoxia induced irreversible cognitive impairment in mice with α -syn phosphorylation and aggregation playing a crucial role. Mechanically, chronic hypoxia promoted the transcriptional expression of *Acer2* by activating HIF-2 α . Increased *Acer2* resulted in the hydrolytic metabolism of ceramide and subsequent decreased activity of PP2A. These changes induced α -syn phosphorylation and abnormal aggregation, resulting in neurological damage. These findings revealed a novel mechanism of cognitive impairment induced by chronic hypoxia. In addition, we found that IHPC significantly alleviated chronic hypoxia-induced α -syn pathology and cognitive impairment in mice, which provided a novel perspective for the intervention of hypoxia-related diseases.

Chronic hypoxia is common in life and includes environmental hypoxia, such as altitude exposure, and hypoxia of the body or organs during aging and disease[25]. The brain is very sensitive to hypoxic stress.

Both environmental hypoxia and body hypoxia cause neuronal damage in the brain[26]. However, the mechanism remains unclear. Our study first highlighted the importance of α -syn phosphorylation and aggregation in chronic hypoxia-induced cognitive impairment, showing that both α -syn knockout and p- α -syn competing interference resisted chronic hypoxia-induced neuronal injury. Furthermore, we systematically revealed the molecular mechanism of chronic hypoxia-induced α -syn pathology and neuronal injury. Compared with the acute hypoxic response mediated mainly by HIF-1 α , HIF-2 α is involved mainly in the chronic hypoxic response[27]. Our study found that chronic hypoxia promoted HIF-2 α rather than HIF-1 α entry into the nucleus, and nucleation of HIF-2 α resulted in transcriptional expression of *Acer2*, a newly discovered downstream target of HIF that was reported in adipocytes[23, 28]. As an important hydrolytic enzyme of ceramides[29], increased *Acer2* promotes the hydrolytic metabolism of ceramides, which is an important reason for the significantly decreased ceramide levels observed after hypoxic treatment. The decrease in ceramides, as the main maintainer of PP2A activity in the brain[22], resulted in the inhibition of PP2A activity. PP2A is the only phosphatase responsible for α -syn dephosphorylation[19]. Therefore, the decreased activity of PP2A induced by chronic hypoxia is an important factor for α -syn hyperphosphorylation and abnormal aggregation. Interestingly, we also found that the hypoxia-induced decline in PP2A activity was reversed when α -syn or p- α -syn was interfered with, suggesting that hypoxia-induced PP2A inhibition and α -syn pathology are mutually reinforcing vicious cycles. This hypothesis has been confirmed by previous studies[30, 31]. In conclusion, our study highlighted the role of α -syn pathology in chronic hypoxia-induced cognitive impairment and revealed a novel underlying mechanism, providing reliable targets for future research.

The α -syn is a protein that is abundant in the brain and is pathologically involved in a variety of neurological diseases. PD and DLB are the most studied diseases of α -syn at present[32, 33]. Their typical pathological feature is the deposition of intracellular Lewy bodies, the main component of which is p- α -syn. A few previous studies have mentioned the relationship between hypoxia and α -syn. For example, hypoxia can increase the expression and aggregation of α -syn in HEK293 cells[15, 34]. There has also been a study showing increased plasma α -syn levels in patients with chronic hypoxia[35]. A recent review has suggested that improving hypoxia adaptation may be an important strategy for PD intervention[36]. In our study, we found that hypoxia significantly promoted phosphorylation of α -syn at ser129, a form of α -syn that was strongly associated with the above diseases. These results suggested that hypoxia stress may play an initiating or promoting role in the pathogenesis of α -syn-related diseases such as PD and DLB. These findings provided a novel perspective for further research and interventions for these diseases.

Compared with persistent hypoxia, intermittent hypoxia is a moderating pattern of hypoxia, and many previous studies have found that IHPC can resist subsequent more severe ischemic hypoxic injury[37, 38]. IHPC has been reported to play protective roles in various neurological diseases[24], it is considered as a potential non-drug intervention[39]. The protective effects of intermittent hypoxia in ischemic stroke have been extensively studied, and IHPC has been shown to reduce the infarct size and improve neurological function in stroke models[40-42]. Several studies have shown that IHPC exerted a neuroprotective effect in models of multiple sclerosis by suppressing inflammation and protecting the blood-brain barrier[43].

IHPC treatment also reduced amyloid pathology and improved cognitive performance in A β PP/PS1 transgenic mice of Alzheimer's disease[44]. In this study, we found that IHPC alleviated chronic hypoxia-induced cognitive impairment in mice and resisted α -syn pathology, which provided a potential direction for the intervention of both hypoxia-related diseases and α -syn-related diseases, such as Parkinson's disease. It is important to note that the IHPC we used was a more moderate model having alternating hypoxia for 5 min and normoxia for 5 min over 10 cycles per day for 14 days. This did not cause additional damage. It is quite different from the intermittent hypoxic conditions associated with some diseases, such as obstructive sleep apnea syndrome.

Conclusions

In summary, our study highlighted the importance of α -syn pathology in hypoxia-related cognitive disorders, elucidated its molecular mechanisms, and provided a reliable molecular target for subsequent research (Supplemental Figure 6). Meanwhile, we also revealed the role of IHPC in resistance to chronic hypoxic injury, which provided a new perspective for the intervention of many hypoxia-related cognitive disorders.

Abbreviations

Acer2, alkaline ceramidase 2; α -syn, alpha-synuclein; CK, creatine kinase; CNS, central nervous system; DLB, dementia of Lewy body; FJC, Fluoro-Jade C; HIF, hypoxia-inducible factor; IHPC, intermittent hypoxia preconditioning; LDH, lactate dehydrogenase; MTT, 3-(4,5)-dimethylthiazolium (-z-y1)-3,5-di-phenyltetrazolium bromide; PD, Parkinson's disease; PI, Propidium Iodide; PLK, polo-like kinase; PP2A, protein phosphatase 2A; RIPA, Radio Immunoprecipitation Assay.

Declarations

- **Ethics approval and consent to participate**

All animal experiments were approved by the Animal Care and Use Committee of the Institute of Animal Management, Capital Medical University, and conducted in accordance with ethical requirements (Num: AEEI-2021-084).

- **Consent for publication**

All authors agree to this manuscript and submit it to Molecular Neurodegeneration.

- **Availability of data and materials**

All data in this study are included in this manuscript and its supplementary information files.

- **Competing interests**

The authors declare that they have no competing interests.

- **Funding**

This manuscript was supported by the National Natural Science Foundation of China (Grant number: 82027802, 32100925), the Beijing Nova Program (Grant number: Z211100002121038), the Beijing

Hundred Thousand and Ten Thousand Talent Project (Grant number: 2019A36), and the Beijing Municipal Health Commission (Grant number: 303–01–005-0019).

- **Authors' contributions**

Gaifen Li, Jia Liu, and Xunming Ji conceived and designed the study. Gaifen Li, Jia Liu, Mengyuan Guo, Yakun Gu, Yuying Guan, Qianqian Shao, Wei Ma, and Xunming Ji performed the experiments and analyzed the data. Gaifen Li, Jia Liu, and Xunming Ji wrote the manuscript. All authors read and approved the final manuscript.

- **Acknowledgements**

We thank Prof. Hui Yang of Capital Medical University for providing the α -syn knockout mice for this project. Thanks also to Beijing Qinglian Biotech Co., Ltd for the detection in target metabolomics.

References

1. Golde TE. Alzheimer's disease - the journey of a healthy brain into organ failure. *Mol Neurodegener.* 2022;17(1):18.
2. Oedekoven C, Egeri L, Jessen F, Wagner M, Dodel R. Subjective cognitive decline in idiopathic Parkinson's disease: A systematic review. *Ageing Res Rev.* 2022;74:101508.
3. Taylor JP, McKeith IG, Burn DJ, Boeve BF, Weintraub D, Bamford C, Allan LM, Thomas AJ, O'Brien JT. New evidence on the management of Lewy body dementia. *Lancet Neurol.* 2020;19(2):157–69.
4. Poh L, Sim WL, Jo DG, Dinh QN, Drummond GR, Sobey CG, Chen CL, Lai MKP, Fann DY, Arumugam TV. The role of inflammasomes in vascular cognitive impairment. *Mol Neurodegener.* 2022;17(1):4.
5. McKenna HT, Murray AJ, Martin DS. Human adaptation to hypoxia in critical illness. *J Appl Physiol (1985).* 2020;129(4):656–63.
6. Burtscher J, Mallet RT, Burtscher M, Millet GP. Hypoxia and brain aging: Neurodegeneration or neuroprotection? *Ageing Res Rev.* 2021;68:101343.
7. Signore AV, Storz JF. Biochemical pedomorphosis and genetic assimilation in the hypoxia adaptation of Tibetan antelope. *Sci Adv.* 2020;6(25):eabb5447.
8. Norris PC, Libreros S, Serhan CN. Resolution metabolomes activated by hypoxic environment. *Sci Adv.* 2019;5(10):eaax4895.
9. Wilson MH, Newman S, Imray CH. The cerebral effects of ascent to high altitudes. *Lancet Neurol.* 2009;8(2):175–91.
10. Mitroshina EV, Savyuk MO, Ponimaskin E, Vedunova MV. Hypoxia-Inducible Factor (HIF) in Ischemic Stroke and Neurodegenerative Disease. *Front Cell Dev Biol.* 2021;9:703084.
11. Koga S, Sekiya H, Kondru N, Ross OA, Dickson DW. Neuropathology and molecular diagnosis of Synucleinopathies. *Mol Neurodegener.* 2021;16(1):83.
12. Cai W, Srivastava P, Feng D, Lin Y, Vanderburg CR, Xu Y, McLean P, Frosch MP, Fisher DE, Schwarzschild MA, et al. Melanocortin 1 receptor activation protects against alpha-synuclein pathologies in models of Parkinson's disease. *Mol Neurodegener.* 2022;17(1):16.

13. Arawaka S, Sato H, Sasaki A, Koyama S, Kato T. Mechanisms underlying extensive Ser129-phosphorylation in alpha-synuclein aggregates. *Acta Neuropathol Commun.* 2017;5(1):48.
14. Anderson JP, Walker DE, Goldstein JM, de Laat R, Banducci K, Caccavello RJ, Barbour R, Huang J, Kling K, Lee M, et al. Phosphorylation of Ser-129 is the dominant pathological modification of alpha-synuclein in familial and sporadic Lewy body disease. *J Biol Chem.* 2006;281(40):29739–52.
15. Chen T, Li J, Chao D, Sandhu HK, Liao X, Zhao J, Wen G, Xia Y. delta-Opioid receptor activation reduces alpha-synuclein overexpression and oligomer formation induced by MPP(+) and/or hypoxia. *Exp Neurol.* 2014;255:127–36.
16. Su Y, Deng MF, Xiong W, Xie AJ, Guo J, Liang ZH, Hu B, Chen JG, Zhu X, Man HY, et al. MicroRNA-26a/Death-Associated Protein Kinase 1 Signaling Induces Synucleinopathy and Dopaminergic Neuron Degeneration in Parkinson's Disease. *Biol Psychiatry.* 2019;85(9):769–81.
17. Zhang X, Witt KE, Banuelos MM, Ko A, Yuan K, Xu S, Nielsen R, Huerta-Sanchez E. **The history and evolution of the Denisovan-EPAS1 haplotype in Tibetans.** *Proc Natl Acad Sci U S A* 2021, 118(22).
18. Waxman EA, Giasson BI. Characterization of kinases involved in the phosphorylation of aggregated alpha-synuclein. *J Neurosci Res.* 2011;89(2):231–47.
19. Lee KW, Chen W, Junn E, Im JY, Grosso H, Sonsalla PK, Feng X, Ray N, Fernandez JR, Chao Y, et al. Enhanced phosphatase activity attenuates alpha-synucleinopathy in a mouse model. *J Neurosci.* 2011;31(19):6963–71.
20. Morita K, He S, Nowak RP, Wang J, Zimmerman MW, Fu C, Durbin AD, Martel MW, Prutsch N, Gray NS, et al. Allosteric Activators of Protein Phosphatase 2A Display Broad Antitumor Activity Mediated by Dephosphorylation of MYBL2. *Cell.* 2020;181(3):702–15 e720.
21. Ho WS, Sizdahkhani S, Hao S, Song H, Seldomridge A, Tandle A, Maric D, Kramp T, Lu R, Heiss JD, et al. LB-100, a novel Protein Phosphatase 2A (PP2A) inhibitor, sensitizes malignant meningioma cells to the therapeutic effects of radiation. *Cancer Lett.* 2018;415:217–26.
22. O'Connor CM, Perl A, Leonard D, Sangodkar J, Narla G. Therapeutic targeting of PP2A. *Int J Biochem Cell Biol.* 2018;96:182–93.
23. Zhang X, Zhang Y, Wang P, Zhang SY, Dong Y, Zeng G, Yan Y, Sun L, Wu Q, Liu H, et al. Adipocyte Hypoxia-Inducible Factor 2alpha Suppresses Atherosclerosis by Promoting Adipose Ceramide Catabolism. *Cell Metab.* 2019;30(5):937–51 e935.
24. Liu J, Gu Y, Guo M, Ji X. Neuroprotective effects and mechanisms of ischemic/hypoxic preconditioning on neurological diseases. *CNS Neurosci Ther.* 2021;27(8):869–82.
25. Yeo EJ. Hypoxia and aging. *Exp Mol Med.* 2019;51(6):1–15.
26. Dunn JF, Isaacs AM. The impact of hypoxia on blood-brain, blood-CSF, and CSF-brain barriers. *J Appl Physiol (1985).* 2021;131(3):977–85.
27. Serocki M, Bartoszewska S, Janaszak-Jasiecka A, Ochocka RJ, Collawn JF, Bartoszewski R. miRNAs regulate the HIF switch during hypoxia: a novel therapeutic target. *Angiogenesis.* 2018;21(2):183–202.

28. Gonzalez FJ, Xie C, Jiang C. The role of hypoxia-inducible factors in metabolic diseases. *Nat Rev Endocrinol.* 2018;15(1):21–32.
29. Xu R, Garcia-Barros M, Wen S, Li F, Lin CL, Hannun YA, Obeid LM, Mao C. Tumor suppressor p53 links ceramide metabolism to DNA damage response through alkaline ceramidase 2. *Cell Death Differ.* 2018;25(5):841–56.
30. Tian H, Lu Y, Liu J, Liu W, Lu L, Duan C, Gao G, Yang H. Leucine Carboxyl Methyltransferase Downregulation and Protein Phosphatase Methylesterase Upregulation Contribute Toward the Inhibition of Protein Phosphatase 2A by alpha-Synuclein. *Front Aging Neurosci.* 2018;10:173.
31. Liu G, Chen M, Mi N, Yang W, Li X, Wang P, Yin N, Li Y, Yue F, Chan P, et al. Increased oligomerization and phosphorylation of alpha-synuclein are associated with decreased activity of glucocerebrosidase and protein phosphatase 2A in aging monkey brains. *Neurobiol Aging.* 2015;36(9):2649–59.
32. Blesa J, Foffani G, Dehay B, Bezard E, Obeso JA. **Motor and non-motor circuit disturbances in early Parkinson disease: which happens first?** *Nat Rev Neurosci* 2021.
33. Arnaoutoglou NA, O'Brien JT, Underwood BR. Dementia with Lewy bodies - from scientific knowledge to clinical insights. *Nat Rev Neurol.* 2019;15(2):103–12.
34. Chen T, Wang Q, Chao D, Xia TC, Sheng S, Li ZR, Zhao JN, Wen GQ, Ding G, Xia Y. delta-Opioid Receptor Activation Attenuates the Oligomer Formation Induced by Hypoxia and/or alpha-Synuclein Overexpression/Mutation Through Dual Signaling Pathways. *Mol Neurobiol.* 2019;56(5):3463–75.
35. Sun HL, Sun BL, Chen DW, Chen Y, Li WW, Xu MY, Shen YY, Xu ZQ, Wang YJ, Bu XL. Plasma alpha-synuclein levels are increased in patients with obstructive sleep apnea syndrome. *Ann Clin Transl Neurol.* 2019;6(4):788–94.
36. Bartscher J, Syed MMK, Lashuel HA, Millet GP. Hypoxia Conditioning as a Promising Therapeutic Target in Parkinson's Disease? *Mov Disord.* 2021;36(4):857–61.
37. Li S, Hafeez A, Noorulla F, Geng X, Shao G, Ren C, Lu G, Zhao H, Ding Y, Ji X. Preconditioning in neuroprotection: From hypoxia to ischemia. *Prog Neurobiol.* 2017;157:79–91.
38. Bartscher J, Mallet RT, Pialoux V, Millet GP, Bartscher M. **Adaptive responses to hypoxia and/or hyperoxia in humans.** *Antioxid Redox Signal* 2022.
39. Brocherie F, Millet GP. Hypoxic exercise as an effective nonpharmacological therapeutic intervention. *Exp Mol Med.* 2020;52(3):529–30.
40. Selvaraj UM, Ortega SB, Hu R, Gilchrist R, Kong X, Partin A, Plautz EJ, Klein RS, Gidday JM, Stowe AM. Preconditioning-induced CXCL12 upregulation minimizes leukocyte infiltration after stroke in ischemia-tolerant mice. *J Cereb Blood Flow Metab.* 2017;37(3):801–13.
41. Guan R, Lv J, Xiao F, Tu Y, Xie Y, Li L. Potential role of the cAMP/PKA/CREB signalling pathway in hypoxic preconditioning and effect on propofol-induced neurotoxicity in the hippocampus of neonatal rats. *Mol Med Rep.* 2019;20(2):1837–45.
42. Monson NL, Ortega SB, Ireland SJ, Meeuwissen AJ, Chen D, Plautz EJ, Shubel E, Kong X, Li MK, Freriks LH, et al. Repetitive hypoxic preconditioning induces an immunosuppressed B cell phenotype

- during endogenous protection from stroke. *J Neuroinflammation*. 2014;11:22.
43. Halder SK, Milner R. Chronic mild hypoxia accelerates recovery from preexisting EAE by enhancing vascular integrity and apoptosis of infiltrated monocytes. *Proc Natl Acad Sci U S A*. 2020;117(20):11126–35.
44. Zhang J, Zhang J, Li XJ, Xiao J, Ye F. Hypoxic Preconditioning Ameliorates Amyloid-beta Pathology and Longterm Cognitive Decline in AbetaPP/PS1 Transgenic Mice. *Curr Alzheimer Res*. 2020;17(7):626–34.

Figures




Figure 1

Chronic hypoxia causes cognitive impairment in mice by promoting hippocampal neurodegeneration. A. Mice received continuous hypoxia-inducing treatment for 0, 1, 3, 7, and 14 days (Con, H1d, H3d, H7d, and H14d), and underwent behavioral tests at different time points. B. Detection and statistical analysis of the novel object recognition tests in mice of each group. C, D. Detection and statistical analysis of the Morris water maze tests in mice of each group. E. Nissl staining was used to evaluate the arrangement of

neurons in the hippocampal (HIPPO) CA1 region of each group. F. Fluoro-Jade C (FJC) staining was used to evaluate the neurodegeneration of the DG and CA1 regions in the hippocampus of each group. Data are expressed as the mean \pm SEM (one-way ANOVA). * P <0.05, ** P <0.01, **** P <0.0001, $n = 10$. Bar = 50 μ m in E, F.




Figure 2

Chronic hypoxia promotes α -syn phosphorylation at ser129 and abnormal aggregation in the mouse hippocampus. A–F. The levels of p- α -syn and α -syn in the RIPA-soluble and RIPA-insoluble fractions (8 M urea dissolved) from the hippocampi of mice in each group (Con, H1d, H3d, H7d, H14d) were detected by western blots with β -actin as the internal reference; Statistical analysis was performed. G. The levels of p- α -syn and the neuronal marker Map2 in the hippocampus of the H14d group and Con group were detected by immunofluorescence. H. The high-power three-dimensional imaging of p- α -syn immunofluorescence and the fluorescence intensity of each channel. Data are expressed as the mean \pm SEM (one-way ANOVA). *P<0.05, **P<0.01, n = 3. Bar = 50 μ m in G.




Figure 3

Mice with α -syn knockout resist chronic hypoxia-induced cognitive impairment. A, B, J. Heterozygous mice with α -syn knockout received continuous hypoxia-inducing treatment for 0, 7, and 14 days (Con, H7d, and H14d), and then underwent behavioral tests at different time points. C. Detection and statistical analysis of the novel object recognition tests in mice of each group. D, E. Detection and statistical analysis of the Morris water maze tests in mice of each group. F–H. The levels of p- α -syn and α -syn in the

hippocampi of mice in the Con, H14d, and H14d+ α -syn^{+/-} groups were detected by western blots with β -actin as the internal reference; Statistical analysis was performed. I. The high-power three-dimensional imaging of p- α -syn immunofluorescence. In C–E, data are expressed as the mean \pm SEM (two-way ANOVA), * P <0.05, ** P <0.01, *** P <0.001, **** P <0.0001, n = 10. In G–H, data are expressed as the mean \pm SEM (one-way ANOVA), * P <0.05, *** P <0.001, n = 3. In J, data are expressed as the mean \pm SEM (unpaired t -test), ** P <0.01, n = 3.




Figure 4

Mice administered with competing peptides targeting ser129 resist chronic hypoxia-induced cognitive impairment. A, B, J. Mice were administered competing peptides targeting ser129 and then received continuous hypoxia-inducing treatment for 0, 7, and 14 days (Con, H7d, and H14d) before undergoing behavioral tests at different time points. C. Detection and statistical analysis of the novel object recognition tests in mice of each group. D, E. Detection and statistical analysis of the Morris water maze tests in mice of each group. F–H. The levels of p- α -syn and α -syn in the hippocampi of mice in the Con, H14d, and H14d+siP-Syn groups were detected by western blots with β -actin as the internal reference; statistical analysis was performed. I. The high-power three-dimensional imaging of p- α -syn immunofluorescence. In C–E, data are expressed as the mean \pm SEM (two-way ANOVA), * P <0.05, ** P <0.01, *** P <0.001, **** P <0.0001, n = 10. In G–H, data are expressed as the mean \pm SEM (one-way ANOVA), * P <0.05, ** P <0.01, *** P <0.001, n = 3. In J, data are expressed as the mean \pm SEM (unpaired t -test), * P <0.05, n = 3.




Figure 5

Chronic hypoxia induces a vicious cycle of decreased PP2A activity and increased p- α -syn. A–C. Enzyme activities of PP2A, PLK, and CK in the hippocampi of mice in each group (Con, H1d, H3d, H7d, H14d) were detected and statistically analyzed. D, G. The levels of p-PP2A and PP2A in the hippocampi of mice of different hypoxic time groups (Con, H1d, H3d, H7d, H14d) were detected by western blots with β -actin as the internal reference; Statistical analysis was performed. E, H. The levels of p-PP2A and PP2A in the hippocampi of mice in the Con, H14d, and H14d+ α -syn^{+/-} groups were detected by western blots with β -actin as the internal reference; Statistical analysis was performed. F, I. The levels of p-PP2A and PP2A in the hippocampi of mice in the Con, H14d, and H14d+ H14d+siP-Syn groups were detected by western blots with β -actin as the internal reference; Statistical analysis was performed. Data are expressed as the mean \pm SEM (one-way ANOVA), *P<0.05, **P<0.01, ***P<0.001, n = 3–5.




Figure 6

Chronic hypoxia promotes ceramide hydrolytic metabolism by activating the HIF-2 α /Acer2 signaling pathway, thereby downregulating PP2A activity. A–E. Metabolomics of ceramides in the hippocampal tissue of mice with hypoxia for different periods (Con, H1d, H3d, H7d, H14d); heat map and statistical analysis of different subtypes of ceramides in each group. F–I. The levels of HIF-1 α , HIF-2 α , and Acer2 in the nuclei of the hippocampi of mice of different hypoxic time groups (Con, H1d, H3d, H7d, H14d) were

detected by western blots with β -actin as the internal reference; Statistical analysis was performed. J. The mRNA levels of *Acer2* in the hippocampi in each group were detected by RT-PCR and statistically analyzed. In B–E, data are expressed as the mean \pm SEM (two-way ANOVA), * P <0.05, ** P <0.01, *** P <0.001, n = 10. In G–J, data are expressed as the mean \pm SEM (one-way ANOVA), * P <0.05, ** P <0.01, **** P <0.0001, n = 3.




Figure 7

Chronic hypoxia-induced α -syn pathology and cognitive impairment in mice remain irreversible after reoxygenation. A. Mice received continuous hypoxia-inducing treatment for 14 days (H14d), reoxygenation for 14 days (H14d+R14d) or 28 days (H14d+R28d), and then underwent behavioral tests at different time points. B. Detection and statistical analysis of the novel object recognition tests in mice of each group. C–F. Detection and statistical analysis of the Morris water maze tests in mice of each group. G–I. The levels of p- α -syn and α -syn in the hippocampi of mice in the Con, H14d, H14d+R2W, and H14d+R4W groups were detected by Western blots with β -actin as the internal reference; statistical analysis was performed. Data are expressed as the mean \pm SEM (one-way ANOVA), * P <0.05, ** P <0.01, *** P <0.001, **** P <0.0001, n = 10 in B–E, n = 3 in G–I.




Figure 8

Intermittent hypoxia preconditioning (IHPC) alleviates chronic hypoxia-induced cognitive impairment in mice by inhibiting α -syn phosphorylation and aggregation. A. Mice were pretreated with intermittent hypoxia (IH) for 7 days by 10 cycles of intermittent hypoxia per day, each of which was 13% O_2 for 5 minutes followed by 21% O_2 for 5 min. After pretreatment, the mice were treated with continuous hypoxic conditions for 7 days and 14 days, and underwent behavioral tests at different time points. B. Detection

and statistical analysis of the novel object recognition tests in mice of each group. C–D. Detection and statistical analysis of the Morris water maze tests in mice of each group. E–G. The differences in mice with α -syn knockout, administered with a competing peptide targeting ser129, and preconditioned with intermittent hypoxia against persistent hypoxic-induced cognitive impairment were compared. H–J. The levels of p- α -syn and α -syn in the hippocampi of mice in the Con, H14d, and H14d+IH groups were detected by Western blots with β -actin as the internal reference; statistical analysis was performed. K. The high-power three-dimensional imaging of p- α -syn immunofluorescence. Data are expressed as the mean \pm SEM (one-way ANOVA), * P <0.05, ** P <0.01, *** P <0.001, **** P <0.0001, n = 10 in B–G, n = 3 in H–K.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Supplementalfigures.docx](#)