

# Do Assessments of Cardiorespiratory and Muscular Fitness Influence Subsequent Reported Physical Activity? A Randomized Controlled Trial.

James T Langland (✉ [langl039@umn.edu](mailto:langl039@umn.edu))

University of Minnesota Medical Center: University of Minnesota Health <https://orcid.org/0000-0001-5188-3331>

Neeraj Sathnur

University of Minnesota Medical Center Fairview: University of Minnesota Health

Qi Wang

University of Minnesota Twin Cities Campus: University of Minnesota

Andrew PJ Olson

University of Minnesota Medical Center: University of Minnesota Health

---

## Research article

**Keywords:** Cardiorespiratory fitness, grip strength, physical activity, exercise

**Posted Date:** January 15th, 2021

**DOI:** <https://doi.org/10.21203/rs.3.rs-143519/v1>

**License:** © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

---

# Abstract

## Background

Clinicians strive to motivate healthy lifestyle behaviors such as exercise. Cardiorespiratory fitness (CRF) and muscular strength are important for good health and their measurement could potentially provide motivation to exercise. We investigated if measurements of CRF and muscular fitness (MF) would influence subsequent self-reported physical activity.

## Methods

Volunteer subjects at a State Fair were randomized in 1:1 parallel fashion to control and intervention groups. The baseline Exercise Vital Sign (EVS) and type of physical activity were obtained from all subjects. The intervention group received estimated maximum oxygen uptake ( $VO_2\text{max}$ ) using a step test and muscular strength using a hand grip dynamometer with age-specific norms for both measurements. All subjects were provided exercise recommendations and follow up surveys during the following year regarding their EVS and physical activity.

## Results

1315 individuals (656 intervention, 659 control) were randomized with one year follow up data obtained from 823 subjects (62.5%). Baseline mean EVS was 213 min/week. No change in EVS was found either group at follow-up ( $p=0.99$ ). Subjects who were less active at baseline ( $EVS < 150$ ) did show an increase in EVS (86 to 146) at 6 months ( $p < 0.05$ ). At 3 months the intervention group increased resistance training (29.1% to 42.8%) compared to controls (26.3% to 31.4%) ( $p < 0.05$ ). Lifestyle physical activity increased in the intervention group at 3 months (27.7% to 29.1%) and 6 months (25%) whereas it declined in the control group at 3 months (24.4% to 20.1%) and 6 months (18.7%) ( $p < 0.05$ ).

## Conclusion

Providing  $VO_2\text{max}$  estimates and grip strength did not produce an increase in overall physical activity. The EVS and exercise recommendations did however result in increased physical activity in less active individuals. In a very active population the  $VO_2\text{max}$  estimate and measured grip strength did increase lifestyle activity and resistance training. Wider adoption of these measures could be effective in promoting physical activity and resistance training.

## Trial Registration Number

clinicaltrials.gov NCT03518931 Registered 05/08/2018 -retrospectively registered

<https://clinicaltrials.gov/ct2/results?cond=&term=NCT03518931&cntry=&state=&city=&dist=>

## Background

Physical activity and regular exercise are important components of a healthy lifestyle. The benefits of physical activity and exercise include lower all-cause mortality, [1][2] reduced cardiovascular disease (CVD) [3], improved blood pressure[4], improved lipids[5], less depression[6], less anxiety[7] and improved cognitive function[8]. Despite the benefits of a regular exercise program, many individuals do not maintain sufficient physical activity and an increasing portion of the adult population engage in no leisure time physical activity[9]. Developing techniques to help motivate individuals to be more physically active can have important public health benefits.

The benefits of exercise are mediated in large part through cardiorespiratory fitness (CRF) and muscular fitness (MF). Although there is a strong genetic component to CRF it can be improved with regular exercise and physical activity[10]. It is well established that increased CRF is associated with better functional ability, improved cardiovascular health and reduced total mortality[11]. Individual measurements of CRF are a more powerful predictor of mortality than more traditional cardiovascular risk factors[12]. CRF is best measured by maximum oxygen uptake ( $VO_2\text{max}$ ) representing one's maximum ability to deliver and consume oxygen during activity. Despite the importance and prognostic significance of CRF and  $VO_2\text{max}$  they are not routinely

measured or estimated in the general population or the clinical setting. Measuring  $VO_2$ max typically involves a maximal exercise session which limits its availability and utility. A self-paced step test has been validated in a primary care setting as a safe and simple method of approximating  $VO_2$ max and CRF[13].

MF and strength also mediate the beneficial effects of exercise. Increased muscular strength is strongly associated with a lower all-cause and CVD mortality[14]. The reduction in all-cause mortality associated with muscular strength has been found to be independent of CRF[15]. Increased muscular strength improves metabolic and cardiovascular risk markers[16] and reduces the risk of developing metabolic syndrome[17]. Exercise, especially resistance training, can increase muscular strength at any age[18][19]. Although MF and strength have important metabolic and health implications they also are not routinely assessed in the general population or in the health care setting. A handgrip dynamometer has been shown to be a cost-effective clinical marker of sarcopenia and to correlate with lower extremity muscle power and poor mobility[20]. A large longitudinal population study found that measurement of grip strength is a simple, inexpensive risk-stratifying method for all-cause death, cardiovascular death, and CVD[21].

Given the major health implications of CRF and MF we hypothesized that providing individuals with an estimate of their CRF and  $VO_2$ max via the step test and MF via grip strength would be effective in motivating increased exercise activity as measured by the Exercise Vital Sign (EVS)[22]. We also secondarily hypothesized that this increase would be through increased cardio and resistance exercise.

## Methods

### Study population:

The study was undertaken at the Minnesota State Fair over 12 days in late August and early September 2014 and 2015. Subjects were self-selected when they volunteered to participate while exploring the University of Minnesota's Driven to Discover Building on the Minnesota State Fairgrounds. Exclusion criteria were age < 18 years, pregnancy, history of heart disease, syncope, chest pain, dyspnea, beta blocker or non-dihydropyridine calcium channel blockers use or evidence of any unstable medical condition. We estimated that we needed to enroll 440 individuals (220 per group) to detect an increase of 15% in EVS with 95% confidence level.

### Enrollment and testing protocol:

Subjects were randomized in a parallel fashion with 1:1 allocation via sealed envelope to either the control or the intervention group. All subjects provided signed written consent and had self-reported weight and height recorded and heart rate and blood pressure measured. The EVS (Product of "On average, how many days per week do you engage in moderate to strenuous exercise like a brisk walk" and "On average how many minutes do you engage in exercise at this level") was recorded for all subjects along with the types of physical activity they performed (none, cardio, sports, resistance exercises, lifestyle activities, balance/flexibility exercises and other). We capped the maximum EVS at 840 by limiting reported exercise time to 120 minutes/day.

All subjects in both groups were provided information on recommended physical activity based on the American College of Sports Medicine Guidance for Prescribing Exercise[23]. Those randomized to the control group did not undergo further testing.

The intervention group had grip strength measured in the dominant arm with a J00105 hand-grip dynamometer (Lafayette Instrument).  $VO_2$ max was estimated using a timed 20 step protocol at "normal" speed as per previously published protocols[13]. No problems or side effects were encountered during the measurements. The measured grip strength and the calculated  $VO_2$ max were provided to subjects along with age specific norms for both the grip strength[24] and  $VO_2$ max[25]. In 2014 participants were given an age specific "good" norm  $VO_2$ max and in 2015 were provided a "superior" norm for  $VO_2$ max to determine if increasing the normative value had a greater impact on subsequent exercise behavior[25].

### Follow-up data collection and analysis:

All study participants in both the control and intervention groups were contacted via email at 3, 6 and 12 months following enrollment to determine their current EVS and type of physical activity after which the trial was stopped and no further contacts were attempted. The change in EVS from baseline for both the intervention and control group was compared at 3, 6, and 12 months. Pre-specified subgroup analysis was also performed based on baseline physical activity levels. Participant baseline characteristics were summarized by group using descriptive statistics. Types of exercise were compared between groups using the Chi-square test. For participants who provided data at all 4 time points, the change in EVS over time were evaluated using linear mixed models. Models included fixed effect of intervention, time, and their interaction, and accounted for correlations among repeated measures. Tukey method was used to adjust for multiple comparisons. Subgroup analysis was conducted to examine change in EVS based on baseline physical activity levels. Analysis was performed using Statistical Analysis Software (version 9.3, SAS Institute Inc., Cary, NC).

## Results

### Study population and baseline measurement:

Total study enrollment was 1315 subjects (776 in 2014 and 539 in 2015) with 659 subjects assigned to the control group and 656 subjects to the intervention group. Mean age was 46.0 (range 18–92), males accounted for 63.0% of the total, and 92.6% of all participants were white. Table 1 shows the demographic characteristics and baseline EVS of the control and intervention groups.

Table 1  
Baseline demographics:

	Overall	Control	Intervention
Number	1315	659	656
Mean Age (range)	46.0(18–92)	44.9 (18–92)	47.1 (18-88.5)
Males	829 (63.0%)	418 (63.2%)	411 (62.3%)
Whites	1218 (92.6%)	607 (92%)	611 (92.6%)
Mean BMI (range)	25.8 (16.0-61.2)	26.0 (16.0-61.2)	25.6 (16.0-56.6)
Mean EVS	213.2 (SD 176.6)	211.7 (SD 168.7)	214.6 (SD 184.1)
BMI = Body mass index EVS = Exercise Vital Sign SD = standard deviation			

The mean EVS at baseline was 213.2 min/week (range 0-840, SD 183.6) with a median of 180 min/week. Nearly all (97.4%) participants reported some regular exercise activity. The reported types of exercise activity performed at baseline and at follow up is reported in Table 2.

Table 2  
Type of Exercise reported by subjects at Baseline and follow-up:

	Baseline Control	Baseline Intervention	3 mo Control	3 mo Intervention	6 mo Control	6 mo Intervention	12 mo Control	12 mo Intervention
Cardio	591 (89.4%)	584 (88.5%)	220 (83.3%)	250 (87.7%)	303 (88.6%)	300 (87.2%)	183 (91.0%)	199 (90.5%)
Sports	100 (15.1%)	109 (16.5%)	39 (14.8%)	33 (11.6%)	58 (17%)	52 (15.1%)	33 (16.4%)	31 (13.1%)
Lifestyle activities	162 (24.5%)	183 (27.7%)	53 (20.1%)	83 (29.1%)*	64 (18.7%)	86 (25.0%)*	58 (28.9%)	82 (37.3%)
Resistance training	174 (26.3%)	192 (29.1%)	83 (31.4%)	122 (42.8%)*	127 (37.1%)	147 (42.7%)	68 (33.8%)	87 (39.5%)
Balance/Flexibility	137 (20.7%)	134 (20.3%)	81 (30.7%)	76 (26.7%)	96 (28.1%)	108 (31.4%)	62 (30.8%)	66 (30.0%)
Other	42 (6.4%)	40 (6.1%)	26 (9.8%)	34 (11.9%)	44 (12.9%)	36 (10.5%)	17 (8.5%)	16 (7.3%)
*Indicates $p < .05$ for difference in change from baseline between study groups (Intervention vs. Control)								

The Intervention group's estimated  $VO_{2max}$  ranged from 11.2 to 77.3 ml/kg/min with a mean of 41.7 (SD 12.1). Maximum grip strength ranged from 10.9–83.2 Kg with a mean of 35.9 Kg (SD 11.9).

#### Follow-up EVS:

During the one year follow-up responses were received from 823 subjects (62.6%) with 262 subjects (20%) responding to all 3 contacts. No additional attempts were made to contact participants if they did not respond to our e-mail. Table 3 shows the follow up EVS values. There was no significant change in the EVS from baseline found in either the control or intervention group ( $p = 0.99$ ).

Table 3  
Change in EVS over time:

	Baseline EVS (N)	3 mo EVS (N)	6 mo mean EVS(N)	12 mo mean EVS (N)
Control	212 min/week (656)	193 min/week (263)	216 min/week (341)	221 min/week (200)
Intervention	215 min/week (659)	215 min/week (283)	219 min/week (343)	250 min/week (219)
N = number of responses, EVS = Exercise Vital Sign				

Among the 262 subjects from both groups who provided data at all 3 follow up intervals 82 had a baseline EVS less than the recommended 150 minutes/week. This group, which includes both control and intervention subjects, and a baseline EVS < 150 exhibited an increase over baseline in mean EVS at each follow up point, reaching statistical significance ( $p < 0.05$ ) at 6 months (Fig. 1). The group meeting or exceeding current recommendations with EVS > 150 at baseline (N = 180) showed no significant change.

#### Follow-up Exercise Type:

Although the follow up EVS showed no significant differences in the total amount of physical activity, there were significant differences in the type of exercise activity performed (Table 2). At 3 months and 6 months the intervention group exhibited significantly increased rates of lifestyle activity ( $p < 0.05$ ). The intervention group also significantly increased resistance exercise at 3 months ( $p < 0.05$ ). This increase in resistance exercise was driven by subjects whose baseline grip strength was less than the norm as they exhibited an increase in resistance training ( $p < 0.05$ ) throughout the one year follow up. Subjects with grip strength at or above norm exhibited a significant increase in resistance training at only 6 months (Table 4).

Table 4  
Subject Reporting Resistance Training related to Grip Strength norms:

	Baseline	3 mo FU	6 mo FU	12 mo FU
Baseline grip strength < norm (N = 69)	22(32%)	36* (52%)	34* (49%)	37* (54%)
Baseline grip strength $\geq$ norm (N = 67)	24(36%)	31 (46%)	34* (51%)	28 (42%)
*indicates $p < 0.05$ for change from baseline, N = number respondents				

### Effect of fitness measurements on subsequent EVS:

No significant influence on the subsequent reported EVS was found in those with a below norm estimated  $VO_{2max}$  ( $p = 1$ ) or grip strength ( $p = 0.75$ ) as seen in Table 5. Providing a higher normative value for  $VO_{2max}$  also had no effect on subsequent reported EVS ( $p = 0.62$ ).

Table 5  
Follow up EVS relative to normative data:

	Baseline (N)	3 mo FU (N)	6 mo FU (N)	12 mo FU (N)
Baseline $VO_{2max}$ < norm (N = 42)	193 (153)	206 (175)	221 (176)	196 (181)
Baseline $VO_{2max}$ $\geq$ norm (N = 100)	272 (194)	244 (183)	257 (179)	261 (191)
Baseline grip strength < norm (N = 69)	264 (203)	224 (175)	246 (211)	235 (202)
Baseline grip strength $\geq$ norm (N = 67)	224 (157)	263 (202)	252 (143)	252 (185)
EVS = Exercise Vital Sign, N = number of respondents				

## Discussion

We did not find that estimates of CRF and MF increased total physical activity over the subsequent year as measured by the EVS. We did find a significant increase in both lifestyle exercise and resistance type exercise in an already very physically active population. Having a below norm grip strength was associated with a significant increase in resistance training throughout the subsequent year. We found that individuals not meeting current recommended physical activity recommendations (EVS < 150 minutes/week) showed a significant increase in their EVS at 6 months of follow up. This study indicates that recording EVS, providing exercise recommendations and estimating CRF and MF could provide both a useful incentive to stimulate greater interest in exercise, lifestyle physical activity and resistance training. To our knowledge no other study has investigated the effect on subsequent physical activity of  $VO_{2max}$  estimates or grip strength.

Recognizing the importance of exercise and physical activity to good health, the Surgeon General and others have called for regular assessments of an individual's physical activity[26][27]. The National Physical Activity Plan asks healthcare systems to prioritize physical activity assessment, advice, and promotion and regularly assess physical activity as a "vital sign" [28]. The EVS has been advocated as a tool to help accomplish this goal[22]. We used the EVS to quantify the exercise activity in our subjects. It is easily calculated with just two questions and corresponds to current exercise guidelines recommending 150

minutes of moderately vigorous physical activity per week[23]. The EVS has been shown to under-report physical activity measured by accelerometer and may best be employed to identify individuals not meeting current physical activity guidelines[29]. Other limitations of using the EVS are the absence of a specific time frame and the inability to differentiate exercise intensity.

Our study population was done in random self-selected volunteers. These volunteers were already very physically active as indicated by the mean EVS of 223.2 with a median value of 180, significantly exceeding current exercise recommendations. The high level of pre-existing physical activity likely attenuated the impact of the fitness measurements on their future physical activity and limited the utility of the EVS as a measurement tool[29]. We did find, however, that the subjects not meeting current physical activity guidelines did exhibit a significant increase in their reported EVS at 6 months. Both the control and intervention groups exhibited this increase indicating that recording the EVS and providing information on current exercise recommendations likely influenced this change. This observation validates calls for recording exercise as a vital sign. It has previously been reported that systematically recording the EVS during outpatient visits was associated with significant changes in exercise-related clinical counseling and documentation (30).

In addition to obtaining the EVS we recorded the types of physical activity both initially and in follow up. We found that the intervention group significantly increased their reported resistance training and lifestyle physical activity relative to controls at 3-months follow-up, despite the much less favorable climate for these activities during winter. The increase in lifestyle activity was sustained at 6 months follow up. A significant increase in resistance training was observed throughout the following year in those individuals with a reported grip strength less than the reported norm. Below norm grip strength appears to stimulate interest in strength training activities in this already very active population. At baseline, 88% of our study population reported participating in some form of cardiovascular exercise but only 28% reported participation in resistance type exercise. This lower level of resistance training is consistent with prior surveys showing only 21.9% of Americans meet muscle strengthening guidelines[30]. This provides greater potential for our assessments and recommendations to have an impact on resistance exercise activity. We did not observe that those subjects with grip strength below the norm increased their EVS even though they did increase their resistance training.

Cardiorespiratory Fitness as measured by  $VO_2$ max is an important indicator of overall health and has significant prognostic implications[11][10][12]. Recognizing the significance of CRF the American Heart Association has called for the inclusion of CRF measurement or estimation in routine clinical practice[31]. Despite its importance it is not typically measured in a clinical encounter. This relates to the difficulty of formal  $VO_2$ max measurements. Other forms of estimating  $VO_2$ max such as maximal or sub-maximal treadmill or bicycle exercise testing are also not suited to routine use.  $VO_2$ max can easily be estimated by several formulas based on demographics and reported exercise habits [32][33]. Estimating CRF from one of these formulas has been associated with CVD and all-cause mortality independent of other risk factors[34]. The estimated CRF from formulas however are significantly influenced by the subjective reporting of exercise activity. We elected to use a step test that had been previously validated in a geriatric population[13]. This test in younger individuals and other populations has been found to be less accurate in the measured  $VO_2$ max yet still felt to be useful in classifying CRF[35]. When this step test has been used to measure CRF to aid in exercise prescription a significant improvement in  $VO_2$ max at 12 months was found compared with baseline measures[36].

We felt that providing a CRF estimate requiring actual physical activity using the step test would positively influence future exercise behavior relative to no measurement. We did not observe significant changes in the EVS in either the control group or the intervention group over one year of follow up. The fitness assessments did not appear to influence this very active population's physical activity as measured by the EVS. We also did not observe an increase in follow up EVS in those individuals who were reported to have an estimated  $VO_2$ max below the provided norm even when the norm was increased from "good" to "superior".

We used a hand-grip dynamometer to estimate muscular strength. This test is inexpensive, convenient and previously demonstrated in multiple studies to be a clinically significant marker of sarcopenia and correlate with lower extremity muscle power and mobility (20). In a large longitudinal population study, measurement of grip strength was found to be a simple,

inexpensive risk-stratifying method for all-cause death, cardiovascular death, and CVD (21). Grip strength is predictive of mortality in both young adults[37] and middle age[38]. Low grip strength has been documented to correlate with increased disability in the elderly[39], greater risk for hospitalization[40], cognitive decline[41] and nutritional status[42]. We used grip strength to estimate MF and felt that this measurement would contribute to increased physical activity and resistance training. We did observe an increase in resistance training but not total exercise time. The increase in resistance exercise was largely driven by those individuals with grip strength below norm. Given the overall high level of baseline physical activity and lower level of resistance training at baseline it appears that having a below norm grip strength shifted physical activity to resistance training from other activities.

The strengths of this study are its size and diversity. The study participants exhibited a wide range of age (18–92 mean 46) and BMI (16–61, mean 25) but were predominately white (92.3%).

The study limitations were the self-selected population that was already very active as exhibited by the high baseline EVS. In this active group the EVS may have been less accurate in measuring their physical activity[29]. This active group may have been more receptive to feedback on their CRF and MF accounting for the short term significant increases in the lifestyle and resistance physical activity but with less potential to observe an increase in EVS over time. The results are also limited by follow up data being provided from only 62.5% of the study population. We also did not perform follow up CRF or MF measurements.

In summary we did not find that our measurements of CRF and MF using a step test and grip strength increased overall exercise or physical activity as measured by the EVS during the ensuing year. The utility of our intervention was likely limited by a self-selected very active population. Less active individuals in both the control and intervention groups (those not meeting current exercise guidelines) did significantly increase their reported exercise activity at 6 months. We found that the fitness measurements appeared to stimulate an increase in lifestyle and resistance training exercise at short term follow up and that the increase in resistance training was largely driven by those having a below norm grip strength. This indicates a potential benefit of recording the EVS and providing current exercise recommendations to less active individuals. Even very active individuals may benefit from measuring grip strength and providing norms to stimulate greater participation in resistance training activities. Given these encouraging improvements in exercise activity it may be useful to more widely record EVS and perform CRF and MF estimates.

## Conclusions

In a very active population providing  $VO_2$ max estimates and grip strength measurement did not produce an increase in overall physical activity however it did shift activity to increased lifestyle physical activity and resistance training. Recording the EVS and providing exercise recommendations did result in a significant increase in overall physical activity in those individuals not meeting current physical activity recommendations. Wider adoption of the EVS and grip strength measurement could be effective in promoting physical activity and resistance training.

## Abbreviations

CVD: Cardiovascular disease

CRF: Cardiorespiratory Fitness

$VO_2$ max: maximum oxygen uptake

MF: Muscular Fitness

EVS: Exercise Vital Sign

## Declarations

The study adheres to CONSORT guidelines and a completed CONSORT checklist has been submitted separately.

**Ethics approval and consent to participate:**

This study and the consent form were approved by the University of Minnesota Institutional Review Board. Signed written consent was obtained from all subjects.

**Consent for publication:**

Not applicable

**Availability of data and materials:**

University of Minnesota RedCap database ([https://redcap.ahc.umn.edu/redcap\\_v10.0.28/index.php?pid=2257&\\_\\_record\\_cache\\_complete=1](https://redcap.ahc.umn.edu/redcap_v10.0.28/index.php?pid=2257&__record_cache_complete=1))

**Competing interests:**

Not applicable

**Funding:**

Not applicable

**Authors' contributions:**

JTL, NS and APJO contributed equally to data collection and manuscript draft, JTL manuscript submission, QW statistical analysis.

**Acknowledgements:**

None


**References**

1. Arem H, Moore SC, Patel A et al. Leisure Time Physical Activity and Mortality. *JAMA Intern Med.* 2015;175: 959-967.
2. Wen CP, Wai JPM, Tsai MK, Chen CH. Minimal amount of exercise to prolong life: To walk, to run, or just mix it up? *J Am Coll Cardiol.* 2014;64: 482–484.
3. Shiroma EJ, Lee I-M. Physical activity and cardiovascular health: lessons learned from epidemiological studies across age, gender, and race/ethnicity. *Circulation.* 2010;122: 743–752.
4. Crump C, Sundquist J, Winkleby MA, Sundquist K. Interactive Effects of Physical Fitness and Body Mass Index on the Risk of Hypertension. *JAMA Intern Med.* 2016;176(2):210-216.
5. Iliam W, Raus EK, Oseph J et al. Effects of the Amount and Intensity of Exercise on Plasma Lipoproteins. *N Engl J Med.* 2002;347: 1483–1492.
6. Cooney GM, Dwan K, Greig CA et al. Exercise for depression ( Review ). *Cochrane Database of Systematic Reviews* 2013, Issue 9. Art. No.: CD004366.
7. Goodwin RD. Association between physical activity and mental disorders among adults in the United States. *Prev Med (Baltim).* 2003;36: 698–703.
8. Zhu W, Wadley VG, Howard VJ et al. Objectively Measured Physical Activity and Cognitive Function in Older Adults. *Med Sci Sports Exerc.* 2017 Jan;49(1):47-53.
9. Ladabaum U, Mannalithara A, Myer PA, Singh G. Obesity, abdominal obesity, physical activity, and caloric intake in US adults: 1988 to 2010. *Am J Med.* 2014;127: 717–727.
10. Lavie CJ, Arena R, Swift DL et al. Exercise and the cardiovascular system: Clinical science and cardiovascular outcomes. *Circ Res.* 2015;117: 207–219.

11. Kaminsky LA, Arena R, Beckie TM et al. The importance of cardiorespiratory fitness in the United States: The need for a national registry: A policy statement from the American Heart Association. *Circulation*. 2013;127: 652–662.
12. Gupta S, Rohatgi A, Ayers CR et al. Cardiorespiratory Fitness and Classification of Risk of Cardiovascular Disease Mortality *Circulation*. 2011;123:1377-1383
13. Petrella RJ, Koval JJ, Cunningham D a, Paterson DH. A self-paced step test to predict aerobic fitness in older adults in the primary care clinic. *J Am Geriatr Soc*. 2001;49: 632–638.
14. Kamiya K, Masuda T, Tanaka S et al. Quadriceps Strength as a Predictor of Mortality in Coronary Artery Disease. *Am J Med*. Elsevier Inc; 2015;128: 1212–1219.
15. Ruiz JR, Sui X, Lobelo F et al. Association between muscular strength and mortality in men: prospective cohort study. *BMJ*. 2008;337: a439
16. Roberts CK, Lee MM, Katiraie M et al. Strength fitness and body weight status on markers of cardiometabolic health. *Med Sci Sports Exerc*. 2015;47: 1211–1218.
17. Sénéchal M, McGavock JM, Church TS et al. Cut points of muscle strength associated with metabolic syndrome in men. *Med Sci Sports Exerc*. 2014;46: 1475–1481.
18. Harber MP, Konopka AR, Udem MK et al. Aerobic exercise training induces skeletal muscle hypertrophy and age-dependent adaptations in myofiber function in young and older men. *J Appl Physiol*. 2012;113: 1495–1504.
19. Peterson MD, Rhea MR, Sen A, Gordon PM. Resistance exercise for muscular strength in older adults: A meta-analysis. *Ageing Res Rev*. Elsevier B.V.; 2010;9: 226–237.
20. Lauretani F, Russo CR, Bandinelli S et al. Age-associated changes in skeletal muscles and their effect on mobility: an operational diagnosis of sarcopenia. *J Appl Physiol*. 2003;95: 1851–60.
21. Leong DP, Teo KK, Rangarajan S et al. Prognostic value of grip strength: Findings from the Prospective Urban Rural Epidemiology (PURE) study. *Lancet*. Elsevier Ltd; 2015;386: 266–273.
22. Coleman KJ, Ngor E, Reynolds K et al. Initial validation of an exercise “vital sign” in electronic medical records. *Med Sci Sports Exerc*. 2012;44: 2071–2076.
23. Garber CE, Blissmer B, Deschenes MR et al. Quantity and quality of exercise for developing and maintaining cardiorespiratory, musculoskeletal, and neuromotor fitness in apparently healthy adults: Guidance for prescribing exercise. *Med Sci Sports Exerc*. 2011;43: 1334–1359.
24. Mathiowetz V, Kashman N, Volland G et al. Grip and pinch strength: normative data for adults. *Arch Phys Med Rehabil*. 1985;66(2):69-74.
25. The Physical Fitness Specialist Certification Manual, The Cooper Institute for Aerobics Research, Dallas TX, revised 1997 printed in *Advance Fitness Assessment & Exercise Prescription*, 3rd Edition, Vivian H. Heyward, 1998.p48
26. Office of the Surgeon General (US). *Step It Up!: The Surgeon General’s Call to Action to Promote Walking and Walkable Communities*. Washington (DC): US Department of Health and Human Services; 2015.
27. Foster C, Hillsdon M, Thorogood M, Kaur A. Interventions for promoting physical activity. *Cochrane Database Syst Rev*. 2014;9: 1–90.
28. National Physical Activity Plan Alliance (2016). U.S. National Physical Activity Plan. Available at: [http://physicalactivityplan.org/docs/2016NPAP\\_Finalforwebsite.pdf](http://physicalactivityplan.org/docs/2016NPAP_Finalforwebsite.pdf)
29. Joseph RP, Keller C, Adams M a, Ainsworth BE. Validity of two brief physical activity questionnaires with accelerometers among African-American women. *Prim Health Care Res Dev*. 2015; 1–12.
30. Carlson SA, Fulton JE, Schoenborn CA, Loustalot F. Trend and prevalence estimates based on the 2008 physical activity guidelines for Americans. *Am J Prev Med*. Elsevier Inc.; 2010;39: 305–313.
31. Ross R, Blair SN, Arena R, Church TS et al. Importance of Assessing Cardiorespiratory Fitness in Clinical Practice: A Case for Fitness as a Clinical Vital Sign: A Scientific Statement From the American Heart Association. *Circulation*. 2016; Dec 13;134(24):e653-e699.

32. Nes BM, Janszky I, Vatten LJ, Nilsen TIL, Aspenes ST, Wisløff U. Estimating  $\dot{V}O_{2peak}$  from a nonexercise prediction model: The HUNT study, Norway. *Med Sci Sports Exerc.* 2011;43: 2024–2030.
33. O'Donovan G, Bakrania K, Ghouri N et al. Nonexercise equations to estimate fitness in White European and South Asian Men. *Med Sci Sports Exerc.* 2016;48: 854–859.
34. Nauman J, Nes BM, Lavie CJ et al. Prediction of Cardiovascular Mortality by Estimated Cardiorespiratory Fitness Independent of Traditional Risk Factors: The HUNT Study. *Mayo Clin Proc.* 2016;92: 1–10.
35. Vidoni ED, Mattlage A, Mahnken J, Burns JM, McDonough J, Billinger SA. Validity of the step test for exercise prescription: No extension to a larger age range. *J Aging Phys Act.* 2013;21: 444–454.
36. Petrella RJ, Lattanzio CN, Shapiro S, Overend T. Improving aerobic fitness in older adults: Effects of a physician-based exercise counseling and prescription program. *Can Fam Physician.* 2010;56:e191-200
37. Ortega FB, Silventoinen K, Tynelius P, Rasmussen F. Muscular strength in male adolescents and premature death: cohort study of one million participants. *BMJ.* 2012;345: e7279.
38. Rantanen T, Masaki K, He Q, Ross GW, Willcox BJ, White L. Midlife muscle strength and human longevity up to age 100 years: A 44-year prospective study among a decedent cohort. *Age (Omaha).* 2012;34: 563–570.
39. Giampaoli S, Ferrucci L, Cecchi F et al. Hand-grip strength predicts incident disability in non-disabled older men. *Age Ageing.* 1999;28: 283–288.
40. Cawthon PM, Fox KM, Gandra SR et al. Do muscle mass, muscle density, strength, and physical function similarly influence risk of hospitalization in older adults? *J Am Geriatr Soc.* 2009;57: 1411–1419.
41. Auyeung TW, Lee JSW, Kwok T, Woo J. Physical frailty predicts future cognitive decline - A four-year prospective study in 2737 cognitively normal older adults. *J Nutr Heal Aging.* 2011;15: 690–694.
42. Norman K, Stobäus N, Gonzalez MC, Schulzke JD, Pirlich M. Hand grip strength: Outcome predictor and marker of nutritional status. *Clin Nutr. Elsevier Ltd;* 2011;30: 135–142.

## Figures



**Figure 1**

Change in EVS for baseline physical activity above and below current guidelines. Legend Figure 1: One year change in Exercise Vital Sign (EVS) for those with baseline below current exercise recommendations of 150 minutes/week (blue) and those

exceeding 150 minutes/week (orange) in combined control and intervention groups. \* signifies  $p < 0.05$  relative to baseline

## Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [CONSORT2010Checklist.Langland.doc](#)