

Estimated frequency of somatic symptom disorder in general practice: cross-sectional survey with general practitioners

Marco Lehmann (✉ ma.lehmann@uke.de)

University Medical Center Hamburg-Eppendorf

Nadine Janis Pohontsch

University Medical Center Hamburg-Eppendorf

Thomas Zimmermann

University Medical Center Hamburg-Eppendorf

Martin Scherer

University Medical Center Hamburg-Eppendorf

Bernd Löwe

University Medical Center Hamburg-Eppendorf

Research Article

Keywords: Persistent somatic symptoms, somatoform disorders, primary health care, general practice, somatic symptom disorder, bodily distress, General Practitioner

Posted Date: April 19th, 2022

DOI: <https://doi.org/10.21203/rs.3.rs-1452786/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Abstract

Background

Somatic symptom disorder (SSD) is the successor diagnosis of somatoform disorder in the 5th Edition of the Diagnostic and Statistical Manual of Mental Disorders (DSM-5). Relevance and frequency of SSD and its clinical symptoms in general practice are still unknown. We estimate frequencies of patients fulfilling the diagnostic criteria of SSD in general practice.

Methods

Mailed and online survey with general practitioners (GP) in Germany using a cross-sectional representative sample from registries of statutory health insurance physicians. GPs estimated percentages of their patients who show the clinical symptoms of SSD according to DSM-5; that is, one or more burdensome somatic symptoms (A criterion), excessive symptom- or illness-related concern, anxiety, or behaviour (B criterion), and persistence of the symptoms over at least 6 months (C criterion). Statistical analysis used means and confidence intervals of estimated patient proportions showing SSD symptoms. Frequency of full-blown SSD was based on the products of these proportions calculated for each GP.

Results

Responses from 1728 GPs were obtained. GPs saw the clinical symptoms of SSD fulfilled (A and B criteria) in 21.5% (95% CI: 20.6 to 22.3) of their patients. They further estimated that in 24.3% (95% CI: 23.3 to 25.2) of patients, symptoms would persist, yielding a total of 7.7% (95% CI: 7.1 to 8.4) of patients to have a full-blown SSD.

Conclusions

GPs see the clinical symptoms of SSD fulfilled in a considerable patient proportion. This indicates a high clinical relevance of the clinical symptoms of SSD in general practice.

Study registration

German Clinical Trials Register (Deutschen Register Klinischer Studien, DRKS)

https://www.drks.de/drks_web/setLocale_EN.do

The date the study was registered: October 2nd 2017

The date the first participant was enrolled: February 9th 2018

DRKS-ID: DRKS00012942

Background

With the introduction of the 5th Edition of the Diagnostic and Statistical Manual of Mental Disorders (DSM-5) in 2013 (1), the diagnosis of somatoform disorder was replaced by somatic symptom disorder (SSD). The clinical symptoms of SSD combine both, physical and psychological symptom burden. That is, besides the A criterion of at least one life-disrupting physical symptom, SSD requires excessive symptom- or illness-related concern, anxiety, or behaviour as it's B criterion – the so-called psychological positive criteria. Finally, SSD assumes symptom

persistence of at least six months as its C criterion. SSD no longer requires the exclusion of a medical cause for physical complaints. The ICD-11 made a similar development from the group of somatoform disorders in ICD-10 to the diagnosis of bodily distress disorder, which is comparable with SSD (2).

Representative and unbiased prevalence studies for SSD are still rare as evidenced in a recent scoping review (3). In a community health centre one study, using a clinical interview to diagnose SSD with a rather small sample of 202 participants, revealed 20.8% of patients diagnosed with somatic symptom disorder (4). Prevalence estimation using a combination of PHQ-15, Whiteley-Index, and BIPQ as a proxy measure for SSD revealed a proportion of 45.5% of SSD in a selected sample of 325 patients with medically unexplained physical symptoms in general practice (5). More gold standard prevalence data is still missing and there is no knowledge about what proportions of patients in clinical practice show symptoms of SSD.

With SSD also came the scholarly debate of the harms and benefits of this new diagnosis in comparison with its predecessor (6). Some argue that SSD was an overdue invention to end dualistic thinking of mind and body in clinical practice (7, 8). Furthermore, clinicians themselves said that they could not assess the former criterion of exclusion of physical disease with certainty anyway (9). Overall, regarding reliability, validity, and clinical usefulness SSD seems to surpass its predecessor (3, 7, 10). Nonetheless, it is unclear of what it means for general practice. It was designed to be viable to implement in general practice (11) and for general practitioners (GPs), who encounter many patients with persistent somatic symptoms of unclear origin. It is uncertain whether GPs evaluate the introduction of SSD as useful and in which proportion of their patients they see the clinical symptoms of SSD. Furthermore, compared with its predecessor somatoform disorder, it is unclear whether GPs see continuity in the diagnoses as they apply to patients showing persistent somatic symptoms. Answering these empirical questions would help to form reasonable expectations about the relevance of SSD in general practice. It would also help to plan for the necessary resources for those in need.

Our research questions were:

(1.) What are the estimated proportions of patients showing the clinical symptoms of somatic symptom disorder and full-blown somatic symptom disorder as viewed by general practitioners?

(2.) Which variables are associated with the estimated proportions of patients showing clinical symptoms of somatic symptom disorder (i. e., proportion of patients with somatoform disorders, characteristics of the general practitioners and the practice setting)?

Methods

We invited a representative sample of GPs to participate in a survey regarding the proportions of patients with somatic symptom disorder in general practice. This study was part of the project '*Identification of barriers and difficulties involved in the process of diagnosing somatic symptom disorders in primary care (BeSSD-GP)*', which was funded by the German Research Foundation (Deutsche Forschungsgemeinschaft, DFG). Details of the project were published in the study protocol (12); and the study method of this cross-sectional representative and anonymous survey is further described elsewhere (13). We administered a questionnaire with GPs in eight federal states of Germany from February 9th to May 15th, 2018 (Registration: October 2nd, 2017, German Clinical Trials Register (DRKS00012942)).

Participants

GPs were randomly sampled (n = 12004) from all working GPs from the German federal states of Schleswig-Holstein, Hamburg, North Rhine, Saarland, Brandenburg, Saxony, Mecklenburg-Vorpommern, and Thuringia. We sampled via publicly accessible contact information through registries of statutory health insurance physicians proportionally to the total number of general practitioners working in the respective federal states. Eligible were physicians working in general practice. Participation was voluntary and anonymous. To comply with common research practice in Germany, we offered no incentive.

Survey

GPs assessed the following items:

- To what extent they evaluate the introduction of SSD as meaningful
- The frequency of occurrence (as percentage of all patients) of fulfilled diagnostic criteria
 - A: One or more somatic symptoms that are distressing or result in significant disruption of daily life.
 - B1: Disproportionate and persistent thoughts about the seriousness of one's symptoms.
 - B2: Persistently high level of anxiety about health or symptoms.
 - B3: Excessive time and energy devoted to these symptoms or health concerns.
 - C: Although any one somatic symptom may not be continuously present, the state of being symptomatic is persistent (typically more than 6 months).
 - Combinations of A- and B-criteria of SSD in their practice.
- Person characteristics
 - Gender
 - Years of practice
- Practice characteristics
 - Practice setting: own practice or collaborative practice
 - Practice region: rural, small town, medium town, large town
 - Number of patients per quarter
- Estimated proportion of patients with DSM-IV-TR somatoform disorders (%) (14).

We captured the subjective meaningfulness of the SSD innovation with a six-point Likert rating scale between *1: does not apply* to *6: applies completely*. We presented the items regarding the subjective frequency of diagnostic categories as Likert rating scales with eleven response options ranging from 0% in steps of 10–100%. Person and practice characteristics used categorical ratings. Years of practice and estimated proportion of patients with somatoform disorder asked for provision of a single number.

Data acquisition

We invited GPs to participate by mail or online. They received a post card announcement and two consecutive mailings of the survey package. Data acquisition commenced February 9th, 2018 with the announcement one week prior the first mailing of the survey, with the second mailing following two weeks later. Data collection was closed on May 15th, 2018. There was also an option for online participation. The invitation to participate consisted of a data safety statement, assurance of anonymous data analysis, and statement that participation was voluntary. Due to the completely anonymous data collection, no written informed consent was necessary in accordance with

German law. GPs were notified that informed consent would be presumed if they returned the completed response form.

Statistical analysis

All analyses were conducted in R (Version 3.6.2). For this data the analysis of responders and non-responders and the comparison of the sample characteristics with official statistics is provided elsewhere (13). We analysed the item of how the GPs appraised the meaning of the diagnostic innovation and the six items regarding the clinical symptoms of SSD in general practice and present means and 95% confidence intervals for these items. An estimate of the frequency of SSD in general practice was calculated using a derived variable, which was the product of the proportions, calculated for each GP, of patients showing the clinical symptoms of SSD (A and B criteria) and the proportion of patients showing symptom persistence (C criterion). Furthermore, we used ordinal regression to model the estimated proportions of patients showing the clinical symptoms of SSD. The regressions used person and practice characteristics and the estimated proportion of patients with DSM-IV-TR somatoform disorders as independent variables. Regressions were calculated with the *ordinal* R package (15). All tests for significance were two-tailed. Multicollinearity between predictors was checked using variance inflation factor from the *car* R package (16).

Results

We obtained responses from 1829 GPs (15.2%). Reasons for non-participation were retirement, death of the GP, comments that the survey method would be inappropriate, and non-specified reasons. After removal of cases with double participation and cases with empty person characteristics, we analysed the data of 1728 general practitioners. The participant and practice characteristics shown in Table 1.

Table 1
Participant characteristics.

	Category	Sample
n		1728
Gender (%)	male	760 (44.8)
	female	938 (55.2)
Years of practice (%)	0–10	455 (27.4)
	11–20	540 (32.5)
	21–30	454 (27.3)
	> 30	211 (12.7)
Practice setting (%)	Own practice	837 (50.1)
	Collaborative practice	834 (49.9)
Practice region (%)	Rural (max. 4.999 Inh.)	321 (19.6)
	Small town (5.000 to 19.999 Inh.)	369 (22.5)
	Medium town (20.000 to 99.999 Inh.)	351 (21.4)
	Large town (above 100.000 Inh.)	600 (36.6)
Number of patients per quarter (%)	< 500	53 (3.4)
	500–1000	654 (42.4)
	1001–1500	647 (41.9)
	> 1500	190 (12.3)
Estimated proportion of patients with DSM-IV-TR* somatoform disorders (%)	0%	210 (15.2)
	10%	524 (37.8)
	20%	324 (23.4)

	Category	Sample
	30%	207 (14.9)
	> 30%	120 (8.7)
*DSM-IV-TR = 4th edition, text revision, Diagnostic and Statistical Manual of Mental Disorders		

The respondents in large part found the replacement of somatoform disorder by somatic symptom disorder meaningful ($M = 4.29$, 95% CI: 4.24 to 4.35). Figure 1 shows the mean proportions of the estimated proportions for the clinical symptoms of SSD. The GPs estimated that 21.5% (95% CI: 20.6 to 22.3) of their patients would show the clinical symptoms of SSD (A and B criteria). In 24.3% (95% CI: 23.3 to 25.2) of their patients, the GPs indicated symptom persistence (C criterion). Regarding the individual diagnostic criteria, the GPs indicated that 33.7% (95% CI: 32.7 to 34.7) of patients would show one or more distressing and significantly impairing somatic symptoms (A criterion). Furthermore, they estimated excessive levels of symptom related thoughts, anxiety, and behaviour in 24.7% (95% CI: 23.8 to 25.6), 25.2% (95% CI: 24.3 to 26.2), and 19.7% (95% CI: 19.0 to 20.6) of their patients, respectively (B criteria). The new variable, consisting of the case wise product of the estimated proportion of patients with clinical symptoms of SSD (A and B criteria), and the estimated proportion of patients with symptoms persisting for more than six months (C criterion), yielded a mean frequency estimate of 7.7% (95% CI: 7.1 to 8.4) of patients in general practice with a diagnosis of full-blown SSD.

For the ordinal regression analyses, to obtain enough counts in all outcome categories, all estimated proportions of diagnostic criteria higher than 40% were aggregated into one category. Variance inflation factors for all predictors in all analyses indicated that multicollinearity was not problematic ($VIF < 1.1$). The regression analyses indicated that the proportion of patients showing a somatoform disorder was the only significant predictor for the proportions of patients showing the clinical symptoms of SSD (Table 2). All other person and practice characteristics were not significantly associated with clinical symptoms of SSD. Significant spearman rank correlations ($r_s = 0.36$ to 0.41 , $p < 0.001$) confirmed that higher estimated proportions of patients showing SSD symptoms were associated with higher estimated proportions of patients showing a somatoform disorder. We further recalculated the analyses as standard linear regressions with quantitative outcomes and the ordinal regressions with data imputations to include higher sample sizes. Both auxiliary analyses yielded the same pattern of results as the initial ordinal regression.

Table 2

Regression coefficients and standard errors of ordinal regression of SSD diagnostic criteria regarding participant characteristics, practice characteristics, and the estimated proportion of patients with somatoform disorder.

Clinical symptoms of somatic symptom disorder						
	Criteria A and B both apply	One or more somatic symptoms (A)	Persistent thoughts about symptoms (B1)	Persistent anxiety about symptoms (B2)	Excessive time and energy spent for symptoms (B3)	Symptom persistence (C)
	B (SE)	B (SE)	B (SE)	B (SE)	B (SE)	B (SE)
Gender female	-0.04 (0.11)	0.05 (0.11)	-0.08 (0.11)	-0.12 (0.11)	-0.03 (0.11)	0.07 (0.11)
11–20 years of practice	-0.15 (0.15)	-0.17 (0.14)	-0.38** (0.14)	-0.14 (0.14)	-0.26 (0.15)	-0.24 (0.14)
21–30 years of practice	-0.11 (0.16)	-0.18 (0.15)	-0.22 (0.15)	0.08 (0.15)	-0.20 (0.16)	-0.29 (0.15)
> 30 years of practice	-0.15 (0.19)	-0.45* (0.18)	-0.35 (0.18)	0.06 (0.19)	-0.21 (0.19)	-0.36 (0.19)
Collaborative practice	-0.09 (0.11)	0.10 (0.11)	0.07 (0.11)	-0.03 (0.11)	-0.15 (0.12)	0.05 (0.11)
Small town (5.000 to 19.999 Inh.)	-0.35* (0.17)	-0.11 (0.17)	-0.07 (0.17)	-0.31 (0.17)	-0.15 (0.17)	-0.16 (0.17)
Medium town (20.000 to 99.999 Inh.)	0.01 (0.17)	0.02 (0.17)	-0.15 (0.17)	-0.16 (0.17)	-0.04 (0.18)	0.08 (0.17)
Large town (above 100.000 Inh.)	-0.10 (0.16)	0.07 (0.16)	-0.04 (0.16)	0.05 (0.16)	0.07 (0.16)	0.08 (0.16)
More than 1000 patients per quarter	0.03 (0.11)	-0.14 (0.11)	0.14 (0.11)	0.22* (0.11)	-0.05 (0.12)	-0.09 (0.11)
Proportion of patients showing a DSM-IV-TR somatoform disorder	2.33*** (0.17)	1.89*** (0.15)	1.91*** (0.16)	2.01*** (0.16)	1.87*** (0.16)	1.96*** (0.16)
Observations	1,131	1,129	1,130	1,130	1,125	1,129
Log Likelihood	-1,579.15	-1,746.65	-1,671.43	-1,673.78	-1,564.65	-1,701.27
<i>Note:</i>	*p < 0.05; **p < 0.01; ***p < 0.001					

Clinical symptoms of somatic symptom disorder

Predictor reference categories: Gender male, 0–10 years of practice, own practice, rural practice region (max. 4.999 Inh.), 0-1000 patients per quarter.

Discussion

Summary

GPs in our study had estimated that about one fifth of their patients show clinical symptoms of SSD – A and B criterion. Moreover, symptom persistence of more than six months was reported for 24.3% of patients – C criterion. The assessments for all diagnostic criteria combined, indicated that a frequency of 7.7% patients in general practice were likely to possess full-blown SSD. Furthermore, GPs estimated that 34% of their patients would show one or more distressing symptoms for the diagnostic A criterion of SSD, and between 20% and 25% of their patients would fulfil any one of the B criteria. The estimated proportion of patients showing a somatoform disorder (according to DSM-IV-TR) best predicted the proportions of clinical symptoms of SSD.

Strengths and limitations

Our study is limited by the still unsatisfactory dissemination of the SSD concept in clinical practice in Germany. In the questionnaire, we introduced the concept of SSD in a short paragraph, so that all participants had at least minimal information about the diagnosis. Furthermore, we used literal translations of the SSD criteria, which may also be intelligible without extra information. However, it was possible that in our questionnaire some participants learned about SSD for the first time. Our study is further limited by the use of subjective estimations regarding patient proportions by the GPs. We did not ask them to search through their actual clinical records for SSD cases, because this would have limited our sample to those patients who were formally diagnosed with SSD.

Comparison with existing literature

Although, our data does not consist of prevalence rates estimated based on standard clinical interviews, our frequency estimation of 7.7% can be compared with estimations from other research. A small sample study in general practice yielded 20.8% as prevalence for SSD (4). In the general population, the prevalence of SSD was estimated as 4.5% (17). About 20% of the general population were at least sometimes concerned about bodily symptoms, had higher levels of anxiety related to symptoms, or showed higher levels of symptom-related behaviour (18). In conclusion, our estimation of SSD frequency is lower than earlier data from general practice, but corresponds with estimations for the general population of about one out of twenty persons to show SSD. Clinicians should be alert for the clinical symptoms of SSD in daily practice.

Conclusions

We found the strongest association between the estimated patient proportions of those showing the clinical symptoms of SSD and those with somatoform disorder. This suggests that GPs may treat the diagnostic criteria of SSD as a continuation of somatoform disorder as it applies to patients showing persistent somatic symptoms and symptom-related concern. So, the results may be due to the actual overlap of both diagnoses in patients in general practice.

It is clear that valid data about the prevalence of SSD in general practice requires representative sampling and a diagnosis based on a clinical interview as gold standard. Until we have these numbers, we have to be content with hopefully converging evidence from different sources, for example, from proxy diagnoses via psychometric tests or from non-random and convenience samples. In this actual state of the evidence, our estimations by GPs about how often the clinical symptoms of SSD occur in general practice are an important component. No matter of what eventually turns out to be the best estimate of the prevalence of SSD, the clinical assessments of GPs are decisive to initiate further treatment of patients. Since many participating general practitioners may not have been familiar with the diagnostic concept of somatic symptom disorder prior to the study, the estimate of 5% of full-blown SSD in general practice is probably a conservative estimate. Nevertheless, this frequency also shows that SSD must be regarded as a frequent and relevant clinical picture in general practice, which should not be underestimated or ignored diagnostically. Thus, it is not enough to introduce the diagnosis, but to disseminate it so that clinicians at gatekeeper positions might take the correct steps for further diagnosis and treatment of the afflicted patients.

Abbreviations

SSD

somatic symptom disorder

DSM-5

Diagnostic and Statistical Manual of Mental Disorders, 5th edition

GP

General practitioner

ICD-11

International Statistical Classification of Diseases, 11th edition

ICD-10

International Statistical Classification of Diseases, 10th edition

DSM-IV-TR

Diagnostic and Statistical Manual of Mental Disorders Text Revision

PHQ-15

Patient Health Questionnaire

BIPQ

Brief Illness Perception Questionnaire

BeSSD-GP

Identification of barriers and difficulties involved in the process of diagnosing somatic symptom disorders in primary care

VIF

variance inflation factor

CI

Confidence interval

Declarations

Ethical approval and consent to participate

We obtained ethical approval from the Ethics Committee of the Hamburg Medical Association, Germany, on April 7th, 2015 (approval number PV4763). This ethics committee is available for all studies located in Hamburg

(Germany), which involve clinical stakeholders. All methods were carried out and are described in accordance with the *Strengthening the Reporting of Observational Studies in Epidemiology* (STROBE) checklist for cross-sectional studies (Version 4, 2007).

The presented study is not medical research involving human subjects. Our results comprise of the opinions of general practitioners regarding aspects of their practice. No clinical data or assessments were involved in this study. Participation of general practitioners was completely anonymous and we obtained no written informed consent. Informed consent was obtained from all subjects by notifying them that if they returned the completed response form by anonymous mail, we would treat this as informed consent.

Consent for publication

Not applicable.

Availability of data and materials

The datasets generated during and analyzed during the current study are not publicly available because this step was not covered by approval of the ethics committee and funding agency, but are available from the corresponding author on reasonable request.

The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

Competing interests

The authors declare that they have no competing interests.

Funding

The study was funded by the Deutsche Forschungsgemeinschaft (DFG), Germany (<http://gepris.dfg.de/gepris/projekt/276028312>; Applicants: Martin Scherer, Astrid Althaus, Bernd Löwe). The sponsor of the study had no role in study design, data collection, data analysis, data interpretation, decision to publish, or writing of the report.

Authors' contributions

BL and MS conceptualised and designed the study. ML and NP were responsible for acquisition of data. ML analysed and interpreted the data. ML drafted the manuscript. BL, NP, TZ and MS revised the manuscript and contributed important intellectual content. BL and MS obtained funding and supervised the study. All authors read and approved the final manuscript.

Acknowledgements

We thank the anonymous general practitioners who volunteered to take part in this study. We thank Meike Mund for technical assistance with data validation. We thank Abigail Smakowski for language revision.

References

1. Diagnostic and Statistical Manual of Mental Disorders (DSM-5). Fifth Edition ed. American Psychiatric A, editor: Arlington, VA; 2013.
2. Gureje O, Reed GM. Bodily distress disorder in ICD-11: problems and prospects. *World Psychiatry*. 2016;15(3):291–2.
3. Löwe B, Levenson J, Depping M, Hüsing P, Kohlmann S, Lehmann M, et al. Somatic symptom disorder: a scoping review on the empirical evidence of a new diagnosis. *Psychol Med*. 2021:1–17.
4. Fergus TA, Kelley LP, Griggs JO. Examining the Whiteley Index-6 as a screener for DSM-5 presentations of severe health anxiety in primary care. *Journal of psychosomatic research*. 2019;127:109839.
5. Claassen-van Dessel N, van der Wouden JC, Dekker J, van der Horst HE. Clinical value of DSM IV and DSM 5 criteria for diagnosing the most prevalent somatoform disorders in patients with medically unexplained physical symptoms (MUPS). *J Psychosom Res*. 2016;82:4–10.
6. Rief W, Martin A. How to use the new DSM-5 somatic symptom disorder diagnosis in research and practice: a critical evaluation and a proposal for modifications. *Annu Rev Clin Psychol*. 2014;10:339–67.
7. Dimsdale JE, Creed F, Escobar J, Sharpe M, Wulsin L, Barsky A, et al. Somatic symptom disorder: an important change in DSM. *J Psychosom Res*. 2013;75(3):223–8.
8. Dimsdale JE, Levenson J. Diagnosis of Somatic Symptom Disorder requires clinical judgment. *J Psychosom Res*. 2013;75(6):588.
9. Lehmann M, Jonas C, Pohontsch NJ, Zimmermann T, Scherer M, Löwe B. General practitioners' views on the diagnostic innovations in DSM-5 somatic symptom disorder - A focus group study. *J Psychosom Res*. 2019;123:109734.
10. Toussaint A, Murray AM, Voigt K, Herzog A, Gierk B, Kroenke K, et al. Development and Validation of the Somatic Symptom Disorder-B Criteria Scale (SSD-12). *Psychosom Med*. 2016;78(1):5–12.
11. Kurlansik SL, Maffei MS. Somatic Symptom Disorder. *Am Fam Physician*. 2016;93(1):49–54.
12. Heinbokel C, Lehmann M, Pohontsch NJ, Zimmermann T, Althaus A, Scherer M, et al. Diagnostic barriers for somatic symptom disorders in primary care: study protocol for a mixed methods study in Germany. *BMJ open*. 2017;7(8):e014157-e.
13. Lehmann M, Pohontsch NJ, Zimmermann T, Scherer M, Löwe B. Diagnostic and treatment barriers to persistent somatic symptoms in primary care - representative survey with physicians. *BMC Fam Pract*. 2021;22(1):60.
14. American Psychiatric Association. *Diagnostic and statistical manual of mental disorders*. 4th ed., text rev. ed2000.
15. Christensen RHB. *ordinal—Regression Models for Ordinal Data*. 2015.
16. Fox J, Weisberg S. *An R companion to applied regression*. Third edition / ed. Los Angeles: SAGE; 2019. 577 p.
17. Häuser W, Hausteiner-Wiehle C, Henningsen P, Brähler E, Schmalbach B, Wolfe F. Prevalence and overlap of somatic symptom disorder, bodily distress syndrome and fibromyalgia syndrome in the German general population: A cross sectional study. *J Psychosom Res*. 2020;133:110111.

18. Kop WJ, Toussaint A, Mols F, Löwe B. Somatic symptom disorder in the general population: Associations with medical status and health care utilization using the SSD-12. *Gen Hosp Psychiatry*. 2019;56:36–41.

Figures

Figure 1

Estimated proportions of patients fulfilling clinical symptoms of somatic symptom disorder.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SupplementaryMaterialIntroductionItemsSSD.docx](#)