

Laboratory colonization by *Dirofilaria immitis* alters the microbiome of female *Aedes aegypti* mosquitoes

Abdulsalam Adegoke

University of Southern Mississippi

Erik Neff

University System of Georgia

Amie Geary

University of Southern Mississippi

Montana Ciara Husser

University of Southern Mississippi

Kevin Wilson

University of Southern Mississippi

Shawn Michael Norris

University of Southern Mississippi

Guha Dharmarajan

University System of Georgia

Shahid Karim (✉ Shahid.Karim@usm.edu)

The University of Southern Mississippi <https://orcid.org/0000-0001-5596-3304>

Research

Keywords: mosquitoes, *Aedes aegypti*, microbiome, metagenome, *Dirofilaria immitis*, Dog heartworm

Posted Date: June 16th, 2020

DOI: <https://doi.org/10.21203/rs.2.23991/v5>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Parasites & Vectors on July 13th, 2020. See the published version at <https://doi.org/10.1186/s13071-020-04218-8>.

Abstract

Background: The ability of blood feeding arthropods to successfully acquire and transmit pathogens of medical and veterinary importance has been shown to be interfered with, or enhanced by, the arthropod's native microbiome. Mosquitoes transmit viruses, protozoan and filarial nematodes, the majority of which contribute to the 17% of infectious disease cases worldwide. *Dirofilaria immitis*, a mosquito transmitted by filarial nematodes of dogs and cats, is vectored by several mosquito species including *Aedes aegypti*.

Methods: In this study, we investigated the impact of *D. immitis* colonization on the microbiome of laboratory reared female *A. aegypti*. Metagenomic analysis of the V3-V4 variable region of the microbial 16S rRNA was used for identification of the microbial differences down to species level.

Results: We generated a total of 1068 OTUs representing 16 phyla, 181 genera and 271 bacterial species. Overall, in order of abundance, Proteobacteria, Bacteroidetes, Actinobacteria and Firmicutes were the most represented phylum with *D. immitis* infected mosquitoes having more of Proteobacteria (71%) than uninfected mosquitoes (56.9%). An interesting finding in this study is the detection of *Klebsiella oxytoca* in relatively similar abundance in infected and uninfected mosquitoes, suggesting a possible endosymbiotic relationship, and has been previously shown to indirectly compete for nutrients with fungi on domestic housefly eggs and larva. While *D. immitis* colonization has no effect on the overall species richness, we identified significant differences in the composition of selected bacteria genus and phylum between the two groups. We also reported distinct compositional and phylogenetic differences in the individual bacteria species when commonly identified bacteria were compared.

Conclusions: In conclusion, this is the first study to the best of our knowledge to understand the impact of a filarial infection on the microbiome of its mosquito vector. Further studies are required to identify bacteria species that could play an important role in the mosquito biology. While the microbiome composition of *A. aegypti* mosquito have been previously reported, our study shows that in an effort to establish itself, a filarial nematode modifies and alters the overall microbial diversity within its mosquito host.

Background

Dirofilaria immitis, responsible for canine respiratory and associated heartworm disease, is a filarial nematode that mainly infects wild and domestic canids, such as dogs and coyotes [1]. It causes serious life-threatening disease of canines and other domestic pets in which adult worms lodge in the cardiopulmonary circulation of infected dogs causing cardiac necrosis which leads to ischemic hypoxia and a total shutdown of the respiratory and cardiovascular system [2]. While primarily a parasite of veterinary importance, human infection has also been reported, albeit rarely [3]. With over 100,000 dogs affected annually, *D. immitis* has been identified as the most important parasite affecting dogs in the United States [4]. Adult female *D. immitis* parasites release microfilariae (mf) into the blood stream of the vertebrate host, usually a dog, which are subsequently taken up by adult female mosquitoes during their

blood meal [1, 5]. Microfilaria in the blood enter the midgut of the mosquito during feeding and subsequently migrate to the Malpighian tubules. In the Malpighian tubules of the mosquito mf develop into first stage larvae and molt twice to reach their infective L3 stage, which migrate to the head and proboscis of the mosquito and can infect the next dog during the next blood meal [1, 5]. The time it takes for the parasite to develop, known as the extrinsic incubation period (EIP), averages 14 days within this mosquito [6].

Beyond the United States, canine dirofilariasis also poses significant problems in dogs in Australia, Europe and Asia [7]. The epidemiology and maintenance of *D. immitis* have been made significant by the abundance of the mosquito vectors with over 70 species of mosquitoes, having been reported to transmit *D. immitis* [8], and 28 of these are found in the United States [9]. The southeastern United States have been shown to have an increased incidence of canine heartworm disease when compared to other states [10], as about half of the mosquito species known to transmit *D. immitis* are found in the Southeastern United States [11], one of which is the *Aedes aegypti* mosquito [12]. In addition to being an important vector of filarial nematodes such as *D. immitis*, mosquitoes and other arthropod vectors also host highly complex and diverse microbial communities. Members of these microbial communities have obligate relationships with mosquitoes and are maintained vertically through mosquito generations [13] or acquired from the environment [14, 15]. The microbiome of mosquitoes have been shown to help maintain normal midgut functions, as well as interfering or aiding vector competence of the mosquito [16, 17]. As previously shown, pathogen susceptibility and vector competence of an arthropod can be modulated by the microbial composition within the arthropod vector [18]. *Aedes aegypti* is the competent vector of many pathogens of humans (Zika, Dengue, Chikungunya Viruses; [19, 20, 21]) and animals (e.g., *D. immitis*; [22]).

Previous studies have demonstrated significant success in reducing vector competence of mosquitoes for pathogenic microbes by taking advantage of the mosquito microbiome and pathogen interactions [23, 24]. It is a fact that pathogens must compete with the native microbial community in their vector host to propagate and facilitate subsequent transmission to a susceptible host [18, 25]. Hence, understanding the pathogen-microbiome interaction in vectors of significant public health importance is important. The prevention and control of canine heartworm disease have primarily focused on the use of therapeutics such as ivermectin and other macrocyclic lactones. While these therapeutics have proven to be effective in killing and preventing the establishment of the third and early fourth-stage larva [26], recent studies have shown significant resistance against these drugs of choice [27, 28]. Understanding the role(s) played by the mosquito microbiome in facilitating or interfering with *D. immitis* colonization and subsequent transmission will add to existing knowledge on the control of mosquito-transmitted filarial nematodes. The current study compared the microbiome changes in *A. aegypti*, infected with *D. immitis* under laboratory conditions. This study provides an exciting opportunity to advance our knowledge of the major changes in the microbiome of the *A. aegypti* mosquito following the acquisition of *D. immitis*. This research seeks to address the question “Does *D. immitis* infection of *A. aegypti* mosquito alters its overall microbial richness and abundances?”

Methods

Study organisms. The mosquito used for investigation was the *A. aegypti*, Liverpool Blackeye strain, a highly susceptible mosquito strain to *D. immitis* used predominately in research for *Aedes sp.* [29]. Although *A. aegypti* is just one of 28 different mosquito species reported to vector *D. immitis*, the relative expertise on this particular species limited us to its use as our study organism.

Mosquito maintenance. *Aedes aegypti*, originally obtained from the Filariasis Research Reagent Resource Center (FR3) [30], were raised under standard laboratory conditions – 27 °C, 80+5% relative humidity, and a 12:12-hour light diurnal cycle (Dharmarajan et al. 2019). Adult female mosquitoes, five days post emergence were blood fed using an artificial membrane feeder. One day prior to membrane feeding 31 female mosquitoes were transferred to ~500 mL plastic containers with mesh tops (henceforth “cages”). Females were starved of sugar for 12 hours and deprived of water for four hours prior to blood feeding. Mosquitoes in each cage (31 each) were allowed to feed for two hours or until repletion on a Parafilm membrane stretched over an inverted water-jacketed glass membrane feeder maintained at 40°C. Each feeder was filled with 200 µL of dog blood infected or uninfected with *D. immitis*. The level of parasitaemia in the infected blood was estimated to be 4,500 mf/mL as previously reported [31]. The dog blood was obtained from FR3.

***Dirofilaria immitis* detection**

Prior to screening for *D. immitis* infection from mosquitoes, DNA from individual mosquito samples was extracted using a commercially available kit (QIAGEN, DNeasy Blood & Tissue Kit) and quality was confirmed using a nanodrop machine (Thermo SCIENTIFIC, NANODROP ONE). All blood fed (infected and uninfected blood) mosquito samples were screened for *D. immitis* infection irrespective of whether they were fed on infected or uninfected blood by amplifying the COI gene (656bp) of the *D. immitis* mitochondrial DNA [32]. Briefly, a 25µl reaction was set up comprising 1µl each of the COI-Forward (5′–TGATTGGTGGTTTTGGTAA–3′) and COI-Reverse (5′–ATAAGTACGAGTATCAATATC–3′) primers, 12.5µl of 2X mastermix (New England Biolabs), 2.5 µl DNA template and 8.5 µl of nuclease free water. For each cycle that was run, a *D. immitis* infected blood sample and nuclease free water were simultaneously included as positive and negative controls respectively. The PCR cycle comprised of an initial denaturation step at 94⁰C for 5 minutes and 40 cycles of 94⁰C for 1 minute, annealing at 50⁰C for 2 minutes and extension at 72⁰C for 3 minutes. A final extension at 72⁰C for 5 minutes and an infinite hold at 4⁰C.

Confirmation of amplification was done by loading the PCR products in a SYBR safe stained gel. Briefly, 2% gel was made by autoclaving a solution of 1X TAE buffer and molecular grade agar. SYBR safe stain (1 µl SYBR safe: 10 mL TAE buffer) was added to the agar solution, poured into a precast gel tray and allowed to cool. To load the samples onto the gel, 6 µl of PCR product was mixed with 4 µl of 6X dye and pipetted into the wells. Lastly, 5 µl of a low molecular weight DNA ladder was loaded onto the gel and the

gel could run for 45 minutes at 100 V. Amplified PCR products were viewed using a Chemidoc gel imager (Supplementary Figure 1).

16S rRNA library preparation and sequencing

Six individual mosquito genomic DNA extracts were pooled to make one biological replicate and five biological replicates each of *D. immitis* infected and uninfected pools were prepared for metagenomic analysis. The hypervariable V1-V3 region of the 16S rRNA gene was PCR amplified using the forward primer 27F; **5'- AGRGTTTGATCMTGGCTCAG-3'**, and reverse primer 519R; **5' – GTNTTACNGCGGCKGCTG - 3'** as outlined by the 16S Illumina's MiSeq protocol (www.mrdnalab.com, Shallowater, TX, USA). Briefly, PCR was performed using the HotStarTaq Plus Master Mix Kit (Qiagen, USA) under the following conditions: 94°C for 3 min, followed by 30-35 cycles of 94°C for 30 s, 53°C for 40 s and 72°C for 1 min, after which a final elongation step at 72°C for 5 min was performed. After amplification, PCR products were electrophoresed in 2% agarose gel to determine the success of amplification and the relative intensity of bands. Multiple samples were pooled together in equal proportions based on their molecular weight and DNA concentrations. Pooled samples were purified using calibrated Ampure XP beads. Then the pooled and purified PCR product was used to prepare Illumina DNA library. Sequencing was performed at MR DNA (www.mrdnalab.com, Shallowater, TX, USA) on a MiSeq following the manufacturer's guidelines.

Sequence analysis

Sequence analysis was done using the Quantitative Insights into Microbial Ecology (QIIME 2) pipeline, unless stated otherwise. Briefly, processing of raw fastq files were demultiplexed. The Atacama soil microbiome pipeline was incorporated for quality control of demultiplexed paired-end reads (Supplementary Figure 2) using the DADA2 plugin as previously described [33]. Sequence alignment and subsequent construction of phylogenetic tree from representative sequences was done using the MAFFT v7 and FasTree v2.1 plugin [34]. Operational taxonomic assignment was done using the qiime2 feature-classifier plugin v7.0 which was previously trained against the **SILVA 132 database** preclustered at 99%. Tables representing operational taxonomic units (OTUs) and representative taxonomy were exported from R and used for diversity metric analysis using the Microbiome Analyst web-based interface [35, 36]. Raw data from this analysis were submitted deposited and assigned the GenBank BioProject number **#PRJNA606536**.

Alpha diversity

To establish whether alpha diversity differs across mosquito samples, reads were transformed and low abundance OTUs were filtered from the datasets. The Observed OTU metric was used to estimate species richness by identifying unique OTUs present across the mosquito groups, while the Shannon index were used to estimate both richness and evenness.

Beta diversity

To compare the differences in the microbiome between mosquito groups, based on measures of distance or dissimilarity, dissimilarity matrix were generated from log-transformed sequence data and ordination of the plots were visualized using both the Principal Coordinates Analysis (PCoA) and the Nonmetric Multidimensional Scaling (NMDS). The matrix used in calculating beta diversity includes the Bray_Curtis and unweighted UniFrac distance matrix.

Statistical analysis

To test if species richness and diversity was significant, the Mann-Whitney/Kruskal-Wallis test was applied to both alpha diversity and classical univariate statistical comparisons analysis, while the significance of beta diversity analysis was determined using the Permutational MANOVA (PERMANOVA) test.

Results

D. immitis effectively colonizes A. aegypti mosquitos under laboratory conditions

A total of 31 mosquitoes were artificially infected to generate *D. immitis* infected mosquito. PCR analysis and confirmation of infection status using gel electrophoresis revealed an infection prevalence of 87% (Supplementary data, Fig 1).

Microbiome composition

Analysis of the demultiplexed paired-end-reads generated a total of 602502 reads which ranged from 31861 to 110235, and an average of 54758 reads per mosquito. Mosquitoes infected with *D. immitis* had the highest number of reads (382714) compared to uninfected mosquitoes with a total of 219788 reads. Taxonomic classification using the SILVA 132 reference database (99% OTUs full-length sequences) identified 268 operational taxonomic units (OTUs), 11 phyla, 16 genera, and 136 species (Supplementary Figure 2 and Table 1).

D. immitis infection alters relative abundance of bacteria taxa in A. aegypti. Taxonomic assignment was done against the SILVA database to observe for the difference in microbiome composition and relative abundance of bacteria species. Both infected and uninfected mosquitoes possess similar composition of bacteria taxa with differences observed in the relative abundances of specific bacteria species. The phylum Proteobacteria and Bacteroidetes were present in higher abundance in both mosquito groups. Mosquitoes infected with *D. immitis* had relatively higher abundance of Proteobacteria (71%) with lower amount of Bacteroidetes (27%), while uninfected mosquitoes had lower amount of Proteobacteria (56.9%) and a higher abundance of Bacteroidetes (37%) (Fig. 1). Among the detected bacteria genera, the genus *Klebsiella* were detected at relatively similar abundances in both infected (36.3%) and uninfected (34.6%) mosquitoes (Fig 2A). The genera *Ochrobactrum*, *Sphingobacterium*, and *Pseudomonas* were present exclusively in uninfected mosquitoes albeit at an abundance of 4.4%, 3.6% and 3% respectively (Fig 2). The genera *Enterobacter* (26.1%) and *Elizabethkingia* (9.4%) were in higher abundance in *D.*

immitis infected mosquitoes (Fig 2A). *Enterobacter hormaechei* (24.4%) and *Elizabethkingia meningoseptica* (9.4%) were more abundant in infected mosquitoes, while *Chryseobacterium indolgenes* (27.4%), and *Grimontella senegalensis* (3.9%), were more abundant in uninfected mosquitoes (Fig.2B). Additional information on the relative bacteria distribution in individual mosquito can be found in the Supplementary figures 4, 5, 6, and 7.

***D. immitis* infection affects species diversity in *A. aegypti*.** Following demultiplexing of paired end reads and quality control by removing chimeric sequences, we normalized OTU counts for individual biological replicates and a rarefaction curve was generated at a depth of 20000 (Fig. 3). Adequate depth coverage was reached as evidenced by the individual curves plateauing out on the rarefaction curve. Our results indicated that phylogenetic diversity, estimated using Shannon index and number of observed OTUs, was reduced in infected mosquitoes compared against uninfected mosquitoes (Fig. 4A and B). Surprisingly, both metrics of alpha diversity used showed no significance when the Kruskal-Wallis test was applied (Observed OTUs: $P=0.053$; (Mann-Whitney) statistic: 3; Shannon Index; $P= 0.309$; (Mann-Whitney) statistic: 7). We also visualized the similarities and differences in the microbial composition of infected and uninfected mosquitoes by carrying out Principal Components Analysis (PCoA) of the unweighted UniFrac and Bray Curtis and distance matrixes (Fig. 5C and D). Figures 5C and 5D shows distinct clustering of *D. immitis* infected mosquito replicates with no outliers. Beta diversity was significantly different when *D. immitis* infected mosquito was compared to those uninfected using the Unweighted UniFrac distance matrix (PERMANOVA, F-value: 1.4043; R-squared: 0.14932; $P=0.286$).

Community profiling and correlation analysis

To assess the extent to which highly abundant bacteria phylum and genus were represented in *R. microplus* ticks, we used a combination of pattern correlation and heat map analysis. A very strong positive correlation was seen between the genus *Ralstonia*, *Francisella*, *Pantoea*, *Elizabethkingia*, *Wolbachia*, *Herbaspirillum*, and *Achromobacter* and *D. immitis* infected mosquitoes (Figure 6A). Heat map analysis and phylogenetic tree of the highly represented and dominant bacteria genera also showed the above identified genera to be well represented in more than one of the *D. immitis* infected mosquito (Supplementary Figure 8, and 9).

To explore how top taxa differed between both infected and uninfected mosquitoes, classical univariate statistical comparisons analysis was applied to identify for phylum and genus that exhibit significant differences (Mann-Whitney test) in their composition. The phylum Actinobacteria ($P=0.036$, FDR; 0.1440), and Firmicutes ($P=0.005$, FDR; 0.14401) has significantly higher abundance in the uninfected mosquitoes (Figure 6B and C). Our analysis also shows that the *D. immitis* mosquito has significantly higher abundance of the genera *Elizabethkingia* ($P=0.015$, FDR; 0.175), and *Wolbachia* ($P=0.011$, FDR; 0.11448), while *Pseudomonas* ($P=0.015$, FDR; 0.12698) was much abundant in the uninfected mosquito (Figure 7A, B, and C).

Discussion

To the best of our understanding, this study is the first to utilize 16S metagenomic analysis to understand the significance of a filarial nematode on the microbiome of a mosquito vector under laboratory conditions. This technique offers the advantage of detecting both culturable and unculturable pathogenic and non-pathogenic microbes from a given DNA or RNA sample. As previously reported from previous metagenome studies on mosquitoes and similar arthropods or insects [37, 38, 39], Figure 1A shows that the phylum Proteobacteria, Bacteroidetes, Actinobacteria and Firmicutes was detected from the mosquito groups tested. Similar studies have also identified the phylum Proteobacteria as one of the dominant phyla in the microbiome of *A. aegypti* mosquitoes [23, 40]. We observed an inverse relationship in the abundances of the phylum Proteobacteria and Bacteroidetes in our study with *D. immitis* infected mosquitoes having a higher abundance of the phylum Proteobacteria when compared to the uninfected mosquitoes and vice-versa (Supplementary data, figure 4). The Proteobacteria group are the largest phylum found in different environments, plants and animals [41] with members ranging from free living commensals to pathogenic microbes. We found that the genus *Klebsiella* was present in relatively similar abundances in both infected and uninfected mosquitoes. Although no known function has been associated to this group in insects or arthropods, *Klebsiella* belongs to a class of gram Proteobacteria known as Gammaproteobacteria with previous detection in field and laboratory raised *Culex quinquefasciatus*, *A. albopictus*, and *A. aegypti* [42, 43, 44].

We also identified the genus *Enterobacter* in *D. immitis* infected mosquito at a relative abundance of approximately 20 fold of the uninfected mosquitoes was also observed (Figure 2). This bacteria genus has been detected in similar microbiome studies of *A. aegypti* where they have been associated with their role in blood meal digestion due to their hemolytic activities [45] which could explain why the genus *Enterobacter* have been commonly associated with females of other mosquito species [46]. The role(s) *D. immitis* colonization and infection density plays in increasing the abundance of *Enterobacter* was beyond the scope of this study, an elegant study by Cirimotich and colleagues showed the inhibition of *Plasmodium* infection in *Anopheles gambiae* mosquitoes mediated by a bacterium designated as *Enterobacter* Esp_Z. The inhibition was due to the production of reactive oxygen species (ROS) by the bacteria. A pro-pathogen role was also recently associated with the *Enterobacter* genus, as it was shown that they produce Enhancins or Enhancins-like proteins, which facilitate pathogen colonization by degrading the peritrophic matrix [44]. If bacteria such as *Enterobacter* can also block *D. immitis* colonization by inducing ROS production will be interesting to see. *Enterobacter hormaechei* is another bacterium that demonstrated increased abundance in *D. immitis* infection as shown in Figure 3. The genus *Enterobacter* have been proposed to aid in blood digestion in hematophagous insects due to their hemolytic activities [47]. Several reports have also identified different mosquito refractoriness or susceptibility to pathogen infection in the presence of different *Enterobacter* species. Cirimotich et al. [48] reported refractoriness of *Anopheles gambiae* to *Plasmodium* infection in the presence of an *Enterobacter* species. Another species of *Enterobacter* (*E. cloacae*) was also reported to express a mucin degrading Enhancin protein that breaks down the mucin component of the *A. aegypti* peritrophic matrix [44] although this wasn't shown to facilitate Dengue virus infection. Another bacteria genus that also

increased with the presence of *D. immitis* infection is *Elizabethkingia*. Previous reports have identified this genus from the midgut of laboratory reared [42] and field collected [40] *A. aegypti*. These studies did not associate any known role to this bacterium. Bacteria in the genus *Chryseobacterium*, *Ochrobactrum*, *Sphingobacterium*, and *Pseudomonas* were all present in higher abundance in uninfected mosquito group. The detection of *Chryseobacterium* from our study agrees with similar detection from previous report in laboratory reared mosquitoes [49, 50]. *Pseudomonas*, a Gram-negative, Gammaproteobacteria was previously shown to have reduced abundance in *Wolbachia* positive *A. aegypti* mosquitoes. A similar observation was also made from our study where the abundance of *Pseudomonas* was inversely correlated with the presence of *D. immitis* infection in the mosquito as shown in Figure 2. Bahia et al. [51] reported blocking of the *Plasmodium* parasite by *Pseudomonas putida* isolated from *A. gambiae*. Unexpectedly, only few bacteria species were differentially altered in *D. immitis* infected and uninfected mosquitoes. *Klebsiella (K.) oxytoca*, a Gammaproteobacteria belonging to the genus *Klebsiella*, was found at relatively similar abundance in infected and uninfected mosquitoes (Fig 3). A study reported the detection of *K. oxytoca* from laboratory reared and field collected *C. quinquefasciatus* and *A. aegypti* [42]. The maintenance of a bacteria by arthropods in both natural and artificial conditions could indicate an important role played by the bacteria. Another study reported that *K. oxytoca* reverse radiation induced loss of copulatory maintenance in male *Ceratitis capitata* [52]. Another study on the fungicidal effects of bacterial colonies found on the domestic housefly eggs revealed that fungal growth is inhibited by the presence of *K. oxytoca* by producing antifungal metabolites and nutrient depletion [53]. Put together, we are proposing *K. oxytoca* as an important bacteria *A. aegypti* with a likely endosymbiotic role, though further studies are still required to understand the specific roles of *K. oxytoca* in the mosquito biology including how it is maintained in the mosquito. Another interesting observation from this study was the inverse correlation between *D. immitis* infection and microbial diversity and richness. In mosquitoes, the innate immune response is activated in the presence of invading pathogens as it was previously shown that activation of the toll pathway and production of reactive oxygen species have been reported in mosquitoes challenged with *Plasmodium* [54] and filarial nematodes [55]. These immune effectors while countering pathogens, could in extension disrupt the normal mosquito microbiome community which could explain the overall reduction in the microbial richness observed from this study as seen in Figure 4B and C. Our observation of reduced microbial richness in infected mosquitoes was in contrast to what was reported in a previous study which shows a more diverse microbiome in *Plasmodium*-infected *Anopheles* mosquitoes [56]. While our current study did not factor the effect blood meal could have on the overall outcome of the microbial richness, quite a number of studies have reported reduction in the bacterial diversity following experimental feeding on host blood in *A. aegypti* [57], *An. gambiae* [50].

Conclusion

This study fills a knowledge gap on the interaction between a mosquito vector and a filarial pathogen of veterinary significance. We were able to show that while some bacterial species were found to be present in both mosquito groups, the relative abundances of individual species changes with the infection status, with infected mosquitoes presenting a reduced microbial richness. This indicates a likely consequence of

the nematode in altering favoring or inhibiting the growth of members of the bacterial community. Ongoing study focuses on the shift in the distribution of culturable bacteria taxa in infected and uninfected mosquitoes, while also comparing the effects of *D. immitis* colonization on the microbial diversity of different tissues of the *A. aegypti* mosquito.

Declarations

Ethics Approval and Consent to participate:

Mosquitoes were blood-fed using an artificial membrane feeder. The protocol for the laboratory was approved by the Institutional Biosafety Committee.

Competing interests

The authors declare that they have no competing interests.

Consent of Publication:

All authors read and approved the manuscript for publication.

Availability of data and material:

The datasets supporting the conclusion of this article are included within the article and its additional files. Raw data are available from the corresponding author upon request.

Funding:

This research was principally supported by a Pakistan-US Science and Technology Cooperation Program award (US Department of State), the National Institutes of General Medical Sciences #P20 and GM103476 to the University of Southern Mississippi and the Department of Energy Office of Environmental Management under Award Number DE-EM0004391 to the University of Georgia Research Foundation. AG, MCH, KW, and SMN were supported the Eagle SPUR award from the USM Drapeau Center for Undergraduate Research.

Acknowledgements:

The following reagents were provided by the NIH/NIAID Filariasis Research Reagent Resource Center through BEI Resources: Uninfected Canine Blood (Not Exposed to Filariae) NR-48922 and *Dirofilaria immitis*, Strain Missouri, *Microfilariae* in Dog Blood (Live) NR-4890. We also thank the University of Southern Mississippi's School of Biological, Environmental, and Earth Sciences Human Parasitology class participants (Fall 2019) for their technical assistance in extracting the genome DNA and pathogen detection.

Author contributions:

Conceived and designed the experiments: GD, SK

Performed the experiments: ASA, EN, AG, MCH, KW, SMN

Analyzed the data: ASA, GD, SK

Contributed reagents/materials/analysis tools: GD, SK

Wrote the paper: ASA, EN, AG, GD, SK

All authors have read and approved the manuscript

Competing Interests:

The authors have declared that no competing interests exist.

References

1. Bowman DD, Atkins CE. Heartworm biology, treatment, and control. *Vet Clin North Am Small Anim Pract.* 2009;39(6):1127-vii.
2. Spence Beaulieu MR, Reiskind MH. Comparative vector efficiency of two prevalent mosquito species for dog heartworm in north Carolina. *J Med Entomol.* 2020;57(2):608-614.
3. Theis JH. Public health aspects of dirofilariasis in the United States. *Vet Parasitol.* 2005;133(2-3):157-180.
4. Bowman, D.D., Liu, Y., McMahan, C.S., Nordone S.K., Yabsley, M.j., Lund, R.B. Forecasting United States heartworm *Dirofilaria immitis* prevalence in dogs. *Vectors.* **9**, 540 (2016).
5. Evans CC, Moorhead AR, Storey BE, Wolstenholme AJ, Kaplan RM. Development of an in vitro bioassay for measuring susceptibility to macrocyclic lactone anthelmintics in *Dirofilaria immitis*. *Int J Parasitol Drugs Drug Resist.* 2013;3:102-108.
6. McCall JW. The role of arthropods in the development of animal - models for filariasis research. *J. Georgia. Ent. Soc.* 1981; 16: 283-293.
7. Morchón R, Carretón E, González-Miguel J, Mellado-Hernández I. Heartworm disease (*Dirofilaria immitis*) and their vectors in Europe - new distribution trends. *Front Physiol.* 2012;3:196.
8. Wang D, Bowman DD, Brown HE, Harrington LC, Kaufman PE, McKay T, et al. Factors influencing U.S. canine heartworm (*Dirofilaria immitis*) prevalence. *Vectors.* 2014;7:264.
9. Ludlam KW, Jachowski LA Jr, Otto GF. Potential vectors of *Dirofilaria immitis*. *J Am Vet Med Assoc.* 1970;157(10):1354-1359.

10. Drake, J., Wiseman, S. Increasing incidence of *Dirofilaria immitis* in dogs in USA with focus on the southeast region 2013–2016. *Vectors*. **11**, 39 (2018).
11. Lok JB, Walker ED, Scoles GA. Filariasis. In: Eldridge BF, Edman JD, editors. *Medical Entomology*. The Netherlands: Kluwer Academic Publishers, Dordrecht; 2000. p. 299-375.
12. Ledesma N, Harrington L. Mosquito vectors of dog heartworm in the United States: vector status and factors influencing transmission efficiency. *Top Companion Anim Med*. 2011;26(4):178-185.
13. Guégan M, Zouache K, Démichel C, Minard G, Van VT, Potier P, et al. The mosquito holobiont: fresh insight into mosquito-microbiota interactions. *Microbiome*. 2018;6(1):49. Published 2018 Mar 20.
14. Engel P, Moran NA. The gut microbiota of insects - diversity in structure and function. *FEMS Microbiol Rev*. 2013;37(5):699-735.
15. Strand MR. The gut microbiota of mosquitoes: diversity and function. In: Wikel S, Aksoy S, Dimopoulos G, editors. *Arthropod Vector: Controller of Disease Transmission*. Vol. 1. Elsevier; 2017. pp. 185–199.
16. Strand MR. Composition and functional roles of the gut microbiota in mosquitoes. *Curr Opin Insect Sci*. 2018;28:59-65.
17. Souza-Neto JA, Powell JR, Bonizzoni M. *Aedes aegypti* vector competence studies: A review. *Infect Genet Evol*. 2019;67:191-209.
18. Bonnet SI, Binetruy F, Hernández-Jarguín AM, Duron O. The tick microbiome: Why non-pathogenic microorganisms matter in tick biology and pathogen transmission. *Front Cell Infect Microbiol*. 2017;7:236.
19. Alto BW, Lounibos LP, Mores CN, Reiskind MH. Larval competition alters susceptibility of adult *Aedes* mosquitoes to dengue infection. *Proc Biol Sci*. 2008;275(1633):463-471.
20. Marchette NJ, Garcia R, Rudnick A. Isolation of Zika virus from *Aedes aegypti* mosquitoes in Malaysia. *Am J Trop Med Hyg*. 1969;18(3):411-415.
21. Martin E, Moutailler S, Madec Y, Failloux AB. Differential responses of the mosquito *Aedes albopictus* from the Indian Ocean region to two chikungunya isolates. *BMC Ecol*. 2010;10:8.
22. Erickson SM, Xi Z, Mayhew GF, Ramirez JL, Aliota MT, Christensen BM, et al. Mosquito infection responses to developing filarial worms. *PLoS Negl Trop Dis*. 2009;3(10):e529.
23. Ramirez JL, Souza-Neto J, Torres Cosme R, Rovira J, Ortiz A, Pascale JM, et al. Reciprocal tripartite interactions between the *Aedes aegypti* midgut microbiota, innate immune system and dengue virus influences vector competence. *PLoS Negl Trop Dis*. 2012;6(3):e1561.

24. Dennison NJ, Jupatanakul N, Dimopoulos G. The mosquito microbiota influences vector competence for human pathogens. *Curr Opin Insect Sci.* 2014;3:6-13.
25. de la Fuente J, Antunes S, Bonnet S, , Domingos AG, Estrada-Peña A, Johnson N, et al. Tick-pathogen interactions and vector competence: identification of molecular drivers for tick-borne diseases. *Front Cell Infect Microbiol.* 2017;7:114.
26. Bourguinat C, Lee AC, Lizundia R, Blagburn BL, Liotta JL, Kraus MS, et al. Macrocyclic lactone resistance in *Dirofilaria immitis*: Failure of heartworm preventives and investigation of genetic markers for resistance. *Vet Parasitol.* 2015;210(3-4):167-178.
27. Bourguinat C, Keller K, Bhan A, Peregrine A, Geary T, Prichard R. Macrocyclic lactone resistance in *Dirofilaria immitis*. *Vet Parasitol.* 2011;181(2-4):388-392.
28. Geary TG, Bourguinat C, Prichard RK. Evidence for macrocyclic lactone anthelmintic resistance in *Dirofilaria immitis*. *Top Companion Anim Med.* 2011;26(4):186-192.
29. Buxton BA, Mullen GR. Comparative susceptibility of 4 strains of *Aedes aegypti* (Diptera, Culicidae) to infection with *Dirofilaria immitis*. *J Med Entomol* 1981; 18: 434-440.
30. Michalski ML, Griffiths KG, Williams SA, Kaplan RM, Moorhead AR. The NIH-NIAID filariasis research reagent resource center. *PLoS Negl Trop Dis.* 2011;5(11):e1261.
31. Dharmarajan G, Walker KD, Lehmann T. Variation in tolerance to parasites affects vectorial capacity of natural asian tiger mosquito populations. *Curr Biol.* 2019;29(22):3946-3952.e5.
32. Murata K, Yanai T, Agatsuma T, Uni S. *Dirofilaria immitis* infection of a snow leopard (*Uncia uncia*) in a Japanese zoo with mitochondrial DNA analysis. *J Vet Med Sci.* 2003;65(8):945-947.
33. Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJ, Holmes SP. DADA2: High-resolution sample inference from Illumina amplicon data. *Nat Methods.* 2016;13(7):581-583.
34. Price MN, Dehal PS, Arkin AP. FastTree 2—approximately maximum-likelihood trees for large alignments. *PLoS One.* 2010;5(3):e9490.
35. Chong J, Liu P, Zhou G, Xia J. Using MicrobiomeAnalyst for comprehensive statistical, functional, and meta-analysis of microbiome data. *Nat Protoc.* 2020;15(3):799-821.
36. Dhariwal A, Chong J, Habib S, King IL, Agellon LB, Xia J. MicrobiomeAnalyst: a web-based tool for comprehensive statistical, visual and meta-analysis of microbiome data. *Nucleic Acids Res.* 2017;45(W1):W180-W188.
37. Mancini MV, Damiani C, Accoti A, Tallarita M, Nunzi E, Cappeli A, et al. Estimating bacteria diversity in different organs of nine species of mosquito by next generation sequencing. *BMC Microbiol.*

2018;18(1):126.

38. Coon KL, Vogel KJ, Brown MR, Strand MR. Mosquitoes rely on their gut microbiota for development. *Mol Ecol.* 2014;23(11):2727-2739.
39. Coatsworth H, Caicedo PA, Van Rossum T, Ocampo CB, Lowenberger C. The composition of midgut bacteria in *Aedes aegypti* (Diptera: Culicidae) that are naturally susceptible or refractory to dengue viruses. *J Insect Sci.* 2018;18(6):12.
40. Audsley MD, Seleznev A, Joubert DA, Woolfit M, O'Neill SL, McGraw EA. *Wolbachia* infection alters the relative abundance of resident bacteria in adult *Aedes aegypti* mosquitoes, but not larvae. *Mol Ecol.* 2018;27(1):297-309.
41. Degli Esposti M, Martinez Romero E. The functional microbiome of arthropods. *PLoS One.* 2017;12(5):e0176573.
42. Gazzoni Araújo Gonçalves G, Feitosa APS, Portela-Júnior NC, de Oliveira CMF, Filho JL de L, Brayner FA, et al. Use of MALDI-TOF MS to identify the culturable midgut microbiota of laboratory and wild mosquitoes. *Acta Trop.* 2019;200:105174.
43. Yadav KK, Adhikari N, Khadka R, Pant AD, Shah B. Multidrug resistant Enterobacteriaceae and extended spectrum β -lactamase producing *Escherichia coli*: a cross-sectional study in National Kidney Center, Nepal. *Antimicrob Resist Infect Control.* 2015;4:42.
44. Wu P, Sun P, Nie K, Zhu Y, Shi M, Xiao C, et al. A gut commensal bacterium promotes mosquito permissiveness to arboviruses. *Cell Host Microbe.* 2019;25(1):101-112.e5.
45. Coon, K.L., Brown, M.R. & Strand, M.R. Gut bacteria differentially affect egg production in the anautogenous mosquito *Aedes aegypti* and facultatively autogenous mosquito *Aedes atropalpus* (Diptera: Culicidae). *Vectors.* **9**, 375 (2016).
46. Minard, G., Mavingui, P. & Moro, C.V. Diversity and function of bacterial microbiota in the mosquito holobiont. *Vectors.* **6**, 146 (2013).
47. Gaio Ade O, Gusmão DS, Santos AV, Berbert-Molina MA, Pimenta PF, Lemos FJ. Contribution of midgut bacteria to blood digestion and egg production in *Aedes aegypti* (diptera: culicidae) (L.). *Vectors.* **4**, 105 (2011).
48. Cirimotich CM, Dong Y, Clayton AM, Sandiford SL, Souza-Neto JA, Mulenga M, et al. Natural microbe-mediated refractoriness to *Plasmodium* infection in *Anopheles gambiae*. *Science.* 2011;332(6031):855-858.
49. Chen S, Zhao J, Joshi D, Xi Z, Norman B, Walker ED. Persistent infection by *Wolbachia* wAlbB has no effect on composition of the gut microbiota in adult female *Anopheles stephensi*. *Front Microbiol.*

2016;7:1485.

50. Wang Y, Gilbreath TM 3rd, Kukutla P, Yan G, Xu J. Dynamic gut microbiome across life history of the malaria mosquito *Anopheles gambiae* in Kenya. PLoS One. 2011;6(9):e24767.
51. Bahia AC, Dong Y, Blumberg BJ, Mlambo G, Tripathi A, Ben-Marzouk-Hidalgo OJ, et al. Exploring *Anopheles* gut bacteria for *Plasmodium* blocking activity. *Environ Microbiol*. 2014;16(9):2980-2994.
52. Ben Ami E, Yuval B, Jurkevitch E. Manipulation of the microbiota of mass-reared Mediterranean fruit flies *Ceratitis capitata* (Diptera: Tephritidae) improves sterile male sexual performance. *ISME J*. 2010;4(1):28-37.
53. Lam K, Thu K, Tsang M, Moore M, Gries G. Bacteria on housefly eggs, *Musca domestica*, suppress fungal growth in chicken manure through nutrient depletion or antifungal metabolites. *Naturwissenschaften*. 2009;96(9):1127-1132.
54. Molina-Cruz A, DeJong RJ, Charles B, Charles B, Gupta L, Kumar S, et al. Reactive oxygen species modulate *Anopheles gambiae* immunity against bacteria and *Plasmodium*. *J Biol Chem*. 2008;283(6):3217-3223.
55. Edgerton EB, McCrea AR, Berry CT, Kwok JY, Thompson LK, Watson B, et al. Activation of mosquito immunity blocks the development of transmission-stage filarial nematodes. *Proc Natl Acad Sci*. 2020;117(7):3711-3717.
56. Bassene H, Niang EHA, Fenollar F, Dipankar B, Doucouré S, Ali E, et al. 16S Metagenomic comparison of *Plasmodium falciparum*-Infected, and Noninfected *Anopheles gambiae* and *Anopheles funestus* microbiota from Senegal. *Am J Trop Med Hyg*. 2018;99(6):1489-1498.
57. Muturi EJ, Dunlap C, Ramirez JL, Rooney AP, Kim CH. Host blood-meal source has a strong impact on gut microbiota of *Aedes aegypti*. *FEMS Microbiol Ecol*. 2019;95(1):10.1093

Figures




Figure 1

Relative distribution of bacteria at phylum taxonomic level in *D. immitis* infected and uninfected *Aedes aegypti* mosquitoes. Bacterial taxa with less than 1% abundances were grouped as others.




Figure 2

Relative distribution of bacteria at A) genus and B) species taxonomic level in *D. immitis* infected and uninfected *Aedes aegypti* mosquitoes. Bacterial taxa with less than 1% abundances were grouped as others.




Figure 3

Rarefaction analysis of biological replicates rarefied at a sequenced depth of 20000. Curves from each samples was allowed to reach a plateau so as to prevent trade off of rarely represented bacterial OTUs.




Figure 4

A), Alpha diversity analysis. A), Observed OTUs (Kruskal-Wallis, $p=0.053173$) distance and B), Shannon diversity index (Kruskal-Wallis, $p=0.30952$) were both used as metrics to measure alpha diversity. Principal coordinate analysis (PCoA) of D), Bray_Curtis and E), Unweighted UniFrac distance matrixes as measures of beta diversity.




Figure 5

Principal coordinate analysis (PCoA) of A), unweighted UniFrac ($p<0.05$) and B), Bray_Curtis ($p<0.05$) distance matrixes as measures of beta diversity. Statistical significance was estimated by PERMANOVA.




Figure 6

Pattern correlation analysis and significant abundance analysis of top taxa identified in between *D. immitis* infected and uninfected mosquito. (A) SparCC correlation of top 25 bacteria Genus showing bacteria with strong positive correlation to those mosquito that were *D. immitis* infected. Box plots analysis of bacteria that shows significant differences in their abundances between infected and uninfected mosquito in the phylum B), Actinobacteria ($p = 0.036145$), and C) Firmicutes ($p = 0.005556$).




Figure 7

Box plots analysis of bacteria that shows significant differences in their abundances between infected and uninfected mosquito in the genus A), Elizabethkingia ($p = 0.01508$), B), Wolbachia, ($p = 0.011925$) C) Pseudomonas ($p = 0.015873$).

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [MosquitoD.immitisMethods.png](#)
- [SupplementaryMaterials.pdf](#)