

Molecular analysis of methicillin-resistant *Staphylococcus aureus* isolates from four *teaching hospitals in Iran: The emergence of Novel MRSA ST-SCCmec clones*

Farzaneh Firoozeh (✉ ffiroozeh@ut.ac.ir)

Alborz University of Medical Sciences

Mitra Omid

Kashan University of Medical Sciences

Mahmood Saffari

Kashan University of Medical Sciences

Hossein Sedaghat

Kashan University of Medical Sciences

Mohammad Zibaei

Alborz University of Medical Sciences

Research

Keywords: MRSA, MLST, SCCmec typing, MSCRAMMs, Genotyping

Posted Date: February 19th, 2020

DOI: <https://doi.org/10.21203/rs.2.24014/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Version of Record: A version of this preprint was published at Antimicrobial Resistance and Infection Control on July 17th, 2020. See the published version at <https://doi.org/10.1186/s13756-020-00777-8>.

Abstract

Background: The global spread of methicillin-resistant *Staphylococcus aureus* (MRSA) infections necessitates the use of validated methods for the identification and typing of this bacterium. This study aimed to determine the distribution of main molecular types of MRSA strain circulating among hospitalized patients in teaching hospitals in Isfahan and Kashan.

Methods: A total of 146 *Staphylococcus aureus* strains were isolated from patients in four teaching hospitals in Isfahan and Kashan during June 2017 to September 2018. The antibiotic resistance patterns of *Staphylococcus aureus* strains were performed by disc diffusion method. The MRSA strains were identified phenotypically and confirmed by PCR assay. The prevalence of microbial surface components recognizing adhesive matrix molecules (MSCRAMMs) genes among MRSA strains was evaluated by multiplex PCR. The genotypes of MRSA strains were determined by multi-locus sequence type and SCC mec typing.

Results: Of 146 *Staphylococcus aureus* isolates 24 (16.4%) isolates identified as MRSA strains. According to antibiotic susceptibility testing the highest resistance rates were seen to erythromycin, cefoxitin and clindamycin. All of *Staphylococcus aureus* isolates were sensitive to vancomycin whereas 3 (2.1%) isolates were resistant to linezolid. Three different SCC mec types were obtained among MRSA strains including 16 (66.7%) SCC mec type V, 3 (12.5%) SCC mec type III and 5 (20.8%) SCC mec type II. Of 24 MRSA isolates 20 (83.3%) carried MSCRAMMs genes including *eno* (70.8%), *fib* (54.1%), *cna* (25.0%), *fnbB* (16.6%), *ebps* 5 (20.8%), and *fnbA*, *bbp* and *clfA* genes were not detected. MLST analysis revealed 11 sequence types among MRSA isolates as follows: ST239, ST239, ST291, ST22, ST861, ST 889, ST8, ST59, ST343, ST772, ST6 and ST1465. Also seven MLST-based clonal complexes (CCs) were identified among MRSA strains including: CC8, CC7, CC398, CC59, CC22, CC1 and CC5.

Conclusions: A relatively high diversity was found in MRSA genotypes in Kashan and Isfahan hospitals, and seven clonal complexes were identified. Pandemic MRSA clones including CC8 and CC22 were the most prevalent clones and the novel ST types including ST1465, ST861, ST 889 and ST772 are reported for the first time in Iran in the present study. In addition the high prevalence of MSCRAMMs genes in MRSA isolates demonstrates the high potential of these strains for pathogenicity.

Background

Staphylococcus aureus (*S. aureus*) is one of the most important human pathogen that causes a wide range of infections. The methicillin-resistant *Staphylococcus aureus* (MRSA) strains are an increasingly danger for public health and can cause outbreaks in hospitals [1]. Fast identification and authentic typing of MRSA strains plays an effective role in control of infection and making therapeutic decisions [2] Although detection of the *mecA* gene is the “gold standard” for identification of methicillin-resistance, specific molecular identification of MRSA strains is based on the detection of *mecA* gene as well as *femA* (factors essential for methicillin-resistance) gene which does not cross-react with other *Staphylococcus* species [2, 3]. In order to carry out epidemiological studies of MRSA strains several typing methods are available [4]. Staphylococcal cassette chromosome *mec* (SCC*mec*) typing is a trustful typing method especially to make difference between hospital-acquired MRSA (HA-MRSA) and community-associated MRSA (CA-MRSA) [4]. MRSA strain generates a low-affinity penicillin binding protein (PBP2a) that is responsible for the resistance to beta-lactam antibiotics [2].

PBP2a is encoded by *mecA* gene, carried by the Staphylococcal cassette chromosome *mec* (SCC*mec*) which as a large mobile genetic element is integrated to a region of the chromosome of MRSA [5]. Till now 12 SCC*mec* types have been documented, and the majority of MRSA strains which isolate from HA-MRSA strains bear SCC*mec* types I, II, and III [5]. Sequencing based typing methods such as multilocus sequence typing (MLST) have recently been introduced and received a lot of attention [4]. MLST typing is a reliable method to study genetic macro- *variation* in *large populations* based on specification of allelic profile of fragments of seven house-keeping genes. [4, 6-8].

aureus produces a large number of extracellular proteins that are considered as important virulence factors and help bacteria to attach to host cells, invade the host, and immune evasion [9]. Staphylococcal infection begins with bacterial adhesion to different kinds of surfaces including epithelial cell, host tissues and medical equipments [10]. Adherence to host cells and proteins is mediated by adhesive molecules which are called microbial surface components recognizing adhesive matrix molecules (MSCRAMMs). These components are attached covalently to peptidoglycan by sortase enzymes [11]. *S. aureus* can express different potential MSCRAMMs genes including *ebpS*, encoding elastin binding protein, *eno* encoding laminin binding protein, *cna* encoding collagen binding protein, *fnbA/fnbB* encoding fibronectin binding proteins A/B, *fib* encoding fibrinogen binding protein, *clfA/clfB* encoding clumping factors A/B, and *bbp* encoding bone sialoprotein-binding protein [12, 13]. In present study we used two reliable typing methods including MLST and SCC*mec* typing to reveal and compare the common genotypes of MRSA isolates circulating in four teaching hospitals in Isfahan and Kashan. Also the prevalence of virulence and MSCRAMMs genes among these MRSA strains was evaluated.

Methods

Bacterial strains and identification

A total of 146 *S. aureus* strains were isolated from clinical samples of patients admitted to teaching hospitals in Isfahan (n=96) and Kashan (n=50) during June 2017 to September 2018. Informed consent from all patients was obtained by survey questionnaire. The samples were cultured on blood agar (Merck, Germany), and plates were incubated at 37 °C for 24 hours. *S. aureus* strains were identified by conventional microbiological methods including gram staining, and biochemical tests such as catalase, coagulase, mannitol fermentation, and DNase tests [14].

Antimicrobial resistance patterns and phenotypic isolation of MRSA strains

The antibiotic resistance patterns of *S. aureus* strains were performed by disc diffusion method according to the Clinical and Laboratory Standards Institute (CLSI) guideline [15]. The antibiotic discs were prepared from MAST Company (MAST, UK) as follows: erythromycin (E; 15µg), clindamycin (CD; 2µg), cefazolin (CZ; 30µg), cefoxitin (FOX; 30µg), linezolid (LZD; 30µg), and trimethoprim sulfamethoxazole (TS; 25µg). MICs for vancomycin were determined by microbroth dilution methodology recommended by the CLSI guideline [16]. The *S. aureus* strain ATCC 25923 was used as the control strain. For confirmation of MRSA strains, cefoxitin

(30µg) and oxacillin (1µg) disks (Mast, UK) and disc diffusion method was performed on Mueller- Hinton agar (Merck, Germany) in accordance with the CLSI [16].

DNA extraction and detection of *femA* and *mecA* genes

DNA extraction was performed by a standard phenol-chloroform method as previously described. All phenotypically isolated MRSA strains were confirmed by polymerase chain amplification (PCR), for this purpose, *femA* and *mecA* genes were amplified. For amplification of *mecA* gene forward and reverse primers were used to detect a 268 bp fragment (Table1) [17]. The PCR *thermocycling program* was: initial denaturation step at 97 °C for 6 min; followed by 30 cycles of 92 °C for 30 sec, 55 °C for 30 sec and 72 °C for 45 sec, which finally ended with final extension step at 72 °C for ten min [18]. The *S. aureus* strain COL which carry *mecA* gene was used as positive control.

The 450 bp amplicon of *femA* gene was detected by PCR assays using specific primers and thermocycler (Eppendorf master cycler®, MA) (Table 1) with *following run program*: initial denaturation at 94 °C for 5 min, 40 cycles of 94 °C for 30 sec, 55 °C for 40 sec and 72 °C for 50 sec and final extension at 72 °C for 10 min [19]. The PCR products were visualized after electrophoresis on 1% agarose gel under UV transilluminator (Bio-Rad, UK).

Multiplex PCR for *SCCmec* typing

Five main *SCCmec* types of MRSA strains were determined using multiplex PCR method with specific primers according to method previously presented by Boye et al. [20, 21]. For this purpose PCR reactions were performed in a final volume of 25 µL. Amplification was done with initial denaturation step (94 °C, 4 min), 30 cycles of denaturation (94 °C, 30 sec), annealing (55 °C, 30 sec), extension (72 °C, 60 sec), and a final extension at 72 °C for 4 min, (Table 1).

Identification of adhesive matrix molecules MSCRAMMs genes

The PCR amplification of MSCRAMMs genes were performed according to Tristan et al. [12] PCR1 was performed to amplify *bbp*, *cna*, *ebpS*, and *eno* genes and PCR2 was applied to amplify *fnbA*, *fnbB*, *fib*, and *clfA*, genes. The thermal cycling condition of multiplex PCR included an initial denaturation step (5 min at 94°C) followed by 25 cycles of denaturation (1 min at 94 °C), annealing (1 min at 55 °C), and extension (1 min at 72 °C). The reaction was terminated with a 10 min incubation step at 72 °C [12]. (Table1).

Multilocus sequence typing (MLST)

The MLST types of 24 identified MRSA strains were determined. Identification of internal fragments of 7 housekeeping genes was performed by PCR amplification according to MLST *websites* (www.mlst.net). [22] (Table 1). The final volume of PCR mixture was 25µL with thermal program including: 95 °C for 15 min followed by 30 cycles of (95° C for 1 min), (51 °C for 1 min), and (72 °C for 1 min) and finally 72 °C for 5 min [22].

Data analysis

Data were analyzed using SPSS software Version 19.0. We were used Chi square or Fisher's exact tests to determine the significance of the differences. A difference was considered statistically significant if the *P*-value was less than 0.05.

Results

This study was performed on 146 clinical samples [Female = 82 (56.2%); Male = 64 (43.8%)] collected from teaching hospitals in Isfahan (Imam Musa Kazem, Amin and Al-Zahra Hospitals) and Kashan (Shahid Beheshti Hospital). Clinical samples were collected of burn wound 56 (38.4%), eye infection 49 (33.6%), Respiratory infections 11 (7.5%), Trauma 6 (4.1%), Diabetic wound 5 (3.4%), Brain abscess 5 (3.4%), Blood 3(2%), Urine 9(6.2), and other infections 2 (1.4%).

The results of PCR amplification *mecA* gene showed among 146 studied *S. aureus* isolates 24 (16.4%) isolates identified as MRSA strains.

According to results obtained from antibiotic susceptibility testing, resistance rates to erythromycin 17 (70.8%), clindamycin 14 (58.3%) and cefazolin 13 (54.2%) was high among MRSA strains and all of *S. aureus* isolates were sensitive to vancomycin (MIC < 2 µg/mL) (Table 2).

SCC*mec* typing

Three different SCC*mec* types were obtained among MRSA strains including 16 (66.7%) SCC*mec* type V, 3 (12.5%) SCC*mec* type III and, 5 (20.8%) SCC*mec* type II (Table 3).

Identification of MSCRAMMs genes

Of 24 MRSA isolates 20 (83.3%) carried MSCRAMMs genes and in 4 (16.6%) of which MSCRAMMs genes was not identified. The prevalence of *eno*, *fib*, *cna*, *fnbB*, *ebps*, genes in MRSA isolates were 17 (70.8%), 13 (54.1%), 6 (25.0%), 4 (16.6%) and 5 (20.8%), respectively. Six, 2, and 4 isolates carried 4, 3 and 2 bands related to MSCRAMMs determinants respectively and *fnbA*, *bbp* and *clfA* genes were not detected in MRSA isolates (Table 3).

In statistical analyses a significant correlation was obtained between MRSA strains and *eno*, *cna* and *fib* genes, ($P < 0.001$).

Multilocus sequence typing (MLST)

MLST analysis revealed 11 sequence types among MRSA isolates. The results of the MLST were as follows: ST239-SCC*mec* type III (2 isolates), ST239-SCC*mec* type II (2 isolates), ST291-SCC*mec* type V (2 isolates), ST291-SCC*mec* type II (1 isolates), ST22-SCC*mec* type II (1 isolate), ST22-SCC*mec* type V (4 isolates), ST861-SCC*mec* type III (1 isolate), ST 889-SCC*mec* type II (1 isolate), ST8-SCC*mec* type V (1 isolate), ST59-SCC*mec* type V (3 isolates), ST343-SCC*mec* type V (1 isolate), ST772-SCC*mec* type V (1 isolate), ST6-SCC*mec* type V (1 isolate) and ST1465-SCC*mec* type V (3 isolates). Also seven MLST-based clonal complexes (CCs) were

identified among MRSA strains including: CC8 (41.7%), CC7 (4.2%), CC398 (12.5%), CC59 (12.5%), CC22 (20.7%), CC1 (4.2%) and CC5 (4.2%) (Table 3).

Discussion

Nowadays, different clones of antibiotic-resistant strains, such as the MRSA strains, are spreading between healthcare centers and the community [8]. This issue is one of the major challenges of health care systems in the world [8]. The goal of this study was to determine the diversity of common clones, prevalence of virulence genes and antibiotic susceptibility patterns of MRSA strains in four University Teaching Hospitals in the center of Iran. It has been documented that MRSA clones are changing, and CA-MRSA are expanding into *healthcare settings* [23]. According to different previous studies, SCC*mec* types I-III and IV-V are commonly responsible for HA-MRSA and CA-MRSA infections, respectively [24]. In the present study, in contrast SCC*mec* type V was the most prevalent SCC*mec* type identified among our clinical isolates of MRSA. However recent studies conducted in Armenia and Iran, SCC*mec* types V and IV were the most types that identified among MRSA isolates from hospitals respectively [8, 25] Also in a study by Hallin et al. [26] in Belgium, most clinical MRSA strains were belonged to SCC*mec* types IV. These results could confirm the rotation of clones between community and hospital. In MLST analysis of 24 MRSA isolates, 11 sequence types were identified and among them the ST22 and ST239 were most prevalent. All our MRSA strains with sequence type ST22 were isolated from hospitals in the city of Isfahan, and majority of them, belonged to SCC*mec* type V except one isolate which was ST22-SCC*mec* type II. These MRSA ST22 types contained limited MSCRAMMs genes, although two isolates from burn wound and eye infection demonstrated multi-drug resistant (MDR) phenotype. In a study conducted by Goudarzi et al. [24] in the city of Tehran, ST22 has been documented as the third most commonly detected MRSA clone though the mentioned strains belonged to ST22-SCC*mec* IV in comparison to our MRSA ST22 which was ST22-SCC*mec* V. These findings highlight the importance of using several typing methods simultaneously to achieve better analysis. In previous studies of MRSA and MSSA strains in the cities of Isfahan and Tehran the majority of MRSA isolates belonged to ST239 [25]. The high prevalence of ST239 has been documented from many Asian countries especially India, Pakistan, Vietnam, Thailand, Taiwan, China, Sri Lanka, and Singapore [27]. In accordance with other studies, our MRSA isolates with ST239 belonged to SCC*mec* types II and III, with similar antibiotic resistance and virulence patterns [22]. Another interesting finding was that these MRSA strains have been isolated from diverse clinical samples in different hospitals in Isfahan Province indicating circulation of these ST clones in clinical settings of our regains. During the time of current study, the third most commonly identified genotypes were ST59-SCC*mec* V, ST291-SCC*mec* V and ST1465-SCC*mec* V. The association of ST291 with HA- MRSA has been reported from Iran [28], and ST59 has been identified among both HA- MRSA and CA- MRSA strains in China [29]. Also ST59- SCC*mec* V has been documented as the predominant CA- MRSA strains in Asia [30]. However, ST1465, ST861, ST 889 and ST772 are reported for the first time in Iran in the present study. Our identified ST types belonged to seven clonal complexes, from which CC8 was the most predominant clone with MDR phenotype and was found in all the studied hospitals. The second most common clonal complex in the present study was CC22 which was not identified in Shahid Beheshti hospital in Kashan. According to MLST genotyping, *S. aureus* strains have been

documented to be highly clonal and the most of the strains belong to a limited number of closely related genotypes. Literature have shown that CC8 (Iberian clone, Brazilian clone, Archaic clone), CC5 (New York/Japanese clone, Pediatric clone) and CC22 (epidemic MRSA-15) belong to epidemic MRSA (EMRSA) clones with global distribution in hospitals [27]. In accordance with our study, CC8-V was found as predominant clone in study conducted in Armenia [8]. Also genotyping methods have been showed that CC22 as the dominant clone are among major clones circulating in Tehran, Iran [21]. ST22 strains which are related to CC22 were determined as the predominant CA-MRSA clone in central of Iran [31]. Furthermore, the ST22 clone has previously been documented as a predominant CA-MRSA clone in Germany [32]. The increasing prevalence of the ST22 clone among HA-MRSA strains in recent studies represents that these strains may have been transferred from community to hospitals. The majority (83.3%) of the MRSA isolates in our study harbored genes of binding factors and prevalence of *eno* and *fib* genes was high. The high prevalence of adhesive protein genes in MRSA isolates indicates the persistence of these strains after colonization and subsequent infection.

Conclusions

According to MLST and SCC*mec* typing results, a relatively high diversity was found in MRSA genotypes in the hospitals of Kashan and Isfahan, and seven clonal complexes were identified. However, it should be considered that only a limited number of isolates were genotyped during current study. Pandemic MRSA clones including CC8 and CC22 were the most prevalent clones and the novel ST types including ST1465, ST861, ST 889 and ST772 are reported in the present study for the first time in Iran. Furthermore, presence of momentous sequence types including ST22-SCC*mec* type II, ST239-SCC*mec* types II and III and ST291-SCC*mec* type II among important clinical samples such as burn wound and blood due to the risk of further spread of these global multi-antibiotic resistant strains is noteworthy. In addition the high prevalence of MSCRAMMs genes in MRSA isolates demonstrates the high potential of these strains for pathogenicity.

Declarations

Acknowledgements

The technical assistance of the staff of Microbiology Laboratory, of the Department of Microbiology and Immunology, School of Medicine, Kashan University of Medical Sciences are gratefully acknowledged. The results described in this paper formed part of a M.Sc. student thesis.

Funding

Nil.

Availability of data and materials

The data analyzed during the study will be available from the corresponding authors upon request.

Authors' contributions

All authors' listed in this research article have been involved and contribute for this study. FF designed the study, collected and analyzed the data, drafted the paper, and supervised the overall work. MO contributed on the data collection and statistical analysis. FF, MO, MS and HS performed the lab works. FF, MS, and MZ analyzed the dataset. Manuscript was prepared by FF, MO, and MZ. All authors read and approved the final manuscript.

Ethics approval and consent to participate

The purpose and procedures of the study were explained to all participants, and a written informed consent was obtained from all of them. The study was approved by the ethical clearance committees of the Kashan University of Medical Sciences. (IR.KAUMS.REC.1396.003).

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

References

1. Omidi M, Firoozeh F, Saffari M, Sedaghat H, Zibaei M, Khaledi A. Ability of biofilm production and molecular analysis of spa and ica genes among clinical isolates of methicillin-resistant *Staphylococcus aureus*. *BMC Res Notes*. 2020;13:19.
2. Francois P, Pittet D, Bento M, Pepey B, Vaudaux P, Lew D, et al. Rapid detection of methicillin-resistant *Staphylococcus aureus* directly from sterile or nonsterile clinical samples by a new molecular assay. *J Clin Microbiol*. 2003;41:254-60.
3. Singh SK, Anwar S, Sharma HK, Chaudhary U, Verma SK. FemA gene in Indian isolates of methicillin resistant *Staphylococcus aureus*—isolation and amplification through real time PCR. *Eur J Exp Biol*. 2014;4:90-4.

4. Huang CC, Ho CM, Chen HC, Li CY, Tien N, Fan HM, et al. Evaluation of double locus (clfB and spa) sequence typing for studying molecular epidemiology of methicillin-resistant *Staphylococcus aureus* in Taiwan. *J Microbiol Immunol Infect*. 2017;50:604-12.
5. Kwon NH, Park KT, Moon JS, Jung WK, Kim SH, Kim JM, et al. Staphylococcal cassette chromosome mec (SCCmec) characterization and molecular analysis for methicillin-resistant *Staphylococcus aureus* and novel SCCmec subtype IVg isolated from bovine milk in Korea. *J Antimicrob Chemother*. 2005;56:624-32.
6. Green SM, Marsh P, Ahmad N, Jefferies JM, Clarke SC. Characterization of community and hospital *Staphylococcus aureus* isolates in Southampton, UK. *J Med Microbiol*. 2010;59:1084-8.
7. Lee CY, Fang YP, Chang YF, Wu TH, Yang YY, Huang YC. Comparison of molecular epidemiology of bloodstream methicillin-resistant *Staphylococcus aureus* isolates between a new and an old hospital in central Taiwan. *Int J Infect Dis*. 2019;79:162-8.
8. Mkrtychyan HV, Xu Z, Yacoub M, Ter-Stepanyan MM, Karapetyan HD, Kearns AM, et al. Detection of diverse genotypes of methicillin-resistant *Staphylococcus aureus* from hospital personnel and the environment in Armenia. *Antimicrob Resist Infect Control*. 2017;6:19.
9. Ghasemian A, Peerayeh SN, Bakhshi B, Mirzaee M. The microbial surface components recognizing adhesive matrix molecules (MSCRAMMs) genes among clinical isolates of *Staphylococcus aureus* from hospitalized children. *Iran J Pathol*. 2015;10:258-64.
10. Kot B, Sytykiewicz H, Sprawka I. Expression of the biofilm-associated genes in methicillin-resistant *Staphylococcus aureus* in biofilm and planktonic conditions. *Int J Mol Sci*. 2018;19:pii: E3487.
11. Joh D, Wann ER, Kreikemeyer B, Speziale P, Höök M. Role of fibronectin-binding MSCRAMMs in bacterial adherence and entry into mammalian cells. *Matrix Biol*. 1999;18:211-23.
12. Tristan A, Ying L, Bes M, Etienne J, Vandenesch F, Lina G. Use of multiplex PCR to identify *Staphylococcus aureus* adhesins involved in human hematogenous infections. *J Clin Microbiol*. 2003;41:4465-7.
13. Walsh EJ, Miajlovic H, Gorkun OV, Foster TJ. Identification of the *Staphylococcus aureus* MSCRAMM clumping factor B (ClfB) binding site in the α C-domain of human fibrinogen. *Microbiology*. 2008;154:550-8.
14. Mahon CR, Lehman DC, Manuselis G. *Textbook of diagnostic microbiology*. 5th ed. Washington DC: Saunders; 2015.
15. Amiri A, Firoozeh F, Moniri R, Zibaei M. Prevalence of CTX-M-Type and PER extended-spectrum β -lactamases among *Klebsiella* spp. isolated from clinical specimens in the teaching hospital of Kashan, Iran. *Iran Red Crescent Med J*. 2016;18:e22260.
16. Clinical and Laboratory Standards Institute: Performance Standards for Antimicrobial Susceptibility Testing. 27th information supplement, M100-S27, Wayne PA, USA 2017.
17. Ito T, Kuwahara-Arai K, Katayama Y, Uehara Y, Han X, Kondo Y, et al. Staphylococcal cassette chromosome mec (SCCmec) analysis of MRSA. *Methods Mol Biol*. 2014;1085:131-48.
18. Pourmand MR, Hassanzadeh S, Mashhadi R, Askari E. Comparison of four diagnostic methods for detection of methicillin resistant *Staphylococcus aureus*. *Iran J Microbiol*. 2014;6:341-4.
19. Jukes L, Mikhail J, Bome-Mannathoko N, Hadfield SJ, Harris LG, El-Bouri K, Davies AP, Mack D. Rapid differentiation of *Staphylococcus aureus*, *Staphylococcus epidermidis* and other coagulase-negative staphylococci and methicillin susceptibility testing directly from growth-positive blood cultures by multiplex real-time PCR. *J Med Microbiol*. 2010;59:1456-61.

20. Boye K, Bartels MD, Andersen IS, Møller JA, Westh H. A new multiplex PCR for easy screening of methicillin-resistant *Staphylococcus aureus* SCCmec types I–V. *Clin Microbiol Infect.* 2007;13:725-7.
21. Goudarzi M, Goudarzi H, Sá Figueiredo AM, Udo EE, Fazeli M, Asadzadeh M, et al. Molecular characterization of methicillin resistant *Staphylococcus aureus* strains isolated from intensive care units in Iran: ST22-SCCmec IV/t790 emerges as the major clone. *PLoS One.* 2016;11:e0155529.
22. D'Souza N, Rodrigues C, Mehta A. Molecular characterization of methicillin-resistant *Staphylococcus aureus* with emergence of epidemic clones of sequence Type (ST) 22 and ST 772 in Mumbai, India. *J Clin Microbiol.* 2010;48:1806-11.
23. Ghanbari F, Saberianpour Sh, Zarkesh-Esfahani F, Ghanbari N, Taraghian A, Khademi F. Staphylococcal cassette chromosome mec (SCCmec) typing of methicillin-resistant *Staphylococcus aureus* strains isolated from community- and hospital-acquired infections. *Avicenna J Clin Microb Infec.* 2017;4:e42244.
24. Goudarzi M, Eslami G, Rezaee R, Heidary M, Khoshnood S, Sajadi Nia R. Clonal dissemination of *Staphylococcus aureus* isolates causing nosocomial infections, Tehran, Iran. *Iran J Basic Med Sci.* 2019;22:238-45.
25. Havaei SA, Halaji M, Vidovic S, Dillon JR, Karbalaei M, Ghanbari F, et al. Prevalence and genotyping of methicillin-resistant and – susceptible *Staphylococcus aureus* strains isolated from patients in a university hospital, Isfahan, Iran. *Jundishapur J Microbiol.* 2017;10:e13571.
26. Hallin M, Denis O, Deplano A, De Mendonça R, De Ryck R, Rottiers S, et al. Genetic relatedness between methicillin-susceptible and methicillin-resistant *Staphylococcus aureus*: results of a national survey. *J Antimicrob Chemother.* 2007;59:465-72.
27. Zafar A, Stone M, Ibrahim S, Parveen Z, Hasan Z, Khan E, et al. Prevalent genotypes of methicillin-resistant *Staphylococcus aureus*: report from Pakistan. *J Med Microbiol.* 2011;60:56-62.
28. Sedaghat H, Esfahani BN, Halaji M, Jazi AS, Mobasherizadeh S, Havaei SR, et al. Genetic diversity of *Staphylococcus aureus* strains from a teaching hospital in Isfahan, Iran: The emergence of MRSA ST639-SCCmec III and ST343-SCCmec III. *Iran J Microbiol.* 2018;10:82-9.
29. Haiying Peng, Dengtao Liu, Yuhua Ma, and Wei Gao. Comparison of community- and healthcare-associated methicillin-resistant *Staphylococcus aureus* isolates at a Chinese tertiary hospital, 2012–2017. *Sci Rep.* 2018;8:17916.
30. Noguchi N, Nakaminami H, Nishijima S, Kurokawa I, So H, Sasatsu M. Antimicrobial agent of susceptibilities and antiseptic resistance gene distribution among methicillin-resistant *Staphylococcus aureus* isolates from patients with impetigo and staphylococcal scalded skin syndrome. *J Clin Microbiol.* 2006;44:2119-25.
31. Japoni-Nejad A, Rezazadeh M, Kazemian H, Fardmousavi N, van Belkum A, Ghaznavi-Rad E. Molecular characterization of the first community-acquired methicillin-resistant *Staphylococcus aureus* strains from Central Iran. *Int J Infect Dis.* 2013;17:e949-54.
32. Linde H, Wagenlehner F, Strommenger B, Drubel I, Tanzer J, Reischl U, et al. Healthcare-associated outbreaks and community-acquired infections due to MRSA carrying the Pantone-Valentine leucocidin gene in southeastern Germany. *Eur J Clin Microbiol Infect Dis.* 2005;24:419-22.

Tables

. The primers of genes used in this study

Target	Gene	Primer sequence (5'-3')	Size of product (bp)	Reference
<i>femA</i>	<i>femA</i>	F: TGCTATCCACCCTCAAACAGG R: AACGTTGTAACCACCCCAAGA	450	[8]
<i>mecA</i>	<i>mecA</i>	F: TGCTATCCACCCTCAAACAGG R: AACGTTGTAACCACCCCAAGA	286	[19]
SCC <i>mec</i>	<i>ccrA2</i>	F: TAAAGGCATCAATGCACAAACACT R: AGCTCAAAGCAAGCAATAGAAT	937	[20]
	<i>ccrC</i>	F: CCTTTATAGACTGGATTATTCAAAATAT R:CGTCTATTACAAGATGTTAAGGATAAT	518	
	<i>IS1272</i>	F: GCCACTCATAACATATGGAA R: CATCCGAGTGAAACCCAAA	415	
	<i>mecA-IS431</i>	F: TATACCAAACCCGACAACACTAC R: CGGCTACAGTGATAACATCC	359	
MSCRAMMs genes	<i>bbp</i>	F: AACTACATCTAGTACTCAACAACAG R: ATGTGCTTGAATAACACCATCATCT	575	[11]
	<i>cna</i>	F: GTCAAGCAGTTATTAACACCAGAC R: AATCAGTAATTGCACCTTTGTCCACTG	423	
	<i>eno</i>	F: ACGTGCAGCAGCTGACT R: CAACAGCATYCTTCAGTACCTTC	302	
	<i>ebpS</i>	F: CATCCAGAACCAATCGAAGAC R:CTTAACAGTTACATCATCATGTTTATCTTTG	186	
	<i>fnbA</i>	F: GTGAAGTTTTAGAAGGTGGAAAGATTAG R: GCTCTTGTAAGACCATTTTTCTTCAC	643	
	<i>fnbB</i>	F: GTAACAGCTAATGGTCGAATTGATACT R: CAAGTTCGATAGGAGTACTATGTTC	524	
	<i>fib</i>	F: CTACAACTACAATTGCCGTC AACAG R: GCTCTTGTAAGACCATTTTTCTTCAC	404	
	<i>clfA</i>	F: ATTGGCGTGGCTTCAGTGCT R: CGTTTCTTCCGTAGTTGCATTTG	292	
	Housekeeping genes	<i>arc</i>	up: TTGATTCACCAGCGCGTATTGTC dn: AGGTATCTGCTTCAATCAGCG	456
<i>aro</i>		up: ATCGGAAATCCTATTTACATTC dn: GGTGTTGTATTAATAACGATATC	456	
<i>glp</i>		up: CTAGGAAGTCAATCTTAATCC dn: TGGTAAAATCGCATGTCCAATTC	465	

<i>gmk</i>	up: ATCGTTTTATCGGGACCATC	429
	dn: TCATTAAC TACAACGTAATCGTA	
<i>pta</i>	up: GTTAAAATCGTATTACCTGAAGG	474
	dn: GACCCTTTTGTGAAAAGCTTAA	
<i>tpi</i>	up: TCGTTCATTCTGAACGTCGTGAA	402
	dn: TTTGCACCTTCTAACAATTGTAC	
<i>ygi</i>	up: CAGCATA CAGGACACCTATTGGC	516
	dn: CGTTGAGGAATCGATACTGGAAC	

Table 2 Antibiotic resistance rate in MRSA and MSSA strains

Antibiotic (%)	MRSA (%) N = 24	MSSA (%) N = 122	Total (%) N = 146
Erythromycin	17 (70.8)	28 (22.9)	45 (30.8)
Clindamycin	14 (58.3)	8 (6.6)	22 (15.1)
Cefazolin	13 (54.2)	1 (1)	14 (9.6)
Linezolid	1 (4.2)	2 (1.6)	3 (2.1)
Trimethoprim Sulfamethaxazole	8 (33.3)	6 (4.9)	14 (9.6)

Table 3 The results of resistance patterns, MSCRAMMs genes, MLST and SCCmec types among MRSA isolates from clinical samples in different hospitals (N = 24)

Source	Hospital	Ward	Antibiotic resistance pattern	MSCRAMMs	SCCmec type	Clonal Complex	ST ^d
Blood	SB ^a	Emergency	E,CD,CZ,TS, FOX	<i>eno, can, fib, FnbB</i>	II	8	239
Burn wound	IMK ^b	Skin	E, CD, CZ, TS, FOX	<i>eno, can, fib, FnbB</i>	III	8	861
	IMK	Skin	E, CD, CZ, FOX	<i>eno, can, fib</i>	III	7	239
	IMK	Skin	CZ, E, FOX	<i>eno</i>	V	8	8
	IMK	Skin	LZD, FOX	<i>eno, ebp, fib, FnbB</i>	V	398	291
	IMK	Skin	E, CD, TS, FOX	<i>Eno, fib</i>	V	59	59
	IMK	Skin	FOX	<i>eno, cna</i>	V	59	59
	IMK	Skin	FOX	-	II	398	291
	IMK	Skin	FOX	-	II	22	22
	IMK	Skin	E, CD, CZ, FOX	<i>eno, cna, ebp, fib</i>	II	8	239
	IMK	Skin	TS, CZ, FOX	<i>eno</i>	V	8	1465
	IMK	Skin	TS, FOX	<i>eno</i>	V	22	22
	IMK	Skin	FOX	<i>eno, cna, ebp, fib</i>	V	1	772
	IMK	Skin	E, CD, CZ, TS, FOX	<i>eno</i>	V	22	22
	IMK	Skin	E, FOX	<i>eno, ebp, fib</i>	V	5	6
	IMK	Skin	E, CD, TS, CZ, FOX	<i>Eno, fib</i>	V	8	1465
Eye infection	SB	Emergency	E, CD, TS, FOX	<i>eno</i>	II	8	889
	Amin	Emergency	E, CD, CZ, FOX	<i>fib</i>	III	8	239
	Amin	Emergency	E, CD, CZ, FOX	-	V	8	1465
	Amin	Emergency	E, CD, CZ, FOX	<i>fib</i>	V	59	59
	Amin	Emergency	CZ, CD, CZ, FOX	-	V	22	22
	SB	Emergency	E, CD, CZ, FOX	<i>eno</i>	V	398	291
Brain abscess	AZ ^c	ICU	CD, E, FOX	<i>eno, ebp, fib, FnbB</i>	V	8	343
Respiratory infection	AZ	ICU	E, FOX	<i>fib</i>	V	22	22

Imam Musa Kazem hospital; ^bShahid Beheshti; ^cAl Zahra; ^dSequence type

