Development of a tool for identifying and addressing prioritised determinants of quality improvement projects led by healthcare professionals: a mixed-methods study

Anne van Tuijl (✉ anne.vantuijl@radboudumc.nl)

Radboudumc <https://orcid.org/0000-0002-5515-3966>

Hub C. Wollersheim

Radboudumc IQ healthcare

Cornelia R.M.G. Fluit

Radboudumc

Petra. J. van Gorp

Radboudumc Afdeling Interne Geneeskunde

Hiske Calsbeek

Radboudumc IQ healthcare

Research

Keywords: Determinants of practice, quality improvement, healthcare professionals, tool development

Posted Date: August 6th, 2020

DOI: <https://doi.org/10.21203/rs.2.24055/v2>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published on October 23rd, 2020. See the published version at <https://doi.org/10.1186/s43058-020-00082-w>.

Abstract

Background: Several frameworks have been developed to identify essential determinants for healthcare improvement. These frameworks aim to be comprehensive, leading to the creation of long lists of determinants that are not prioritised based on being experienced as most important. Furthermore, most existing frameworks do not describe the methods or actions used to identify and address the determinants, limiting their practical value. The aim of this study is to describe the development of a tool with prioritised facilitators and barriers supplemented with methods to identify and address each determinant. The tool can be used by those performing quality improvement initiatives in healthcare practice.

Methods: A mixed-methods study design was used to develop the tool. First, an online survey was used to ask healthcare professionals about the determinants they experienced as most facilitating and most hindering during the performance of their quality improvement initiative. A priority score was calculated for every named determinant, and those with a priority score ≥ 20 were incorporated into the tool. Semi-structured interviews with implementation experts were performed to gain insight on how to analyse and address the determinants in our tool.

Results: The 25 healthcare professionals in this study experienced 64 facilitators and 66 barriers when performing their improvement initiatives. Of these, 12 facilitators and nine barriers were incorporated into the tool. Sufficient support from management of the department was identified as the most important facilitator, while having limited time to perform the initiative was considered the most important barrier. The interviews with 16 experts in implementation science led to various inputs for identifying and addressing each determinant. Important themes included maintaining adequate communication with stakeholders, keeping the initiative at a manageable size, learning by doing and being able to influence determinants.

Conclusions: This paper describes the development of a tool with prioritized determinants for performing quality improvement initiatives with suggestions for analysing and addressing these determinants. The tool is developed for those engaged in quality improvement initiatives in practice, so in this way it helps to bridging the research to practice gap of determinants frameworks. More research is needed to validate and develop the tool further.

Contributions To The Literature

- In the existing literature, there are various determinant models to understand the influence of barriers and facilitators on implementation outcomes.
- Although the literature agrees to a large extent on which determinants are important, it is less clear which determinants are experienced as most important by those engaged in quality improvement in the field, and how to find methods to analyse and address these determinants.

- This study describes a tool of prioritised facilitators and barriers for healthcare quality improvement initiatives, with suggestions for analysing and addressing them. This tool contributes to bridging the research to practice gap by taking a serious look at how practical research-centric determinant frameworks are for those engaged in quality improvement projects in the field.

Background

The importance of contextual determinants for the success of quality improvement (QI) initiatives is widely accepted(1, 2). The increased interest in the context where QI initiatives are implemented has resulted in the publication of several frameworks of determinants in the literature (e.g., Tailored Implementation for Chronic Disease (TICD(2)), Consolidated Framework for Implementation Research (CFIR(3)), Model for Understanding Success in Quality (MUSIQ(4)) and Measurement Instrument for Determinants of Innovations (MIDI(5)). These frameworks describe general classes of determinants, which are independent variables that prevent or enable implementation outcomes, such as changes in healthcare professional behaviours(2, 6). These determinants are also described as barriers and enablers, barriers and facilitators or problems and incentives(2). The aim of these frameworks is to help implementation researchers and people responsible for QI initiatives in healthcare to identify determinants, enabling them to design and execute more effective implementation strategies to support implementation of the QI initiative (6).

In this article, we use QI to refer to improvements in healthcare related to more effective, safe, efficient, timely and patient-centered care (7). QI initiatives are defined as e.g. new procedures, technologies, guidelines, protocols, programs that are firmly established as being able to contribute to more effective, safe, efficient, timely and patient-centered care(8). With the term implementation we refer to a planned process and systematic introduction of these initiatives; the aim being that these are given a structural place in professional practice, in the functioning of organizations or in the health care structure(7). Implementation strategies in healthcare can be defined as methods or techniques used to enhance the adoption, implementation, and sustainability of a clinical practice or programme(9).

Most of the existing generic frameworks of determinants aim to be comprehensive, including all domains and determinants (2). These frameworks are based on the idea that implementation is a multidimensional phenomenon taking place in complex settings with multiple interacting influences within and across different types of determinants(6). The downside is that these frameworks lead to long lists of determinants (e.g., TICD contains 57 determinants,), which can be challenging to use in practice; it would be difficult to analyse and address all of these determinants during implementation (10). For healthcare professionals planning to implement an improvement initiative , a shorter tool of determinants that are prioritized based on experienced importance by those engaged in QI in the field would be useful; however, to our knowledge, no such tool yet exists

In most general determinant frameworks, it is unclear whether the barriers and facilitators are the most important determinants within a specific context and for a specific population. A systematic review of 12

available frameworks and taxonomies of determinants of practice revealed that most were based on literature reviews or developed using brainstorming and consensus processes(2). A recent scoping review(11) of determinant frameworks showed that many were developed based on earlier frameworks, which could lead to a narrow approach to exploring and understanding determinants. Although some frameworks are based on the author's own implementation experiences(11), it is unclear to what extent these determinants are experienced as important by other healthcare professionals responsible for the implementation of healthcare improvements.

The use of determinant frameworks in practice is also restricted by the limited guidance available on how to analyse determinants and how to match determinants with implementation strategies(8, 12). Most frameworks of implementation strategies (e.g., Expert Recommendations for Implementing Change (ERIC)(13)) have not linked their implementation strategies to determinants. This can lead to a mismatch between identified determinants and strategies(14), making the effect of implementation strategies variable(8). A systematic review of 32 studies(15) showed that strategies tailored to determinants were more effective than those that were not; however, the methods used to select implementation strategies were often not well described(15). Enhancing the link between identified determinants and implementation strategies is therefore a priority for implementation science(16).

To our knowledge, there is no relatively short general determinant tool with prioritized determinants based on the experienced importance according to healthcare professionals who are responsible for QI initiatives in their own practice. Such a tool could support these professionals to have a structured discussion regarding which determinants are most important to consider during different moments in the implementation process of the QI initiative and to find ways to analyse and address these determinants.. The aim of this study was therefore to develop a tool of prioritized determinants based on the experienced importance according to healthcare professionals leading QI initiatives in their own workplace. In this way, this tool helps to bridging the research to practice gap by taking a serious look at how practical research-centric determinant frameworks are for those engaged in QI initiatives in the field. Therefore, the tool was developed to be practical in the field rather than comprehensive, resulting in a relatively short list of determinants with suggestions for how to analyse and address them. In this article, we describe the development of this tool.

Methods

Setting

Healthcare professionals and implementation experts participated in this study. Both groups were involved in a two-year half-time post-initial scientific master's programme (subsequent to an initial master's) on QI in healthcare. The program trains healthcare professionals with an academic background to become leaders in the evidence-based improvement of healthcare quality and safety. Professionals are enrolled in 12 learning modules (see Figure 1, which shows the core elements of the programme). The implementation experts are teachers and supervisors in these modules and have different specialties,

such as leadership or patient involvement in QI . Professionals work intensively on their personal (leadership) development in QI through portfolios and coaching. Together, these educational interventions support professionals to perform a QI initiative at their workplace during their master's programme. Because the initiatives are carried out in the context of a teaching programme, they are to some extent standardised (e.g., all professionals receive methodological support, all initiatives are performed in a hospital setting and are led by healthcare professionals, and all theses are evaluated using the Standards for Quality Improvement Reporting Excellence (SQUIRE) guidelines(17)), making them valuable sources for the development of this tool.

Design

We used a mixed-methods design(18) to develop the tool. First, we performed a cross-sectional study using an online survey among healthcare professionals from the master's programme, in which they were asked about the determinants they experienced as most facilitating and hindering during the performance of their QI initiative. After including the determinants in our tool, we performed semi-structured interviews with implementation experts about how healthcare professionals can analyse and address the determinants identified in their initiative . We used the Strengthening of Reporting of Observational Studies in Epidemiology guidelines (STROBE) and the Consolidated Criteria for Reporting Qualitative Research guidelines (COREQ) when designing our study. For more details, see additional files 1 and 2 [see Additional file 1 and 2].

Part 1: Survey

Development

Two cohorts of healthcare professionals following the master's programme (N = 41) were asked to fill in an online survey in which they were retrospectively asked to express the five most facilitating and five most hindering determinants they experienced during the performance of their QI initiative. s. See additional file 3 for the complete survey (in Dutch) [see Additional file 3].

To support healthcare professionals with which determinants to name in their top 5, we provided a pre-specified list with an overview of determinants that are described in the literature. This list was based on existing determinants models. After reviewing and considering the literature on determinants models, we based our pre-specified list on two models: : MUSIQ(4) and the TICD model(2). To the best of our knowledge, MUSIQ is the only model that includes determinants related to team-level QI , while the TICD model is based on a systematic review of other determinant frameworks (including for example the Consolidated Framework For Implementation Research), not including MUSIQ. We therefore assume that the combination of both models provides an overview of all unique determinants.

First, the researcher (AvT) combined the two models to gain insight into the unique and overlapping determinants. For the MUSIQ model, we used the questionnaire that was developed to measure contextual determinants(19). For the TICD model, we used the TICD tool that was developed to facilitate

the use of the model(2). The researcher (AvT) and one member of the research team (HW) scored all determinants (57 TICD determinants, 39 MUSIQ determinants) by three criteria: relevance (the determinant should be of relevance for performing a QI initiative in healthcare), applicability (the determinant should be applicable across various relevant settings and types of improvement initiatives) and recognition (the determinant must be easily understood by healthcare professionals)(2). Of the selected determinants, 32 came from the MUSIQ questionnaire, 22 from the TICD tool and five were identified in both tools. Two determinants were formulated by discussion with the research group based on their relevance to our context. We formulated all the included determinants as both facilitators and barriers, resulting in two lists of 61 determinants each. A flowchart of the selection process can be found in Figure 2.

The translation and adaptation of the determinants for the pre-specified list was completed by researcher AvT. The list was extensively discussed with the entire research group. The determinants were organised according to the MUSIQ model levels: those of the external environment, organisation, microsystem and QI team. We also added two levels from the TICD model: the level of the patient and of the QI initiative itself.

Procedure

In 2014–2018, a total of 46 healthcare professionals with backgrounds in various medical and nursing specialties participated in the first two cohorts of the master's programme. We included all healthcare professionals who started performing a QI initiative during their master's. Professionals who dropped out of the programme and did not begin an initiative and those who explicitly stated that they did not want to participate were not recruited. In total, 41 healthcare professionals were recruited by e-mail to participate.. Participation in this survey was voluntary. Informed consent was implied by the overt action of completing the online questionnaire after reading the information letter.

We administered the survey after the professionals, whether or not successfully, implemented their QI initiative. In the survey, professionals were retrospectively asked to name the five most facilitating and five most hindering determinants s they experienced at any point during the performance of their QI initiative. Professionals could select determinants in two ways; choosing from the pre-specified list of facilitators and barriers and/or naming self-experienced determinants that were not included in this list. The survey could be completed in approximately 10 minutes.

Analyses

For the analyses, the participants' self-experienced determinants were carefully evaluated by the researcher (AvT) to assess whether they could be merged with determinants from the pre-specified list or with determinants reported by other participants. Self-experienced determinants with the same meaning as a prespecified determinant were accommodated within this determinant. Self-experienced determinants with the same meaning as another self-experienced determinant were combined,

sometimes leading to a reformulation of the self-experienced determinant. The combined determinants were discussed with another member of the research team until consensus was reached (HW and/or HC).

A priority score was calculated for each of the top-five determinants named by the respondents. This score consisted of the sum of the ranking in the top five (i.e., a determinant ranked in first place scored five points, a determinant ranked fifth place scored one point), multiplied by the number of times a determinant was placed in a top five by professionals. Determinants with a priority score ≥ 20 were included in our tool.

Part 2: Interviews

Questions

After prioritizing each determinant and including the determinants with a priority score ≥ 20 in our tool, we performed semi-structured interviews with implementation experts. The purpose of these interviews was to gain insight on how to analyse and address the determinants in our tool, not further triangulation of which determinants are important. The interview guide consisted of open-ended questions about how healthcare professionals could be assisted to analyse determinants, use facilitators and confront barriers when performing a QI initiative. We also asked for general comments on the tool. The interview guide was pilot-tested with a member of the research team (HC) and adapted where necessary.

Procedure

We recruited 29 implementation experts from eight Dutch universities. They all had a teaching role in the master's programme and possessed different expertise regarding implementation science. We used purposive sampling to select implementation experts with varied expertise in this field. All 29 implementation experts were approached by e-mail to participate, and were provided with information about the study. Oral consent was recorded. The primary researcher interviewed all implementation experts. All interviews were audio-taped and field notes were made.

During the interview, implementation experts were asked to choose two facilitators and two barriers from the tool, on which they wanted to provide input. Input was based on the implementation experts' theoretical knowledge and practical experiences with QI. As an increasing number of interviews were held, it became clear that some determinants had received more input than others. Therefore, as we conducted more interviews, we asked implementation experts if they could also advise on the determinants which had previously received less input.

Analyses

All interviews were transcribed verbatim by the primary researcher (AvT), and from each transcript an extensive summary was made to facilitate the analyses. The transcript summaries were analysed in pairs using principles from thematic analysis(20). The primary researcher (AvT) analysed all summaries and two researchers (HW and HC) with implementation expertise also analysed half of the summaries each.

This involved carefully reading the summaries before discussing the meaning of the input to identify and develop themes, which were reformulated into suggestions for analysing and addressing the determinants in the tool.

Results

Survey

Of the 41 professionals who received the survey, 28 responded (68% participation rate). Most participants had a professional background as a physician (50%), followed by a background as a nurse (36%). Table 1a and 1b show characteristics of the respondents.

(Insert Table 1a and 1b) We found 130 unique determinants named by respondents, consisting of 64 facilitators (53%) and 66 barriers (47%). Facilitators related to the QI team were most commonly reported, while the most commonly reported barriers were related to healthcare organisations. Of the 64 facilitators, 32 (50%) were self-reported by the professionals and thus outside our pre-specified list of determinants. Of the 66 barriers, 34 (52%) were self-reported. Additional file 4 provides all determinants that were ranked in the professionals' top five lists [see Additional file 4].

Twelve facilitators were found to have a priority score ≥ 20 (min. 24, max. 288). Most facilitators were related to the level of the department. The availability of management support was identified as the most important facilitator. We found nine barriers with a priority score ≥ 20 (min. 20, max. 767). Most barriers were experienced at the organisation level. Healthcare professionals feeling they had insufficient time to perform the QI initiative was identified as the most important barrier. Table 2 provides a list of the 12 facilitators and nine barriers with their priority scores.

Interviews

We interviewed 16 implementation experts (response rate = 55%). Reasons for non-participation were needing to meet other deadlines and overlapping expertise. Five interviews were conducted face to face at the professional's workplace, while the remainder were held over the telephone. The interviews took on average 34 minutes (range 20–57 minutes). Table 3 shows the areas of expertise of the implementation experts.

For each determinant with a priority score ≥ 20 , we identified several suggestions on 1) how to assist healthcare professionals in analysing the determinant and 2) how to use a facilitator or tackle a barrier. The suggestions are included in the tool under the headings 'analyse determinant' and 'address determinant'; for example, Table 4 shows the suggestions included in the tool for analysing and addressing a facilitator and a barrier.

In addition to providing input on how to analyse and address the determinants in our tool, experts frequently made general comments about performing QI initiatives. These comments led to the

identification of four themes: communication, keep it small, influence and concern, and learning by doing. Box 1.1 provides a description of each theme.

Box 1.1 Themes identified from interviews with implementation experts on how to analyse and address determinants for performing QI initiatives

Communication: It seems important to communicate about the initiative to different stakeholders at different levels both inside and outside the organisation. Depending on the stakeholder, this communication differs in terms of how much information the project leader provides, which information is provided and the way in which the information is provided. Experts talked about *'speaking the language of the stakeholder'*. It is important to express what is expected from stakeholders, to periodically inform them about progress and to listen carefully to what they say about the project. Creating room for stakeholders to ask questions about the initiative is also considered important. A communication plan can facilitate the structuring of this communication.

Keep it small: Experts emphasised the importance of keeping an initiative small, both in terms of the goal(s) and the participants and departments to be included. Project leaders should look at the initiative as one part of the whole QI process in their institution and try to improve something in their microsystem. Successful implementation in this context can lead to future spread and sustainability.

Influence and concern: This theme relates to focusing on determinants that can be influenced by healthcare professionals as improvement leaders, rather than determinants that lie beyond their influence. Determinants that can only be addressed with disproportionate effort will result in a disbalance between effort and result, which can lead to an energy leak.

Learning by doing: Experts frequently noticed that the implementation of an QI initiative is a process that is equally or even more important than the beneficial outcomes. One expert said: *'the journey itself is much more important than your end goal. Projects can fail in terms of outcomes, but you can learn so much of that by putting value on the process'*. By seeing the implementation process as a learning process, healthcare professionals can learn for future projects.

All facilitators and barriers with a priority score >20, together with suggestions for analysing and addressing the determinants, were combined in a practical tool. Based on the major themes that emerged from interviews with implementation experts, a distinction is made between determinants that lie within the control of healthcare professionals (proximal) and determinants that lie beyond their control (distal). The full tool can be found in additional file 5 [see Additional file 5].

Discussion

In this study, we developed a tool of 12 prioritised facilitators and nine prioritised barriers based on the experienced importance according to healthcare professionals performing QI initiatives in practice. For every determinant in the tool, , suggestions on how to analyse and address these determinants are

provided . This tool can be used before, during and after the implementation of a QI initiative to guide discussion on which determinants are important to consider. The tool therefore helps healthcare professionals to learn from failures and successes, which can be used in future initiatives . Most facilitators in our tool are at the level of the department, while most barriers are at the organisational level. We identified support from the departmental management staff as the most important facilitator, while lacking the proper time to perform an initiative was the most important barrier. Methods for analysing and addressing each determinant are provided, based on interviews with implementation experts. Although not every determinant can be directly influenced by professionals, good communication with stakeholders, keeping the project small and learning from implementation are important overall recommendations for performing QI initiatives .

Differences with other determinant models

In contrast to most determinant models in the literature, our tool makes an explicit distinction between barriers and facilitators. We assume that the presence of a determinant during implementation does not necessarily have to be equally as helpful as its absence would be prohibitive, and vice versa. By analysing the professionals' top five most important experienced facilitators and barriers, we conclude that this assumption is valid because we found that determinants that are experienced as most important facilitators were different than determinants that are experienced as most important barriers. Our tool therefore contains separate lists of prioritised facilitators and barriers. We hope that this distinction supports professionals in making more informed choices on which facilitators to use and which barriers to address during the implementation of an QI initiative.

Because identifying determinants is the first step to challenge or use determinants, we not only include a list of prioritised facilitators and barriers but also provide practical suggestions on how to analyse determinants, use facilitators and address barriers. Current determinant models = often fail to provide a link between determinants and the strategies to address them(8, 15, 16). A recent study of which implementation strategies in the ERIC model would best address contextual barriers from the CFIR found that respondents had varying opinions regarding which implementation strategies best addressed each contextual barrier(12). This result can be explained by the fact that professionals chose implementation strategies without fully understanding the determinant and that most implementation strategies are limited in their specification, poorly described and 'package' approaches consisting of multiple poorly understood elements(21). Our tool first helps to address these concerns by including suggestions how to analyse the determinants. By applying those analysing methods, there will be a better understanding of the determinant leading to more informed choices on selecting implementation strategies. Thereby, the practical suggestions in our tool to address the determinants are different to those of other implementation strategies in that they are concrete and focus on one specific determinant. Future research should strengthen our suggestions to address the determinants by enhancing the evidence for these suggestions. Also linking specific behavioural change strategies, based on theoretical constructs, to the determinants in our tool would be a valuable future step. . .

Reflection on the barriers and facilitators in the tool

We identified receiving sufficient support from management as the most important facilitator. Support from management helps to formally confirm an initiative, for example by integrating it in the policy statements of the department(22) or by providing resources(19). In an exploratory analysis of the MUSIQ model, researchers found that microsystem determinants (e.g., department-level factors) have direct effects on the success of QI initiatives(19). Microsystem leadership (similar to our determinant 'management support of department') was not found to have a direct influence on success; however, this determinant was found to be directly influenced by QI team leadership, which in turn had one of the strongest direct effects on measures of success. Our tool is based on the experiences of QI team leaders, who, it could be argued, are highly influenced by their departmental management team.

Analysis of the MUSIQ model also found that most determinants related to the QI team had a direct effect on success because this team is responsible for guiding the implementation(19). Our results also show that facilitators at the level of the QI team were most often reported in professionals' top 5 but were not included in our tool due to their priority score (< 20). This result could mean that the various individual determinants related to the QI team are independently not experienced as most important but that these determinants together could make the total synergistic importance of this domain very high. Future research on this tool should study how and why the QI team is important in the performance of a QI initiative.

By providing a tool of prioritised facilitators and barriers with practical suggestions for how to analyse and address determinants, we intend to help healthcare professionals to have structured discussions regarding which determinants are important to consider during different moments in the implementation process of the QI initiative. Using the tool prior to the implementation can help to identify potential barriers and facilitators to implementation, adapt the initiative before implementation and consider how learning can be supported. During implementation, our tool can serve to monitor the implementation and to identify and address determinants. After implementation, the tool can help to reflect upon which determinants influenced the implementation. Although we recommend to use our tool during different phases of the implementation, we did not take into account that the experienced importance of determinants will vary throughout the lifetime of a QI initiative in the development of the tool. We recommend future research to ask professionals at different moments during the implementation of the initiative which determinants they experience as most important.

Our determinants are prioritised based on the experiences of healthcare professionals who were leading a QI initiative in the hospital setting. It is assumed that the higher the priority score, the more important this determinant is in the implementation of the initiative. However, as every initiative is unique and every context is different, it's not definite that the determinants we found will also be reflected in other contexts or initiatives. It's therefore important that QI teams have a structured discussion about the determinants in our tool to discuss whether the determinants are reflected in their context.

Although the science of measuring determinants is immature⁽¹⁾, the column 'analysis' in the tool will be helpful for this task. During those discussions professionals should also think about other possible determinants that are important in their context and add them to this tool. Additional file 4 which provides all determinants that were ranked in professionals' top five, could support these discussions.

Knowing which determinants could potentially influence the implementation does not mean that the determinant can be addressed by healthcare professionals.; interviews with implementation experts showed that some determinants lie beyond the control of healthcare professionals. An organisational culture supportive of QI is an example of a facilitator, but is difficult to influence directly and in the short term by healthcare professionals. However, this does not imply that these determinants should be fully ignored by healthcare professionals. We recommend professionals to give priority on addressing the determinants that lie within their control yet at the same time do not lose sight on the determinants beyond their control⁽³³⁾. First addressing the determinants within practitioners' control will help to build motivation and commitment to the QI initiative. These 'early wins' could also help to address or use the determinants that are beyond their influence by stimulating the organization to influence those determinants. Also, these determinants that lie beyond practitioners' control should be kept on the horizon, knowing that they won't change quickly but by the use of adaptive leadership behaviours and other strategies small steps in the right direction can be obtained

Because our tool is one of the first that includes the sphere of influence, we recommend more research on which determinants can be influenced by whom and under which circumstances.

Limitations

Our study has several limitations. The first limitation has to do with the context dependency of our tool, limiting the generalizability. The QI initiatives s at the basis of this study were performed within the context of an educational programme, making the results susceptible to selection bias. It is possible that healthcare professionals performing a QI initiative outside the context of an educational programme will experience different determinants. Although this possibility, our initiatives are performed using the principles from experiential learning which means that real-life experiences in the context relevant to learners own future career are used, making the difference with initiatives performed outside the educational context relatively small^(34, 35). Thereby, all of the included improvement initiatives are performed within the Dutch hospital setting. We propose future research to test this tool in other contexts, to see whether the determinants we found are also experienced as most important in those contexts. Also, our tool is based on a small number of QI initiatives (N = 28), which may further limit its generalisability. The initiatives were performed in a wide range of hospitals across the Netherlands however, including all academic hospitals in the country. Another limitation is that there could be a potential bias in the participants 'top 5 barriers and facilitators. Respondents could have identify the easier well-known determinants more easy and overlooked the less known determinants that could potentially have a greater impact. However , we tried to minimize this bias by letting participants choose

determinants from our pre-specified list of determinants which included also the less obvious determinants.

Finally, our tool has not been prospectively validated. Further evaluation and modification is needed, including feedback from a broad range of healthcare professionals about their experiences with the tool. This tool is not a finished product and will benefit from further adjustments and developments.

Conclusion

To our knowledge, this is the first tool of prioritised determinants based on the experiences of healthcare professionals in practice. Our tool is not a substitute for the comprehensive well-developed frameworks that exists but is complementary to these frameworks in that it helps bridging the research to practice gap by including those determinants that are experienced as most importance during the performance of a QI initiative by those engaged in QI in the field. The tool consists of nine barriers and 12 facilitators for the performance of QI initiatives. For each of these barriers and facilitators, suggestions on how to analyse and address the determinant are given based on input from implementation experts. This tool will facilitate healthcare professionals in the systematic reflection on determinants for QI initiatives s; however, additional research is needed to further develop and validate our tool.

Declarations

Ethics declarations

Ethics approval and consent to participate

Research ethics approval was sent to the Research Ethics Committee of the Radboud university medical centre. The committee judged that ethical approval was not required under Dutch national law (2018-4918). Participation in this study was fully voluntary and anonymity was guaranteed by the use of non-identifiable data. For the survey, informed consent was implied by the overt action of completing the online questionnaire after reading the information letter. Oral consent was recorded from the participants of the interviews.

Consent for publication

Not applicable

Availability of data and materials

The datasets that support the findings of this study are not publicly available due to information that could compromise research participant consent and privacy but can be made available from the corresponding author (AvT) with appropriate precautions and upon reasonable request. *Competing interests*

HW, HC and PG have roles in the master's programme on which the participants for this study are employed; HW was the former director of the master's, HC is co-ordinator of the programme and PG is current director of the master's. The master's programme is partly financed by the Dutch Federation of University Medical Centers, which is the same organisation that financed this study. To ensure confidentiality of participants, identifying data about the individuals were only available to the main researcher AvT, who has no role in the master's programme nor any association with the Dutch Federation of University Medical Centers.

Funding

This work was carried out under the auspices of the Consortium Quality of Care of the Dutch Federation of University Medical Centers (NFU) and the National Health Care Institute.

Authors contributions

AvT collected the data, analysed the data and drafted the manuscript. HW analysed the data and contributed to early drafting. HC also analysed the data. All researchers participated in the design of the study and made important contributions to the theoretical approach and interpreting insights. All authors read, revised and approved the final manuscript.

Acknowledgements

We thank colleagues of the Dutch Federation of University Medical Centers and the National Health Care institute for their input and agreement to develop this tool and their ongoing participation.

Abbreviations

QI: Quality improvement

TICD: Tailored Implementation for Chronic Disease

CFIR: Consolidated Framework for Implementation Research

MUSIQ: Model for Understanding Success in Quality

MIDI: Measurement Instrument for Determinants of Innovations

ERIC: Expert Recommendations for Implementing Change

SQUIRE: Standards for Quality Improvement Reporting Excellence

References

1. Lau R, Stevenson F, Ong BN, Dziedzic K, Treweek S, Eldridge S, et al. Achieving change in primary care—effectiveness of strategies for improving implementation of complex interventions: systematic review of reviews. *BMJ open*. 2015;5(12):e009993.
2. Flottorp SA, Oxman AD, Krause J, Musila NR, Wensing M, Godoycki-Cwirko M, et al. A checklist for identifying determinants of practice: a systematic review and synthesis of frameworks and taxonomies of factors that prevent or enable improvements in healthcare professional practice. *Implementation Science*. 2013;8(1):35.
3. Damschroder LJ, Aron DC, Keith RE, Kirsh SR, Alexander JA, Lowery JC. Fostering implementation of health services research findings into practice: a consolidated framework for advancing implementation science. *Implementation science*. 2009;4(1):50.
4. Kaplan HC, Provost LP, Froehle CM, Margolis PA. The Model for Understanding Success in Quality (MUSIQ): building a theory of context in healthcare quality improvement. *BMJ Qual Saf*. 2012;21(1):13-20.
5. Fleuren M, Paulussen T, Van Dommelen P, Van Buuren S. Measurement instrument for determinants of innovations (MIDI). Leiden: TNO. 2014.
6. Nilsen P. Making sense of implementation theories, models and frameworks. *Implementation science*. 2015;10(1):53.
7. Grol R, Wensing M, Eccles M, Davis D. Improving patient care: the implementation of change in health care: John Wiley & Sons; 2013.
8. Grol R, Bosch M, Wensing M. Development and selection of strategies for improving patient care. *Improving patient care: The implementation of change in health care*. 2013:165-84.
9. Curran GM, Bauer M, Mittman B, Pyne JM, Stetler C. Effectiveness-implementation hybrid designs: combining elements of clinical effectiveness and implementation research to enhance public health impact. *Medical care*. 2012;50(3):217.
10. Krause J, Van Lieshout J, Klomp R, Huntink E, Aakhus E, Flottorp S, et al. Identifying determinants of care for tailoring implementation in chronic diseases: an evaluation of different methods. *Implementation Science*. 2014;9(1):102.
11. Nilsen P, Bernhardsson S. Context matters in implementation science: a scoping review of determinant frameworks that describe contextual determinants for implementation outcomes. *BMC health services research*. 2019;19(1):189.
12. Waltz TJ, Powell BJ, Fernández ME, Abadie B, Damschroder LJ. Choosing implementation strategies to address contextual barriers: diversity in recommendations and future directions. *Implementation Science*. 2019;14(1):42.
13. Powell BJ, Waltz TJ, Chinman MJ, Damschroder LJ, Smith JL, Matthieu MM, et al. A refined compilation of implementation strategies: results from the Expert Recommendations for Implementing Change (ERIC) project. *Implementation Science*. 2015;10(1):21.
14. Bosch M, Van Der Weijden T, Wensing M, Grol R. Tailoring quality improvement interventions to identified barriers: a multiple case analysis. *Journal of evaluation in clinical practice*. 2007;13(2):161-

8.

15. Baker R, Camosso-Stefinovic J, Gillies C, Shaw EJ, Cheater F, Flottorp S, et al. Tailored interventions to address determinants of practice. *Cochrane Database of Systematic Reviews*. 2015(4).
16. Powell BJ, Fernandez ME, Williams NJ, Aarons GA, Beidas RS, Lewis CC, et al. Enhancing the impact of implementation strategies in healthcare: a research agenda. *Frontiers in public health*. 2019;7:3.
17. Davidoff F, Batalden P, Stevens D, Ogrinc G, Mooney SE. Publication guidelines for quality improvement studies in health care: evolution of the SQUIRE project. *BMJ (Clinical research ed)*. 2009;338:a3152.
18. Johnson RB, Onwuegbuzie AJ. Mixed methods research: A research paradigm whose time has come. *Educational researcher*. 2004;33(7):14-26.
19. Kaplan HC, Froehle CM, Cassedy A, Provost LP, Margolis PA. An exploratory analysis of the model for understanding success in quality. *Health care management review*. 2013;38(4):325-38.
20. Fereday J, Muir-Cochrane E. Demonstrating rigor using thematic analysis: A hybrid approach of inductive and deductive coding and theme development. *International journal of qualitative methods*. 2006;5(1):80-92.
21. Proctor EK, Powell BJ, McMillen JC. Implementation strategies: recommendations for specifying and reporting. *Implementation Science*. 2013;8(1):139.
22. Fleuren M, Paulussen T, Van Dommelen P, Van Buuren S. Meetinstrument voor determinanten van innovaties (MIDI). Leiden: TNO. 2012:16.
23. Cochrane LJ, Olson CA, Murray S, Dupuis M, Tooman T, Hayes S. Gaps between knowing and doing: understanding and assessing the barriers to optimal health care. *Journal of continuing education in the health professions*. 2007;27(2):94-102.
24. Sadeghi-Bazargani H, Tabrizi JS, Azami-Aghdash S. Barriers to evidence-based medicine: a systematic review. *Journal of evaluation in clinical practice*. 2014;20(6):793-802.
25. Schofield J. Time for a revival? Public policy implementation: a review of the literature and an agenda for future research. *International Journal of Management Reviews*. 2001;3(3):245-63.
26. Sabatier PA. Top-down and bottom-up approaches to implementation research: a critical analysis and suggested synthesis. *Journal of public policy*. 1986;6(1):21-48.
27. Verdonschot S, Van Rooij M, editors. Validation of a set of design principles to promote knowledge productivity and innovation. 10th European Conference on Creativity and Innovation, Copenhagen; 2007.
28. Verdonschot SG. Methods to enhance reflective behaviour in innovation processes. *Journal of European Industrial Training*. 2006;30(9):670-86.
29. Pronovost PJ. Navigating adaptive challenges in quality improvement. *BMJ Publishing Group Ltd*; 2011.
30. Moore GF, Audrey S, Barker M, Bond L, Bonell C, Hardeman W, et al. Process evaluation of complex interventions: Medical Research Council guidance. *BMJ (Clinical research ed)*. 2015;350:h1258.

31. Proctor E, Silmere H, Raghavan R, Hovmand P, Aarons G, Bunger A, et al. Outcomes for implementation research: conceptual distinctions, measurement challenges, and research agenda. *Administration and Policy in Mental Health and Mental Health Services Research*. 2011;38(2):65-76.

32. Reed JE, Kaplan HC, Ismail SA. A new typology for understanding context: qualitative exploration of the model for understanding success in quality (MUSIQ). *BMC health services research*. 2018;18(1):584.

33. Goldman J, Kuper A, Wong BM. How Theory Can Inform Our Understanding of Experiential Learning in Quality Improvement Education. *Academic medicine : journal of the Association of American Medical Colleges*. 2018;93(12):1784-90.

34. Kolb DA. *Experiential learning: Experience as the source of learning and development*: FT press; 2014.

Tables

Table 1a: Characteristics of healthcare professionals participating in the survey (N=28)

Characteristics		N (%)
Professional background		
	Physician	14 (50%)
	Nurse	10 (36%)
	Pharmacist	2 (8%)
	Healthcare scientist	1 (4%)
	Healthcare jurist	1 (4%)
Gender		
	Female	18 (%)
	Male	10 (%)
Hospital setting of their quality improvement initiative		
	Academic medical center	
	Teaching hospital	21 (75%)
	General hospital	6 (21%)
		1 (4%)
Medical department of their quality improvement initiative		
	Surgery	
	(Pediatric) intensive care	6 (21%)
	Internal medicine	6 (21%)
	Anesthesiology	2 (8%)
	Emergency	1 (4%)
	Geriatrics	1 (4%)
	Gastrointestinal liver	1 (4%)
	Neonatology	1 (4%)
	Neurology	1 (4%)
	Oncology	1 (4%)
	Orthopedics	1 (4%)
	Pediatrics	1 (4%)
	Radiology	1 (4%)
	Rehabilitation	1 (4%)
	Rheumatology	1 (4%)
	Transcending departments	1 (4%)
	Urology	1 (4%)
		1 (4%)

Table 1b. Improvement goals of healthcare professionals' quality improvement initiatives
(N=28)

Improvement goal
Reduction of medication administration errors
Improving early detection of deterioration amongst children
Improving resuscitation and non-technical team skills of nurses
Improving professional skills of nurses in using echography
Improving the use of the intra-hospital checklist by transporting patients
improving the employability and well-being of healthcare professionals
Improving the use of the 'Utrecht Symptoom Dagboek'(Utrecht Symptoms Diary)
Improving hand hygiene compliance
Improving the amount of information patients remember during consultation
Improving the collection of patient experiences
Reduction of (extreme) pain of children below the age of 10
Improving shared decision making between parents and healthcare professionals
Improving patient transfer from intensive care
Reduction of the amount of unplanned readmissions
Reduction of acetylcysteine and/or salbutamol without a strict clinical indication
Improving shared decision making
Improving sleep quality on intensive care
Improving the medical handover in complex cases
Improving safety of preoperative agreements
Reduction of deep postoperative wound infections
Improving system awareness, communication skills, empathy and professional identity
Improving patient satisfaction by substitution of care to outreach clinics
Improving patient identification and verification procedure
Reduction of intravenous medication errors
Improving the performance of STOP moments during the perioperative process
Improving wound care after hand operations
Improving nurses as role model in evidence based practice
Reduction of complications in admitted older patients

Table 2. Facilitators and barriers with a priority score ≥ 20 , grouped by category

Facilitators	Priority score^a
<i>External environment</i>	
Incentives or pressure (financial, legal or politica)	64
<i>Organisation</i>	
Sufficient support of expertise in the field of quality improvement	95
Culture of improvement	60
Sense of urgency ^b	42
Sufficient available time	90
<i>Department</i>	
Sufficient support of management	288
Employee support ^b	203
Bottom-up project approach ^b	96
Enthusiastic and supportive department head ^b	76
Workforce is motivated about the improvement project	33
<i>Quality improvement team</i>	
Sufficient participation in the decision-making process by team members	24
<i>Intervention of the project</i>	
Intervention fits in with current workflow	120
Barriers	
<i>Organisation</i>	
Insufficient available time	767
Data infrastructure	120
Insufficient support of the Executive Board for the project	110
Opponents of the project	30
Other organisational changes (reorganisation, merger)	27
Insufficient integration of quality improvement	27
<i>Intervention</i>	
Lack of evidence in literature of the effects of intervention	24
<i>Department</i>	
Insufficient motivation among the workforce	21
Experiencing one's competencies needed for the intervention as insufficient	20

^a The priority score consists of 1) the sum of the ranking in the top five (i.e., a determinant ranked first place in the top five got five points, while a determinant ranked fifth place in the top five received one point) multiplied by 2) the number of times a determinant was placed in the top five by professionals.

^b Self-experienced determinant listed by professionals (not included in the pre-specified list of determinants used in the survey).

Table 3. Area of expertise implementation experts participating in interviews

Area of expertise	N implementation experts ^a
Qualitative research	1
Governance of quality and safety in healthcare	2
Patient safety and teamwork	2
Quality and safety advisor	4
Quality improvement in patient care and education	1
Evidence based healthcare	2
Implementation improvement projects in patient care	2
Implementation as learning	2
Implementation in the field of perioperative patient safety	3
Evaluation of quality improvement	1
Patient involvement	3

^aSome experts were interviewed from different areas of expertise

Table 4 . Example for analysing and addressing a facilitator and barrier

Determinant	Analysis	Approaches
<i>Facilitator</i>		<i>Ways to use facilitator when performing QIPs</i>
Sufficient support of management	<ol style="list-style-type: none"> 1. Conduct a stakeholder analysis 2. Does management meet its agreements? Are you involved in matters relating to the initiative? Do you receive information about the project? 3. Are a member of the Executive Board and a department head the sponsor of the initiative? Do they actively support it and provide resources? 	<ol style="list-style-type: none"> 1. Seek support from management before the start of the QI initiative. 2. At least once every 9 months, according to an agreement with the management, provide periodic information about the progress of the initiative. This is part of a detailed communication plan. 3. Explicitly ask the department head and a member of the Executive Board to sign the project plan.
<i>Barrier</i>		<i>Ways to tackle barrier when performing QIPs</i>
Insufficient support of the Executive Board for the project	<ol style="list-style-type: none"> 1. Conduct a stakeholder analysis 2. Has the project plan been signed by a member of the Executive Board? 	<ol style="list-style-type: none"> 1. Make sure your initiative fits the mission or the strategic principles of the Executive Board and show this explicitly in the title of your initiative . 2. Take someone else from your project team to a meeting with the Executive Board so that you can complement each other.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Additionalfile2COREQchecklist.docx](#)
- [Additionalfile4Tableswithfacilitatorsandbarriers.docx](#)
- [Additionalfile1.STROBEchecklist.docx](#)
- [Additionalfile3DeterminantsurveyDutch.docx](#)
- [Additionalfile5TooldeterminantsQlinitiativesrevised.docx](#)