

Molecular mechanism and effects of urocortin II on rat adventitial fibroblast calcification induced by calcified medium

Xusheng Zhang

Department of Cardiology, The First Central Clinical College of Tianjin Medical University, 300070 Tianjin, China

Zhanjun Huang

Department of Cardiology, Longgang People's Hospital of Shenzhen, 518172 Shenzhen, Guangdong, China

Xiaorong Fan

Department of Cardiology, Longgang People's Hospital of Shenzhen, 518172 Shenzhen, Guangdong, China

Xiaoqing Tan

Department of Cardiology, Longgang People's Hospital of Shenzhen, 518172 Shenzhen, Guangdong, China

Chengzhi Lu (✉ 5020200072@nankai.edu.cn)

Department of Cardiology, The First Central Clinical College of Tianjin Medical University, 300070 Tianjin, China

Jianshe Yang (✉ yangjs@impcas.ac.cn)

Shanghai Tenth People's Hospital, Tongji University, Shanghai 200072, China

Research Article

Keywords: Urocortin II, vascular calcification, rat adventitial fibroblasts, Wnt/ β -catenin signaling pathway

Posted Date: March 18th, 2022

DOI: <https://doi.org/10.21203/rs.3.rs-1463803/v1>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

The present study aimed to assess the role of urocortin II (UII) in the process of vascular calcification *in vitro* by using a calcification model, and to detect the changes in the mRNA and protein levels of associated markers in rat adventitial fibroblasts (AFs) during their phenotypic transformation to osteoblast cells, and clarify the main signal transduction pathway of UII responsible for regulating vascular calcification and AF phenotypic transformation of osteoblast cells and prove that UII was an endogenous factor promoting vascular calcification, so as to provide an effective experimental basis for the clinical regulation of related diseases caused by vascular calcification. Finally we successfully constructed the calcified cell model, found that UII was an endogenous substance regulating vascular calcification, regulated the vascular calcification by promoting apoptosis and inhibiting autophagy through up- and down-regulated Bax and Bcl-2/Beclin-1 level, and the Wnt/ β -catenin signaling pathway was involved.

Introduction

Vascular calcification is a common condition noted in aging, atherosclerosis, hypertension, diabetes, dyslipidemia, valvular heart disease, and chronic kidney disease [1, 2]. It is caused by heterotopic deposition of hydroxyapatite on vascular walls [3]. Vascular calcification increases the risk of heart disease, atherosclerotic plaque rupture, and stroke. It is an important risk factor for the increased incidence of cardiovascular disease-associated morbidity and mortality [4, 5]. Therefore, vascular calcification is attracting increased attention worldwide. At present, various research studies have contributed to intima calcification, middle membrane calcification, and heart valve calcification. Previous clinical studies demonstrated that adventitia calcification is also possible. However, research studies on adventitia calcification are limited. Therefore, additional studies are required to elucidate the pathogenesis of vascular calcification.

Originally, vascular calcification was considered to be the result of passive degeneration of blood vessels, marking the aging of blood vessels. However, a growing number of studies have shown that vascular calcification is an active process regulated by a variety of molecular signaling pathways [4–7], which are highly similar to those involved in bone formation [7, 8]. Boström et al. demonstrated that during calcification, the morphology of rat vascular mesenchymal cells (VSMCs) was altered from the original contractile phenotype to the osteoblast-like phenotype. The expression levels of the original contractile markers, such as α -SMA were decreased, while the expression levels of Runx2, BMP-2, and other osteoblast-like markers were increased, which enabled the deposition of bone matrix in the blood vessels [9, 10]. Apoptosis and autophagy are two pathways that induce cell death, which are closely associated with vascular calcification. Previous studies have shown that VSMCs release apoptotic bodies during the induction of apoptosis to promote cellular calcification. Autophagy has also been shown to regulate cellular calcification through multiple mechanisms of action [11].

Urocortin II (Ull) is a neurocyclic peptide composed of 11 amino acid residues, which is currently considered to be a vasoactive peptide [12]. Immunohistochemical and quantitative PCR analyses were used to measure the expression levels of Ull in human atherosclerotic lesions. The data indicated that the expression levels of Ull were increased [13]. Rakowski et al. demonstrated that the use of the Ull antagonist SB-611812 against Ull could reduce the thickness of new intima and increase the lumen flow in a rat carotid artery angioplasty restenosis model [14]. These studies suggested that Ull played an important role in vascular dysfunction and remodeling. However, the mechanism of Ull in the calcification of adventitia fibroblasts remains unclear.

The purpose of the present study was to investigate whether Ull promoted adventitia fibroblast calcification by inducing apoptosis and inhibiting autophagy and to explore the possible mechanism of action. We hypothesized that Ull promoted AF calcification by inducing apoptosis and inhibiting autophagy. To verify this hypothesis, we constructed an *in vitro* calcification model using AFs and investigated the effects of Ull on the calcification of AFs and its possible mechanism of action.

Methods And Materials

Cell culture

AFs were isolated from the thoracic aortas of 4 to 6-week-old male Wistar-Kyoto rats weighing 120–180 g and cultured according to the tissue adherence method [15]. Animal experiments for AFs acquirement were carried out at the Molecular Biology Center of the First Central Clinical College of Tianjin Medical University, China. Briefly, for the rats, intravenous anesthesia was performed with 3% sterile saline solution of sodium pentobarbital according 30mg/kg body weight, finally sacrificed by medullotomy after the experiments. These operations were strictly complied with the Helsinki Declaration, and approved by the Ethics Committees of the Tianjin Medical University (TMU-2020-LLSC-0106). The cells were cultured in Dulbecco's modified Eagle's medium (DMEM) (Cat#11965092, Gibco, NY, USA) supplemented with 10% fetal bovine serum (Gibco, NY, USA) and maintained in a humidified atmosphere of 5% CO₂ at 37°C. AFs were cultured at 80–90% confluence and passaged with 0.125% trypsin-EDTA. Cells from passages three to six were used.

The following reagents were used: Fetal bovine serum (FBS), high glucose DMEM, penicillin, streptomycin (Cat#85886, Sigma Aldrich, USA), alizarin red solution, β-Glycerol phosphate, ascorbic acid, Dexamethasone (Sigma, USA), antibodies against GAPDH (Cat#AP0063, Bioworld, USA), antibodies against Bcl-2, Bax, Beclin1 and LC3 (Cell signaling), alkaline phosphatase assay kit and calcium (Ca) test kit (Nanjing Jiancheng Biological Engineering Institute). The primers for the RT-PCR assay were synthesized by BGI Genomics.

Grouping

The cells were seeded into 6-well plates at a density of 1×10^5 cells/l and maintained in DMEM containing 10% FBS. Following 24 h of incubation, the cells were divided into the following 7 groups: 1)

Normal control group, which contained cells maintained in DMEM without other treatments, 2) calcification group, including cells that were treated with 10 mmol/l β -glycerophosphate + 0.05 mmol/l ascorbic acid + 100 mmol/l dexamethasone, 3) calcification + Ull group, which included cells treated with Ull at different concentrations (10^{-7} , 10^{-8} and 10^{-9} mol/l) in the presence of the calcified medium, and 4) inhibitor group, which contained calcified medium and Ull (10^{-9} mol/l), and cells treated with or without DKK-1 (50 ng/ml). The medium was replaced with fresh medium every 3 days. Following 15 days, the medium was removed and the cells were used for different experiments.

Measurement of alkaline phosphatase activity

The cells were seeded in 6-well plates. The medium was removed and the cells were washed thrice with PBS. The ALP activity was measured according to the manufacturer's instructions with a kit (A0592, Nanjing Jiancheng Biological Engineering Institute). In brief, 300 μ l Triton X-100 (0.1%) was added to each well, followed by incubation at 4°C for 40 min. The lysed cells were transferred to a centrifuge tube. The cells were centrifuged and the supernatant was used for detection. The incubation was performed at 37°C. The ALP activity was measured by using an atomic absorption spectrophotometer at 520 nm. The hydrolysis of para-nitrophenyl phosphate, which is used as a chromogenic substrate, within 1 min may produce 1 μ mol p-nitrophenol, which requires 1 unit of ALP activity.

Measurement of calcium content

The cells were seeded into 6-well plates. The medium was removed and the cells were washed thrice with PBS. The calcium content was measured according to the manufacturer's instructions with a kit (C004-2-1 Nanjing Jiancheng Biological Engineering Institute). For decalcification, 300 μ l 0.6 M HCl was added to each well, followed by incubation at room temperature for 24 h. The supernatant was used for detection. The calcium content was measured using an atomic absorption spectrophotometer at 610 nm.

Alizarin red staining

The cells were seeded into 6-well plates. The medium was removed, and the cells were washed in PBS twice and fixed in 4% paraformaldehyde (1 ml/well) for 30 min. Following removal of formalin, the cells were washed in PBS twice, followed by incubation with 0.1% alizarin red (1 ml/well) for 15 min. Following washing with PBS, the cells were observed under a light microscope in order to detect the calcified nodules. Three samples were selected from each group, and one field was randomly selected from each sample at $\times 100$. The number of calcified nodules was compared between the groups. The calcified nodules were orange in color.

RT-PCR

The cells were harvested and total RNA was extracted with TRIzol[®] reagent (Invitrogen). The mRNA expression levels were determined by an SYBR green-based real-time PCR assay using a standard protocol as follows: Predenaturation at 95°C for 5 min, and a total of 35 cycles of denaturation at 95°C for 45 sec, annealing at 58°C for 45 sec, and extension at 72°C for 50 sec, followed by a final extension at

72°C for 10 min. The experiments were performed thrice and GAPDH was used as an internal control. The relative mRNA expression levels of each gene were determined. The primers used for PCR are shown in Table 1.

Table 1
Primer sequence list

Gene		Primer
ALP	Forward	5'-TGCAGTACGAGCTGAACAGG
	Reverse	5'-GTCAATTCTGCCTCCTTCCCA
OPN	Forward	5'-TCAGCTGGATGACCAGAGTG
	Reverse	5'-TTGGGGTCTACAACCAGCAT
Runx2	Forward	5'-TGG CTT TGG TTT CAG GTT AGG-3'
	Reverse	5'-TGG AGA TGT TGC TCT GTT CG-3'
BMP-2	Forward	5'-TGA GGA TTA GCA GGT CTT TGC-3'
	Reverse	5'-TCT CGT TTG TGG AGT GGA TG-3'
GAPDH	Forward	5'-GGC TGC CCA GAA CAT CAT-3'
	Reverse	5'-CGG ACA CAT TGG GGG TAG-3'

Western blotting

The cells were harvested with lysis buffer. The lysates were collected, and the protein concentration was determined using the BCA Protein Assay Kit (A045-4, Nanjing Jiancheng Biological Engineering Institute). The protein samples were separated on 12% SDS-PAGE gels, transferred to polyvinylidene fluoride membranes (Millipore, MA, US), and blocked with 5% nonfat milk/TBST for 2 h at room temperature. The membranes were subsequently incubated overnight at 48°C with primary antibodies. The membranes were washed with TBST three times and incubated with secondary antibodies for 1 h at room temperature. Immunoreactive bands were visualized by enhanced chemiluminescence (ECL) (Millipore, MA, US). The density of the bands was quantified with the Quantity One image analysis software (Bio-Rad). The optical density (OD) of each band was determined and subsequently normalized to that of GAPDH.

All of the above experiments

Data analysis

All data are expressed as mean \pm SEM. The mean differences between the groups were analyzed by one-way ANOVA or t-test using SPSS 22.0 statistical software. $P < 0.05$ was considered to indicate a statistically significant difference.

Results

AFs can induce calcification *in vitro*

It has been previously reported that the *in vitro* culture of VSMC in the presence of calcified medium can induce their calcification. To assess whether high calcium and phosphorus can induce calcification of rat AFs, the cells were cultured in a calcified medium, and the ALP activity and intracellular calcium contents were detected at different time points. By extending the incubation time, the ALP activity was gradually increased and reached its peak on the 9th day of culture. The results were significantly different compared with the control group ($P < 0.05$). The intracellular calcium content was significantly increased following 9 days of induction compared with that of the control group ($P < 0.05$), and the increase noted in ALP activity levels and in the calcium content was time-dependent (Fig. 1A). The highest calcium content was detected on the 15th day of induction. Therefore, the induction time selected for subsequent experiments was 15 days. Alizarin red staining was performed on the cells following 15 days of induction of calcification. The results indicated that the cells in the control group were long and spindle-shaped, while the cells in the calcification group were diamond-shaped (Data not shown). Following alizarin red staining, a large number of orange calcium nodules were observed in the calcified group (Fig. 1B), whereas these nodules were not present in the control group. The results demonstrated that AFs could induce calcification *in vitro* following incubation with the calcified medium.

Ull increases the calcification of AFs

In order to explore whether Ull exerts an effect on the calcification of AFs, different concentrations of Ull were added in the calcified medium to induce calcification. ALP activity and intracellular calcium content were detected following 15 days of incubation. The data indicated that 10^{-9} , 10^{-8} , and 10^{-7} M Ull significantly increased ALP activity ($P < 0.05$) (Fig. 2A). The intracellular calcium content was significantly higher than that of the calcified group ($P < 0.05$) following 15 days of cell incubation with 10^{-9} , 10^{-8} , and 10^{-7} M Ull in the medium (Fig. 1A). Increased vascular calcification is associated with increased expression levels of osteoblast-specific markers, such as BMP-2 and Runx2 [16]. RT-PCR results indicated that the mRNA expression levels of the osteoblast-specific markers BMP-2 and Runx2 were significantly increased in the calcified group compared with those of the control group ($P < 0.05$). Compared with the calcified group, the mRNA expression levels of BMP-2 and Runx2 were further increased in the presence of 10^{-9} M Ull, and the differences were statistically significant ($P < 0.05$) (Fig. 2B).

Ull promotes calcification by inducing apoptosis and inhibiting autophagy In order to further explore the mechanism of Ull in promoting cell calcification, the changes in the expression levels of apoptotic and autophagic proteins were assessed by western blot analysis. Compared with the control group, the expression levels of the pro-apoptotic protein Bax were upregulated, while those of the antiapoptotic protein Bcl-2 were downregulated in the calcified group. The increase in the concentration of Ull caused an upregulation in the expression levels of the Bax protein and downregulation in the expression levels of the Bcl-2 protein in a concentration-dependent manner (Fig. 3A). Compared with the control group, the

expression levels of the autophagy-related protein Beclin1 and the ratio of LC3-II/LC3-I were increased in the calcification group, indicating that intracellular autophagy levels were enhanced following incubation with the calcified medium. However, the expression levels of the autophagy-related protein Beclin1 and the ratio of LC3-II to LC3-I were decreased following the addition of Ull to the calcium content in a concentration-dependent manner (Fig. 3B). In conclusion, Ull inhibited autophagy during calcification.

Ull promotes calcification by activating the Wnt/ β -catenin pathway Activation of the Wnt/ β -catenin pathway is one of the key factors involved in cell calcification. In order to investigate whether Ull promotes the calcification of AFs via the Wnt/ β -catenin pathway, the cells were treated with DKK-1, which is a Wnt/ β -catenin pathway inhibitor. The results of the western blot analysis indicated that the expression levels of the Bax protein were upregulated, whereas the expression levels of the Bcl-2 protein were downregulated following incubation of the cells with calcified medium for 15 days. The expression levels of Bax were further upregulated following the addition of Ull in the calcium content, whereas the expression levels of Bcl-2 were downregulated. The data indicated that DKK-1 reversed the pro-apoptotic effect of Ull (Fig. 4A). Figure 3B indicated that Ull inhibited the expression levels of the autophagy-related protein Beclin1, whereas it could decrease the LC3-II/LC3-I ratio. DKK-1 intervention caused upregulation of the expression levels of the autophagy-related protein Beclin1 and an increase in the LC3-II/LC3-I ratio, which partially restored the autophagy levels of the cells, suggesting that Ull may promote calcification in AF cells through the Wnt/ β -catenin pathway.

To further confirm that the Wnt/ β -catenin inhibitor DKK-1 can inhibit Ull-induced cell calcification, the cells were simultaneously treated with Ull + DKK-1, and the formation of the intracellular calcium nodules was detected following 15 days of culture by alizarin red staining (Fig. 4C). The data indicated that following DKK-1 intervention, the number of calcium nodules induced by Ull was significantly reduced.

Discussion

Vascular calcification refers to the abnormal deposition of calcium and phosphorus crystals on the vascular wall, which is divided into the three following types: Intima calcification, middle membrane calcification, and outer membrane calcification [17]. Vascular calcification is one of the complications of cardiovascular disease and has increased in incidence and severity in the past years. It is considered to be an important factor leading to the high mortality of cardiovascular disease. Currently, no effective treatment is available for vascular calcification. Therefore, the exploration of the pathogenesis of vascular calcification and the identification of effective diagnostic and prevention targets at the molecular level are urgently required.

Previous studies have shown that TGF- β 1 can induce calcification of rat extravascular membrane cells in the calcification medium [17]. However, the present study demonstrated that significant calcification of AF cells could be induced *in vitro* by 10 mmol/l β -glycerophosphate + 0.05 mmol/l ascorbic acid + 100 mmol/l dexamethasone without the addition of exogenous cytokines. This was determined by alizarin red S staining, and measurement of ALP activity levels and intracellular calcium content. In addition, the

experimental results indicated that Ull could further increase ALP activity and intracellular calcium content.

Certain studies have implied that in the process of vascular calcification, VSMCs transform into osteoblast-like cells and express specific osteogenic markers, resulting in the deposition of bone matrix, which is an important feature and mechanism of vascular calcification [18]. The calcification conditions of AF cells used in the present study are consistent with those reported in previous studies for VSMCs, suggesting that the mechanism of AF cell calcification may be partially consistent with that of VSMC calcification. It has been reported that during the calcification process of VSMCs, the cells express osteogenic transcription factors, such as Runx2, which promote the expression levels of downstream bone-related proteins, such as bone morphogenetic protein BMP-2 and facilitate the active differentiation of cells into osteoblast-like cells [19]. This mediated the deposition of the bone matrix in the blood vessels [19]. In the present study, a similar mechanism was observed. The mRNA expression levels of BMP-2 and Runx2 were significantly increased in AFs following incubation of the cells in the calcified medium for 15 days. Ull could further promote the mRNA expression levels of BMP-2 and Runx2. In the present study, calcium salt deposition and phenotypic transformation of osteoblasts suggesting that AFs could be calcified by calcium-rich medium, which extended the classification of vascular calcification.

The mechanism of vascular calcification is complex, involving various signaling pathways, such as autophagy and apoptosis, activation of the Wnt/ β -catenin signaling pathway, and induction of ER stress. As an adaptive response to cellular stress, autophagy is critical in maintaining vascular structure and function. Previous studies have shown increased levels of autophagy during vascular calcification [20, 21]. Autophagy can inhibit vascular calcification through a variety of mechanisms. In *in vitro* experiments, phosphorus increased the levels of autophagy in VSMCs and the number of autophagosomes, thereby inhibiting the apoptotic and calcification processes [22]. Additional studies have shown that autophagy can inhibit oxidative stress of VSMCs and the inflammatory response of vascular endothelial cells, which regulates lipid metabolism and reduces vascular calcification [19, 23]. LC3 and Beclin1 are two typical markers of autophagy. The results of the western blot analysis indicated that the LC3-II/LC3-I ratio and the expression levels of Beclin1 protein were increased in AF cells incubated with calcified medium for 15 days, while Ull treatment decreased LC3-II/LC3-I ratio and Beclin1 expression in a concentration-dependent manner.

Apoptosis involves the induction of vascular calcification, and its inhibition can in turn inhibit the calcification process [20, 21]. Shi et al. demonstrated that fibroblast growth factor 21 could regulate the CHOP and caspase-12 signaling pathways by endoplasmic reticulum stress so as to reduce vascular smooth muscle cell apoptosis and inhibit vascular calcification [22]. In addition, the increased concentration of Pi or Ca^{2+} in the medium induced the formation and release of matrix vesicles, such as the apoptotic bodies, from the plasma membrane of VSMCs, leading to the calcification of the extracellular matrix, which may be the nucleation site for vascular calcification [24]. The findings demonstrated that the intracellular apoptotic levels were increased during the calcification induced by the calcified medium in AFs, which was consistent with previous studies. Concomitantly, intracellular

autophagy levels were also increased, which may be the result of the cells regulating themselves against calcification. However, Ull increased the apoptotic levels and inhibited autophagy, which aggravated the calcification of AF cells *in vitro*.

Previous studies have shown that the Wnt/ β -catenin pathway is an important signaling pathway involved in the process of osteogenesis and vascular calcification. Activation of the Wnt/ β -catenin pathway can promote the expression levels of the calcification factors and ultimately initiate and promote the occurrence and development of vascular calcification by upregulating ALP activity. In contrast to this mechanism of action, inhibition of the Wnt/ β -catenin pathway has been reported to protect against vascular calcification [25–27]. To determine whether the Wnt/ β -catenin pathway plays a role in AF calcification, the cells were treated with DKK-1, which is a Wnt/ β -catenin pathway inhibitor. The results indicated that inhibition of the Wnt/ β -catenin pathway inhibited apoptosis and promoted autophagy of AFs, which is consistent with the report of Liu et al. [25]. Concomitantly, alizarin red staining was used to detect the formation of intracellular calcium nodules following induction for 15 days. The data indicated that following DKK-1 intervention, the number of calcium nodules induced by Ull was significantly reduced. These results further demonstrated that Ull promoted AF calcification partially by activating the WNT/ β -catenin pathway.

In conclusion, the present study demonstrated that the calcium content could induce the calcification of rat AF cells, whereas Ull could aggravate the degree of calcification by promoting apoptosis and inhibiting calcification. Moreover, inhibition of the WNT/ β -catenin pathway could alleviate the calcification induced by Ull.

Conclusions

The culture of AF cells for 15 days in the calcified medium could successfully construct the calcified cell model. Ull can increase calcium deposition and ALP activity in AF calcified cells in a concentration-dependent manner, suggesting that it is an endogenous substance regulating vascular calcification. Following Ull intervention, the protein levels of Bax were significantly upregulated, while the protein levels of Bcl-2 and Beclin-1 were downregulated. The LC3-II/LC3-I ratio was upregulated, suggesting that Ull regulated vascular calcification by promoting apoptosis and inhibiting autophagy. The Wnt/ β -catenin signaling pathway blocker DKK-1 inhibited apoptosis and autophagy induced by Ull, and reduced the number of calcium nodules, suggesting that the ability of Ull to promote cell calcification may be related to the activation of the Wnt/ β -catenin signaling pathway.

Abbreviations

Ull, urocortin II; AFs, adventitial fibroblasts; ALP, alkaline phosphatase; Runx2, Runt-related transcription factor 2; DKK-1, Dickkopf-related protein-1; VSMC, vascular mesenchymal cells.

Declarations

Authors' contributions

XZ, ZH, XF, and XT carried out the studies, participated in collecting data, and drafted the manuscript. XZ performed the statistical analysis and participated in its design. CL, XZ, and JY participated in acquisition, analysis, or interpretation of the data and in drafting the manuscript. All authors read and approved the final version of the manuscript.

Ethics approval and consent to participate

The present study protocol was approved by the Ethics Committees of the Tianjin Medical University (TMU-2020-LLSC-0106).

Acknowledgement

Not applicable.

Funding

This work was supported by research grants from the National Natural Science Foundation of China (grant number 81970303).

Conflict of interest

The authors declare no conflict of interest.

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

References

1. Shanahan CM, Crouthamel MH, Kapustin A, Giachelli CM. Arterial calcification in chronic kidney disease: key roles for calcium and phosphate. *Circ Res.* 2011;109(6):697–711.
2. Ho CY, Shanahan CM. Medial Arterial calcification: An overlooked player in peripheral arterial disease. *Arterioscler Thromb Vasc Biol.* 2016;36(8):1475–1482.
3. Nicoll R, Henein MY. The predictive value of arterial and valvular calcification for mortality and cardiovascular events. *Int J Cardiol Heart Vessel.* 2014;3:1–5.

4. Johnson RC, Leopold JA, Loscalzo J. Vascular calcification: pathobiological mechanisms and clinical implications. *Circ Res*. 2006;99(10):1044–1059.
5. Yamada S, Giachelli CM. Vascular calcification in CKD-MBD: Roles for phosphate, FGF23, and Klotho. *Bone*. 2017;100:87–93.
6. Zhang X, Zhou X, Huang Z, Fan X, Tan X, Lu C, et al. Aldosterone is a possible new stimulating factor for promoting vascular calcification. *Front Biosci (Landmark Ed)*. 2021;26(11):1052–1063.
7. Durham AL, Speer MY, Scatena M, Giachelli CM, Shanahan CM. Role of smooth muscle cells in vascular calcification: implications in atherosclerosis and arterial stiffness. *Cardiovasc Res*. 2018;114(4):590–600.
8. Lin ME, Chen TM, Wallingford MC, Nguyen NB, Yamada S, Sawangmake C, et al. Runx2 deletion in smooth muscle cells inhibits vascular osteochondrogenesis and calcification but not atherosclerotic lesion formation. *Cardiovasc Res*. 2016;112(2):606–616.
9. Boström KI, Rajamannan NM, Towler DA. The regulation of valvular and vascular sclerosis by osteogenic morphogens. *Circ Res*. 2011;109(5):564–577.
10. Speer MY, Yang HY, Brabb T, Leaf E, Look A, Lin WL, et al. Smooth muscle cells give rise to osteochondrogenic precursors and chondrocytes in calcifying arteries. *Circ Res*. 2009;104(6):733–741.
11. Proudfoot D, Skepper JN, Hegyi L, Bennett MR, Shanahan CM, Weissberg PL. Apoptosis regulates human vascular calcification in vitro: evidence for initiation of vascular calcification by apoptotic bodies. *Circ Res*. 2000;87(11):1055–1062.
12. Eyre HJ, Speight T, Glazier JD, Smith DM, Ashton N. Urotensin II in the development and progression of chronic kidney disease following 5/6 nephrectomy in the rat. *Exp Physiol*. 2019;104(3):421–433.
13. Bousette N, Patel L, Douglas SA, Ohlstein EH, Giaid A. Increased expression of urotensin II and its cognate receptor GPR14 in atherosclerotic lesions of the human aorta. *Atherosclerosis*. 2004;176(1):117–123.
14. Rakowski E, Hassan GS, Dhanak D, Ohlstein EH, Douglas SA, Giaid A. A role for urotensin II in restenosis following balloon angioplasty: use of a selective UT receptor blocker. *J Mol Cell Cardiol*. 2005;39(5):785–791.
15. An SJ, Boyd R, Zhu M, Chapman A, Pimentel DR, Wang HD. NADPH oxidase mediates angiotensin II-induced endothelin-1 expression in vascular adventitial fibroblasts. *Cardiovasc Res*. 2007;75(4):702–709.
16. Zeadin M, Butcher M, Werstuck G, Khan M, Yee CK, Shaughnessy SG. Effect of leptin on vascular calcification in apolipoprotein E-deficient mice. *Arterioscler Thromb Vasc Biol*. 2009;29(12):2069–2205.
17. Li N, Cheng W, Huang T, Yuan J, Wang X, Song M. Vascular Adventitia Calcification and Its Underlying Mechanism. *PLoS One*. 2015;10(7):e0132506.
18. Leopold JA. Vascular calcification: Mechanisms of vascular smooth muscle cell calcification. *Trends Cardiovasc Med*. 2015;25(4):267–274.

19. Byon CH, Javed A, Dai Q, Kappes JC, Clemens TL, Darley-Usmar VM, et al. Oxidative stress induces vascular calcification through modulation of the osteogenic transcription factor Runx2 by AKT signaling. *J Biol Chem.* 2008;283(22):15319–15327.
20. Liu Q, Luo Y, Zhao Y, Xiang P, Zhu J, Jing W, et al. Nano-hydroxyapatite accelerates vascular calcification via lysosome impairment and autophagy dysfunction in smooth muscle cells. *Bioact Mater.* 2022;8:478–493.
21. Liang J, Huang J, He W, Shi G, Chen J, Huang H. β -Hydroxybutyric Inhibits Vascular Calcification via Autophagy Enhancement in Models Induced by High Phosphate. *Front Cardiovasc Med.* 2021;8:685748.
22. Ciceri P, Elli F, Cappelletti L, Tosi D, Braidotti P, Bulfamante G, et al. A new in vitro model to delay high phosphate-induced vascular calcification progression. *Mol Cell Biochem.* 2015;410(1–2):197–206.
23. Ouimet M, Franklin V, Mak E, Liao X, Tabas I, Marcel YL. Autophagy regulates cholesterol efflux from macrophage foam cells via lysosomal acid lipase. *Cell Metab.* 2011;13(6):655–667.
24. Reynolds JL, Joannides AJ, Skepper JN, McNair R, Schurgers LJ, Proudfoot D, et al. Human vascular smooth muscle cells undergo vesicle-mediated calcification in response to changes in extracellular calcium and phosphate concentrations: a potential mechanism for accelerated vascular calcification in ESRD. *J Am Soc Nephrol.* 2004;15(11):2857–2867.
25. Liu Y, Wang WM, Zhang XL, He HQ, Sun XL, Zeng H, et al. AGE/RAGE promotes the calcification of human aortic smooth muscle cells via the Wnt/ β -catenin axis. *Am J Transl Res.* 2016;8(11):4644–4656.
26. Cai T, Sun D, Duan Y, Wen P, Dai C, Yang J, et al. WNT/ β -catenin signaling promotes VSMCs to osteogenic transdifferentiation and calcification through directly modulating Runx2 gene expression. *Exp Cell Res.* 2016;345(2):206–217.
27. Jin X, Rong S, Yuan W, Gu L, Jia J, Wang L, et al. High Mobility Group Box 1 Promotes Aortic Calcification in Chronic Kidney Disease via the Wnt/ β -Catenin Pathway. *Front Physiol.* 2018;9:665.

Table 1

Table 1. Primer sequence list

Gene		Primer
ALP	Forward	5'-TGCAGTACGAGCTGAACAGG
	Reverse	5'-GTCAATTCTGCCTCCTTCCCA
OPN	Forward	5'-TCAGCTGGATGACCAGAGTG
	Reverse	5'-TTGGGGTCTACAACCAGCAT
Runx2	Forward	5'-TGG CTT TGG TTT CAG GTT AGG-3'
	Reverse	5'-TGG AGA TGT TGC TCT GTT CG-3'
BMP-2	Forward	5'-TGA GGA TTA GCA GGT CTT TGC-3'
	Reverse	5'-TCT CGT TTG TGG AGT GGA TG-3'
GAPDH	Forward	5'-GGC TGC CCA GAA CAT CAT-3'
	Reverse	5'-CGG ACA CAT TGG GGG TAG-3'

Figures




Figure 1

Rat extravascular outer membrane cells induce calcification *in vitro*. A: By using the calcification-induced medium, the ALP activity and the intracellular calcium content were increased following the time of culture; B: In culture, intracellular calcium nodules were clearly observed for 15 days with cissarin-red staining (100×).




Figure 2

Ull aggravates the calcification of vascular outer membrane cells in rats. A: The incubation of the cells with different concentrations of Ull demonstrated that high concentrations of Ull could increase ALP activity and intracellular calcium content; B: The expression levels of specific osteogenesis-related genes were detected using the RT-PCR assay and the data indicated that the levels all four osteogenesis-related genes were upregulated at 15 days following induction of calcification. Ull caused upregulation in the expression levels of osteogenesis-related genes.




Figure 3

U II promotes cell calcification by inducing apoptosis and inhibiting autophagy. WB analysis indicated that following 15 days of calcification, the expression levels of the apoptotic protein Bax were increased, whereas those of the antiapoptotic protein Bcl-2 were decreased. The expression levels of the autophagy-related proteins Beclin1 and LC3 were upregulated. U II caused further upregulation of the expression levels of the apoptotic protein Bax and downregulation of the expression levels of the apoptotic protein

Bcl-2. The expression levels of the autophagy-related proteins Beclin1 and LC3 were downregulated. The data indicated that Ull exacerbated cell calcification by promoting apoptosis and inhibiting autophagy.




Figure 4

DKK-1, the inhibitor of the Wnt/ β -catenin pathway inhibits the effect of Ull. A, B: WB analysis indicated that treatment of the cells with the Wnt/ β -catenin inhibitor DKK-1 reversed the effects of Ull, causing

downregulation of the apoptotic protein Bax, upregulation of the antiapoptotic protein Bcl-2, and upregulation of the autophagy-related proteins Beclin1 and LC3; C: Sissarin-red staining demonstrated that significant calcium nodules were observed in the intracellular fraction following 15 days of induction with calcified media. The simultaneous addition of the Ull culture indicated significantly denser intracellular calcium nodules. The increase in the concentration levels of DKK-1 resulted in significantly smaller orange calcium nodules being present in the cells.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [supplementfileswesternblot.docx](#)