Novel Mumps Virus Epitopes Reveal Robust Cytotoxic T Cell Responses After Natural Infection but Not After Vaccination

Patricia Kaaijk (✉ patricia.kaaijk@rivm.nl)

National Institute for Public Health and the Environment

Maarten Emmelot

National Institute for Public Health and the Environment

Hugo Meiring

Intravacc (Netherlands)

Cecile van Els

National Institute for Public Health and the Environment

Jelle de Wit

National Institute for Public Health and the Environment

Research Article

Keywords: Mumps virus, T cell epitopes, T cell response, cytotoxic CD8+ T-cells, cellular immunity, MHC class I, mumps virus infection, measles-mumps-rubella vaccination

Posted Date: January 22nd, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-147373/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Mumps is nowadays re-emerging despite vaccination. The contribution of T cell immunity to protection against mumps has not been clearly defined. Previously, we described a set of 41 peptides that were eluted from human leukocyte antigen (HLA) class I molecules of mumps virus (MuV)-infected cells. Here, we confirmed immunogenicity of five novel HLA-B*07:02- and HLA-A*01:01-restricted MuV T cell epitopes from this set of peptides. High frequencies of T cells against these five MuV epitopes could be detected *ex vivo* in all tested mumps patients. Moreover, these epitope-specific T cells derived from mumps patients displayed strong cytotoxic activity. In contrast, only marginal T cell responses against these novel MuV epitopes could be detected in recently vaccinated persons, corroborating earlier findings. Identifying which MuV epitopes are dominantly targeted in the mumps-specific CD8⁺ T- response is an important step towards better understanding in the discrepancies between natural infection or vaccination-induced cell-mediated immune protection

Introduction

The implementation of routine measles-mumps-rubella (MMR) vaccination has led to a dramatic reduction in incidence of these infectious diseases, however, mumps is re-emerging nowadays ^{1,2}. Especially young adults that were vaccinated during childhood can be affected by mumps. Waning of vaccine-induced immunity in the absence of natural boosters by circulating mumps virus (MuV) has been suggested as major cause ³⁻⁶. Various studies have shown a decline of antibody levels in time following childhood vaccination ^{5,7,8}. However, the finding that antibodies alone do not fully correlate with protection against MuV infection indicates that, apart from the humoral response, other immune compartments play a role in the immunity against MuV ^{6,9,10}. T cell responses against MuV have been detected in mumps cases as well as in vaccinated healthy persons ⁹⁻¹². Previously, we showed that strong MuV-specific CD8⁺ T cells dominate the T cell response in mumps patients. In the same study, we showed that the CD8⁺ T cell response in children that were recently vaccinated with the MMR vaccine was considerably less vigorous than after natural infection ⁹. Several scientists have emphasized the importance of fully understanding the T cell immune responses to MuV after vaccination, which is needed to be able to develop strategies to improve the quality and durability of vaccine-induced immunity ^{2,13-15}. For the detailed characterization of the T cell immune response to MuV and the understanding of its role in natural infection or vaccination, the identification of MuV-specific T cell epitopes is indispensable. Furthermore, to be able to investigate the cross-reactive potency of the T cells induced by vaccination towards the circulating strains, data on the T cell response at the epitope level is also crucial.

Recently, a set of 41 unique viral peptides was eluted from HLA class I molecules of MuV-infected cells (typing: HLA-A*01:01, A*02:01, B*07:02, B*40:01, C*03:04, C*07:02) and characterized by mass spectrometry. From this panel, we confirmed the first six HLA-A*02:01-restricted MuV T cell epitopes to be able to induce a T-cell response ¹⁶. To expand our knowledge on the CD8⁺ T cell repertoire of MuV, we here searched for novel T cell epitopes among the earlier identified panel of naturally processed peptides

with high predicted binding scores to the two other common HLA-I molecules, HLA-B*07:02 and HLA-A*01:01. We describe five novel HLA-B*07:02- and HLA-A*01:01-restricted MuV epitopes that reveal for the first time robust ongoing cytotoxic T cell responses in all mumps patients tested, with only low frequencies of epitope-specific T cells in recently vaccinated persons.

Results

In the present study, peptides from a previously characterized set of 41 naturally processed MuV epitopes¹⁶, predicted to bind best to HLA-B*07:02 (n = 12) and HLA-A*01:01 molecules (n = 10) were selected for immunogenicity assessment. Selected epitopes were tested for their capacity to recall a specific T cell response, by using expanded CD8⁺ T-effector cells isolated from three HLA-B*07:02 positive and three HLA-A*01:01 positive mumps patients (Fig. 1).

Three T cell epitopes, i.e., APIQGTNLL (phosphoprotein, aminoacid positions 27–35) V/P/I_{27–35}; KPRTSTPVTEF, V/P/I_{142–152}; IPNARANL, (nucleoprotein) NP_{115–122}, were identified that all elicit a T cell response in 3 out of 3 HLA-B*07:02 positive mumps patients, as measured by IFN- γ and TNF production. In addition, three HLA-A*01:01-restricted epitopes, i.e., YSDPNNHEVY, (large protein) L_{1336–1343}; YSDPNDHEVY, L_{1336–1343}; VTDSNLIY, L_{1983–1992}, induced a T cell response. The YSDPNNHEVY peptide, and its deamination variant, YSDPNDHEVY (in which the asparagine (N) on position 6 was deaminated to aspartic acid (D)), both induced a T cell response in 3 out of 3 HLA-A*01:01 positive mumps patients. The other HLA-A*01:01-restricted epitope (VTDSNLIY) induced IFN- γ and TNF production by CD8⁺ effector T cells from 2 out 3 HLA-A*01:01 positive mumps patients (Fig. 1). Subsequently, APIQGTNLL, KPRTSTPVTEF, IPNARANL (HLA-B*07:02 restricted epitopes), and YSDPNNHEVY and VTDSNLIY (HLA-A*01:01 restricted epitopes) were selected for custom-made (p:HLA) dextramer synthesis to investigate the *ex vivo* occurrence of CD8⁺ T cells against these epitopes in mumps patients. YSDPNNHEVY and not its post-translationally modified deamination variant, YSDPNDHEVY, was selected because it was more abundantly presented based on elution from MHC-I molecules from MuV-infected cells¹⁶ and had a better predictive binding to HLA-A*01:01 (Fig. 1).

Using p:HLA dextramers, high frequencies of MuV epitope-specific T cells were observed in all three B*07:02- and three HLA-A*01:01-positive patients, even exceeding 2% of the total CD8⁺ T cell pool (Fig. 2A). The frequencies of the epitope-specific CD8⁺ T cells and the dominance of the response per epitope varied between patients. Further analyses showed that the majority peptide-specific T cells were within the T-effector cell subset (CD3⁺CD8⁺CD45RO⁺CCR7⁺). Functional characterization revealed that, upon stimulation with peptides, the epitope-specific T cells expressed the activation marker CD137 and displayed a cytotoxic phenotype by expression of CD107a. In addition to pro-inflammatory cytokines IFN- γ , TNF and IL-17A, epitope-specific T cells produced granzyme-A/B, granulysin, and to a lesser extent sFas(L), and perforin that are involved in cell-mediated cytotoxicity (Fig. 2B). Finally, we showed that epitope-specific T cells were able to efficiently lyse peptide-pulsed antigen-presenting target cells in a flow cytometry-based cytotoxicity assay (Fig. 2C).

In contrast to the high frequencies of MuV epitope-specific T cells observed in mumps patients, only very low frequencies of MuV epitope-specific T cells could be detected in young adults that were recently vaccinated with a third dose of the MMR vaccine (Fig. 3). Despite the fact that epitope-specificity varied between individuals, in all tested PBMC samples of HLA-B7- and HLA-A1-positive vaccinated persons, small amounts of MuV epitope-specific CD8⁺ T cells could be visualized with the p:HLA dextramers.

Discussion

Identification of more functional CD8⁺ T cell epitopes that cover a broad repertoire of T cells against mumps virus (MuV) is important for monitoring and functionally characterizing the T cell response following mumps infection or vaccination. Based on HLA-I binding prediction, most of the set of 41 MuV peptides, from an earlier identified panel ¹⁶, bound well to the HLA-B*07:02 molecule, followed by the HLA-A*02:01 and HLA-A*01:01 molecules, i.e. belonging to the most common HLA-I alleles ¹⁷. In addition to the previously identified HLA-A*02:01 epitopes, we now confirmed three HLA-B*07:02-restricted T cell epitopes (i.e. APIQGTNLL (V/P/I₂₇₋₃₅), KPRTSTPVTEF (V/P/I₁₄₂₋₁₅₂), IPNARANL (NP₁₁₅₋₁₂₂)), each inducing a strong T cell response in expanded effector T cells obtained from three different HLA-B*07:02-positive mumps patients. Interestingly, the first described MuV-specific CD4⁺ T-helper cell epitope, GTYRLIPNARANLTA (NP₁₁₀₋₁₂₄), which we recently discovered ¹⁸, contains the newly identified HLA-B*07:02-restricted CD8⁺ T cell epitope IPNARANL (NP₁₁₅₋₁₂₂). This opens the possibility of using synthetic peptides, containing both epitope sequences, to stimulate CD4⁺ and CD8⁺ T cell responses simultaneously. In addition to the HLA-B*07:02-restricted epitopes, HLA-A*01:01-restricted T cell epitopes (i.e. YSDPNNHEVY (L₁₃₃₆₋₁₃₄₃) and VTDSNLIY (L₁₉₈₃₋₁₉₉₂)) showed to induce a T cell response in HLA-A*01:01 positive mumps patients. Four out of the five novel MuV epitopes of the genotype G outbreak strain (MuVi/Utrecht.NLD/40.10) appeared to be conserved among the two Jeryl Lynn vaccine strains (JL2 and JL5), and therefore are useful for monitoring cellular immune responses after natural infection with the genotype G outbreak strain and after vaccination. Interestingly, one of the HLA-B*07:02-restricted epitopes, i.e. APIQGTNLL (V/P/I₂₇₋₃₅) showed to have sequence differences among various circulating mumps strains, including genotype G, H and C strains, and also differ from the peptide sequences in the two Jeryl Lynn vaccine strains (JL2 and JL5). Future experiments are planned to explore the impact of this mismatch on the T cell recognition of the various MuV strains, i.e. vaccine *versus* circulating MuV strains. Using peptide-HLA class I (p:HLA-I) dextramers, high frequencies of T cells against all five MuV epitopes were *ex vivo* detected in PBMCs of all three HLA-B*07:02- and three HLA-A*01:01-positive mumps patients. Functional characterization of these specific T cells revealed a cytotoxic phenotype with expression of degranulation marker CD107a, and secretion of granzyme-A/B, granulysin, perforin, sFas(L). With a flow cytometry-based cytotoxicity assay, epitope-specific T cells showed indeed to be able to efficiently lyse MuV peptide-pulsed target cells. Thus, we for the first time describe that MuV epitope-specific CD8⁺ effector T cells isolated from mumps patients have a cytotoxic phenotype which is a critical feature for mediating viral clearance following acute MuV infection. Although T cells against the novel MuV epitopes were also detectable in recently vaccinated persons, only very low frequencies of

these specific CD8⁺ T cells were observed, which is in line with our previous findings⁹. Live-attenuated vaccines, such as MMR vaccines, are likely to be able to induce CD8⁺ T cells, but it might be that the vaccine-induced CD8⁺ T cell response is low due to attenuated replication of the vaccine virus. As a result, the mumps-specific CD4/CD8 T cell ratio may be lower after vaccination than after natural infection, as confirmed in our previous study⁹. Both CD4⁺ and CD8⁺ T cells are involved in protection against viral disease. Prevention of infection can be achieved by induction of CD4⁺ T cells providing help to B cells to produce protective virus-neutralizing antibodies, whereas vaccine-induced CD8⁺ T cells support disease attenuation and protection against complications after infection¹⁹.

In order to be able to accurately measure the low CD8⁺ T cell response after vaccination, we propose that not a single peptide but rather a pool of confirmed immunogenic peptides representing a broad HLA-I coverage should be used. Such a peptide pool would provide a useful tool to monitor and functionally characterize the CD8⁺ T cell response to MuV to better understand its role after natural infection and vaccination.

Methods

Subjects and blood sample handling

Peripheral blood samples of mumps patients were collected within the first month after disease as part of the clinical study VAC-263 (NL37852.094)²⁰. In addition, peripheral blood samples were used that were collected from young adults (18–30 years) four weeks after they received a third dose of the MMR vaccine as part of our MMR3 clinical study⁶ or from our blood donor system²¹. Written informed consent was obtained for all participants. The clinical studies were approved by the medical ethical committee (METC Noord-Holland) and performed in accordance with the Declaration of Helsinki.

HLA class I typing (B*07:02 and HLA-A*01:01) was performed by high resolution next-generation sequencing (University Medical Center Utrecht, the Netherlands) or by flow cytometry using anti-HLA class I A1 antibody (8.L.104, biotin-labeled) or anti-HLA class I B7 antibody (BB7.1, PE-labeled (both Abcam) utilizing LSRFortessaX20 (BD) and analyzed with FlowJo (V10/Treestar). Peripheral blood mononuclear cells (PBMCs) were isolated from the blood samples by Ficoll-Hypaque gradient centrifugation and cryopreserved. Epstein-Barr virus-transformed B-lymphoblastoid cell lines (BLCL) as antigen-presenting cells, were generated from PBMCs as described before²¹.

Prediction of binding of MuV peptides to HLA-A*01:01 and B*07:02 molecules

The 41 MuV peptides, that were previously eluted from HLA class I molecules of MuV-infected cells¹⁶, were tested for predicted binding to the (common) HLA-A*01:01 and B*07:02 molecules using the NetMHCpan 4.0 server. A threshold for selection of best-binding peptides was set on rank < 2%²².

Generation of effector CD8⁺ T cell lines

In order to generate effector CD8⁺ T cell lines, PBMCs from three HLA-B*07:02 positive and three HLA-A*01:01 positive mumps patients were thawed. Cells were stained with fluorochrome-conjugated antibodies anti-CD3-APC/R700, anti-CD4-BV711, anti-CD8-BV786, anti-CD45RO-PE/Cy7 (all from BD Biosciences), and CCR7-PerCP/Cy5.5 (BioLegend). A subset of the effector T cells (CD3⁺, CD8⁺, CCR7⁻, CD45RO⁻) was sorted using FACSMelody sorter (BD) in order to increase MuV-specific T cell frequencies for peptide immunogenicity testing. Sorted T cells were further expanded in AIM-V medium (Gibco) supplemented with 2% human AB serum, with irradiated (3000 rad) allogeneic PBMCs and irradiated (5000 rad) allogeneic BLCL cells, 1 µg/ml phytohemagglutinin (Sigma), and 5 ng/ml IL-2. Every 3–4 days IL-2 (5 ng/ml) was added. Cells were frozen when they reached a total of $\geq 2 \times 10^6$.

Immunogenicity testing for CD8⁺ T cell epitope identification

Thawed CD8⁺ T-effector cell lines were cultured for 6 hours in AIM-V containing 2% human AB serum in the presence of one of the selected best-binding peptides (1 µM; synthesized by JPT Peptide Technologies Inc). During the last 4 hours, Brefeldin A and Monensin (BD) were added. Stimulated cells were stained with fluorochrome-conjugated antibodies, anti-CD3-BUV395, anti-CD4-BV711, anti-CD8-BV786, and Fixable Viability Stain 780 (all BD Biosciences). For intracellular staining, cells were fixed, permeabilized and stained with anti-IFN γ -APC (BD) and anti-TNF-FITC (eBioscience/ThermoFisher), using the Foxp3/transcription factor staining buffer set (eBioscience/ThermoFisher). Cells were acquired on an LSRFortessaX20 (BD) and analyzed using FlowJo (V10/Treestar).

Ex vivo detection of MuV epitope-specific CD8⁺ T cells from mumps patients and vaccinees

Five immunogenic peptides were selected that were able to induce a T cell response. Subsequently, fluorochrome-labeled peptide-HLA class I (p:HLA-I)-dextramers each loaded with one of these five peptides were synthesized: B*0702-APIQGTNLL (FITC-labeled), B*0702-KPRTSTPVTEF (PE-labeled), B*0702-IPNARANL (APC-labeled), A*0101-YSDPNNHEVY (PE-labeled), and A*0101-VTDSNLIY (APC-labeled) (all Immudex). In order to detect *ex vivo* MuV-specific CD8⁺ T cells, 4×10^6 PBMCs from three HLA-B*0702- and three HLA-A*0101-positive mumps patients and from two HLA-B7- and two HLA-A1-positive young adults that recently received a third dose of the MMR vaccine, were stained with a combination of the different p:HLA-I-dextramers and incubated for 10 minutes at room temperature. Subsequently, cells were stained with anti-CD3, anti-CD4, anti-CD8, anti-CD45RO and anti-CCR7 for 15 minutes at 4 °C. Cells were acquired on an LSRFortessaX20.

Expansion of MuV epitope-specific CD8⁺ T cells from mumps patients

Epitope-specific CD8⁺ T cell lines were generated by sorting dextramer-positive cells (FACSMelody sorter (BD)) from HLA-B*0702- and HLA-A*0101-positive mumps patients and subsequent further expanded as described above for effector CD8⁺ T cell lines.

Functional characterization of epitope-specific CD8⁺ T cells

Cytokine secretion analysis

Cytokine release of epitope-specific T cell lines was measured after stimulation with one of the five selected peptides (5 µM) or medium as negative control, and performed in triplicate. Cell-free culture supernatant was collected to determine quantitating levels of IL-2, IL-4, IL-6, IL-10, IL-17A, TNF, IFN-γ, granzyme A (GrA), granzyme B (GrB), perforin, soluble Fas ligand (sFasL), soluble Fas (sFas), and granulysin in a multiplex bead-based assay (LEGENDplex human CD8/NK panel; BioLegend) according to the manufacturers' instructions and using FACSCantoII (BD).

Testing for expression of functional markers

Epitope-specific cell lines were stained with anti-CD3, anti-CD4, anti-CD8, anti-CD45RO and anti-CCR7 for 15 minutes at 4 °C. Anti-CD107a-BV421 (BioLegend) was added during culture. For intracellular staining, cells were fixed, permeabilized, and stained with anti-CD137-PE, anti-IFN-γ and anti-TNF using the Foxp3/transcription factor staining buffer set. Cells were acquired on an LSRFortessaX20.

Testing for cytotoxicity capacity

A flow cytometry-based killing assay was used to determine the cytotoxic capacity of the various epitope-specific T cell lines¹⁸. For this purpose, BLCL as antigen-presenting cells were labeled with CellTrace violet (Life Technologies) and pulsed with one of the five selected peptides (5 µM) or medium as negative control. Subsequently, epitope-specific T cell lines and BLCL were co-cultured at different effector/target (E:T) ratios in AIM-V supplemented with 2% human AB serum. After 4 hours, cells were stained with Fixable Viability Stain 780 (BD) and fixated. Cells were acquired using a LSRFortessa X20. Measurements were made in triplicate and averaged.

Declarations

Acknowledgements

We are grateful to Jacqueline van Gaans for helping with peptide elution experiments.

Author information

Affiliations

Centre for Infectious Disease Control, National Institute for Public Health and the Environment, Bilthoven, the Netherlands

Patricia Kaaijk, Maarten E. Emmelot, Cécile A.C.M. van Els, Jelle de Wit¹

Intravacc (Institute for Translational Vaccinology), Bilthoven, The Netherlands

Hugo D. Meiring,

Contributions

P.K. conceived the project; P.K., J.W. and C.E. designed the study; M.E., H.D.M., P.K. and J.W. planned experiments; M.E. and H.D.M. performed experiments; M.E., P.K. and J.W. analyzed data; M.E. and P.K. prepared figures; P.K. wrote first draft of the manuscript and all authors contributed to the final version of the paper.

Corresponding author

Correspondence to Patricia Kaaijk

Ethics Declaration

Competing interests

The authors declare no competing interests.

Financial support

This work was supported by the Dutch Ministry of Health, Welfare and Sport.

Data availability

NetMHCpan-4.0 server²² was used to predict binding of the set of 41 mumps virus (MuV) peptides to the HLA-A*01:01 and B*07:02 molecules. This server is available via <https://services.healthtech.dtu.dk/service.php?NetMHC-4.0>

References

- 1 Bankamp, B., Hickman, C., Icenogle, J. P. & Rota, P. A. Successes and challenges for preventing measles, mumps and rubella by vaccination. *Curr Opin Virol***34**, 110-116, doi:10.1016/j.coviro.2019.01.002 (2019).
- 2 Rubin, S., Kennedy, R. & Poland, G. Emerging Mumps Infection. *Pediatr Infect Dis J***35**, 799-801, doi:10.1097/INF.0000000000001182 (2016).
- 3 Dayan, G. H. *et al.* Recent resurgence of mumps in the United States. *N Engl J Med***358**, 1580-1589 (2008).

- 4 Cohen, C. *et al.* Vaccine effectiveness estimates, 2004-2005 mumps outbreak, England. *Emerg Infect Dis***13**, 12-17 (2007).
- 5 Kontio, M., Jokinen, S., Paunio, M., Peltola, H. & Davidkin, I. Waning antibody levels and avidity: implications for MMR vaccine-induced protection. *J Infect Dis***206**, 1542-1548, doi:10.1093/infdis/jis568 (2012).
- 6 Kaaijk, P. *et al.* A Third Dose of Measles-Mumps-Rubella Vaccine to Improve Immunity Against Mumps in Young Adults. *J Infect Dis***221**, 902-909, doi:10.1093/infdis/jiz188 (2020).
- 7 Smits, G. *et al.* Seroprevalence of mumps in The Netherlands: dynamics over a decade with high vaccination coverage and recent outbreaks. *PLoS One***8**, e58234, doi:10.1371/journal.pone.0058234 (2013).
- 8 Kaaijk, P. *et al.* Dynamics of the Antibody Response After a Third Dose of Measles-Mumps-Rubella Vaccine Indicate a Slower Decline Compared With a Second Dose. *Open Forum Infect Dis***7**, ofaa505, doi:10.1093/ofid/ofaa505 (2020).
- 9 de Wit, J. *et al.* Mumps infection but not childhood vaccination induces persistent polyfunctional CD8(+) T cell memory. *The Journal of allergy and clinical immunology***141**, 1908-1911 e1912, doi:10.1016/j.jaci.2017.11.047 (2018).
- 10 Jokinen, S., Osterlund, P., Julkunen, I. & Davidkin, I. Cellular immunity to mumps virus in young adults 21 years after measles-mumps-rubella vaccination. *J Infect Dis***196**, 861-867, doi:JID38157 [pii] 10.1086/521029 (2007).
- 11 Hanna-Wakim, R., Yasukawa, L. L., Sung, P., Arvin, A. M. & Gans, H. A. Immune responses to mumps vaccine in adults who were vaccinated in childhood. *J Infect Dis***197**, 1669-1675, doi:10.1086/588195 (2008).
- 12 Vandermeulen, C. *et al.* Evaluation of cellular immunity to mumps in vaccinated individuals with or without circulating antibodies up to 16 years after their last vaccination. *J Infect Dis***199**, 1457-1460, doi:10.1086/598482 (2009).
- 13 Rasheed, M. A. U. *et al.* Decreased humoral immunity to mumps in young adults immunized with MMR vaccine in childhood. *Proc Natl Acad Sci U S A***116**, 19071-19076, doi:10.1073/pnas.1905570116 (2019).
- 14 Connell, A. R., Connell, J., Leahy, T. R. & Hassan, J. Mumps Outbreaks in Vaccinated Populations-Is It Time to Re-assess the Clinical Efficacy of Vaccines? *Front Immunol***11**, 2089, doi:10.3389/fimmu.2020.02089 (2020).
- 15 Ramanathan, R., Voigt, E. A., Kennedy, R. B. & Poland, G. A. Knowledge gaps persist and hinder progress in eliminating mumps. *Vaccine***36**, 3721-3726, doi:10.1016/j.vaccine.2018.05.067 (2018).

- 16 de Wit, J. *et al.* Identification of Naturally Processed Mumps Virus Epitopes by Mass Spectrometry: Confirmation of Multiple CD8+ T cell Responses in Mumps Patients. *J Infect Dis***221**, 474-482, doi:10.1093/infdis/jiz480 (2020).
- 17 Gonzalez-Galarza, F. F. *et al.* Allele frequency net database (AFND) 2020 update: gold-standard data classification, open access genotype data and new query tools. *Nucleic Acids Res***48**, D783-D788, doi:10.1093/nar/gkz1029 (2020).
- 18 de Wit, J. *et al.* The Human CD4(+) T Cell Response against Mumps Virus Targets a Broadly Recognized Nucleoprotein Epitope. *J Virol***93**, doi:10.1128/JVI.01883-18 (2019).
- 19 Siegrist, C. A. *Vaccine Immunology (Chapter 2)*. In: Plotkin SA, Orenstein WA, *Vaccines*. 7th edn, (Elsevier Inc. Philadelphia, 2018).
- 20 Kaaijk, P. *et al.* Dynamics of the serologic response in vaccinated and unvaccinated mumps cases during an epidemic. *Hum Vaccin Immunother***11**, 1754-1761, doi:10.1080/21645515.2015.1040967 (2015).
- 21 Han, W. G. *et al.* Development of an IFNgamma ELISPOT for the analysis of the human T cell response against mumps virus. *J Immunol Methods***431**, 52-59, doi:10.1016/j.jim.2016.02.010 (2016).
- 22 Jurtz, V. *et al.* NetMHCpan-4.0: Improved Peptide-MHC Class I Interaction Predictions Integrating Eluted Ligand and Peptide Binding Affinity Data. *J Immunol***199**, 3360-3368, doi:10.4049/jimmunol.1700893 (2017).

Figures

Figure 1

Predicted binding affinity of the candidate epitopes to HLA and capability to induce cytokine production by T cells is presented. Mumps virus (MuV) candidate epitopes were selected based on their predicted binding (threshold rank <2%) to HLA-B*07:02 molecule (left panel; n=12 peptides) and HLA-A*01:01 molecule (right panel; n= 10 peptides). Selected peptides were used to stimulate peripheral blood mononuclear cells (PBMC) of respectively three HLA-B*0702 positive and three HLA-A*01:01 positive mumps patients. CD8+ T effector cell lines from these patients were stimulated with the selected peptides and production of interferon (IFN)- γ and tumor necrosis factor (TNF) by CD8+ T cells was measured by flow cytometry. Responding T cells were identified as a 2-fold increase of IFN γ + /TNF+ cells within the CD3+/CD4-/CD8+ live gate. Peptides that were tested positive, induced a response in all three HLA-B*07:02 and HLA-A*01:01 positive mumps patients. Filled bars (gray) indicate the peptides that tested positive.

Figure 2

Detection and functional characterization of epitope-specific T cells from mumps patients. (A) Flow cytometric ex vivo detection of epitope-specific T cells using p:HLA-I dextramers in T cellPBMC samples of three HLA-B*07:02 patients (#03, #08 and #20) and three HLA-A*01:01-positive mumps patients (#04, #14 and #25). (B) Cytokine analysis of epitope-specific T cell lines derived from HLA-B*07:02- or HLA-A*01:01-positive mumps patients measured with a multiplex bead-based assay quantitating levels of interleukin-2 (IL-2), IL-4, IL-6, IL-10, tumor necrosis factor (TNF), interferon gamma (IFN-γ), granzyme A (GrA), granzyme B (GrB), perforin, soluble Fas ligand (sFasL), soluble Fas (sFas), and granulysin. Concentrations of measured cytokines are all expressed as means of triplicates in pg/mL. (C) Cytotoxic activity of epitope-specific T cell lines isolated from HLA-B*07:02- or HLA-A*01:01-positive mumps patients measured with a flow cytometry-based killing assay. Specific lysis, presented as mean of triplicates $\pm$ SD, was calculated as follows: % specific lysis = $[(\% \text{ dead target cells} - \% \text{ spontaneous dead target cells}) / (100 - \% \text{ spontaneous dead target cells})] \times 100\%$

Figure 3

Detection of epitope-specific T cells from recently vaccinated young adults. (A) Flow cytometric ex vivo detection of epitope-specific T cells using p:HLA-I dextramers from PBMC samples of two HLA-B7-positive (#IIV-291-034 and VAC-076) and two HLA-A1-positive vaccinees (#IIV-291-002 and #025).