Prevalence of Neonatal Mortality and Its Associated Factor among Neonates Who Admitted In Neonatal Intensive Care Unit At Debre Tabor General Hospital, South Gondar, Amhara, Ethiopia

Solomon kebede Demis (✉ solomondemis@gmail.com)

Debre Tabor University <https://orcid.org/0000-0003-0537-7571>

Tigabu Munye

Debre Tabor University

Biniam Munye

Debre Tabor University

Research

Keywords: Neonatal Mortality, NICU, Debre Tabor General Hospital, South Gondar

Posted Date: January 19th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-147885/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background

Neonatal mortality is the death of newborn babies from the time of birth to 28 completed days of life which are the most vulnerable time for a child's survival. About one million of them passed away on their first day of life, and more than two thirds (38%) of the deaths were in sub-Saharan Africa where Ethiopia is one of the countries with the highest neonatal mortality in the world which accounts for 29 deaths per 1,000 live births.

Objective

To assess the prevalence and associated factors of neonatal mortality among neonates admitted in Debre Tabor General Hospital in South Gondar, Ethiopia.

Methods

Institutional based retrospective cross-sectional study design was conducted from November 1, 2018, up to January 30, 2019, in Debre Tabor General Hospital. A Structured interviewer-administered pre-tested questionnaire was used to collect data. The collected data were entered into Epi data version 4.2 and then exported into SPSS window version 24. Bivariate and multivariate analysis was undertaken and information was presented by using simple frequency tables, graphs, and pie charts.

Result

the prevalence of neonatal mortality was found to be 12.3%. Gestational age group 28–32 weeks (AOR = 9.5, 95% CI: 2.39–37.97), Gestational Age beyond 42 weeks (AOR = 4.6, 95% CI: 6.3–33.8), and forceps delivery (AOR = 0.18, 95% CI: 0.05–0.68) were found to be statistically significant.

Conclusion and recommendation

Neonatal mortality was higher than the national with independently associated factors of prematurity and post maturity while forceps delivery as a preventive factor. Therefore, this might be essential to the hospital NICU to plan for managing prematurity and post maturity in better quality as well as providing quality ANC and identifying most predisposing factors for prematurity.

Introduction

Neonatal morbidity and mortality are the major global public health challenges, specifically for the first 28 days of life. Globally, each year over 4 million neonates died within 28 days of birth among 130 million births. Every minute, twenty under-five children die, leading to 8 million deaths before they reach their fifth birthday due to the conditions which could be either avoided or treated. Neonatal mortality accounts for two-thirds of deaths of infants, and nearly two-fifths of all deaths in under-five children. Many babies die

nameless and unrecorded, indicating the perceived inevitability of their deaths. Most neonatal deaths (99%) occur in low and middle-income countries, where about half of the deaths occur at home [2].

Ethiopia, one of the countries with the highest neonatal mortality in the world, is responsible for 29 deaths per 1,000 live births which was over 9 times more than that of highly developed countries, where the rate is 29 per 1,000 live births [1].

The health of future societies depends on the health of the children of today and their mothers. The neonatal period is considered as the highest risk period. Childhood mortality is often used as a broad indicator of social development or a specific indicator of the health conditions of a country. However, child health programs were given low attention, especially neonatal health [5].

Every year an estimated 4 million babies die in the first 4 weeks of life (the neonatal period). Three-quarters of neonatal death happens in the first week, the highest risk of death is on the first day of life. The highest number of neonatal death was in South Asian countries and the highest rate is generally in sub-Saharan Africa. Preventing death in newborn babies has not been a focus of child survival or safe motherhood programs. While we neglect these challenges, 450 newborn die every hour, mainly from the preventable cause which is unjustifiable in the 21 century [2].

Neonatal mortality accounts for 44% of under-five mortality in 2014. Average death during the neonatal period is 30 times higher compared to the rest of under-five children.

Ten countries account for 67% of neonatal mortality globally by which Ethiopia accounts for 4% of global neonatal mortality [6].

Neonatal mortality has three main causes in low and middle-income countries. The complication of preterm, asphyxia, and neonatal infection together contribute to 85% of newborn death.

In 2013, 35% of the global neonatal deaths were caused by complications of preterm birth, 24% by intrapartum related complications, and 25% by infection. The rest of the death is caused by congenital malformations [7, 8].

In Ethiopia, the rate of under-five mortality (U5MR) decreased by 60% from the year 2000 to 2016 which implies from 123 to 67 per 1000 of live births. However, the neonatal mortality rate (NMR) decreased only by 40% from 49/1000 live births to 29 per 1000 live births from 2000 to 2016. Because of this, the share of neonatal mortality in under-five mortality has been increased from 29.5% to 43% [3].

Neonatal health problems are usually seen as similar to an older child's health problem. But, the causes of neonatal mortality and intervention to improve neonatal health are different from that of other under-five children. Neonatal survival is the most important indicator of improved health care during childbirth. There is a gap in integrating the need for care of neonates in the NICU, the neonatal health problems, and health programs for maternal and child health. As a result, there is still slow progress to decrease

neonatal mortality in the majority part of the country by which they contribute to the national burden of neonatal mortality in Ethiopia.

Even though there were some endeavors in identifying major causes of neonatal death, in Ethiopia, studies that identify the prevalence and associated factors of neonatal mortality were limited specifically for hospital admitted neonates. Having data on the prevalence of the problem is crucial to take prioritized actions. Therefore, this study has done to identify the prevalence of neonatal mortality and associated factors among NICU admitted neonates.

Methods

Study Design and period

An Institutional based cross-sectional study was conducted from November 1, 2018, to January 30, 2019, in Debre Tabor General Hospital in NICU among admitted neonates.

Study setting

Debre Tabor General Hospital is found in the South Gondar *zone* in the Amhara region of Ethiopia. It was officially commenced its function in 1917 and currently, it delivers the health care services for more than 2.3 million populations through medical, surgical, gynecological, pediatrics, NICU, ophthalmological wards, and 14 OPD with a total of 182 beds and 444 staffs.

Annually, nearly more than 1250 neonates were admitted to the neonatal intensive care unit with different health problems. It has 22 beds, 3 pediatricians, and 20 nurses in which 2 are neonatal nurses.

Study population

All neonates who were admitted at NICU in Debre Tabor General Hospital from November 1, 2018, up to January 30, 2019.

Sample size estimation

The sample size was calculated using single population proportion formula and by taking into the following consideration: prevalence (P) of neonatal mortality 13.29% [17] confidence level (CL) 95%, the margin of error (d) 5%, and by adding 10% for non-response rate. The final sample size came up to 195. However, to make a statistical analysis of logistic binary logistic regression, a 50% proportion (p) was used to get the final sample size to be 422.

Data collection and procedure

Before data collection, both the data collectors and supervisors have trained by the principal investigator for one day. Secondary data was collected from Neonatal ICU treatment registers by using a structured questionnaire and checklist designed in the way that it could collect pieces of information about all the

relevant variables. Then it was pre-tested with individuals equivalent to 5% of the calculated sample size among Debre tabor General hospital. Data were collected by two Bsc nurses and one Msc nurse for the supervisor. The data has collected among admitted neonates on a specified period from NICU. A simple random sampling method was used to select neonates admitted in NICU from medical records. This was done by sampling frame 1 up to N [1; 1000] admitted neonates in NICU, then choose 422 samples were chosen every 2 intervals of medical record systematically, so it was taken the sample from a patient chart and register which have full information.

Data quality control

Data quality was assured by proper designing of the questionnaire in Amharic the local language to prevent misinterpretation, apply pre-test on 5% of the sample in Debre Tabor General Hospital. The data have also coded, entered, cleaned, and made close supervision during data collection by the principal investigator.

Data analysis

Then data have coded, entered, and cleaned using Epi-data version 4.2 software and finally exported into SPSS version 24 for analysis. Bivariate analysis, crude odds ratio with 95% CI, was used to see the association between each Independent variable and the outcome variable by using binary logistic regression. Independent Variables with a p-value of ≤ 0.05 were included in the multivariable analysis to control Confounding factors. Adjusted odds ratios with 95% CI were estimated to identify the prevalence and associated factors of neonatal mortality using multivariable logistic regression analysis. The level of statistical significance was declared at p-value ≤ 0.05 .

Ethical consideration

Ethical clearance was obtained from Debre Tabor University, college of health sciences, Institutional Health Research Ethics Review Committee (IHRERC). A formal letter for permission and support was written from Debre Tabor University to Debre Tabor General Hospital. Finally, informed consent was obtained from NICU coordinators.

Results

Sociodemographic characteristics

From a total of 928 selected neonatal charts during study periods, 422 neonatal charts were reviewed. The majority of the maternal age group were 20-34 and the least were age group 20 and below which accounts 284 (67.3%) and 16 (3.7%) respectively. The majority 392 (92.90%) were born from married parents while 219 (51.8%) were from rural family while 203 (48.2%) from urban; and 172 (40.5%) born from mothers with no formal education and 41 (9.7%) were from mothers with completed primary educational level. The majority of neonates were from housewife mothers and the least was from other occupations which account for 223 (52.8%) and 80 (19 %) respectively (**Table 1**).

Prevalence of neonatal mortality

The prevalence of neonatal mortality was found to be 12.3% (**figure 1**).

Neonatal Characteristics

Out of 422 neonates, female, male, and ambiguous genitalia were 214 (50.7 %), 207 (49%), and 1(0.3%) respectively. The majority of neonates were admitted at age less than or equal to 7 days 409 (96.92%) and the least 13 (3.08%) was 8 days or beyond.

Most neonates died at early neonatal age and late neonatal age which accounts for 45 (87%) and 7(13) respectively. Sepsis 203 (48.2%), asphyxia 95 (22.56%), and others 71 (16.92%) were the leading cause of neonatal admission from first to third respectively whereas the majority 209 (53.85%) were low birth weight and the least were very low birth weight which accounts 10 (2.56 %). The majority of neonates were born from mothers whose membrane rupture from 1 to 12 hours 395 (93.58%) and those with 12 to 24 hours duration were 26 (6.15%).

Pregnancy and obstetric characteristics

The majority of maternal factors were duration of labor lasts from one to twelve hours 218 (56.41%), and the least was above twenty-four hours 19 (5.13%). Regarding labor delivery, the majority were through SVD, and least were forceps which accounts for 296 (70.26%) and 33 (7.69%) respectively.

The majority were delivered at health institution 389 (92.28 %) and the majority of presentations were cephalic which accounts for 315 (81.03%). The majority of neonates were term (37-42 weeks) 232 (54.80%) and the least were both post-term (≥ 42 weeks) and unknown gestational age which accounts for 13 (3.10%). The majority 385 (91.28%) had ANC follow up visits whereas 37(8.72%) did not have ANC follow up in which majority 192 (45.64%) had four ANC visits (**Table 2**).

Maternal health problems during pregnancy

The majority, 303 (71.9%) women did not have health problems during pregnancy while the rest 119 (28.2%) had different health problems at the latest pregnancy (**Figure 2**).

Types of health problem during pregnancy

Among 119 women who had a health problem, 86 (72.27%) were with hypertensive disorders during pregnancy and those with other disorders were 2 (1.68%) (**Figure 3**).

Factors associated with neonatal mortality

Binary and multiple logistic regression has been used to identify associated factors. There was no marked variation by mode of delivery, presentation of labor, maternal health problems during pregnancy,

and maternal educational level even though it appeared associated factors of neonatal mortality by bivariate analysis.

Neonatal gestational age group (28-32 weeks) and beyond 42 weeks were risk factors for neonatal mortality while forceps delivery was a protective factor for neonatal mortality.

Neonates with (28-32 weeks) age group were 9.5 times more likely to die than (37-42 weeks) age group. And those with gestational age group > 42 weeks were 4.6 times more likely to die than the gestational age group (37-42 weeks). Those neonates who delivered by forceps were 18% less likely to die than SVD (**Table 3**).

Discussion

In this study, the prevalence of neonatal mortality was 12.3%. The prevalence was higher than the national neonatal mortality with is 29 per 1000 live births (1). The study conducted at Jimma specialized hospital and Felege Hiwot Referral Hospital Ethiopia showed that neonatal mortality was 35.4% which was nearly three times higher than the prevalence in Debre Tabor Hospital and 13.29% which was comparable to this study respectively (14, 17). And this study finding was higher than the study conducted in Dire Dawa Ethiopia which was 11.44% (20). The study conducted in Pakistan showed that neonatal mortality was 27% which was higher than this study (10).

This variation might be due to the different socioeconomic status of the mother and the level of differences among hospitals in which the studies conducted.

The previous studies showed that mode of delivery, educational status of secondary school, marital status, household wealth, male sex and ANC visit, preterm gestational age, and early neonatal age were associated factors for neonatal mortality (17, 18). Similarly, in this study, forceps delivery and gestational age group (28-32 weeks) had a significant association for neonatal mortality. Besides, the gestational age group beyond 42 weeks was associated factors

The neonate who was delivered through forceps were 18% less likely to die than those who delivered by spontaneous vaginal delivery. But, the study in Sudan showed that neonates delivered in C/S were more likely to die than spontaneous vaginal delivery (11). This might be due to the timely management of labor which in turn decreases labor complications.

Gestational age at delivery was significantly associated with neonatal mortality. The preterm newborns (28-32) weeks were 9.5 times more likely to die than those who were delivered at term (37-42) weeks which was 4 times higher than the study conducted at Bahir Dar Felege Hiwot Referral Hospital (17). This might be due to differences in hospital care levels like ANC, institutional delivery, and quality of maternal health service.

Limitation

Secondary data was used that makes it difficult to collect full data from registration and charts due to missing of some treatment outcomes and other relevant data.

Conclusion And Recommendation

The prevalence of neonatal mortality was higher compared to the national figure while Factors associated with neonatal mortality were prematurity (28-32 weeks) of gestational age, post-term (>42 weeks) of gestational age, and preventive forceps delivery. And, the way forwarded, using this stud as input would be providing client-based individualized ANC care to decrease prematurity, enhanced pre term care at NICU, and studies on causes of prematurity as it was the leading factor for neonatal mortality in this study.

References

1. WHO, U. and C. Mathers, Global strategy for women's, children's and adolescents' health (2016-2030). The organization, 2017. 2016(9).
2. Lawn, J.E., et al., 4 million neonatal deaths: when? Where? Why? *Lancet*, 2005. 365(9462): p. 891-900.
3. Tarekegn, S.M., L.S. Lieberman, and V. Giedraitis, Determinants of maternal health service utilization in Ethiopia: analysis of the 2011 Ethiopian Demographic and Health Survey. *BMC pregnancy and childbirth*, 2014. 14(1): p. 161.
4. Antonucci, R., A. Porcella, and M.D. Pilloni, Perinatal asphyxia in the term newborn. *Journal of Pediatric and Neonatal Individualized Medicine (JPNIM)*, 2014. 3(2): p. e030269.
5. Van Lerberghe, W., *The World Health Report 2005: Make every mother and child count*. 2005: World Health Organization.
6. Lawn, J.E., et al., Every Newborn: progress, priorities, and potential beyond survival. *The Lancet*, 2014. 384(9938): p. 189-205.
7. Wang, H., et al., Global, regional, and national levels of neonatal, infant, and under-5 mortality during 1990–2013: a systematic analysis for the Global Burden of Disease Study 2013. *The Lancet*, 2014. 384(9947): p. 957-979.
8. Bhutta, Z.A., et al., Can available interventions end preventable deaths in mothers, newborn babies, and stillbirths, and at what cost? *The Lancet*, 2014. 384(9940): p. 347-370.
9. Tran, H.T., et al., A systematic review of the burden of neonatal mortality and morbidity in the ASEAN Region. *WHO South-East Asia journal of public health*, 2012. 1(3): p. 239.
10. Ayaz, A. and S. Saleem, Neonatal mortality and prevalence of practices for newborn care in a squatter settlement of Karachi, Pakistan: a cross-sectional study. *PLoS One*, 2010. 5(11): p. e13783.
11. Bashir, A.O., et al., Neonatal mortality in Sudan: analysis of the Sudan household survey, 2010. *BMC Public Health*, 2013. 13(1): p. 287.

12. Gizaw, M., M. Molla, and W. Mekonnen, Trends and risk factors for neonatal mortality in Butajira District, South Central Ethiopia,(1987-2008): a prospective cohort study. BMC pregnancy and childbirth, 2014. 14(1): p. 64.
13. Yaya, Y., et al., Maternal and neonatal mortality in south-west Ethiopia: Estimates and socio-economic inequality. PloS one, 2014. 9(4): p. e96294.
14. Debelew, G.T., M.F. Afework, and A.W. Yalew, Determinants and causes of neonatal mortality in Jimma zone, southwest Ethiopia: a multilevel analysis of prospective follow up study. PLoS One, 2014. 9(9): p. e107184.
15. Yirgu, R., et al., Perinatal mortality magnitude, determinants and causes in west Gojam: a population-based nested case-control study. PloS one, 2016. 11(7): p. e0159390.
16. Kolola, T., et al., Determinants of Neonatal Mortality in North Shoa Zone, Amhara Regional State, Ethiopia. PloS one, 2016. 11(10): p. e0164472.
17. Tewabe, T., et al., Neonatal mortality in the case of Felege Hiwot referral hospital, Bahir Dar, Amhara Regional State, North West Ethiopia 2016: a one-year retrospective chart review. Italian journal of pediatrics, 2018. 44(1): p. 57
18. Yared Mekonnen, Biruk Tensou, Daniel S Telake, Ted babe Degeffie and Abeba Bekele Neonatal mortality in Ethiopia: trends and determinants JSI Research & Training Institute, Inc 2013, 13:483. Page 5 of 14
19. Mekonnen al. BMC Public Health Bekele, Neonatal mortality in Ethiopia: trends and Determinants <http://www.biomedcentral.com/1471-2458/13/4832013>, 13:483 P. 6 of 14.
20. Roba AA and Diro DH. Morbidities, Rate, and Time Trends of Neonatal Mortality in Dilchora Referral Hospital, Dire Dawa, Ethiopia, 2012-2017. Austin Med Sci. 2017; 2(2): 1019.

Tables

Table 1. Maternal sociodemographic characteristics of prevalence and associated factors of neonatal mortality among admitted neonates in NICU, Debre Tabor Hospital, 2019.

Characteristics		Frequency	Percentage (%)
Age of mothers	< 20 years	16	3.7
	20-34 years	284	67.3
	>=35 years	122	28.9
Marital status	Married	392	92.90%
	Not married	28	6.63%
	Others	2	0.47
Residence	Rural	219	51.8%
	Urban	203	48.2%
Educational status	No formal education	172	40.5%
	Can read and write	91	21.6%
	Primary school	41	9.7%
	Secondary school	56	13.3%
	College and above	63	14.9%
Occupation	House wife	223	52.8%
	Employers	119	28.2%
	Others	80	19.0%

Table 2. Maternal and neonatal characteristics for prevalence and associated factors of neonatal mortality among admitted neonates in NICU, Debre Tabor General Hospital, 2019.

Variables		
Cause of admission	Frequency (n=422)	Percent (%)
Asphyxia	95	22.56
Sepsis	203	48.20
Jaundice	20	4.62
Pneumonia	20	4.62
Birth trauma	13	3.08
Others	71	16.92
Duration of labor(DOL)	N=385	Percent
1-12 hours	218	56.41
12-24 hours	148	38.46
Above 24 hours	19	5.13
Mode of delivery	N=422	Percent (%)
SVD	296	70.26
Forceps	33	7.69
Vacuum	39	9.23
C/S	54	12.82
Presentation at labor	N=389	Percent (%)
Cephalic	315	81.03
Breach	66	16.92
Others	8	2.06
Gestational age at birth	N=389	Percent (%)
28-32 weeks	45	10.80
32-36 weeks	119	28.20
37-42 weeks	232	54.80
Above 42 weeks	13	3.10
Unknown	13	3.10
Birth weight	N=386	Percent (%)
VLBW	10	2.56

LBW	209	53.85
NBW	167	43.59
ANC visit during pregnancy	N= 422	Percent (%)
Yes	385	91.28
No	37	8.72
Duration of rupture of membrane (ROM)	N=389	Percent (%)
≤ 12 hours	365	93.84
>12 hours	24	6.16

Table 3. The factors for prevalence and associated factors among admitted neonates in NICU, Debre Tabor General Hospital (n=422), 2019.

Variables	Outcome of admission		95% confidence interval	
Mode of delivery	Discharged	Died	COR	AOR
SVD	388 (92%)	34 (8%)	1	1
Forceps	366 (86.7%)	56 (13.3%)	1.76 (0.35-8.82)	0.18(0.05-0.68)*
Vacuum	304 (72.2%)	118 (27.8%)	4.40(1.32-14.64)	0.44(0.07-2.82)
C/S	321 (76%)	101 (24%)	3.61(1.19-10.92)	1.49 (0.41-7.86)
Presentation at labor^a				
Cephalic	352(90.5%)	37(9.5%)	1	1
Breach	295(75.8%)	94(24.2%)	3.05(1.17-7.94)	0.15(0.0-4.61)
Gestational age at birth				
28-32 wks	276(71.4%)	96(24.8%)	3.88(1.23-12.24)	9.5(2.39-37.97)*
32-36 wks	350(90.2%)	38(9.8%)	1.05(0.36-3.06)	2.0(0.60-6.7)
Above 42 wks	259(66.7%)	128(33.3%)	4.85(0.78-29.87)	4.6(6.3-33.8)
37-42 wks	353(90.7%)	35(9.3%)	1	1
Problem during pregnancy^a				
Yes	338(80.8%)	84(20%)	2.44(1.02-5.84)	2.41(0.72-7.92)
No	380(90.7%)	38(9.3%)	1	1
Educational status of mothers^a				
No formal education	375(88.9%)	42(10.1%)	0.43(1.3-1.37)	0.17(0.07-1.55)
Can read and write	338(81%)	80(19%)	0.90(0.27-2.94)	0.99(0.19-5.0)
Primary school	397(94.7%)	21(5.3%)	0.21(0.23-1.93)	0.58(0.03-8.62)
Secondary school	405(96.2%)	16(3.8%)	0.15(0.01-1.37)	0.16(0.01-2.27)
Above secondary	333(79.3%)	87(20.7%)	1	1

Note: ^a variables significant at p <0.2 in bi variable analysis, * significant with p≤0.001, 1= reference.

Figures

Prevalence of neonatal mortality

Figure 1

Prevalence of neonatal mortality in NICU, Debre Tabor General Hospital, 2019.

Health problem

Figure 2

Maternal health problems during pregnancy of prevalence and associated factors of neonatal mortality among admitted neonates in NICU, Debre Tabor General Hospital, 2019.

Figure 3

Types of maternal pregnancy health problems of prevalence and associated factors of neonatal mortality among admitted neonates in NICU, Debre Tabor General Hospital, 2019.