

# Prevalence and determinants of test anxiety among medical students in Addis Ababa Ethiopia

Light Tsegaye Girmay (✉ [leilatsegay@gmail.com](mailto:leilatsegay@gmail.com))

Aksum University

Shegaye Shumet

University of Gondar

Woyanabeba Damene

Amanuel Mental Specialized Hospital

Gebrselassie Gebreegzihabhier

Amanuel Mental Specilized Hospital

Getinet Ayano

Amanuel Mental specialized Hospital

---

## Research article

**Keywords:** Test anxiety, prevalence, medical students, Ethiopia

**Posted Date:** October 18th, 2019

**DOI:** <https://doi.org/10.21203/rs.2.10507/v4>

**License:**  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---

**Version of Record:** A version of this preprint was published on November 14th, 2019. See the published version at <https://doi.org/10.1186/s12909-019-1859-5>.

# Abstract

Background Worldwide, problematic test anxiety is a common health problem among medical students. The magnitude of problematic test anxiety ranges from 25 to 40% in undergraduate medical students and has a detrimental effect on academic achievement and success of students. However, data on the prevalence of test anxiety among medical students is limited. Thus, the study aimed to assess the prevalence and associated factors of test anxiety among medical students. Methods In this cross-sectional study, a stratified random sampling technique was used to select the participants. The level of test anxiety was determined by the Westside Test Anxiety Inventory (WTAI). We utilized logistic regression to explore the association between test anxiety and the potential sociodemographic/student-related characteristics among medical students. Results The study included 423 medical students. Our study resulted the prevalence of problematic test anxiety among medical students to be 52.30% (95% CI 47.40-57.30). The prevalence of test anxiety was remarkably higher in women (79.75%) than in men (33.62%) students. Female sex [AOR=3.25, 95% CI: (1.54, 6.89)], having low grade [AOR=0.11,95% CI: (0.044,0.288)], being first year [AOR=10.55,95% CI: (1.4,76.7)], excessive course load [AOR=6.128,95% CI: (2.675,14.039)], and taking oral examination [AOR=2.89,95% CI: (1.42,5.84)] were determined as some of the predicting factors of test anxiety among medical students. Additionally, lack of systemic study plan [AOR=2.4,95% CI: (1.25,4.59)], poor social support [AOR=3.6,95% CI: (1.56,8.29)], moderate social support [AOR=3.39,95% CI: (1.56,7.4)], psychologically distressed [AOR=2.68,95% CI: (1.37,5.27)] independently predicts test anxiety among medical students. Conclusion Findings suggest that a substantial percentage of medical students had problematic test anxiety in Ethiopia (52.30%). This study also showed a significant association between test anxiety and female sex, having poor grade point average, being the first year, excessive course load, oral examination, lack of study plan, poor social support, moderate social support, and having psychological distress. Problematic test anxiety, which is found to be common among medical students, deserves more attention.

## Background

Test anxiety is referred to as the set of psychological and behavioral responses that accompany concern about likely negative consequences or failure of an exam or similar evaluation situations (1). Test anxiety is a situation-specific trait that refers to the anxiety states and worries conditions that are happened during examinations. This type of anxiety appeared abruptly or gradually. Sometimes it is persistent, and sometimes or ended within a few hours (2-4).

Literature shows that most students experience test anxiety during exam but when the anxiety interferes with the student's capacity to perform in exam adequately and express their knowledge on examinations, it becomes a problem (5). In fact, test anxiety is a great obstacle in the way of many individuals to reach their real academic destination (6)

Test anxiety can be diagnosed using the Diagnostic and Statistical Manual-IV, under the classification of social phobia and has emotional and cognitive components (7). Cognitive test anxiety is the worry part of

cognitive reaction or internal dialogue in the lead up to, during and after evaluative situations, which is correlated with academic performance. Individuals who have high levels of cognitive test anxiety often report feelings of comparing self-performance to peers, worries over the consequences of failure, low levels of confidence in their own abilities as well as too much worry over evaluation. Additionally, the emotional component (sometimes called tension) refers the sharp physiological symptoms stemming from arousal of the autonomic nervous system and related affective responses that students have during the test, which is physiological manifestations include increased galvanic skin response and heart rate, muscle tension dizziness, nausea, or feelings of panic (3, 8).

According to educational psychologists and experts in education, an average level of anxiety is useful as an effective motivational factor can enhance one's performance for more effort. However, excessive anxiety can result in different effects like disturbance of mental processes (4).

Evidence also suggests that medical school programs are characterized by challenging classes and high credit loads in each academic term, which in turn is associated with higher rates of test anxiety, resulting poor academic performance and higher school dropout rates (1). Epidemiologic studies shows that 25 to 40% of undergraduate students experience test anxiety (9, 10).

Several studies revealed that the major factors associated with test anxiety were female sex, inability to manage time, and lacking study skills, low previous grade (GPA), psychological distress, poor self-esteem, and poor social support (3, 11-16). Excessive course load, lack of revision time before the exam (5, 16). In addition, there is variation across faculty, having low academic competence and test competencies were the significant predictors of test anxiety (5, 16).

To the best of our knowledge, there are no previous studies that explored the prevalence and associated factors of test anxiety in undergraduate medical students in Ethiopia. Therefore, this is the first study to assess the prevalence of test anxiety and its determinants in undergraduate medical students in Ethiopia. Additionally, the study is important for a better understanding of the level of test anxiety and its predictors, which enables the responsible stakeholders to intervene, based on the findings.

## **Methods**

### **Study design and population**

This was an institutional-based cross-sectional study conducted from May to May 2018 at Addis Ababa University (AAU). Of 1650 medical students, 390 of them were randomly selected and completed the required questionnaires. All undergraduate medical students were eligible to participate in the current study.

### **Sample Size determination and Sampling techniques**

The sample size (n) to estimate the prevalence of test anxiety among medical students was determined by a single population proportion formula. 95% confidence level was considered to estimate the sample

size and since there are no previous similar studies conducted in Ethiopia, we assumed the prevalence of test anxiety 50% and a precision of 5% between the sample and the parameter was taken.  $\alpha = 0.05$  (95%)  $= 1.96$

We employed the following sample size calculation formula for a single proportion: (See Formula 1 in Supplementary Files)

Additionally, we calculated sample size based on the potential associated factors using STATA 14.2 for sample size calculations, assuming the proportion of test anxiety 50% (no similar study in the area) and taking the odds ratio of 2. In addition, we assumed a significant level of 5% and 80% power. The total sample size was 296 (148 per groups). Adding 10% non-response rate the sample was 320 for each associated factors. However, we used the larger sample size (calculated based on proportion,  $n=423$ ) for this study.

Concerning the sampling technique, in this study, a total of 390 participants, were randomly selected by using a stratified random sampling technique. As suggested, we used a proportionate stratified random sampling to allocated sample for students in each year (year 1 to 5) and we employed a simple random sampling method to select each participant. **(See figure 1).**

## Measures

### Measures for the dependent variable (test anxiety)

Data was collected by a self-administrated questionnaire. Westside Test Anxiety inventory developed by Richard Driscoll (WTAI), is a self-reported questionnaire of 10 statements to which respondents are asked to report how often they experience anxiety symptoms before, during, and after taking tests. Each statement response is scored with a 5-point Likert scale (1–5) yielding a total test anxiety score ranging from 10 to 50 points. Participant were attributed to five different levels of test anxiety according to WTAI score: score of 1.0-1.9 comfortably low-test anxiety, a score between 2.0–2.5 Normal or average test anxiety, 2.5–2.9 High normal test anxiety, 3.0–3.4 Moderately high ,3.5–3.9 High test anxiety ,4.0–5.0 Extremely high anxiety and the cut point for problematic test anxiety is  $\geq 30$ . The Cronbach alpha was alpha of .78, split-half reliability of .77 in a Nigerian sample(17, 18).

Therefore, In this study, those students who scored below 30 were considered as having no test anxiety and those score 30 and above were classified as having test anxiety(18). In our sample, the internal consistency of WTAI was very high (Cronbach's alpha =0.94).

### Measures for the predictor variables

#### Study skills

Study skills referred to as the ability to effectively use the specific skills (learning and planning study, library use, note-taking, course participation, preparation for exams, motivation, preparation for courses,

effective reading, writing, health and nutrition, and listening skills). Time Management (TM) includes six items inventory was used in this study to measure the time management of students it has 6 questions and 5 points Likert scale <20 suggest not effective time management skill (19).

To assess the study skills we used approaches and a study skills inventory for students (ASSIST) (19). The inventory consists of 51 items 5-point Likert scale. It measures the scores on six subscales regarding study habits and skills. These subscales are Textbook Reading, Note-Taking, Memory, Test Preparation, Concentration and Time Management. The domain of Time Management (TM) includes six items of the inventory. The convergent validity of the tool is high and consistency reliability is 0.96 which is

### **Self-esteem**

Rosenberg Self-esteem scale (RSES) was used to measure levels of self-esteem among the study participants (20). Students were scored based on a 4-point Likert scale, scoring of the scale "Strongly Disagree" 1 point, "Disagree" 2 points, "Agree" 3 points, and "Strongly Agree" 4 points. The sums of the scores for all ten items and keep them on a continuous scale. Higher scores indicate higher self-esteem. The scale ranges from 0-30. Scores between 15 and 25 are within normal range; scores below 15 suggest low self-esteem. Internal Validity is 0.78-0.84 And reliability is 0.94 which is high (20).

### **Psychological Distress**

Psychological distress was measured using the Kessler Psychological Distress Scale (K-10). It is a simple measure of psychological distress, the K10 scale involves 10 questions about emotional states each with a five-level response scale. According to the scale, a score 10 - 19 Likely to be well, 20 - 24 Likely to have a mild disorder, 25 - 29 Likely to have a moderate disorder, 30 - 50 Likely to have severe distress. The scale has a sensitivity of 70% and specificity 67% (21).

### **Social support**

We also measured social support of the students and individual who were scored 3-8 poor, 9-11 moderate and 12-14 strong social support on Oslo 3-item social support scale (OSSS-3) (22). Evidence suggests that the OSSS-3 is a very brief and economic scale which has satisfactory internal consistency with an  $\alpha = .640$  (23). Although OSSS-3 is not validated in Ethiopia, it is extensively used in previous studies to assess social support in different population groups (24-26).

### **The sociodemographic and other variables**

We collected self-reported data on sociodemographic and other characteristics of the participants using self-reported and structured questioners.

### **Data collection procedure**

In this study to maintain the quality of the data high, before they engage in the actual data collection activities the data collectors received adequate training on data collection procedure and protocol. Data

was collected by using an English version of self-administrated questionnaire. It had components which assess a different aspect of the participant. The 1<sup>st</sup> part assessed a socio-demographic characteristic of participants. The 2<sup>nd</sup> part assessed the level of test anxiety by using the WTAI scale. The 3<sup>rd</sup> part was on psychosocial factors (psychological distress, self-esteem, and social support) which affect the level of test anxiety and measured by K-10 scale, RSES and oslo3-social support scale respectively. The 4<sup>th</sup> part was on questions assessing behavioral factors using SSI scale to assess time management. The fifth part assessed the academic-related factors that affect test anxiety using yes or no questions. This questionnaire disseminated to 423 eligible medical students.

### **Data quality assurance**

The questionnaire was designed and modified appropriately. The self-administrated English version was disseminated. The training was given for data collectors and supervisor. Pre-test conducted two days before the actual data collection on saint Paulo's medical school and the Pre-test was conducted among 22 samples (5%) of saint Paulo's medical students; the result was not included in the main survey.

Based on the finding from the pre-test, the questioner revised and helped to estimate the time needed for data collection. The data collectors were supervised daily and assist the students to fill the questionnaire and checked daily by the supervisors and principal investigator. The solution to problems during data collection was given immediately by discussing with the supervisors and the data collectors.

### **Data processing and analysis**

First, the data were checked for completeness and consistency and then coded and entered in the computer using EPI DATA software for cleaning, storing and recording and then imported to SPSS version 25 for analysis. Descriptive statistic was used to explain the study participants in relation to study variables. A p-value of less than 0.05 was considered statistically significant. Bivariate and multivariate logistic regression analysis was conducted to identify associated factors of test anxiety among medical students and the strength of the association was presented by odds ratio with a 95% confidence interval.

### **Ethical consideration**

This cross-sectional study was reviewed and approved by the Institutional Review Board (IRB) of the University of Gondar and Amanuel Mental Specialized Hospital. Informed consent from each participant was obtained after clearly explaining the objectives as well as the significance of the study for each study participant. We advised the study participants about the right to participate as well as refuse or discontinue participation at any time they want and the chance to ask anything about the study. The participants were also advised that all data collected would remain confidential.

## **Result**

## **Socio-demographic characteristics of respondents**

Of the total 423 participants reached by random sampling, 390 participants were involved in this study with a response rate of 92.2%. The population was (40.5%) female and (59.5%) male and with a mean of 21.8 years with SD ( $\pm 1.72$ ) ranging 18 to 26 years.

More than half of the participants 245 (62.8%) were Orthodox by religion and most of the respondents were single 385(98.7%) and regarding education level of the respondent's majority of the participant were 3<sup>rd</sup> year 92(23.6%) followed by 2<sup>nd</sup> year 80(20.5%). **(Table 1)**

## **Academic characteristics**

Out of 390 students, 192(49.2%) of them reported maternal educational status as tertiary and above, (75.6%) of them reported history of anxiety that increased during oral exam and (24.4%) reported history of anxiety that increased during a written exam, (43.1%) faced excessive course load, and (73.8%) of them lack enough time to revise before exam. **(Table 2)**

## **Psychosocial characteristics of participants**

In this study, 186 (47.7%) of the participant had psychologically distressed. Regarding social support 138 (35.38%) and 166 (42.57%) had poor and strong social support respectively. In this study, 337 (86.41% )of the participant had high self-esteem. **(Table 3)**.

## **Behavioral characteristics of the participant**

The results of this study found that about 10 (2.6%) of the participant had attempted suicide in their lifetime. Regarding time management, 267(68.5%) of students had poor time management and 24.1% of the participant used the internet more than 5 hours per day and 22.8% used more 30 minutes to one hour per day. **(Table 4)**.

## **Prevalence of test anxiety among medical students**

In this study, more than half (52.30% (95%CI 47.40-57.30)) of medical students had problematic test anxiety. The prevalence of test anxiety in females (79.75%) was remarkably higher than in males (33.62%).

## **Factors associated with the level of test anxiety among medical students**

In this study, female sex, having a poor average grade, being the first year, excessive course load, oral examination, lack of study plan, poor social support, moderate social support, and psychological distress were significantly associated with test anxiety in our final multivariable logistic regression. The final model has the Cox and Snell R-square of 0.523 and 0.79 Neglekerke R-square. The Hosmer and Lemeshow test was not statistically significant (P=0.289).

The odds of having positive test anxiety for a female is 3.25-fold higher than male [AOR=3.25, 95% CI: (1.54, 6.89)]. Regarding grade of students, when the average grade point increase in a unit, the odds of developing severe test anxiety decreased by 88.8% [AOR=0.11,95% (0.044,0.288)].

We also found that the risk of developing positive test anxiety for first-year students is 10 times higher than the fifth-year students [AOR=10.55, 95 % (1.4, 76.7)]. Having excessive course load was 6 times higher risk to develop positive test anxiety [AOR=6.128,95% (2.675,14.039)]

Furthermore, the odds of having positive test anxiety for an oral examination is 2 times higher than that of written examination. [AOR=2.89,95(1.42,5.84)]. The odds of having severe test anxiety for students who did not have a systemic study plan is 2.4 higher than those who had a study plan [AOR=2.4,95% (1.25,4.59)]. The odds of having severe test anxiety for those who have poor social support is 3.6 higher than those who have strong social support [AOR=3.6,95% (1.56,8.29)].

Additionally, the odds of having positive test anxiety for those who have moderate social support is 3.39 higher than those who have strong social support [AOR=3.39,95% (1.56,7.4)]. The odds of having severe test anxiety for those psychologically distressed students is 2.68 times higher than those not distressed [AOR=2.68,95% (1.37,5.27)]

## Discussion

### Main findings

To the best of our knowledge, this is the first study that assessed the prevalence of problematic test anxiety and determinants among medical students in Ethiopia. Findings from the present study demonstrated that more than half of the medical students had problematic test anxiety (52.30%). This prevalence is consistent with the reported magnitude of test anxiety from Saudi Arabia (53%) (27), the USA (55%) (28), and Turkey(48%) (29).

However, the result of the current study was higher than the study conducted at Iran ((43.4%) **(30)**, Malaysia (32.5%) **(2)**, and India (32.3%) **(31)**). This discrepancy might be due to the sampling size difference, the methodological differences including the instrument used to measure test anxiety and differences in the study in the characteristics of the population in each country.

On the contrary, the result of this study is lower than the study conducted in Saudi Arabia and Kenya with the prevalence of (65%)(32) 68.1%) (33) respectively. This difference may be due to the variations in several factors that have an impact on anxiety, such as different course contents, educational environment, test conditions, types of test questions and other factors.

In this study, the prevalence of test anxiety was found to be more than two-fold higher in female students as compared to male students. This result is supported by a study carried out in Sudan and Pakistan (34, 35). The possible reason for this strong association can be due to the reason that all night studying before exams is significantly higher among female students, as compared to male, this can create fatigue

and overall exertion among students which may lead to a lower performance in examinations or due to females over-report their problems than males (36).

On the other hand, the present finding does not agree with the results the study conducted in Iran (10) where no significant association between test anxiety and gender was reported and a study in India (42) where test anxiety is higher in males compared to female students. The discrepancy might be due to variation in the study area and the characteristics and type of the study participants because in the above study the study participants were high school students and the study involved greater number of male than female participants. Biological (including defense mechanisms) (37-40) and environmental factors such as parental pressure (41), previous poor performance, fear of failure and procrastination, expectations, and preparations (42, 43) as well as the nature and the characteristics of the test (43, 44) are the possible explanations for the higher magnitude of test anxiety among the medical students. Additionally, as described by Yerkes-Dodson law, the higher level of test anxiety could be due to an excessive level of arousal which potentially leaving them nervous and unable to concentrate on the test (45).

Regarding the associated factors, this study showed that the score of the students was found to be a significant predictor for test anxiety among medical students. Higher-grade scores was associated with increased odds of developing test anxiety among medical students. This result is supported by studies conducted in Saudi Arabia, Sri Lanka and Nigeria (14, 46, 47) where higher-grade scores were associated with greater risks of test anxiety. On the other hand, a study conducted in Iran showed no significant association between GPA and test anxiety (48). The discrepancy could be due to differences between universities educational environment, teaching and evaluation methods, and systems of rewards and punishments resulting from test results play a role in this regard.

Furthermore, the year of students was also found to be a significant predictor of test anxiety among medical students. The odds of developing test anxiety was 7.0 times higher among first-year medical students as compared to fifth-year students. This might be due to substantial stress at the beginning of the course with more concern and uncertainty about their academic performance at the early stages of the study. The finding of this study is in agreement with studies done in India (31). Conversely, a study conducted in Saudi Arabia among female medical students showed a greeter odds of test anxiety for higher year students as compared with first-year students (11) test. The variation could be due to the study was conducted on female medical students only or due to curriculum difference.

In the present study, test anxiety for an oral examination was 5.5 times higher as compared with a written examination. This finding is in line with the results from studies conducted in Germany and India (49, 50). The possible reason might be due to oral exams require additional skills such as language, social interaction, and communication skills as compared to written exams, which possibly increase the anxiety.

In the current study, the level of test anxiety for students who faced extensive course, the load is four times higher than those who did not face excessive course load. This is in line with studies conducted in Pakistan and Saudi Arabia (32, 51) where extensive course load contributes 90.5% and 23.4% for exam

anxiety, it can be due to the reason that since the medical course is so vast and examination process is so lengthy, students' perception of course load and their ability to manage time with their course work is associated with exam anxiety (32).

Moreover, in this study, test anxiety for students who did not have a systemic study plan was two times higher than those who had a study plan. This results is in line with the reported result from a study conducted in Pakistan (52). The possible reason can be due to Ineffective study habits leads to poor preparation and students encode and store the material inadequately, as a result, they are unable to recall poorly learned material during the examination.

The odds of having high-test anxiety for those who had poor social support is three times higher than those who had strong social support. This finding is in agreement with the findings from studies conducted in Turkey and Iran (53, 54). Having someone around when needed and participating in social activities decrease students' emotional tension and release energy, which in turn decrease anxiety.

Finally, the odds of developing test anxiety was three times higher among students who had psychological distress as compared to those students who had no psychological distress. This result is in line with the reported results from studies conducted in Saudi Arabia (11, 55) where a significant and positive association between psychological distress and test anxiety was identified.

## **Conclusion**

This study showed that the average prevalence of test anxiety was high among undergraduate medical students (52.30%). Female sex, being the first year, lower grade point average, oral exam, excessive course load, lack of systemic study plan, lower social support and psychological distress were significantly associated with the level of test anxiety. Early screening and interventions of test anxiety among medical students were warranted.

## **Abbreviations**

ASSIST: Approaches and Study Skills Inventory for Students

IRB: Institutional Review Board

K-10: Kessler Psychological Distress Scale

RSES: Rosenberg Self-esteem scale

SPSS: Statistical Software for Social Science

TM: Time Management

WTAI: Westside Test Anxiety inventory

# Declarations

## Acknowledgments

We are very grateful to Amanuel mental specialized hospital for funding the study. We are also very grateful to the study participants for their cooperation in providing the necessary information.

## Funding

This research work is funded by the Amanuel mental specialized hospital. However, the funding body has no role in the design of the study and collection, analysis, as well interpretation of data and in writing the manuscript.

## Competing interests

The authors declare that they have no competing interests.

## Consent for publication

Not applicable

## Availability of data and materials

The datasets used and analyzed during the current study are not publically available due to ethical restriction and personal data protections but are available from the corresponding author on reasonable request.

## Ethics approval and consent to participate

This cross-sectional study was reviewed and approved by the Institutional Review Board (IRB) of the University of Gondar and Amanuel Mental Specialized Hospital. Informed consent from each participant was obtained after clearly explaining the objectives as well as the significance of the study for each study participant. We advised the study participants about the right to participate as well as refuse or discontinue participation at any time they want and the chance to ask anything about the study. The participants were also advised that all data collected would remain confidential.

## Authors' contributions

LT conceived the study and was involved in the study design, reviewed the article, analysis, report writing and drafted the manuscript. SS, WD, GZ, and GA were involved in the study design, analysis and drafted the manuscript. All authors read and approved the final manuscript.

# References

1. Kassim MA, Hanafi S, Hancock DRJ. Test anxiety and its consequences on academic performance among university students. 2007;15:17-37.
2. Rajiah K, Coumaravelou S, Ying OW. Relationship of test anxiety, psychological distress and academic motivation among first year undergraduate pharmacy students. *International Journal of Applied Psychology*. 2014;4(2):68-72.
3. Durakua ZH. Factors Influencing Test Anxiety among University Students. 2016.
4. Arbabisarjou A, Zare S, Shahrakipour M, Ghoreishinia G. Analyzing Test Anxiety among Medical Sciences Students of Zahedan in 2015. *INTERNATIONAL JOURNAL OF MEDICAL RESEARCH & HEALTH SCIENCES*. 2016;5(7):334-7.
5. Javanbakht N, Hadian M. The Effects of Test Anxiety on Learners' Reading Test Performance. *Procedia - Social and Behavioral Sciences*. 2014;98:775-83.
6. Malik M, Akhter M, Fatima G, Safder M. Emotional intelligence and test anxiety: A case study of unique school system. *Journal of Elementary Education*. 2013;23(2):49-56.
7. Diagnostic and statistical manual of mental disorders-IV-TR. Washington, DC: American Psychiatric Association. American Psychiatric Association, 2000.
8. Karatas H, Alci B, Aydin H. Correlation among high school senior students' test anxiety, academic performance and points of university entrance exam. *Educational Research and Reviews*. 2013;8(13):919.
9. Alghamdi AR. Test Anxiety: Concept and Implication.
10. Saravanan C, Kingston R, Gin M. Is Test Anxiety a Problem Among Medical Students: A Cross Sectional Study on Outcome of Test Anxiety among Medical Students? *International journal of psychological studies*. 2014;6(3):24.
11. Aziz N, Serafi AH. Increasing Levels of Test Anxiety and Psychological Distress with Advancing Years of Medical Education.
12. Sangsiry SS, Bhosle M, Dutta AP. Predictors of test anxiety in doctor of pharmacy students: An empirical study. *Pharmacy Education*. 2005;5.
13. Kurt AS, Balci S, Kose D. Test anxiety levels and related factors: Students preparing for university exams. *JPM The Journal of the Pakistan Medical Association*. 2014;64(11):1235-9.
14. Gunawardena S, de Zoysa P, Jayasinghe S, Manathunge A, Alles H, Shenoy V, et al. Selected correlates associated with test anxiety among 14-16 year olds in a Colombo district school. *Sri Lanka journal of child health*. 2017;46(2).
15. Gerwing TG, Rash JA, Allen Gerwing AM, Bramble B, Landine J. Perceptions and Incidence of Test Anxiety. *Canadian Journal for the Scholarship of Teaching and Learning*. 2015;6(3):3.
16. Cipra C, Müller-Hilke B. Testing anxiety in undergraduate medical students and its correlation with different learning approaches. *PloS one*. 2019;14(3):e0210130.
17. Driscoll R. Westside Test Anxiety Scale Validation. Online submission. 2007.

18. Rajiah K, Saravanan C. The effectiveness of psychoeducation and systematic desensitization to reduce test anxiety among first-year pharmacy students. *American journal of pharmaceutical education*. 2014;78(9):163.
19. Gadelrab HF. Factorial structure and predictive validity of approaches and study skills inventory for students (ASSIST) in Egypt: A confirmatory factor analysis approach. 2011.
20. Westaway MS, Jordaan ER, Tsai J. Investigating the psychometric properties of the Rosenberg Self-Esteem scale for South African residents of Greater Pretoria. *Evaluation & the health professions*. 2015;38(2):181-99.
21. Andersen L, Grimsrud A, Myer L, Williams D, Stein D, Seedat S. The psychometric properties of the K10 and K6 scales in screening for mood and anxiety disorders in the South African Stress and Health study. *International Journal of Methods in Psychiatric Research*. 2011;20(4):215-23.
22. Dalgard OS DC, Lehtinen V, Vazquez-Barquero JL, Casey P, Wilkinson G, Ayuso-Mateos JL, Page H, Dunn G; ODIN Group Negative life event ,social support and gender difference in depression. *social psychiatry and psychiatric epidemiology*. 2006; 42(6):444-51.
23. Kocalevent R-D, Berg L, Beutel ME, Hinz A, Zenger M, Härter M, et al. Social support in the general population: standardization of the Oslo social support scale (OSSS-3). *BMC Psychology*. 2018;6(1):31.
24. Kebede MA, Anbessie B, Ayano G. Prevalence and predictors of depression and anxiety among medical students in Addis Ababa, Ethiopia. *International journal of mental health systems*. 2019;13:30-.
25. Yohannes K, Gebeyehu A, Adera T, Ayano G, Fekadu W. Prevalence and correlates of post-traumatic stress disorder among survivors of road traffic accidents in Ethiopia. *International journal of mental health systems*. 2018;12:50-.
26. Aberha M, Gebeyehu A, Ayano GJJJoP. Prevalence and factors associated with anxiety among patients with hypertension on follow up at Menelik-II Referral Hospital, Addis Ababa Ethiopia. 2016;19(4):378.
27. Nusrat Aziz , Serafi AH. Increasing Levels of Test Anxiety and Psychological Distress with Advancing Years of Medical Education. *British Journal Of Medical and Health Research*. 2019.
28. Moore B. *Test Anxiety and Nursing Students*. 2013.
29. Kavakci O, Semiz M, Kartal A, Dikici A, Kugu N. Test anxiety prevalence and related variables in the students who are going to take the university entrance examination. *The Journal of Psychiatry and Neurological Sciences*. 2014;27:301-7.
30. Asayesh H, Hosseini MA, Sharififard F, Taheri Kharameh Z. The relationship between self-efficacy and test anxiety among the Paramedical students of Qom University of Medical Sciences. *Journal of Advances in Medical Education*. 2018;1(3):8-12.
31. Patil SG, Aithala MR. Exam anxiety: Its prevalence and causative factors among Indian medical students. *National Journal of Physiology, Pharmacy and Pharmacology*. 2017;7(12):1323-8.

32. Khoshhal KI, Khairy GA, Guraya SY, Guraya SS. Exam anxiety in the undergraduate medical students of Taibah University. *Medical teacher*. 2017;39(sup1):S22-S6.
33. Ndirangu G, Muola JM, Kithuka M, Nassiuma D. An investigation of the relationship between test anxiety and academic performance in secondary schools in Nyeri District, Kenya. *Global journal of educational research*. 2009;8(1-2).
34. Afzal H, Afzal S, Siddique SA, Naqvi S. Measures used by medical students to reduce test anxiety. *JPMA The Journal of the Pakistan Medical Association*. 2012;62(9):982-6.
35. Hashmat S, Hashmat M, Amanullah F, Aziz S. Factors causing exam anxiety in medical students. *JOURNAL-PAKISTAN MEDICAL ASSOCIATION*. 2008;58(4):167.
36. Morales-Casasola A. *Assessment of Test Anxiety on the OSU Lima Campus: Prevalence, Intensity, and Coping Methods: The Ohio State University*; 2018.
37. Liu Y, Lu Z. The relationship between MAOA gene polymorphism and test anxiety. *Twin research and human genetics : the official journal of the International Society for Twin Studies*. 2013;16(6):1103-6.
38. Waqas A, Rehman A, Malik A, Muhammad U, Khan S, Mahmood N. Association of Ego Defense Mechanisms with Academic Performance, Anxiety and Depression in Medical Students: A Mixed Methods Study. *Cureus*. 2015;7(9):e337.
39. Schamess G, Miller RR. *Ego Mechanisms of Defense: A Guide for Clinicians and Researchers* George E. Vaillant. Washington, DC: American Psychiatric Press, 1992. 306 pp. \$38.50 hardback. *Social Work*. 1994;39(2):239-.
40. Vaillant GE, Sobowale NC, McArthur C. Some psychologic vulnerabilities of physicians. *The New England journal of medicine*. 1972;287(8):372-5.
41. Putwain DW, Woods KA, Symes W. Personal and situational predictors of test anxiety of students in post-compulsory education. 2010;80(1):137-60.
42. Salend SJ. Teaching Students Not to Sweat the Test. *Phi Delta Kappan*. 2012;93(6):20-5.
43. Putwain DW. Situated and Contextual Features of Test Anxiety in UK Adolsecent Students. *School Psychology International*. 2009;30(1):56-74.
44. Encandela J, Gibson C, Angoff N, Leydon G, Green M. Characteristics of test anxiety among medical students and congruence of strategies to address it. *Medical education online*. 2014;19:25211.
45. Cohen RA. Yerkes–Dodson Law. In: Kreutzer JS, DeLuca J, Caplan B, editors. *Encyclopedia of Clinical Neuropsychology*. New York, NY: Springer New York; 2011. p. 2737-8.
46. Sideeg A. Test anxiety, self-esteem, gender difference, and academic achievement: the case of the students of medical sciences at Sudanese universities: a mixed methods approach. *British Journal of Arts and Social Sciences*. 2015;9(11):39-59.
47. Balogun AG, Balogun SK, Onyencho CV. Test anxiety and academic performance among undergraduates: The moderating role of achievement motivation. *The Spanish journal of psychology*. 2017;20.

48. Kashfi SM, Jeihooni AK, Kashfi SH, Yazdankhah M. The relationship between test anxiety and educational performance among the students at school of health and nutrition, shiraz university of medical sciences in 2011. *J Contemp Med Edu.* 2014;2(3):158-62.
49. Hahn H, Kropp P, Kirschstein T, Rücker G, Müller-Hilke B. Test anxiety in medical school is unrelated to academic performance but correlates with an effort/reward imbalance. *PloS one.* 2017;12(2):e0171220.
50. Laurin-Barantke L, Hoyer J, Fehm L, Knappe S. Oral but not written test anxiety is related to social anxiety. *World journal of psychiatry.* 2016;6(3):351.
51. Khan AN, Rasool SA, Sultan A, Tahira I. Prevalence of examination related anxiety in a private medical college. *Journal of Ayub Medical College Abbottabad.* 2013;25(1-2):113-5.
52. Numan A, Hasan SS. Effect of Study Habits on Test Anxiety and Academic Achievement of Undergraduate Students. *Journal of Research & Reflections in Education (JRRE).* 2017;11(1).
53. Kurt AS, Balci S, Kose D. Test anxiety levels and related factors: Students preparing for university exams. *Journal of Pakistan Medical Association.* 2014;64:1235-9.
54. Bayani AA. The Effect of Self-Esteem, Self-Efficacy and Family Social Support on Test Anxiety in Elementary Students: A Path Model. *International Journal of School Health.* 2016;3(4).
55. Lyndon MP, Strom JM, Alyami HM, Yu T-C, Wilson NC, Singh PP, et al. The relationship between academic assessment and psychological distress among medical students: a systematic review. *Perspectives on medical education.* 2014;3(6):405-18.

## Tables

**Table 1:** Socio demographic characteristics of medical students at AAU, Tikur Anbessa specialized Hospital, Addis Ababa, Ethiopia, 2018 (n=390)

Variable	Category	Frequency	Percentage
Age	<b>Mean</b>	Standard deviation	
	<b>21.8</b>	±1.72	
<b>Sex</b>			
	Female	158	40.5
	Male	232	59.5
<b>Religion</b>			
	Orthodox	245	62.8
	Muslim	38	9.7
	Protestant	89	22.8
	Others	18	4.6
<b>Marital status</b>			
	Single	385	98.7
	Married	5	1.3
<b>Educational status</b>			
	First year	74	19.0
	Second year	80	20.5
	Third year	92	23.6
	Fourth year	73	18.7
	Fifth year	71	18.2

N.B. others

\* Catholic, ova witness and seventh Adventist

Table 2: Academic characteristics of undergraduate medical students at Tikur Anbessa Specialized Hospital Addis Ababa, Ethiopia, 2018 (n=390)

Variable	Category	Frequency	Percentage
Grade Point average	Mean	Standard deviation	
	3.38	<b>0.48</b>	
<b>Maternal educational status</b>			
	Less than primary	40	10.3
	primary	73	18.7
	secondary	85	21.8
	tertiary and above	192	49.2
<b>Examination type</b>			
	oral exam	295	75.6
	written exam	95	24.4
<b>Excessive course load</b>			
	Yes	168	43.1
	No	222	56.9
<b>Enough revision time</b>			
	No	73.8	73.8
	Yes	26.2	26.2
<b>Study plan</b>			
	No	297	76.2
	Yes	93	23.8

Table 3: Psychosocial characteristics of undergraduate medical students at Tikur Anbessa Specialized Hospital Addis Ababa, Ethiopia, 2018 (n=390)

Variable	Category	Frequency	Percentage
Psychological distress	Yes	186	47.7
	No	204	52.3
Social support	poor	138	35.38
	moderate	86	22.05
	strong	166	42.57
Self-esteem	low	8	2.05
	average	45	11.54
	high	337	86.41

Table 4: Behavioral and clinical characteristics of undergraduate medical students at Tikur Anbessa Specialized Hospital Addis Ababa, Ethiopia, 2018.

Variable	Category	Frequency	Percentage
Suicidal attempt	Yes	10	2.6
	No	380	97.4
Time management	Poor	267	68.5
	Good	123	31.5
Internet use habit	more than 5hr	94	24.1
	3hr-4hr	121	31.0
	1.30min-2hr	86	22.1
	30min-1hr	89	22.8

Table 5. Factors associated with level of test anxiety among undergraduate medical students at Tikur Anbessa Specialized Hospital (Bivariate and multivariate analysis) (n=390), Addis Ababa, Ethiopia, 2018.

Variable	Test anxiety		COR(95%CI)	AOR(95%CI)
	No	Yes		

	Age		Sex	COR(95%CI)	AOR(95%CI)
	mean	standard deviation			
	21.18	1.719		0.83(0.73,0.93)	1.084(0.71,1.64)
Female	32	126		7.77(4.8, 12.4)	3.25(1.54, 6.89)
Male	154	78		reference	

Grade point average	mean	standard deviation
---------------------	------	--------------------

3.38	0.48	0.03(0.02,0.05)	0.11(0.044,0.288)
------	------	-----------------	-------------------

Educational level
-------------------

1 <sup>st</sup> year	20	54	3.69(1.83,7.4)	10.55(1.4,76.7)
2 <sup>nd</sup> year	37	43	1.58(0.83,3.02)	1.81(0.339,9.7)
3 <sup>rd</sup> year	49	43	1.19(0.64,2.23)	1.68(0.49,5.76)
4 <sup>th</sup> year	39	34	1.19(0.61,2.3)	0.5(0.168,1.49)
5 <sup>th</sup> year	41	30	reference	

Maternal educational status
-----------------------------

< Primary	16	24	1.5(0.75,2.99)	1.89(0.628,5.73)
Primary	35	38	1.08(0.63,1.86)	1.07(0.42,2.67)
Secondary	39	46	1.17(0.7,1.96)	1.63(0.69,3.86)
Tertiary	96	96	reference	

Exam type
-----------

Oral 5.84)	108	187	7.94(4.46, 14.1)	2.89, (1.42,
Written	78	17	reference	

Course load
-------------

Yes	42	126	5,53(3.55,8.64)	6.128,(2.675,14.039)
No	144	78	reference	

Study plan
------------

No	110	187	7.6(4.27,13.5)	2.4(1.25,4.59)
Yes	76	17	reference	

Social support

Poor	20	118	19.2(10.6,34.8)	3.6(1.56,8.29)
Moderate	39	47	3.92(2.25,6.84)	3.39(1.56,7.4)
Strong	127	39	reference	

Internet use

30min-1hr	45	44	reference	
1.30min-2hr	38	48	1.29(0.71,2.34)	0.98(0.37,2.59)
3-4hr	53	68	1.31(0.75,2.27)	1.85(0.702,4.88)
>5hr	50	44	0.9(0.5,1.6)	1.72 (0.62,2.5)


Time management

Poor	101	166	3.67(2.33,5.79)	1.25(0.62,2.5)
Good	85	38	reference	

Psychological distress


Distressed	49	137	5.71(3.68,8.85)	2.68(1.37,5.27)
Not distressed	137	67	reference	

## Figures



**Figure 1-schematic presentation of sampling technique (n=390)**

Figure 1



**Figure 2:** Prevalence of test anxiety among undergraduate medical students at Addis Ababa University, Addis Ababa, Ethiopia, 2018.

¶

Figure 2

## Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Formula1.jpg](#)