

# Expression of mitochondrial protein genes encoded by nuclear and mitochondrial genomes correlate with energy metabolism in dairy cattle

Jigme Dorji (✉ [jigme.dorji@agriculture.vic.gov.au](mailto:jigme.dorji@agriculture.vic.gov.au))

AgriBio Centre for Agriculture Bioscience <https://orcid.org/0000-0003-4057-8918>

**Christy J. Vander Jagt**

Agriculture Victoria, AgriBio, Centre for AgriBioscience, Bundoora

**Josie B. Garner**

Agriculture Victoria, Ellinbank Dairy Centre, Ellinbank, Victoria, 3822

**Leah C. Marett**

Agriculture Victoria, Ellinbank Dairy Centre, Ellinbank, Victoria

**Brett Mason**

Agriculture Victoria, Ellinbank Dairy Centre, Ellinbank, Victoria

**Coralie M. Reich**

Agriculture Victoria, AgriBio, Centre for AgriBioscience, Bundoora, Victoria

**Ruidong Xiang**

Agriculture Victoria, AgriBio, Centre for AgriBioscience, Bundoora, Victoria

**Emily L. Clark**

The Roslin Institute and Royal (Dick) School of Veterinary Studies, University of Edinburgh, Edinburgh, Scotland

**Benjamin J. Cocks**

Agriculture Victoria, AgriBio, Centre for AgriBioscience, Bundoora, Victoria

**Amanda J. Chamberlain**

Agriculture Victoria, AgriBio, Centre for AgriBioscience, Bundoora, Victoria

**Iona M. MacLeod**

Agriculture Victoria, AgriBio, Centre for AgriBioscience, Bundoora, Victoria

**Hans D. Daetwyler**

Agriculture Victoria, AgriBio, Centre for AgriBioscience, Bundoora, Victoria

---

## Research article

**Keywords:** mitochondria, energy metabolism, differential gene expression, gene co-expression, cattle

**Posted Date:** February 24th, 2020

**DOI:** <https://doi.org/10.21203/rs.2.24317/v1>

**License:** © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

---

**Version of Record:** A version of this preprint was published on October 19th, 2020. See the published version at <https://doi.org/10.1186/s12864-020-07018-7>.

## Abstract

Background Mutations in the mitochondrial genome have been implicated in mitochondrial disease, often characterised by impaired cellular energy metabolism. Cellular energy metabolism in mitochondria involves mitochondrial proteins (MPs) from both the nuclear (Nu MP) and mitochondrial (Mt MP) genomes. The expression of MP genes in tissues may be tissue specific to meet varying specific energy demands across the tissues. Currently, the characteristics of MP gene expression in tissues of dairy cattle are not well understood. In this study, we profile the expression of MP genes in 29 adult and six foetal tissues in dairy cattle using RNA sequencing and gene expression analyses: particularly differential gene expression and co-expression network analyses.

Results MP genes were differentially expressed (DE; over-expressed or under-expressed) across tissues in cattle. All 29 tissues showed DE Nu MP genes in varying proportions of over-expression and under-expression. On the other hand, DE of Mt MP genes was observed in <50% of tissues and notably Mt MP genes within a tissue was either all over-expressed or all under-expressed. A high proportion of Nu MP (up to 60%) and Mt MP (up to 100%) genes were over-expressed in tissues with expected high metabolic demand; heart, skeletal muscles and tongue, and under-expressed (up to 45% of Nu MP, 77% of Mt MP genes) in tissues with expected low metabolic rates; leukocytes, thymus, and lymph nodes. These tissues also invariably had the expression of all Mt MP genes in the direction of dominant Nu MP genes expression. The Nu MP and Mt MP genes were highly co-expressed across tissues and co-expression of genes in a cluster were non-random and functionally enriched for energy generation pathways. The differential gene expression and co-expression patterns were validated in independent cow and sheep datasets.

Conclusions This study demonstrated the biological interaction of MP genes from the mitochondrial and nuclear genomes and their over-expression in tissues with high energy demand. This highlights the importance of considering MP genes from both genomes in future studies related to mitochondrial functions and traits related to energy metabolism.

## Background

There is growing evidence that mitochondrial dysfunction arise from variations in the mitochondrial genome and that their interplay with the nuclear genome has a role in mitochondrial diseases in humans, including metabolic disorders and diabetes [1–3]. Mitochondria and mitochondrial functions are critical for tissues with high energy requirement [4]. Energy is produced in mitochondria through a process of oxidative phosphorylation (OXIPHOS). Besides energy production, mitochondria mediate programmed cell death (apoptosis), aging, calcium homeostasis, and signalling as reviewed in [5–7].

Mitochondrial proteins (MP) are the proteins localized in mitochondria and are key component to mitochondrial functions [8]. There are an estimated 1500 proteins in mitochondria of rats [9], participating as components of electron transport chain, metabolic pathways, and factors for replication, initiation and regulation in transcription and translation. To date, 1158 MPs stand verified in human [10] and almost all MPs (> 99%) are of nuclear origin (NuMP) and imported into the mitochondria [11, 12] with the exception for 13 proteins (< 1%), which originate from the mitochondrial genome (MtMP). Mitochondria have their own genome, which is inherited maternally [13–15]. The cattle mitochondrial genome is haploid with a small circular structure (~ 16.4 kb) with a coding region encoding for 37 genes (13 proteins, 22 tRNAs and 2 rRNAs) and a non-coding control region [16]. Mitochondrial DNA mutations in cattle have previously been shown to be associated with fertility and productivity [17–20], and environmental adaptability to high altitudes in yaks [21, 22]. Unlike nuclear DNA, mitochondrial genomes occur in multiple copies, and their numbers are relatively constant within a cell type and development stage but vary considerably among cell types [23–25].

Gene expression is referred to as one of a series of processes from gene activation to mature protein function that contributes to the expression of cellular phenotypes [26]. The expression of a gene is often specific to tissue types, and a notable example is the dominance of major milk protein transcripts in the bovine lactating mammary gland [27]. Gene expression is commonly studied using RNA sequencing (RNAseq) where the number of reads mapping to a gene (counts) is used to measure gene expression.

The characterization of gene expression, identification of gene function, gene-disease or gene-production associations from genome-wide gene expression [28] employs differential gene expression and co-expression network analyses. Differential gene expression compares the gene expression in a sample with another sample or group of samples. Gene co-expression analysis measures the correlation between the expression levels of genes and associates gene clusters with biological processes and facilitates prediction of gene function of previously unknown genes [29]. At a very local level, co-expression of small groups of genes results from being in close proximity [30, 31] and in chromosomal domains characterized by frequent internal DNA-DNA interactions known as topological association domains (TADs)[32].

Most RNAseq based gene expression analyses to date have focused on nuclear genes rather than genes from the mitochondrial genome [33]. Nonetheless, a comprehensive examination of MP genes from both genomes is central to understanding genome-genome interactions, their role in meeting specific energy demand, and development of mitochondrial diseases. Metabolic profiles and energy demands vary widely across organs and tissue types [34–36]. The varying demand for energy across tissues is possibly in part facilitated through tissue specific and differential expression of MP genes. Currently, a comprehensive study on the expression of MP genes (both NuMPs and MtMP) across tissues is lacking in bovine, although the expression of individual or groups of MtMP genes has been published as part of larger gene sets [33, 37]. Therefore, our study aimed to characterize MP gene expression across both adult and foetal tissues in dairy cattle. We used RNAseq of 35 tissues from two adult cows and two foetuses (29 adult and six foetal tissues) to investigate differential gene expression and gene co-expression. We validated our findings using publicly available RNAseq data for an additional dairy cow and three sheep.

## Methods

### Data

This study utilizes three RNAseq datasets: i) two Holstein cows and their foetuses in this experiment (Main Cows), ii) one Holstein cow (Validation Cow) [27], and iii) gene expression counts from three sheep (Validation Sheep) [37] from previously published studies. The Main Cows dataset had RNAseq from 29 tissues from two adult cows and six tissues from two 16 weeks old foetuses. The Validation Cow data consisted of RNAseq reads from 18 tissues, and the Validation Sheep data were gene expression counts for a subset of tissues (15 tissue types) of three Texel x Blackface adult females. The tissue-specific gene expression patterns in the Main Cows dataset were validated using the validation datasets.

### Ethics, animals and tissue sampling

The ethical approval, including the permission to animals of the Main Cows datasets were obtained from the Department of Jobs, Precincts and Regions Animal Ethics Committee (Application No. 2014-23). Two lactating and pregnant Holstein cows and their two foetuses at 16 weeks of gestation from the Agriculture Victoria Research dairy herd at Ellinbank, Victoria, Australia (38°14' S, 145°56' E) were chosen for the study. Cow 2181 had a male fetus (2181F), and cow 6819 had a female fetus (6819F), both foetuses were from the same sire (half-sibs).

Blood samples were drawn from the coccygeal vein by venipuncture and processed following the blood fractionation and white blood cell stabilization protocols of the RiboPure™ blood kit (Ambion by Life Technologies). Other tissues were sampled following euthanasia of the animals. The cows were euthanised individually by a trained veterinarian and not within line of sight of another deceased animal to minimise stress. The cow was restrained in a crush and given an intravenous injection of 600 mg of xylazine IV adequate to cause moderate sedation. The cow was immediately released from the crush, and once the sedation had taken effect and the cow was sitting down, 300 mg of ketamine was given intravenously. Once the cow laid down, 1 litre of 25% magnesium sulphate solution was administered intravenously until pronounced deceased by the veterinarian. Once pronounced dead, all tissue types were dissected from the animal. Connective tissue was removed, and the samples were dissected into 1 cm squares, sealed in a 5 ml tube and flash-frozen in liquid nitrogen. Blood (on ice) and tissues samples (in liquid nitrogen canisters) were then moved to the main laboratory and stored at -80 °C. The metadata and RNAseq reads for all 40 tissues are available at EMBL-EBI European Nucleotide Archive (ENA) under study accession ERP118133. For this study, we generated data for 35 samples (29 tissues from adult cows and six from the foetuses) (Table 1).

Table 1

List of 35 organ-tissue sampled from the two adult cows and two foetuses in the Main Cows dataset

Tissues/organs	Tissues/organs	Tissues/organs
Adrenal gland cortex	Lung	Spleen
Adrenal gland medulla	Mammary gland (Mammary)	Thymus
Omental fat pad (Omental fat)	Heart	Thyroid gland (Thyroid)
Subcutaneous fat	Brain cerebellum	Tongue
Kidney cortex	Brain stem	Blood leukocytes (Leukocytes)
Kidney medulla	Brain caudal lobe	
Longissimus thoracic muscle (Loin muscle)	Pituitary gland	Foetal brain (fBrain)
Semimembranosus muscle (Leg muscle)	Placenta	Foetal kidney (fKidney)
Intercostal muscle (Rib muscle)	Liver	Foetal lung (fLung)
Latissimus dorsi muscle (Msub)	Skin black	Foetal heart (fHeart)
Mediastinal lymph node (Mediastinal LN)	Skin white	Foetal liver (fLiver)
Lymph node	Ovary	Foetal leg muscle (fLeg muscle)

## RNA extraction and sequencing

RNA from blood leukocytes was extracted using the RiboPure Blood Kit (Ambion) according to the manufacturer instructions. For tissues, 100 mg of tissue was ground in a TissueLyserII (Qiagen) with liquid nitrogen, and then ~ 30 mg of ground tissue was used to extract RNA using Trizol (Invitrogen) following standard procedures. RNA was passed through a PureLink RNA Mini column (Qiagen) and eluted in 30 µl RNase free water. RNA Integrity Numbers (RIN), which indicates the RNA quality, were determined using Agilent TapeStation (Agilent) and RNAseq libraries were prepared from all samples with RIN > 6 using the SureSelect Strand-Specific RNA Library Prep Kit (Agilent) as instructed by the manufacturer. Libraries were barcoded uniquely, assigned randomly to one of two pools and sequenced on a HiSeq™ 3000 (Illumina) in a 150-cycle paired-end run. One hundred and fifty bases paired-end reads were called with bcltfastq and output in fastq format. The quality of the libraries and alignment are as presented in Additional file 1. Poor-quality bases were filtered, and sequence reads trimmed using an in-house script. Bases with a quality score of < 20 were trimmed from the 3' end of reads. Reads with a mean quality score < 20, >3 N's, or final length < 50 bases were not included. Only paired reads were retained for alignment.

## Read alignment and gene counting

For each library, paired-end reads were mapped to Ensembl bovine genome UMD3.1 reference and annotated using STAR version 2.5.3ab [38]. Aligned reads were checked for quality using Qualimap 2 [39], and unique mapping reads for samples (Additional file 2). The R package featureCounts [40] was used to generate a count matrix of read counts per gene for every sample.

## Mitochondrial protein genes

Mitochondrial protein genes in the current study were based on the list of MP identified in humans, available in Mitocarta 2.0 [10]. The official gene names of 1158 MP genes were directly converted to bovine ensemble gene IDs using a gene ID conversion function in the software DAVID (Database for Annotation, Visualization, and Integrated Discovery) version 6.8 [41, 42]. This translated into 1054 bovine MP ensemble gene IDs (1,041 NuMPs and 13 MtMP), which were used as the final list of MP genes for further analysis in this study (Additional file 3). Additionally, 24 non-protein coding genes from the mitochondrial genome (22 tRNAs and 2 rRNAs) were also included in the analysis.

## Differential gene expression analysis

The lowly expressed genes were filtered out using function `filterByExpr` of `edgeR` package for differential expression analysis in R [43]. Differential expression of genes was analysed using the `glmQLFit` function. A model was fitted to the data with a design matrix of an overall mean of gene expression counts across all other tissues as the intercept and tissue as a fixed effect, i.e. differential expression is relative to the average expression of the same gene across all other tissues. The `glmQLTest` method was used to identify DE genes, specifically up or down expressed. A list of DE genes, along with their fold changes, was generated and summarized for each tissue. A gene was considered as differentially expressed (DE) in tissue if its expression was significantly higher than the mean expression of same gene across all other tissues (i.e.  $\geq |0.6| \log_2$  fold changes (LFC) = 1.5-fold difference, FDR < 0.01). The sign + and - of LFC values of the DE gene was used to deduce the expression as either over-expression or under-expression respectively compared to the mean of expression across all other tissues. Further,  $\log_2$ CPM (counts per million) values of all NuMP and MtMP genes (including non-DE MP genes) across tissues were visualized as heatmap using R package `pheatmap` [44]. In addition, we looked at the number of DE MP genes by genome (i.e. NuMP and MtMP), their direction of expression and the proportion of DE genes to the total genes in category. The foetal tissues were analysed separately following the procedures implemented for the adult cows.

## Co-expression network analysis across tissues

The functionally associated genes tend to be co-expressed, and this is used to infer novel function as well as to identify candidate genes in diseases and their prediction [28]. To study the co-expression pattern in tissues, we used a similarity network based on a Pearson correlation coefficient of gene expression ( $>|0.95|$ ), executed using a plugin `ExpressionCorrelation` [45] in `Cytoscape 3.6.1` [46], to construct gene co-expression clusters. We analysed the co-expression cluster involving MP genes for;

- i. biological significance of the cluster using functional enrichment analysis and composition of the genes,
- ii. whether the co-expression of NuMP genes in NuMP-MtMP cluster was greater than random expectations using Chi-square goodness of fit ( $\chi^2$ );

$$\chi^2 = \sum \frac{(O_i - E_i)^2}{E_i}$$

Where,

$O_j$  = observed frequency of genes ( $j$ ) in the NuMP-MtMP cluster ( $j$  = NuMP, Non-NuMP gene)

$E_j$  = expected frequency of genes ( $j$ ) from the overall clusters ( $j$  = NuMP, Non-NuMP gene)

- iii. the effect of TAD in co-expression of NuMP genes in NuMP-MtMP cluster considering TAD as one of the several factors potentially influencing the co-expression of small group of genes. Briefly, we mapped the co-expressed bovine genes across the clusters (3,643) to the putative bovine TADs derived from the IMR90hg18 [47] and 3,022 genes mapped to 1,286 TADs. Similarly, within the NuMP-MtMP cluster, 651 of 813 co-expressed genes mapped to 484 TADs. Of this, 282 co-expressed genes were distributed in groups of 2 or more per TAD. We compared this to averages from 100 random samples of 651 genes from TAD mapping genes across all clusters (3,022). The averages for number of TADs of the random samples and genes found in group of 2 or more within a TAD were 472 ( $\pm 10$ ) and 116 ( $\pm 10$ ) respectively.

## Functional enrichment analysis

The DAVID software was used to investigate the functional enrichment of differentially expressed genes within a tissue and co-expressed genes across tissues: up to 3000 genes (maximum permissible in DAVID) were selected and analysed for over-representation in KEGG (Kyoto Encyclopedia of Genes and Genomes) pathways [48]. Up to top 10 pathways with adjusted  $p < 1e-5$  are discussed in this study.

## Validation

The patterns of MP gene expression in tissues of Main Cows were validated using two previously published datasets from a lactating Holstein cow consisting of 18 tissues [27] (i.e. Validation Cow) and three adult female Texel x Scottish Blackface sheep (i.e. Validation Sheep) (15 tissues) from the sheep gene expression atlas project [37]. The RNAseq reads of the Validation Cow were processed, aligned, gene counts generated and analysed following the protocols for Main Cows. Similarly for sheep, the raw gene counts [49] were normalized and subjected to standard processing and analyses for differential expression and co-expression. In sheep, 823 MP genes were identified as overlapping the Mitocarta 2.0 Human database, using the same approach as in cattle (Additional file 4). The pattern of MP gene expression across tissues was visualized with a heatmap and co-expression networks as described for Main Cows. One of the purposes of validation was to look at the consistency of gene expression patterns across datasets. To evaluate the consistency of differential expression of MP gene expression in a tissue across the datasets, a scatterplot of the LFC values of DE NuMP genes (in common between the datasets) and their coefficient of determination ( $R^2$ ) was used to indicate correlation between datasets. For consistency in co-expression of MP genes, the NuMP-MtMP co-expression cluster was further examined for the composition and commonality of genes among the datasets.

## Results

### Differential expression of mitochondrial protein genes

#### Main Cows: Adult

In total, 16,166 genes including 1,041 MP genes were available for analysis after filtering. A gene was considered as differentially expressed (DE) in a tissue if the expression was significantly different from the average expression across all other tissues (LFC  $>|0.6|$ , FDR  $< 0.01$ ). Across all genes, 13 to 40% of 24,616 genes in total were DE in one or more tissues and as high as 50% each of the DE genes were over-expressed or under-expressed (Fig. 1). Table 2 summarizes the number of DE genes by category across tissues. The highest overall numbers of DE genes among the tissues were in blood leukocytes, brain caudal lobe, loin muscle, and brain cerebellum, and the lowest in the ovary, omental fat pad, and mediastinal lymph node. The DE genes of heart, skeletal muscles and tongue were significantly enriched for OXIPHOS, metabolic pathways and neurodegenerative diseases pathways while liver and kidney cortex were enriched for metabolic pathways (Table 3).

Tissue	MG <sup>l</sup> (37)			NuMP* (1,041)			All <sup>l</sup> (24,616)			MtMP <sup>l</sup>	Mt	Mt
	Over-exp	Under-exp	Total DE	Over-exp	Under-exp	Total DE	Over-exp	Under-exp	Total DE	(13)	tRNA <sup>y</sup>	rRNA <sup>l</sup>
Adrenal cortex	0	0	0	66	157	223	2018	2532	4550	-	-	-
Adrenal medulla	2	0	2	124	89	213	1620	2260	3880	-	-	2
Brain caudal lobe	2	0	2	190	151	341	3354	4150	7504	-	-	2
Brain cerebellum	1	0	1	120	187	307	3282	3879	7161	-	-	1
Brain stem	0	0	0	95	161	256	1911	3294	5205	-	-	-
Heart	17	0	17	576	53	629	2185	2833	5018	13	2	2
Kidney cortex	5	0	5	315	30	345	1465	2851	4316	<i>ATP8, CYB, COX2</i>	-	2
Kidney medulla	0	0	0	93	58	151	1618	2444	4062	-	-	-
Latissimus d muscle	14	0	14	481	74	555	2668	2970	5638	13	1	-
Leg muscle	14	0	14	456	80	536	2773	3083	5856	13	1	-
Leukocytes	0	9	9	99	368	467	5270	3948	9218	<i>ATP8, ND3, COX3, COX2, ATP6, CYB, COX1, ND1, ND2</i>	-	-
Liver	0	0	0	360	82	442	2914	3124	6038	-	-	-
Loin muscle	14	0	14	514	114	628	3853	3707	7560	13	1	-
Lung	0	5	5	31	430	461	1793	2565	4358	<i>ND3, ATP6, COX3, CYB, ND1</i>	-	-
Lymph node	0	7	7	38	306	344	2649	2562	5211	<i>ND1, ND3, ATP6, ND4, ND4L, CYB, ND2</i>	-	-
Mammary	0	0	0	86	250	336	1805	1864	3669	-	-	-
Mediastinal LN	0	0	0	46	77	123	993	2432	3425	-	-	-
Omental fat	0	2	2	78	163	241	1107	1901	3008	-	-	2
Ovary	0	0	0	58	70	128	985	2018	3003	-	-	-

Pituitary gland	0	5	5	73	251	324	2522	3033	5555	<i>ND1, ND3, CYB, ATP6, ATP8</i>	-	-
Placenta	0	10	10	110	232	342	2842	3267	6109	<i>ND3, ND1, ATP6, COX3, ND2, CYB, ND4L, ND4, COX2, ND5</i>	-	-
Rib muscle	7	0	7	500	70	570	2325	2933	5258	<i>ND1, ND3, COX2, ATP6, CYB, ND4L, ND2</i>	-	-
Skin black	0	0	0	79	176	255	1814	2574	4388	-	-	-
Skin white	0	0	0	79	205	284	1780	2554	4334	-	-	-
Spleen	2	0	2	47	260	307	2003	2449	4452	-	-	2
Subcutaneous fat	0	0	0	105	95	200	1299	2238	3537	-	-	-
Thymus	0	10	10	63	363	426	3509	3208	6717	<i>ND3, ATP6, ND1, CYB, ND4, COX3, COX1, ND4L, ATP8, COX2</i>	-	-
Thyroid	0	2	2	33	329	362	2335	2335	4670	<i>ND1, ND3</i>	-	-
Tongue	14	0	14	413	30	443	1369	2365	3734	13	1	-
<p>( ) total number of genes in a category; Over-exp Over-expression, under-exp under-expression; MG<sup>Ⓜ</sup> Genes from mitochondrial genome including tRNA and rRNAs, <i>NuMP</i>* Mitochondrial protein genes encoded by the nuclear genome, <i>MtMP</i><sup>Ⓜ</sup> Mitochondrial protein genes encoded by mitochondrial genome, <i>Mt</i> tRNA<sup>Ⓜ</sup> Mitochondrial transfer RNA, <i>Mt</i> rRNA<sup>Ⓜ</sup> Mitochondrial ribosomal RNA, All<sup>Ⓜ</sup> all genes from nuclear and mitochondrial genomes</p>												

Table 2. Number of differentially expressed (DE) genes in tissues by gene categories averaged for two cows in the Main Cows dataset

Table 3

KEGG functional annotation of overall differentially expressed genes of selected tissues with the largest number of genes averaged across two cows in the Main Cows dataset

Tissues	Enrichment	No. of genes (Overlap NuMP*)	Adj. p	Other pathways
Heart	Oxidative phosphorylation	101 (93)	3.3e <sup>-53</sup>	Parkinson's disease, Alzheimer's disease, Huntington's disease, NAFLD, carbon metabolism, Metabolic pathways, cardiac muscle contraction, TCA cycle,
Leg muscle	Oxidative phosphorylation	83 (78)	1.0e <sup>-32</sup>	Parkinson's disease, Alzheimer's disease, NAFLD, Huntington's disease, Carbon metabolism, metabolic pathways, Proteasome, cardiac muscle contraction Biosynthesis of antibiotics
Loin muscle	Oxidative phosphorylation	89(86)	5.8e <sup>-39</sup>	Parkinson's disease, Alzheimer's disease, NAFLD, Huntington's disease, Huntington's disease, carbon metabolism, Proteasome, Cardiac muscle contraction
Rib muscle	Oxidative phosphorylation	86 (81)	1.8e <sup>-35</sup>	Parkinson's disease, Alzheimer's disease, NAFLD, Huntington's disease, Metabolic pathways, carbon metabolism, Proteasome
Tongue	Oxidative phosphorylation	98(91)	3.0e <sup>-50</sup>	Parkinson's disease, Alzheimer's disease, NAFLD, Huntington's disease, Metabolic pathways, carbon metabolism, cardiac muscle contraction,
Kidney cortex	Metabolic pathways	312(118)	3.9e <sup>-28</sup>	Biosynthesis of antibiotics, Carbon metabolism, Valine, leucine and isoleucine degradation, Glycine, serine and threonine metabolism, tryptophan metabolism, Fatty acid metabolism
Kidney medulla	Focal adhesion	67(0)	3.0e <sup>-10</sup>	Tight junction, calcium signalling pathway, Gastric acid secretion. ECM-receptor interaction, cGMP-PKG signalling pathway, Gastric acid secretion, Dilated cardiomyopathy
Liver	Metabolic pathways	387(123)	4.1e <sup>-49</sup>	Biosynthesis of antibiotics, Peroxisome, valine, leucine and isoleucine degradation, complement and coagulation cascades, fatty acid degradation, tryptophan metabolism, carbon metabolism
Brain caudal lobe	Axon guidance	57(0)	8.0e <sup>-18</sup>	Glutamatergic Synapse, Domaminergic synapse, MAPK signalling pathway, Adrenergic signalling in cardiomyocytes, Retrograde endocannabinoid signalling, cAMP signalling pathway, Synaptic vesicle cycle, GABAergic synapse, Morphine addiction, Glutamatergic synapse, Circadian entrainment, Dopaminergic synapse
Brain cerebellum	Glutamatergic synapse	49(2)	3.1e <sup>-14</sup>	GABAergic synapse, Retrograde endocannabinoid signalling, Morphine addiction, Circadian entrainment, Dopaminergic synapse, cAMP signalling pathway, Adrenergic signalling in cardiomyocytes, axon guidance,
Brain stem	GABAergic synapse	36(2)	3.1e <sup>-9</sup>	Glutamatergic synapse, Morphine addiction, Retrograde endocannabinoid signalling, Dopaminergic synapse, Circadian entrainment
Leukocytes	Chemokine signalling pathways	74	4.7e <sup>-15</sup>	Focal adhesion, leukocyte transendothelial migration, Rap1 signalling pathway, natural killer cell mediated cytotoxicity, regulation of actin cytoskeleton, B cell receptor signalling pathway
*we show the number of genes in the top enriched pathway that overlap mitochondrial proteins				

More than 99% of the MP genes originate from the nuclear genome (NuMP). The proportion of DE NuMP genes across the tissues varied from 12 to 60% with higher proportions (> 50%) in heart and skeletal muscles. The proportion of under or over-expressed DE NuMP genes within tissues varied considerably. A relatively greater proportion of NuMP genes were over-

expressed in heart, kidney cortex, skeletal muscles, and tongue, and under-expressed in blood leukocytes, lymph nodes, placenta, lungs, mammary, and thymus (Fig. 2). The expression of NuMP genes was similar between animals in the Main Cows as indicated by the clustering together of same tissues, with the exception of five tissues (Fig. 2; adipose, ovary, kidney cortex, and leukocytes).

In contrast to NuMP, differential expression of MtMP genes were observed in less than 50% of tissues (14 out of 29 tissues). The proportion of DE MtMP genes within tissues ranged widely from 0 (no genes) to 100% (all 13 MtMP genes). Specifically, MtMP genes were 100% DE in heart, leg muscle, latissimus dorsi muscle, loin muscle, and tongue, and ranged between 50–75% in other tissues (leukocytes, placenta, thymus, rib muscle, and lymph node). Unlike NuMP genes, all DE MtMP genes were expressed in a single direction (i.e. either all over-expressed or all under-expressed) meaning every DE MtMP gene was over-expressed in heart, tongue, muscles and kidney cortex, and under-expressed in blood leukocytes, placenta, lymph node, pituitary, thymus, and thyroid (Fig. 3). Further, there were similarities between the expression of DE MtMP and NuMP genes within a tissue. For instance, every tissue showing over-expression of DE MtMP genes invariably showed predominant over-expression of NuMP genes and similarly for under-expression. In addition to MtMP genes, some of the non-protein coding genes from the mitochondrial genome were also DE in several tissues (16 s rRNA, 12 s rRNA, tRNA-Pro and tRNA-Ser).

Within groups of tissues with either over-expression of MP genes (heart, skeletal muscles, liver and kidney cortex) or under-expression (leukocytes, thymus, placenta and lymph node), we examined all overlapping genes and their functional enrichment. In heart and skeletal muscles, there were 1,088 over-expressed genes in common including 320 NuMP and seven MtMP genes. Altogether across these 1,088 genes, there was significant enrichment for OXIPHOS, metabolic pathways and neurodegenerative disease pathways as in these individual tissues. Similarly, liver and kidney cortex had 1,249 over-expressed genes in common including 223 NuMP genes (0 MtMP genes) and these were significantly enriched for metabolic pathways and peroxisome, valine, leucine and isoleucine degradation. In contrast, the DE genes in common for tissues in the under-expression group was low (63) with only 20 NuMP genes (0 MtMP genes). Across all 63 genes, there was enrichment for adrenergic signalling in cardiomyocytes, dilated cardiomyopathy, cardiac muscle contraction and hypertrophic cardiomyopathy. Altogether, these results indicated a significant role of the over-expressed MP genes contributing to the enriched pathways in the over-expression tissue group, while this pattern was not observed in the under-expression group.

## **Main Cows: Foetuses**

The analysis for functional enrichment of overall DE genes in six foetal tissues showed significant enrichment of OXIPHOS and metabolic pathways only in heart and lungs but not in leg muscle (Additional file 5). The NuMP genes were over-expressed in the heart and under-expressed in the remaining tissues, including leg muscles (Additional file 6). Similarly, the MtMP genes were prominently over-expressed in heart, under-expressed in the lungs and not significant in the remaining tissues (Additional file 7). Between the two foetuses, liver of the male fetus showed a relatively higher expression of NuMP genes compared to the female.

## **Co-expression network analysis of mitochondrial protein genes**

The gene co-expression network constructed based on the affinity matrix from genes correlated in expression  $> |0.95|$  had altogether 3643 genes clustered into four major network clusters I-IV (Fig. 4). The NuMP genes were concentrated in two main clusters (I and IV) indicating co-expression among NuMP genes and the remaining NuMP genes were sparsely scattered across all other clusters. Similarly, MtMP genes were all grouped in cluster I. Clusters I and IV containing subgroups of highly co-expressed NuMP and MtMP and NuMP genes are referred to as NuMP-MtMP and NuMP clusters respectively. Within the NuMP-MtMP cluster, the MP genes from the respective genomes were highly co-expressed. The NuMP-MtMP cluster was significantly enriched for OXIPHOS, metabolic pathways and mitochondrial diseases' pathways. Similarly, the NuMP cluster (cluster IV) was over-represented for signalling pathways, contraction, metabolic pathways and myopathies related to the heart (Table 4). The gene functions of the non-mitochondrial protein genes (Non-MP) in the NuMP-MtMP cluster were associated with heart and muscle functioning, signalling, and contraction (Additional file 8,9).

Table 4

Summary of gene, composition, and functional enrichment of KEGG pathways of genes in co-expression clusters (FDR < 1e-05) in the Main Cows dataset

Cluster	No. of MtMP genes	No. of NuMP Genes	No. of Non-MP genes	Total No. of genes	Enrichment of pathways
I (NuMP-MtMP Cluster)	13	216	584	813	Parkinson's disease, Oxidative phosphorylation, Alzheimer's diseases, Huntington diseases, Non-alcohol fatty liver diseases, metabolic pathways, Citrate cycle, carbon metabolism, Cardiac muscle contraction, Proteasome
II	0	10	871	881	Retrograde endocannabinoid signaling, GABAergic synapse, Nicotine addiction, Morphine addictions, Glutamatergic synapse, Dopaminergic synapse, Synaptic vesicle cycle, Neuroactive ligand-receptor interaction
III	0	12	923	935	Cell adhesion molecules (CAMs), Staphylococcus aureus infection, intestinal immune network for IgA production, Leishmaniasis, Antigen processing and presentation, viral myocarditis, Allograft rejection, primary immunodeficiency, Hematopoietic cell lineage, Natural killer cell-mediated cytotoxicity
IV (NuMP cluster)	0	79	466	545	Chemical carcinogenesis, Complement and coagulation cascade, Drug metabolism – cytochrome p450 metabolism, steroid hormone biosynthesis, retinol metabolism, Metabolic pathways, Complement and coagulation cascades, bile secretion, primary bile acid biosynthesis, tryptophan metabolism, carbon metabolism, fatty acid metabolism

We tested if the co-expression of NuMP genes in the NuMP-MtMP cluster was due to random chance using a Chi-square goodness of fit test. The frequency of NuMP genes in the cluster was significantly higher than random ( $\chi^2 = 307.6$ ,  $p < 0.01$ ), supporting that the cluster was enriched with co-expressed MPs.

Further, we investigated the effect of TAD on the co-expression by comparing the number of 651 TAD mapped genes in the NuMP-MtMP cluster with the mean from 100 randomly generated samples of 651 genes from 3022 TAD mapping genes across the clusters. It showed involvement of NuMP-MtMP genes in a similar number TADs ( $472 \pm 10$  vs 484), but NuMP-MtMP were more likely to be present in groups of two or more within TADs. The total number of genes occurring in a two or more in a TAD was 282 and 116 ( $\pm 6$ ) in NuMP-MtMP cluster and random samples respectively. This indicated that co-expression in the NuMP-MtMP cluster was enriched within TADs.

## Validation of patterns of mitochondrial protein gene expression

The key findings on the expression of MP genes from the Main Cows dataset were validated using two independent datasets (Validation Cow and Validation Sheep). Both validation sets confirmed the general trends of MP gene expression and coexpression in tissues.

Firstly, both validation datasets confirmed the over-expression of MP genes in heart and skeletal muscles, and under-expression in blood leukocytes as in the adult tissues of the Main Cows dataset (Additional file 10–17). Further, expression of MP genes within tissues, as indicated by LFC values between Main Cows and Validation Cow (Fig. 5), were highly correlated ( $R^2$  0.67–0.96) except for thyroid ( $R^2$  0.013). Similarly, the correlation of LFC values between Main Cows and Validation Sheep was high ( $R^2$  0.6–0.87), except for mammary and lungs ( $R^2$  0.34, 0.37) (Additional file 18). We investigated the poor correlation of gene expression in thyroid between the Main Cows and the Validation Cow. At least 35 DE NuMP genes in common between the datasets were expressed in opposite directions. These genes were mainly enriched for metabolic pathways, pyruvate

metabolism and synthesis of antibiotics. Interestingly, the expression of NuMP genes between Validation Cow and Validation Sheep were moderately correlated in the thyroid ( $R^2$  0.59) (Additional file 19).

Secondly, the 'either all over-expression or all under-expression' of DE MtMP genes within tissues was supported by findings from both validation datasets. Further, the expression of MtMP genes in the direction of the dominant DE NuMP genes also remained evident across datasets.

Thirdly, the co-expression of MtMP and NuMP genes in a cluster were reproduced in the Validation Cow (Additional file 20), and to some extent in Validation Sheep (Additional file 21). The co-expression of MP genes in the NuMP-MtMP cluster in the Validation Cow was more than expected by random chance ( $\chi^2=207.847$ ,  $p < 0.01$ ) indicating that this cluster was enriched for co-expression of MP genes.

Finally, the overlap of genes in NuMP-MtMP clusters across the Main Cow and validation datasets was higher than would be expected if genes were randomly allocated to clusters. In particular, the occurrence of MtMP genes were almost coincidental (13/13) between cow datasets and 12/13 genes in common between cow and sheep datasets. Similarly, a considerable proportion of NuMP genes and also non-mitochondrial protein genes, were in common across datasets (Fig. 6).

## Discussion

This study described and validated specific patterns of differential expression for over 1000 mitochondrial protein genes, encoded by the nuclear and mitochondrial genomes, in bovine across different tissues. The study also presented strong evidence of co-expression between NuMP and MtMP genes.

## Differential expression of mitochondrial protein genes

Overexpression of MP gene in more metabolically active tissues

The observed patterns of differential expression of MP genes within tissue, where the proportion of DE MP genes exceeded 40%, appears to correlate with the known metabolic demand of tissues. MP genes were over-expressed in tissues with high reported metabolic demand (heart, skeletal muscles, tongue, and kidney cortex: Table 5), and under-expressed in tissues with low reported energy demand (adipose tissue: Table 5 and blood leukocytes). In humans, about 60–70% of the total resting energy expenditure is accounted for by kidney, brain, liver, and heart, which altogether constitute less than 6% of adult body weight, while skeletal muscle (40–50% of body weight) accounts for 20–30% of resting energy expenditure [4, 50], which altogether account up to 80% of energy expenditure.

The heart meets almost the entire energy demand through the mitochondrial OXIPHOS pathway (95%) [52]. A higher expression of selected MtMP genes (ND1,ND5,ATP6, CYTB) were reported in the heart compared to other tissues (brain, kidney, liver and skeletal muscle) in mice [53], which supports heart as the tissue with the highest MtMP gene expression. Similarly, skeletal muscles, which has a low resting energy demand, are capable of spiking by almost 1000-fold during short intensive exercise [54, 55]. OXIPHOS is highlighted as an important pathway for generating energy during the exercise/muscular activity in both short intensive as well as prolonged exercise [56]. The observed a higher expression of MP genes in the tongue seems likely because the tongue is a muscular organ. Furthermore, results from the heart and skeletal muscles group reinforced the OXIPHOS and Metabolic pathways as important for energy metabolism in these tissues.

A high expression of MP genes specifically in kidney cortices may be attributed to energy generation taking place at the proximal and distal convoluted tubules, which are also the site for active reabsorption of metabolites [57, 58]. In kidney and liver, the enrichment for Metabolic pathways and not for OXIPHOS pathway, despite their high energy demand, is suggestive of non-OXIPHOS pathways predominating for energy metabolism.

## Tissues with under-expression of mitochondrial protein genes

Among the tissues with under-expression of MP genes, only adipose tissue in human has been published. In keeping with our observed low MP gene expression in adipose, the metabolic rate of human adipose tissues was low (3.2–4.6 kcal/kg/day) [59]. On the other hand, leukocytes (and other tissues with under-expression for MP) have mainly non-energy related mitochondrial functions, such as redox signalling and controlling apoptosis [60], which, in part, could explain the incidence of under-expression of MP genes in blood leukocytes. Further, results from the analysis of group of tissues showing MP gene under-expression revealed a low number of DE genes in common across these tissues and no enrichment for energy pathways supports a diminished role of mitochondrial energy function.

As for the adult cows, the highest expression of MP genes in the foetal heart tissue was expected considering the early foetal development and establishment of the heartbeat occurs as early as three weeks in the bovine foetus. In contrast to adult cows, the low expression of MP genes in foetal leg muscle was likely attributable to only partial development and non-functionality of the muscle. Skeletal muscle development, mainly secondary myogenesis, is initiated in the foetal stage from 9 weeks post-fertilization to parturition [61] and our foetal calves were around 16 weeks old. Generally, the remaining foetal tissues measured in this study are reported to be functionally inert in the foetal development stage, including lungs [62] and explains the under-expression of the MP genes.

In general, the expression profiles of MP genes in a tissue were consistent as indicated by clustering of same tissue of two or more animals within the dataset. Nonetheless, some tissues were exceptions, including foetal livers may be attributed to the sampling and cellular heterogeneity of the samples because each cell type may have a specific expression profile.

To sum up, the expression of MP genes in this study concurs with the energy demand of tissues (where known) implying that the increased energy demand may be met through increased expression of MP genes. Furthermore, previous studies report that there is a specific correlation between mRNA and protein quantity across tissues [63, 64].

## **Mitochondrial genome encoded mitochondrial protein (MtMP) gene expression**

Besides, energy demand in tissues as the basis of increased transcription rates of MP genes, high MtMP gene expression could also result from increased mitochondrial genome copy numbers. Mitochondrial DNA copy number differs considerably across tissue types but remains closely regulated within a tissue type [23]. Studies in humans indicate that mitochondrial genome copy numbers are aligned with tissue energy demands: for example heart, skeletal muscle, omental fat, and blood leukocytes had 6970, 3650, 400–600 and 91 copies per diploid nuclear genome respectively [65–67]. Studies comparing copy number and gene expression of all MtMP genes across tissues are scarce. A study in striated muscles (cardiac, type 1 skeletal muscle and type 2 skeletal muscles) of rabbit [68] demonstrated that the expression of MtMP gene (CYTB) was proportional to copy number. Thus, it is plausible that the varying energy requirements of tissue types are at least partly modulated through their mitochondrial DNA copy number.

## **Direction of expression of differentially expressed mitochondrial protein genes**

There were two interesting aspects of the direction of DE MtMP genes; first, the 'all over-expression or all under-expression' of DE MtMP genes within tissues and second, the directional consistency of DE MtMP genes expression in the dominant direction of DE NuMP genes. The first phenomenon of occurrence of DE MtMP genes in single direction has not been previously reported to the best of our knowledge. A possible explanation of this phenomenon rests in the mechanism of transcription because the entire mitochondrial genome is transcribed as a near-complete polycistronic unit [69, 70], so that almost all mitochondrial genes are transcribed as one unit. The initiation of transcription particularly at HSP2 promotor site generates a near-complete polycistronic unit [71]. The second trend showing the common direction of DE MtMP and NuMP genes was observed in all tissues exhibiting significant DE of MtMP genes. This suggests DE NuMP and MtMP in these tissues are co-regulated.

## **Coexpression of mitochondrial protein genes**

The co-expression of mitochondrial protein genes was a prominent finding in the current study. Co-expression of MP gene in MtMP-NuMP cluster was further tested to be non-random and non-random co-expression of genes are previously reported across species [72]. Further, the significant enrichment of NuMP-MtMP co-expression cluster for OXIPHOS and Metabolic supports co-functional coexpression of genes [29, 73]. Similarly, results from MP gene expression study in humans showed a significant correlation between MtMP and NuMP gene expression within tissues [74], suggesting close coordination between nuclear and mitochondrial genomes in relation to energy demand. The functional enrichment of our NuMP-MtMP cluster for the OXIPHOS pathway and non-MP genes in the cluster for heart myopathies, contraction and signalling, emphasize their role in energy metabolism and supporting systems. The NuMP cluster was enriched for metabolic pathways which is another important energy metabolism component of mitochondria.

The investigation of involvement of TAD on the co-expression demonstrated that the co-expressed genes in NuMP-MtMP cluster occurring in two or more within a TAD compared to the random sample. This indicated the potential role of TADs in co-expression of mitochondrial protein gene in our study. As such, the intra-TAD gene co-expression was not different from random for most chromosomes in another study [75].

## Validation

Overall, there were high correlation and consistency evident in the expression (differential expression and co-expression) of mitochondrial protein genes in tissue across the datasets. However, we have not taken into account for the physiological states, number tissues sampled and sequencing platforms employed in our validation study. Firstly, a notable difference in expression profile of NuMP genes in the thyroid between the Main Cows and the Validation Cow, and Main Cows and Validation Sheep is potentially related to pregnancy of the Main Cows, as the Validation Cow and the Validation Sheep were not pregnant. The activity of thyroid and thyroid hormone synthesis are reportedly increased during pregnancy in human [76] and thyroid hormones are known to regulate metabolism [77]. The interaction of thyroid and MP function in metabolism is an area of interest for further investigation but beyond the scope of current work. Secondly, we based the differential gene expression of a gene in tissue to the mean expression across all other tissues where both the number of tissues and tissue types were not completely identical across the datasets (29 tissues in Main Cows, 18 tissues in Validation Cow and 15 tissues in Validation Sheep). Thereby, expression in tissue across the datasets has been compared to the mean expression of different sample sizes, which might vary across the datasets. Thirdly, the sequencing platforms used were different for each dataset: the Main Cows dataset were sequenced on HiSeq™ 3000 (Illumina), the Validation Cow was sequenced on HiSeq2000 sequencer (Illumina) and Validation Sheep were sequenced on Illumina HiSeq 2500.

## Conclusions

Mitochondrial protein genes were differentially expressed across tissues. Tissues with high energy demand showed over-expression and under-expression was observed in tissues with low energy requirements, which suggests a link between mitochondrial protein gene expression and the energy demand of each tissue. Furthermore, mitochondrial protein genes from both genomes (NuMP and MtMP) were significantly co-expressed and enriched for co-functionality. This implies that it is necessary to consider mitochondrial protein genes from both genomes in studies related to mitochondrial function. Mitochondrial protein gene expression analysis may be extrapolated to production traits such as feed efficiency, heat tolerance, adaptability to cold climate, to further elucidate their role in relation to energy metabolism.

## Abbreviations

<b>CPM</b>	Counts Per Million
<b>DAVID</b>	Database for Annotation, Visualization, and Integrated Discovery
<b>DE</b>	Differentially expressed
<b>EDTA</b>	Ethylenediaminetetraacetic acid
<b>FDR</b>	False discovery rate

<b>KEGG</b>	Kyoto Encyclopedia of Genes and Genomes
<b>LFC</b>	Log2 fold change
<b>MP</b>	Mitochondrial protein
<b>MPG</b>	Mitochondrial protein genes
<b><i>Mt</i></b>	Mitochondrial
<b><i>MtMP</i></b>	Mitochondrial genome encoded mitochondrial proteins
<b>NAFLD</b>	Non-alcohol fatty liver disease
<b><i>NuMP</i></b>	Nuclear genome encoded mitochondrial proteins
<b>OXIPHOS</b>	Oxidative phosphorylation
<b>PBS</b>	Phosphate Buffer Saline
<b>RIN</b>	RNA integrity number
<b>RNA</b>	Ribonucleic acid
<b>rRNA</b>	Ribosomal ribonucleic acid
<b>TAD</b>	Topologically associating domain
<b>TCA</b>	Tricarboxylic acid
<b>tRNA</b>	Transfer ribonucleic acid

## Declarations

Ethics approval and consent to participate

Use of animals and ethical approval for this experiment was granted by the Department of Jobs, Precincts and Regions Animal Ethics Committee (Application No. 2014-23).

Consent to publish

Not applicable

Availability of data and materials

RNAseq datasets for two cows are available at EMBL EBI European Nucleotide Archive (ENA) under study accession ERP118133 at <https://www.ebi.ac.uk/ena/data/search?query=ERP118133>, and for Validation Cow in fastq format under study accession number ERP107617 at <https://www.ebi.ac.uk/ena/data/search?query=ERP107617>. The raw gene counts from three adult female Texel X Blackface (1, 2 and 3) are from the sheep gene expression atlas dataset available at <https://doi.org/10.7488/ds/2616>. The data generated and analysed in our study supporting the conclusions of the article are included in the Additional files.

## Competing interests

The authors declare that they have no competing interests.

## Funding

The study was funded by DairyBio project (a joint venture between Agriculture Victoria, Dairy Australia, and the Gardiner Foundation). JD is a recipient of Full Fee Remission Scholarship from La Trobe University (LUFFRS). However, funders did not have direct role in the study.

## Authors' contributions

The conception of the experiment: JD, IMM, BGC, AJC, CJV, HDD

Sampling, RNA sequencing and processing: AJC, CJV, JBG, LM, BM, CMR, JD

Data analysis JD, IMM, RX, CJV, AJC, ELC, HDD

Manuscript preparation JD, IMM, CJV, AJC, HDD

Manuscript revision JD, IMM, BGC, LM, CMR, JBG, HDD

Management IMM, BGC, HDD

All authors have read and approved the final manuscript.

## Acknowledgements

The authors are grateful to the Agriculture Victoria staff at the Ellinbank and Bundoora sites that were involved in sample collection. The authors thank the DairyBio project (a joint venture between Agriculture Victoria, Dairy Australia, and the Gardiner Foundation) for funding. JD receives a La Trobe University fee remission scholarship.

## References

1. Taylor RW, Turnbull DM: Mitochondrial DNA mutations in human disease. *Nature Reviews Genetics* 2005, 6(5):389-402.
2. Gorman GS, Chinnery PF, DiMauro S, Hirano M, Koga Y, McFarland R, Suomalainen A, Thorburn DR, Zeviani M, Turnbull DM: Mitochondrial diseases. *Nature Reviews Disease Primers* 2016, 2:16080.
3. Wallace DC: Mitochondrial Diseases in Man and Mouse. *Science* 1999, 283(5407):1482.
4. Elia M, Livesey G: Energy expenditure and fuel selection in biological systems: the theory and practice of calculations based on indirect calorimetry and tracer methods. *World review of nutrition and dietetics* 1992, 70:68-131.
5. Wang C, Youle RJ: The role of mitochondria in apoptosis\*. *Annu Rev Genet* 2009, 43:95-118.
6. Sun N, Youle RJ, Finkel T: The Mitochondrial Basis of Aging. *Molecular cell* 2016, 61(5):654-666.
7. Contreras L, Drago I, Zampese E, Pozzan T: Mitochondria: The calcium connection. *Biochimica et Biophysica Acta (BBA) - Bioenergetics* 2010, 1797(6):607-618.
8. Fox TD: Mitochondrial protein synthesis, import, and assembly. *Genetics* 2012, 192(4):1203-1234.
9. Lopez MF, Kristal BS, Chernokalskaya E, Lazarev A, Shestopalov AI, Bogdanova A, Robinson M: High-throughput profiling of the mitochondrial proteome using affinity fractionation and automation. *ELECTROPHORESIS* 2000, 21(16):3427-3440.

10. Calvo SE, Clauser KR, Mootha VK: MitoCarta2.0: an updated inventory of mammalian mitochondrial proteins. *Nucleic Acids Research* 2016, 44(Database issue):D1251-D1257.
11. Dudek J, Rehling P, van der Laan M: Mitochondrial protein import: Common principles and physiological networks. *Biochimica et Biophysica Acta (BBA) - Molecular Cell Research* 2013, 1833(2):274-285.
12. Chacinska A, Koehler CM, Milenkovic D, Lithgow T, Pfanner N: Importing mitochondrial proteins: machineries and mechanisms. *Cell* 2009, 138(4):628-644.
13. Margulis L: Origin of eukaryotic cells : evidence and research implications for a theory of the origin and evolution of microbial, plant, and animal cells on the Precambrian earth. New Haven: New Haven : Yale University Press; 1970.
14. Giles RE, Blanc H, Cann HM, Wallace DC: Maternal inheritance of human mitochondrial DNA. *Proceedings of the National Academy of Sciences* 1980, 77(11):6715.
15. Hutchison CA, Newbold JE, Potter SS, Edgell MH: Maternal inheritance of mammalian mitochondrial DNA. *Nature* 1974, 251(5475):536-538.
16. Anderson S, de Bruijn MHL, Coulson AR, Eperon IC, Sanger F, Young IG: Complete sequence of bovine mitochondrial DNA conserved features of the mammalian mitochondrial genome. *Journal of Molecular Biology* 1982, 156(4):683-717.
17. Sutarno, Cummins JM, Greeff J, Lymbery AJ: Mitochondrial DNA polymorphisms and fertility in beef cattle. *Theriogenology* 2002, 57(6):1603-1610.
18. Holyoake AJ, McHugh P, Wu M, O'Carroll S, Benny P, Sin IL, Sin FYT: High incidence of single nucleotide substitutions in the mitochondrial genome is associated with poor semen parameters in men. *International Journal of Andrology* 2001, 24(3):175-182.
19. Chan CC, Liu VW, Lau EY, Yeung WS, Ng EH, Ho PC: Mitochondrial DNA content and 4977 bp deletion in unfertilized oocytes. *Molecular human reproduction* 2005, 11(12):843-846.
20. Schutz MM, Freeman AE, Lindberg GL, Koehler CM, Beitz DC: The effect of mitochondrial DNA on milk production and health of dairy cattle. *Livestock Production Science* 1994, 37(3):283-295.
21. Wang J, Shi Y, Elzo MA, Dang S, Jia X, Lai S: Genetic diversity of ATP8 and ATP6 genes is associated with high-altitude adaptation in yak. *Mitochondrial DNA Part A* 2017:1-9.
22. Shi Y, Hu Y, Wang J, Elzo MA, Yang X, Lai S: Genetic diversities of MT-ND1 and MT-ND2 genes are associated with high-altitude adaptation in yak. *Mitochondrial DNA Part A* 2017:1-10.
23. Robin ED, Wong R: Mitochondrial DNA molecules and virtual number of mitochondria per cell in mammalian cells. *Journal of Cellular Physiology* 1988, 136(3):507-513.
24. Goffart S, Wiesner RJ: Regulation and Co-Ordination of Nuclear Gene Expression During Mitochondrial Biogenesis. *Experimental Physiology* 2003, 88(1):33-40.
25. Clay Montier LL, Deng JJ, Bai Y: Number matters: control of mammalian mitochondrial DNA copy number. *Journal of Genetics and Genomics* 2009, 36(3):125-131.
26. San Segundo-Val I, Sanz-Lozano CS: Introduction to the Gene Expression Analysis. In: *Molecular Genetics of Asthma*. Edited by Isidoro García M. New York, NY: Springer New York; 2016: 29-43.
27. Chamberlain AJ, Vander Jagt CJ, Hayes BJ, Khansefid M, Marett LC, Millen CA, Nguyen TTT, Goddard ME: Extensive variation between tissues in allele specific expression in an outbred mammal. *BMC Genomics* 2015, 16:993.
28. van Dam S, Vosa U, van der Graaf A, Franke L, de Magalhaes JP: Gene co-expression analysis for functional classification and gene-disease predictions. *Briefings in bioinformatics* 2018, 19(4):575-592.
29. Eisen MB, Spellman PT, Brown PO, Botstein D: Cluster analysis and display of genome-wide expression patterns. *Proceedings of the National Academy of Sciences of the United States of America* 1998, 95(25):14863-14868.
30. Michalak P: Coexpression, coregulation, and cofunctionality of neighboring genes in eukaryotic genomes. *Genomics* 2008, 91(3):243-248.
31. Hurst LD, Pal C, Lercher MJ: The evolutionary dynamics of eukaryotic gene order. *Nat Rev Genet* 2004, 5(4):299-310.

32. Nora EP, Lajoie BR, Schulz EG, Giorgetti L, Okamoto I, Servant N, Piolot T, van Berkum NL, Meisig J, Sedat J *et al*: Spatial partitioning of the regulatory landscape of the X-inactivation centre. *Nature* 2012, 485(7398):381-385.
33. Harhay GP, Smith TPL, Alexander LJ, Haudenschild CD, Keele JW, Matukumalli LK, Schroeder SG, Van Tassel CP, Gresham CR, Bridges SM *et al*: An atlas of bovine gene expression reveals novel distinctive tissue characteristics and evidence for improving genome annotation. *Genome Biology* 2010, 11(10):R102-R102.
34. Wang ZM, O'Connor TP, Heshka S, Heymsfield SB: The Reconstruction of Kleiber's Law at the Organ-Tissue Level. *J Nutr* 2001, 131.
35. Berg JM, Tymoczko JL, L S: Biochemistry, 5th edition edn. New York : Basingstoke New York: New York, NY, USA: W. H. Freeman; 2002.
36. Wang Z, Zhang J, Ying Z, Heymsfield SB: Organ-Tissue Level Model of Resting Energy Expenditure Across Mammals: New Insights into Kleiber's Law. *ISRN Zoology* 2012, 2012:9.
37. Clark EL, Bush SJ, McCulloch MEB, Farquhar IL, Young R, Lefevre L, Pridans C, Tsang HG, Wu C, Afrasiabi C *et al*: A high resolution atlas of gene expression in the domestic sheep (*Ovis aries*). *PLoS Genetics* 2017, 13(9):e1006997.
38. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S: STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* 2013, 29.
39. Okonechnikov K, Conesa A, García-Alcalde F: Qualimap 2: advanced multi-sample quality control for high-throughput sequencing data. *Bioinformatics* 2016, 32(2):292-294.
40. Liao Y, Smyth GK, Shi W: featureCounts: an efficient general purpose program for assigning sequence reads to genomic features. *Bioinformatics* 2014, 30(7):923-930.
41. Huang da W, Sherman BT, Lempicki RA: Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. *Nature protocols* 2009, 4(1):44-57.
42. Huang da W, Sherman BT, Lempicki RA: Bioinformatics enrichment tools: paths toward the comprehensive functional analysis of large gene lists. *Nucleic Acids Res* 2009, 37(1):1-13.
43. Robinson MD, McCarthy DJ, Smyth GK: edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics* 2010, 26(1):139-140.
44. Kolde R: Pheatmap: pretty heatmaps. . In., R Package Version 61 edn; 2012.
45. Hui S, Sander C, Potylitsine E, Whitaker W, Bader G, Morrison L, Lopes C: ExpressionCorrelation. In: *Makes a similarity network where nodes are genes, and edges denote highly correlated genes*. Version 1.1.0 edn; 2015.
46. Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, Amin N, Schwikowski B, Ideker T: Cytoscape: A Software Environment for Integrated Models of Biomolecular Interaction Networks. *Genome Research* 2003, 13(11):2498-2504.
47. Wang M, Hancock TP, Chamberlain AJ, Vander Jagt CJ, Pryce JE, Cocks BG, Goddard ME, Hayes BJ: Putative bovine topological association domains and CTCF binding motifs can reduce the search space for causative regulatory variants of complex traits. *BMC Genomics* 2018, 19(1):395.
48. Kanehisa M, Goto S: KEGG: kyoto encyclopedia of genes and genomes. *Nucleic acids research* 2000, 28(1):27-30.
49. Bush SJ, Hume DA, Clark EL: Unnormalised raw count estimates for the sheep gene expression atlas dataset. In. Edited by Edinburgh Uo; 2019.
50. Gallagher D, Belmonte D, Deurenberg P, Wang Z, Krasnow N, Pi-Sunyer FX, Heymsfield SB: Organ-tissue mass measurement allows modeling of REE and metabolically active tissue mass. *The American journal of physiology* 1998, 275(2):E249-258.
51. Wang Z, Ying Z, Bosy-Westphal A, Zhang J, Schautz B, Later W, Heymsfield SB, Müller MJ: Specific metabolic rates of major organs and tissues across adulthood: evaluation by mechanistic model of resting energy expenditure. *The American journal of clinical nutrition* 2010, 92(6):1369-1377.
52. Stanley WC, Chandler MP: Energy Metabolism in the Normal and Failing Heart: Potential for Therapeutic Interventions. *Heart Failure Reviews* 2002, 7(2):115-130.
53. Herbers E, Kekäläinen NJ, Hangas A, Pohjoismäki JL, Goffart S: Tissue specific differences in mitochondrial DNA maintenance and expression. *Mitochondrion* 2018.

54. Glaister M: Multiple sprint work : physiological responses, mechanisms of fatigue and the influence of aerobic fitness. *Sports medicine (Auckland, NZ)* 2005, 35(9):757-777.
55. Spriet LL: Anaerobic metabolism in human skeletal muscle during short-term, intense activity. *Canadian Journal of Physiology and Pharmacology* 1992, 70(1):157-165.
56. Hargreaves M: Skeletal muscle metabolism during exercise in humans. *Clinical and experimental pharmacology & physiology* 2000, 27(3):225-228.
57. Forbes JM: Mitochondria–Power Players in Kidney Function? *Trends in Endocrinology & Metabolism* 2016, 27(7):441-442.
58. Bhargava P, Schnellmann RG: Mitochondrial energetics in the kidney. *Nature Reviews Nephrology* 2017, 13:629.
59. Wang Z, Ying Z, Bosy-Westphal A, Zhang J, Schautz B, Later W, Heymsfield SB, Muller MJ: Specific metabolic rates of major organs and tissues across adulthood: evaluation by mechanistic model of resting energy expenditure. *Am J Clin Nutr* 2010, 92(6):1369-1377.
60. Kramer PA, Ravi S, Chacko B, Johnson MS, Darley-Usmar VM: A review of the mitochondrial and glycolytic metabolism in human platelets and leukocytes: implications for their use as bioenergetic biomarkers. *Redox biology* 2014, 2:206-210.
61. Yan X, Zhu M-J, Dodson MV, Du M: Developmental programming of fetal skeletal muscle and adipose tissue development. *J Genomics* 2013, 1:29-38.
62. Strang LB: The Lungs at birth. *Arch Dis Childh* 1965, 40 575.
63. Edfors F, Danielsson F, Hallström BM, Käll L, Lundberg E, Pontén F, Forsström B, Uhlén M: Gene-specific correlation of RNA and protein levels in human cells and tissues. *Molecular Systems Biology* 2016, 12(10):883.
64. Schwanhäusser B, Busse D, Li N, Dittmar G, Schuchhardt J, Wolf J, Chen W, Selbach M: Global quantification of mammalian gene expression control. *Nature* 2011, 473:337.
65. Miller FJ, Rosenfeldt FL, Zhang C, Linnane AW, Nagley P: Precise determination of mitochondrial DNA copy number in human skeletal and cardiac muscle by a PCR-based assay: lack of change of copy number with age. *Nucleic Acids Research* 2003, 31(11):e61-e61.
66. Lindinger A, Peterli R, Peters T, Kern B, von Flue M, Calame M, Hoch M, Eberle AN, Lindinger PW: Mitochondrial DNA content in human omental adipose tissue. *Obesity surgery* 2010, 20(1):84-92.
67. Svendsen AJ, Tan Q, Jakobsen MA, Thyagarajan B, Nygaard M, Christiansen L, Mengel-From J: White blood cell mitochondrial DNA copy number is decreased in rheumatoid arthritis and linked with risk factors. A twin study. *Journal of Autoimmunity* 2019, 96:142-146.
68. Williams RS: Mitochondrial gene expression in mammalian striated muscle. Evidence that variation in gene dosage is the major regulatory event. *Journal of Biological Chemistry* 1986, 261(26):12390-12394.
69. Aloni Y, Attardi G: Expression of the mitochondrial genome in HeLa cells. II. Evidence for complete transcription of mitochondrial DNA. *J Mol Biol* 1971, 55(2):251-267.
70. Gelfand R, Attardi G: Synthesis and turnover of mitochondrial ribonucleic acid in HeLa cells: the mature ribosomal and messenger ribonucleic acid species are metabolically unstable. *Molecular and Cellular Biology* 1981, 1(6):497-511.
71. Shokolenko IN, Alexeyev MF: Mitochondrial transcription in mammalian cells. *Frontiers in bioscience (Landmark edition)* 2017, 22:835-853.
72. Michalak P: Coexpression, coregulation, and cofunctionality of neighboring genes in eukaryotic genomes. *Genomics* 2008, 91(3):243-248.
73. Lee JM, Sonnhammer EL: Genomic gene clustering analysis of pathways in eukaryotes. *Genome Res* 2003, 13(5):875-882.
74. Mercer Tim R, Neph S, Dinger Marcel E, Crawford J, Smith Martin A, Shearwood A-Marie J, Haugen E, Bracken Cameron P, Rackham O, Stamatoyannopoulos John A *et al*: The Human Mitochondrial Transcriptome. *Cell* 2011, 146(4):645-658.
75. Soler-Oliva ME, Guerrero-Martínez JA, Bachetti V, Reyes JC: Analysis of the relationship between coexpression domains and chromatin 3D organization. *PLOS Computational Biology* 2017, 13(9):e1005708.
76. Fantz CR, Dagogo-Jack S, Ladenson JH, Gronowski AM: Thyroid Function during Pregnancy. *Clinical Chemistry* 1999, 45(12):2250.

77. Mullur R, Liu Y-Y, Brent GA: Thyroid hormone regulation of metabolism. *Physiol Rev* 2014, 94(2):355-382.

## Additional Files

Additional file 1

Table S1. Quality of library preparation.xls

Additional file 2

Table S2. Read alignment quality check.xls

Additional file 3

Table S3. List of Mitochondrial protein genes derived from Mitocarta in cattle.xls

Additional file 4:

Table S4. List of Mitochondrial protein genes derived from Mitocarta in Sheep.xls

Additional file 5:

Table S5. Number of differentially expressed (DE) genes by gene categories averaged for two foetuses in the Main Cows

Additional file 6:

Figure S1. Heatmap of expression of nuclear genome encoded mitochondrial protein (NuMP) in tissues of foetuses 6819F and 2181F in Main Cows

Additional file 7:

Figure S2. Heatmap of mitochondrial genome encoded mitochondrial protein (MtMP) genes in tissues of foetuses 6819F and 2181F in Main Cows.

Additional file 8:

Table S6. List of non-mitochondrial protein (Non-MP) genes clustering with the mitochondrial protein genes in cluster I (NuMP-MtMP cluster) in Main Cows

Additional file 9:

Table S7. KEGG pathway enrichment of the non-mitochondrial protein (Non-MP) genes in NuMP-MtMP cluster in Main Cows

Additional file 10:

Figure S3. The proportion of differentially expressed gene in each gene category in 18 tissues in a Validation Cow (All=All genes encoded by nuclear and mitochondrial genome, Nu=Mitochondrial protein genes encoded by nuclear genome (NuMP), Mt=Mitochondrial protein genes encoded by mitochondrial genome (MtMP))

Additional file 11:

Figure S4. Heatmap of expression of nuclear genome encoded mitochondrial (NuMP) gene in a Validation Cow

Additional file 12:

Figure S5. Heatmap of expression of mitochondrial genome encoded mitochondrial protein (MtMP) genes in the Validation Cow.

Additional file 13:

Figure S6. The proportion of differentially expressed gene in each gene category and direction of gene regulation in 15 tissues in Validation Sheep (All=All genes encoded by nuclear and mitochondrial genome, Nu=Mitochondrial protein genes encoded by nuclear genome (NuMP), Mt=Mitochondrial protein genes encoded by mitochondrial genome (MtMP))

Additional file 14:

Figure S7. Heatmap of nuclear genome encoded mitochondrial protein genes (NuMP) in Validation Sheep (three adults Texel x Blackface female sheep AF1, AF2, and AF3)

Additional file 15:

Figure S8. Heatmap of mitochondrial genome encoded mitochondrial protein genes (MtMP) genes in Validation Sheep (three adults Texel x Blackface females AF1, AF2, and AF3)

Additional file 16:

Table S8. Number of differentially expressed gene (DEG)s and their direction in tissues by gene categories in a Validation Cow

Additional file 17:

Table S9. Number of differentially expressed gene (DEG)s and their direction in tissues by gene categories in Validation Sheep

Additional file 18:

Figure S9. Scatter plot of log fold changes of Main Cows against the log-fold changes of Validation Sheep for mitochondrial protein gene expression in tissues

Additional file 19:

Figure S10. Scatter plot of log fold changes of Validation Cow against the log-fold changes of Validation Sheep for mitochondrial protein gene expression in thyroid

Additional file 20:

Figure S11. Gene co-expression network constructed based similarity matrix computed using Person Correlation Co-efficient of gene expression at  $r > |0.95|$  across tissues of the Validation Cow.


Additional file 21:

Figure S12. Gene co-expression network constructed based similarity matrix computed using Person correlation coefficient of gene expression at  $r > |0.95|$  across tissues of Validation Sheep (three Texel x blackface adult female sheep AF1, AF2 and AF3).

Additional file 21:


Figure S12. Gene co-expression network constructed based similarity matrix computed using Person correlation coefficient of gene expression at  $r > |0.95|$  across tissues of Validation Sheep (three Texel x blackface adult female sheep AF1, AF2 and AF3).

## Figures




**Figure 1**

Percentage of differentially expressed genes by gene categories for 29 tissues in the Main Cows dataset. m.= muscle, LN=lymph node; Gene category: All=All protein coding genes from nuclear and mitochondrial genomes, Nu=Mitochondrial protein coding genes from the nuclear genome (NuMP), Mt=Mitochondrial protein coding genes from the mitochondrial genome (MtMP)




**Figure 2**

Heatmap of nuclear genome encoded mitochondrial protein (NuMP) gene expression the Main Cows dataset. 6819 and 2181 represents Cow No. 6819 and Cow No. 2181 respectively.




**Figure 3**

Heatmap of mitochondrial genome encoded mitochondrial protein (MtMP) gene expression in the Main Cows dataset. Based on the log2 counts per million of MtMP genes across 29 tissues from cows 6819 and 2181.




**Figure 4**

Gene co-expression network clusters across tissues in the Main Cows based on similarity matrix computed using Pearson correlations  $> |0.95|$ .



**Figure 5**

Scatterplot of log2 fold change values of differentially expressed mitochondrial protein genes from nuclear genome between the Main Cows against the Validation Cow.



**Figure 6**

Venn diagram showing the number of genes in common among the NuMP-MtMP co-expression clusters. a. Mitochondrial protein genes encoded by the nuclear genome (NuMP genes), b. Mitochondrial protein genes encoded by the mitochondrial genome (MtMP genes) and c. non-mitochondrial protein genes from nuclear genomes in common between Main Cows and Validation Cow

## Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Additionalfile4.xlsx](#)
- [Additionalfile1.xlsx](#)
- [Additionalfile12.tif](#)
- [Additionalfile2.xlsx](#)
- [ARRIVEGuidelinesChecklistfilled.pdf](#)
- [Additionalfile8.xlsx](#)
- [Additionalfile3.xlsx](#)
- [Additionalfile21.png](#)
- [Additionalfile19.tiff](#)
- [Additionalfile11.tif](#)
- [Additionalfile6.tif](#)
- [Additionalfile20.tif](#)
- [Additionalfile18.tiff](#)
- [Additionalfile10.tif](#)
- [Additionalfile7.tif](#)
- [Additionalfile17.xlsx](#)
- [Additionalfile16.xlsx](#)
- [Additionalfile14.tif](#)
- [Additionalfile5.xlsx](#)

- [Additionalfile9.xlsx](#)
- [Additionalfile15.tif](#)
- [Additionalfile13.tif](#)

Table 5  
Specific metabolic rates of organs and tissues across species (kcal/kg/day)

Species	Heart	Kidney	Brain	Liver	Skeletal muscle*	Adipose	Reference
Cattle	429	412	185	130	-	-	[36]
Sheep	588	496	255	200	-	-	[36]
Human	440	440	240	200	10–15	4.5	[4, 51]
*Rate for resting muscle							