Prevalence of hypertension and its risk factor among textile workers in low-and-middle-income countries: A systematic review protocol

Naureen Ali (✉ naureenalimeghani@gmail.com)

Aga Khan University Hospital

Anam Feroz

Aga Khan University Medical College Pakistan

Protocol

Keywords: Blood pressure, Hypertension, Risk-factors, Prevalence, Adult workers, Low and Middle Income Countries, Systematic Review, Textile workers

Posted Date: February 24th, 2020

DOI: <https://doi.org/10.21203/rs.2.24338/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background

Textile workers are exposed to various hazards in the textile industry that might result in different ailments including hypertension (HTN). However, few attempts have been made to systematically review the prevalence of hypertension and its risk factor among textile workers in low-and-middle-income countries (LMICs). This review aims to identify the factors and burden of hypertension; considering that modifiable factors can be prevented in this high risk population.

Methods

We will carry out the systematic literature search on the major electronic databases including PubMed, CINAHL Plus, Science Direct, and Cochrane between January 2000 and December 2019, to identify the prevalence and incidence along with risk factors of hypertension in adults' textile workers in LMIC. Two researchers will independently search the databases. The search strategy will be piloted to ensure sufficient specificity and sensitivity. We will limit the search findings to human studies conducted in the adult workers.

Discussion

This review will highlight the proportion of hypertension along with its risk factor among textile workers in LMICs. Given that, some of the factors can be prevented in this high-risk population, we will call on health, experts to prioritize policies and commission and conduct programs to support the improvement in their health. The findings of this review will be made publicly available and the results will be disseminated via presentations and peer-reviewed publications

Conclusion

The review provides detailed information about the prevalence and risk factors of hypertension that will help to progress the health of textile workers in LMICs.

Background

Globally, hypertension (HTN) is identified as one of the leading causes of mortality and disability(1). It has led to 7.5 million deaths and accounted for 57 million disability adjusted life years (DALYS) or 3.7% of total DALYS(2). Hypertension is characterized as constant rise of systolic blood pressure and/or diastolic blood pressure of more than or equivalent to 140 mmHg and/or 90 mmHg respectively(3). Currently, around 1.13 billion individuals are suffering from hypertension globally and two-third of them resides in low- and middle-income countries (LMICs)(4). The number of hypertensive people is projected to grow by 70 million individual from 2000 to 2025 in high income countries (HIC) while it is estimated to grow by >500 million in LMICs over the same period(5). Moreover, worldwide high blood pressure is one of the major risk factors for cardiovascular diseases (CVD) that refers to diseases of the heart and blood

vessels(6, 7),stroke, coronary artery disease and peripheral vascular disease(7). The burden of CVD across the world is very high and in LMICs, 47% of mortality secondary to CVDs and 44% of CVDs burden are attributable to HTN/high blood pressure (BP) (7, 8). Generally, it is asymptomatic and the primary presentation can be seen with a devastating event like stroke or heart disease(9). It is estimated that 45% of deaths with heart disease, and 51% of deaths with stroke are related with hypertension (10). This is the reason that it is referred to as the “silent killer” (9).

Worldwide, the textile industry has more than 60 million employees(11) that are exposed to various hazards in textile industry that might results in different ailments(12, 13). Studies have revealed that oxidative stress is found in high concentration in cotton textile workers (14)leading to the development of essential hypertension(15, 16). A study conducted in Kannur district in India found out that the prevalence of hypertension was 22.3% among textile workers which was equivalent with the prevalence of general population(17). Furthermore, increasing age, alcohol consumption, family history of hypertension, BMI >25 kg/m² and high waist-hip ratio were identified as a strong factor of hypertension among cotton workers(17). Similarly, a study conducted in Dhaka, Bangladesh notified that both general and central obesity were found to be significantly associated with hypertension(18). Additionally, factors such as work related stress, noise pollution, use of earplug, improper diet, tobacco smoking, long hours and night shifts contributes towards the development of hypertension among the cotton textile workers (9, 19-22). However, the association between dust and increased blood pressure may not be strong, as confounding and bias impact could not be excluded(23). These factors have negative impact on worker’s cardiovascular health that will in turn effect their job performance(9).

Chronic diseases make workers less productive, more prone to injury that leads to high suffering and adds huge financial and service challenges on health care systems (18, 24-26). Furthermore, it is witnessed that collective monetary losses related with the hypertension, & DM, in low and middle-income countries are estimated to be over \$7 trillion during the years 2011–2025 that move millions of individuals underneath the poverty line(27). Besides, cardiovascular disease and its risk conditions like hypertension are chronic conditions that also adds a substantial burden on the individual and on the company as it requires investment of time and resources(9). Therefore, early detection of hypertension by identifying its risk factor is essential among high risk population particularly in LMIC. Therefore, this systematic review aims to highlight potential leading factors of hypertension among textile workers. Additionally, the review will also notify the magnitude of this disease in this occupational group.

Methods

The protocol for the systematic review is registered in the International Prospective Register for Systematic Reviews (PROSPERO) [167175]. The protocol has been designed and reported based on the guidelines Preferred Reporting Items for Systematic Reviews and Meta-analyses Protocols (PRISMA-P) checklist (28). (Figure. 1).

Exclusion and inclusion criteria:

All observational studies (cross-sectional, case-control or follow-up studies) published between 1 January, 2000 to 31 December, 2019 which reported a prevalence or incidence along with risk factors of hypertension in adults textile workers in LMICs will be included. Original articles, conference proceedings or abstracts reporting the outcome of interest will also be eligible for inclusion.

Articles will be excluded if the adult workers are from other than cotton textile industry. Non-English articles will not be included as the authors are not proficient in other languages. Moreover, commentaries, editorials, symposium proceedings, and systematic reviews will also be excluded.

The inclusion and exclusion criterion is shown in Table 1.

Information sources and search strategy

We will carry out the systematic literature search on the major electronic databases including PubMed, CINAHL Plus, Science Direct, and Cochrane because they are usually considered as good databases for systematic reviews. Furthermore, gray literature (non-published, non-reviewed papers internal, or online repositories) will also be searched. In addition, the bibliography of the included records will be assessed to identify relevant studies. Besides, cross-referencing of identified systematic review will also be performed to include pertinent studies. Two researchers will independently search the data bases (NA, AF).

The search term will be group into following categories of interest: population (textile, workers, factory workers, and mill workers), exposure (risk-factors, factors, predictors, prevalence, prevalence and factors) outcome (hypertension, blood pressure, systolic blood pressure, diastolic blood pressure, hypertension, high blood pressure and HTN), and settings (LMICs). Additionally, indexed keywords in the Medical Subject Headings (MeSH) will be used in order to ensure uniform search terms. The search strategy will be piloted to ensure sufficient specificity and sensitivity. The preliminary search strategy is illustrated in Table 2. We will limit the search findings to human studies conducted in the adult workers.

Study selection

Citation management system (Endnote software) will be used to manage the search results from the different databases(29). Firstly, the titles will be screened based on eligibility criteria using Endnote software. Secondly, the abstract will be screened within the short-listed studies guided by the inclusion and exclusion criteria. Lastly, the full text of potentially analyzed studies will be retrieved. For the reliability of selected studies among the two reviewers, a pre-defined screening tool will be prepared and pilot testing will be done as per the inclusion criteria. Two reviewer will depict outcome measures subsequently after reviewing the studies to confirm the relevance of the included articles. All studies will be given as a unique study ID to identify the data for this review. Each reviewer will give strong justification for excluding studies. Any discrepancies between the both reviewers will be resolved by mutual consensus. If it still persists, the third reviewer will be referred to conclude the final decision about

including the study. The Preferred Reporting Items for Systematic Reviews and Meta-Analysis (PRISMA) flow diagram will be used to report the study selection process as given in figure 1.

Data collection process

Data will be extracted onto a customized sheet in Excel that will be pilot tested prior to initiating the data extraction process. This sheet will be completed by two independent reviewers (NA, AF) for the eligible studies. Data extraction tables filled by two reviewers will be compared to confirm that all main findings are included. During the data extraction process a third assessor will be involved if any discordant data is witnessed. The primary data extraction table is shown in Table 3. Besides, existing studies on this research domain have been revised to assess items in the extraction form. The form included the study's title, primary author, year of publication, study setting, country of study, objective, study design, study population, reviewer name, BP measurement method, prevalence of hypertension, risk-factors of hypertension co-morbidities, study limitations, included/excluded, reason for exclusion, and quality appraisal of included studies.

Evaluation of study quality

The quality of the included studies will be evaluated by standardized quality assessment tool which will be conducted by two authors (NA, AF). Mixed Methods Appraisal Tool (MMAT) tool will be used to evaluate the methodological quality of all non-randomized studies(30). This tool is used to assess several aspects like selection of study participant, study tool, exposure period, missing data, measurement in outcomes, and selection of the reported result. Two reviewer (NA, AF) will rate each study as critical, serious, moderate, or low risk of bias via judgment of the gathered information. If there are limited information then the risk of bias will be categorized as "no information" or the reviewer will contact corresponding study authors for complete information.

Synthesis of included studies

The findings of the studies will be synthesized narratively. Firstly, we will conduct a descriptive analysis of all the selected studies to note down their main characteristics that includes study title, study authors, publication year, study aim, study methods, sampling strategy, characteristics of study participants and study outcomes. After that narrative synthesis will be conducted in which included studies will be grouped under outcomes. Both reviewers (NA, AF) will read each included study several times to extract data and group-related results. Finally, the reviewers will highlight potential risk factors that are associated with blood pressure among textile workers. For the quantitative studies, we will conduct a sub-group analysis under different categories including gender and textile department. Measures of association i.e. relative risks, odds ratio and prevalence ratios will be reported. Furthermore, we will also provide a narrative of confounders or effect modifiers being adjusted in different quantitative studies.

Discussion

This review will highlight the proportion of hypertension along with its risk factor among textile workers in LMICs. Given that, some of the modifiable factors can be prevented in this high risk population, we will call on health experts to prioritize policies and commission and conduct programs to support the improvement in their health. Besides, this review will highlight an urgent need for further research on hypertension in textile workers that fill in some of the data gaps such as the trends and determinants of hypertension. The findings of this review will be made publicly available and the results will be disseminated via presentations and peer-reviewed publications.

List Of Abbreviations

HTN: Hypertension; DALYS: disability adjusted life years; CVD: cardiovascular disease; LMICs: Lower-Middle-Income Countries; PRISMA: Preferred Reporting Items for Systematic Reviews and Meta-analyses

Declarations

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Availability of data and materials

Not applicable

Competing interests

The authors declare that they have no competing interests.

Funding

Not applicable

Authors' contributions

NA will conceive and design the study. NA will draft the manuscript and is the guarantor of the systematic review. AF, NA will develop the search strings and perform searching the studies, selection of the studies, data extraction, and synthesis. AF, NA, will extensively review the manuscript and incorporated intellectual inputs. Both authors will read, provide feedback, and approve the final version of the manuscript.

References

1. Forouzanfar MH, Liu P, Roth GA, Ng M, Biryukov S, Marczak L, et al. Global burden of hypertension and systolic blood pressure of at least 110 to 115 mm Hg, 1990-2015. *Jama*. 2017;317(2):165-82.
2. World Health Organization. Global Health Observatory (GHO) data. Raised blood pressure, Situation and trends. https://www.who.int/gho/ncd/risk_factors/blood_pressure_prevalence_text/en/.
3. Chobanian AV, Bakris GL, Black HR, Cushman WC, Green LA, Izzo Jr JL, et al. The seventh report of the joint national committee on prevention, detection, evaluation, and treatment of high blood pressure: the JNC 7 report. *Jama*. 2003;289(19):2560-71.
4. World Health Organization (WHO, 2019). Hypertension; Key-facts. <https://www.who.int/news-room/fact-sheets/detail/hypertension>.
5. Perkovic V, Huxley R, Wu Y, Prabhakaran D, MacMahon S. The burden of blood pressure-related disease: a neglected priority for global health. *Hypertension*. 2007;50(6):991-7.
6. Chioloro PBA. Prevalence and control of hypertension. *THE LANCET*. 2018; 392(10155):1305-6.
7. Roth GA, Johnson C, Abajobir A, Abd-Allah F, Abera SF, Abyu G, et al. Global, regional, and national burden of cardiovascular diseases for 10 causes, 1990 to 2015. *Journal of the American College of Cardiology*. 2017;70(1):1-25.
8. Horsa BA, Tadesse Y, Engidawork E. Assessment of hypertension control and factors associated with the control among hypertensive patients attending at Zewditu Memorial Hospital: a cross sectional study. *BMC research notes*. 2019;12(1):152.
9. Ofori SN, Obosi J. Prevalence of hypertension among office workers in a multi-national company in the Niger-Delta with the 2017 American College of Cardiology/American Heart Association Blood Pressure Guidelines. *Preventive medicine reports*. 2019;15:100899.
10. World Health Organization. A global brief on hypertension. WHO/DCO/ WHD/2013.2. Geneva: WHO 2013. Available from http://www.who.int/cardiovascular_diseases/publications/global_brief_hypertension/en/.
11. Forstater M, Implications of the global financial and economic crisis on the textile and clothing sector. International Labour Organization. International Labour Office, Geneva, 2009.
12. Ali NA, Nafees AA, Fatmi Z, Azam SI. Dose-response of Cotton Dust Exposure with Lung Function among Textile Workers: MultiTex Study in Karachi, Pakistan. *The international journal of occupational and environmental medicine*. 2018;9(3):120.
13. ZIA S, JAVEED S, JAN F. Pathophysiological Effects of Cotton Dust Pollution on Blood Pressure in Textile Workers. *Cell*.300:6617901.
14. Suryakar A, Katkam R, Dhadke V, Bhogafe R. A study of oxidative stress in cotton industry workers from Solapur city. *Biomedical Research*. 2010;21(3):260-4.
15. Rodrigo R, Prat H, Passalacqua W, Araya J, Guichard C, Bächler JP. Relationship between oxidative stress and essential hypertension. *Hypertension Research*. 2007;30(12):1159.
16. Briones AM, Touyz RM. Oxidative stress and hypertension: current concepts. *Current hypertension reports*. 2010;12(2):135-42.

17. Ismail IM, Binub K. Prevalence of hypertension and its associated factors among cotton textile workers of Kannur, Kerala. *Menoufia Medical Journal*. 2016;29(4):991.
18. Bhowmik B, Afsana F, Ahmed T, Akhter S, Choudhury HA, Rahman A, et al. Obesity and associated type 2 diabetes and hypertension in factory workers of Bangladesh. *BMC research notes*. 2015;8(1):460.
19. Rizi HAY, Dehghan H. Effects of occupational noise exposure on changes in blood pressure of workers. *ARYA Atherosclerosis*. 2013:183-6.
20. Starzyński Z, Wilczyńska U, Kubasiewicz M, Szymczak W. Incidence of arterial hypertension in the population of workers in the textile industry in Łódź. *Medycyna pracy*. 1985;36(2):131-8.
21. Tiwai R, Pathak M, Zodpey S, Babar V. Hypertension among cotton textile workers. *Indian journal of public health*. 2003;47(1):34-6.
22. Sumardiyono S, Hartono H, Probandari A, Setyono P. The Association between Risk Factors and Blood Pressure in the Textile Industry Workers. *Global Medical & Health Communication*. 2017;5(3):228-35.
23. Lin J, Wang H, Yan F, Tang K, Zhu H, Weng Z, et al. Effects of occupational exposure to noise and dust on blood pressure in Chinese industrial workers. *Clinical and Experimental Hypertension*. 2018;40(3):257-61.
24. Luckhaupt SE, Cohen MA, Li J, Calvert GM. Prevalence of obesity among US workers and associations with occupational factors. *American journal of preventive medicine*. 2014;46(3):237-48.
25. Safety A, Council C. Overweight and Obesity: Implications for Workplace Health and Safety and Workers' Compensation: Australian Safety and Compensation Council; 2008.
26. Di Lorenzo L, De Pergola G, Zocchetti C, L'Abbate N, Basso A, Pannacciulli N, et al. Effect of shift work on body mass index: results of a study performed in 319 glucose-tolerant men working in a Southern Italian industry. *International journal of obesity*. 2003;27(11):1353.
27. Bloom DE, Chisholm D, Jané-Llopis E, Prettner K, Stein A, Feigl A. From burden to "best buys": reducing the economic impact of non-communicable disease in low-and middle-income countries. Program on the Global Demography of Aging, 2011.
28. Moher D, Shamseer L, Clarke M, Ghersi D, Liberati A, Petticrew M, et al. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement. *Systematic reviews*. 2015;4(1):1.
29. Xiehang CDL. Evaluation and prospect of reference management software—a case study of EndNote and NoteExpress. *New Technology of Library and Information Service*. 2009:Z1.
30. Sterne JA, Hernán MA, Reeves BC, Savović J, Berkman ND, Viswanathan M, et al. ROBINS-I: a tool for assessing risk of bias in non-randomised studies of interventions. *bmj*. 2016;355:i4919.

Tables

Table 1: Eligibility criteria

Attribute	Inclusion Criteria	Exclusion Criteria
Population	Studies involving adult cotton textile workers of both genders	Studies involving target groups of adult workers working in another factory mill (oil mill, steel mill and coal mill)
Type of studies	Observational studies (survey, descriptive, cross-sectional cohort and case control study and brief communication studies having full-text article	RCT and quasi-experimental study, Qualitative formative studies, systematic reviews case studies, protocols, commentaries, editorials, unpublished articles, symposium proceedings, conference abstracts and irretrievable documents
Outcome	Studies demonstrating outcomes related to prevalence, incidence along with risk factor of hypertension	Studies demonstrating outcomes in workers having different comorbidity
Setting	Studies conducted in LMICs	Studies conducted elsewhere
Language	Studies available in the English Language	Studies in other languages
Time period	Studies published between the period of 1 January, 2000 to 31 December, 2019	Studies published before the period of 2000 to 2019

Table 2: Search strategy

Population	('textile* [Mesh] OR 'workers*' OR 'factory worker*' OR 'mill worker'* OR 'male worker*'OR ' cotton worker*' [Mesh]) AND
Exposure	('Risk-factors' OR 'factors' OR 'predictors' OR 'prevalence' OR 'incidence' 'risks', 'prevalence and factors') AND
Outcome	Outcome ('hypertension' OR 'blood pressure' OR 'systolic blood pressure' OR 'diastolic blood pressure' OR hypertension* OR high blood pressure OR HTN) AND
Setting	Studies conducted in LMIC
Filters	Publication date from 1 January, 2000 to 31 December, 2019; Humans

Table 3: Data Extraction Table

S.no	Characteristics	Study 1	Study 2	Study 3	Study 4	Study 5	Study 6	Study 7	Study 8
	Study Title								
	Study Author & Year								
	Setting/Country								
	Study Aim								
	Study Design								
	Study Population								
	Blood pressure Measurement Apparatus								
	Prevalence								
	Risk-Factors								
	Co-Morbidities								

Figures

Figure 1

PRISMA Flow Diagram for Database Search of Studies