Attitude and Self Care Practice on Hypertension among Antihypertensive Medication Users in a Tertiary Care Hospital Nepal

Jyoti Shrestha

Jyoti Shrestha

Sabina Sanhi

Pharmaceutical Sciences Program, School of Health and Allied Sciences, Pokhara University, Kaski Nepal

Rajendra lamichhane

4Department of Public Health, Asian College for Advance Studies, Lalitpur, Nepal

Nabin Raj Marasini

5Department of Public Health, La Grande International College, Kaski, Nepal

Nirmal Raj Marasine (✉ nirmalmarasine@gmail.com)

Pharmaceutical Sciences Program, School of Health and Allied Sciences, Pokhara University, Kaski Nepal

Research Article

Keywords: Attitude, Hypertension, Hypertensive patients, Self-care, Practice

Posted Date: January 18th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-149687/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background: Hypertension a “silent killer” is a serious global health problem, whose prevalence is increasing in Nepal. This study aimed to determine the attitude and practice of hypertension among antihypertensive medication users in a tertiary care teaching hospital in western Nepal.

Methodology: A hospital-based cross-sectional study was conducted among 136 hypertensive patients under medication, aged ≥ 30 years, and visiting medical outpatient department, medical ward, and geriatric ward, using a non-probability convenience sampling technique. Semistructured questionnaires were used for data collection through interviews. Descriptive and inferential statistics were used, and a p -value < 0.05 was considered statistically significant.

Results: The maximum number of patients (36; 26.5%) was in the age range of 50-60 years with a mean (SD) of 56.74 (12.58). Most of 77(56.6%) were male, Hindu, 95(69.9%), upper caste people (40; 29.4%), illiterate (30; 22.1%), house maker (37; 27.2%), and monthly family income of 21,000 and above (66; 48.5%). Almost half of the patients (69; 50.7%) had a positive attitude and more than half (71; 52.2%) had performed adequate self-care practice. Selected variables such as educational status and dietary pattern and attitude were significantly associated, whereas no association was found between sociodemographic variables and self-care practice. Attitude and self-care practices were found strongly associated with one another ($p=0.002$).

Conclusion: Hypertensive patients were found with better level of attitude and adequate self-care practice. Educational interventions and awareness programs on dietary aspects should be focused for improving the attitude and practice of all the patient groups.

Introduction

Hypertension, also known as “silent killer”, is a serious public health problem globally with an age-standardized prevalence of 24.1% in men and 20.1% in women, respectively [1]. By year 2025, approximately 1.56 billion global population are expected to suffer from this disease [2] and two-thirds of those living in economically developing countries. Hypertension is the leading cause of death and disability, which is responsible for 75 million disability-adjusted life-year gains (DALYs) and approximately 7.5 million deaths (12.8 of all-cause deaths) worldwide [3]. Untreated or poorly controlled high blood pressure can result in large medical, economic, and human costs, with an estimated 10% of health care spending directly related to increased blood pressure and its complications[4]. In Nepal, the prevalence of hypertension is 24.5% [5]. Despite of a high prevalence, blood pressure control rates among hypertensive patients are reported to be very low, i.e., only 30%- 34% in developing and 33%-38% in developed countries [6]. Hypertension is an important risk factor for cardiovascular disease, renal, and eye diseases [7]. According to the World Health Organization report, complications of hypertension account for 9.4 million deaths each year, which is greater than that caused by all other infectious diseases in

combination. It is also responsible for approximately 45% of deaths due to heart disease and 51% of deaths due to stroke [3, 4].

Evidence suggests that attitude and self-care practices are an important aspects to control blood pressure and related heart and renal complications. Self-care practice includes taking prescribed medications, consumption of low salt and low fat diet, quit smoking, limiting alcohol, reducing stress and weight, regular physical exercise, self-monitoring of blood pressure, and regular healthcare visits [8]. Consumption of a combination of diets such as fruits, vegetables, and low-fat dairy products lowers systolic blood pressure (SBP) by 5.5 mmHg and diastolic blood pressure (DBP) by 3 mmHg [9]. Salt intake not more than 2.4 g per day lowers systolic blood pressure by 2-8 mmHg [10]. Similarly, regular aerobic exercise for 30 min reduces SBP by 4-9 mmHg and alcohol consumption reduction by 50% reduces SBP by 2-7 mmHg and DBP by 2-5 mmHg [11, 12]. Moreover, weight reduction to normal lowers SBP by 3-7 mmHg and DBP by 3-9 mmHg [13].

People often do not implement the recommended self-care practices and results with uncontrolled blood pressure. Despite the high prevalence, studies examining self-care practices among hypertensive patients are scarce in Nepal. Information from such studies will help to modify treatment options, develop interventional programs aimed at increasing disease self-management behaviors, reducing adverse outcomes, and provide better healthcare services. To fill this gap, this study aimed to assess the attitude and self-care practices on hypertension among hypertensive medication users in a tertiary care teaching hospital in western Nepal.

Methods

Study design and population: A hospital-based cross-sectional study was conducted from October to December 2017 among 136 patients, using the nonprobability convenience sampling technique at Western Regional Hospital (WRH), Pokhara-10, Kaski, Nepal. All the new and old hypertensive patients under antihypertensive medication, aged ≥ 30 years, both sex, visiting medical outpatient department, medical ward, and geriatric ward, and those willing to participate were included in the study. Pregnant women and unresponsive participants were excluded.

Study variables: Attitude and self-care practices of hypertensive patients were our dependent variables, whereas non-modifiable factors such as social determinants (age, gender, educational status, monthly income, occupation, family history of hypertension) and metabolic disorders (diabetes, kidney diseases and chronic artery diseases) and modifiable factors such as lifestyle related factors (smoking, alcoholism, unhealthy dietary pattern, and physical exercise) and confounding variables (previous information about hypertension and availability of health facilities) were our independent variables.

Data Collection: The data on attitude and self-care practices of 136 patients were collected using a semi-structured questionnaire through an interview method. A standard proforma was designed to collect data on sociodemographic information. There were 13 questions for assessing sociodemographic characteristics, 10 statements for assessing attitude (5 positive and 5 negative), and 22 questions for

assessing self-care practice. The English form of the questionnaire was translated into Nepali and pilot testing was done on 10% of the total sample size, i.e., 14 hypertensive patients who met our inclusion and exclusion criteria. The reliability of attitude of hypertensive patients and self-care practice using Karl Pearson's correlation coefficient formula by split-half method, was 0.81 and 0.84, respectively. The patients involved in the pilot test were not included in the main study.

Data measurements: For attitude, where the total score was 50, firstly the median of the total result was calculated and score >41 was defined as positive attitude and score ≤ 41 as negative attitude. Likewise, for self-care practice with 20 a total score, the median was calculated and a score >16 was regarded as adequate practice and ≤ 16 as inadequate practice.

Statistical analysis: The data were entered in Microsoft Excel version 13 and analyzed using IBM-SPSS 20.0 (IBM Corporation, Armonk, NY, USA). The descriptive data were analyzed using mean, median, frequency, and percentage. Inferential statistics were used to assess the relationship between selected variables and attitude and self-care practice. $P < 0.05$ was considered statistically significant.

Ethics: Ethical approval for this study was obtained from the Institutional Review Committee (IRC) of Pokhara University Research Center (PURC) (Ref.No:85/073/74), and prior data collection permission was taken from the Western Regional Hospital, Pokhara, Nepal. The patients were fully informed about the nature and purpose of the study in the Nepali language, and their written consent was obtained prior to data collection. The personal details of the patients were kept confidential and anonymity was maintained.

Results

Of the 136 hypertensive patients, 77 (56.6%) were male. The principal characteristics of the study population are described in Table 1. They had a mean (SD) age of 56.74 (± 12.58) years; the majority (26.5%) were in the age range of 30-40 years. The majority (95; 69.9%) of the hypertensive patients were Hindus and very fewer (1; 0.7%) were of other religions like Shikh. Regarding ethnicity, upper caste patients were predominant (40; 29.4%) and only 6 (4.4%) patients belonged to religious minorities. Similarly, a higher proportion (95; 69.9%) of patients were married and very few (4; 2.9%) were divorced.

Most of the patients (30; 22.1%) were illiterate and only 11 (8.1%) of them had a bachelor and above level of education. Twenty-two (16.2%) of the patients had a family income of less than 7000 and 66 (48.5%) of them had an income range of 21,000 and above. Thirty-seven (27.2%) were homemakers, 20 (14.7%) were involved in agriculture, 26 (19.1%) were self-employed, 14 (10.3%) were employed in the government sector, 10 (7.4%) were employed in the private sector, 22 (16.2%) were unemployed, and 7 (5.1%) were retired.

Likewise, sixty-nine (50.7%) patients had reported a family history of hypertension, whereas 67 (49.3%) had reported not having any such history. Among the 69, 29 (42.03%) told that

their father or mother had hypertension, 16(23.18) told their son or daughter had hypertension, 7(10.15%) told their grandparents had hypertension, and 17(24.64%) complained that their husband or wife had hypertension. Among 136 patients, 56(41.2%)had no any comorbid condition, but 32(23.5%) complained of having diabetes mellitus. The majority (107; 78.7%) were nonvegetarian, whereas 29(21.3%) of them were vegetarian. Moreover, 134(98.5%) of the patients had easy access to healthcare services, and only 2(1.5%) of them complained of nonavailability of health service facilitieswithin 30 minutesof walking distance.

Table 1: Socio-demographic determinants of patients (n=136)

Characteristics	Categories	n (%)
Age	30-80	136 (100.0)
Sex	Male	77 (56.6)
Religion	Hindu	95 (69.9)
	Buddhist	28 (20.6)
	Christian	2 (1.5)
	Muslim	10 (7.4)
	Others	1 (0.7)
Ethnicity	Dalit	26 (19.1)
	Disadvantaged non-dalit tarai caste	11 (8.1)
	Religious minorities	6 (4.4)
	Relatively disadvantaged janajati	22 (16.2)
	Relatively advantaged janajati	31 (22.8)
	Upper caste people	40 (29.4)
Marital status	Unmarried	9 (6.6)
	Married	95 (69.9)
	Divorced	4 (2.9)
	Widow/Widower	28 (20.6)
Educational status	Illiterate	30 (22.1)
	Informal schooling	24 (17.6)
	Primary	25 (18.4)
	Secondary	25 (18.4)
	Higher secondary	21 (15.4)
	Bachelor and above	11 (8.1)
Monthly family income	Less than 7,000	22 (16.2)
	7,000-14,000	23 (16.9)
	14,000-21,000	25 (18.4)
	21,000 and above	66 (48.5)
Occupation	House maker	37 (27.2)

	Agriculture	20 (14.7)
	Self employed	26 (19.1)
	Government sector	14 (10.3)
	Private sector	10 (7.4)
	Unemployed	22 (16.2)
Family history of hypertension	Retired	7 (5.1)
	Present	69 (50.7)
	Absent	67 (49.3)
Family member having hypertension(n=69)	Parents	29 (42.03)
	Offspring	1 (23.18)
	Grandparents	7 (10.15)
	Partner	17 (24.64)
Presence of comorbidities	No any	56 (41.2)
	Diabetes mellitus	32 (23.5)
	Kidney diseases	10 (7.4)
	Asthma	21 (15.4)
	Other diseases	17 (12.5)
Dietary pattern	Vegetarian	29 (21.3)
	Non-vegetarian	107 (78.7)
Availability of health services at 30 minutes walking distance	Available	134 (98.5)
	Not available	2 (1.5)

Others: Shikh, Other diseases: Skin problems, psychological disorders, gastritis, reproductive problems

The majority (69; 50.7%) of the hypertensive patients were found to have a positive attitude regarding hypertension, whereas 67(49.3%) had a negative attitude. Likewise, 71 (52.2%) patients were found to perform adequate self-care practice and only 65(47.8%) were found to perform inadequate self-care practice, as depicted in Table 2.

Table 2: Attitude and self-care practice scoring of hypertensive patients (n=136)

Variables	Category	n (%)
Attitude	Negative (≤ 40)	67 (49.3)
	Positive (>40)	69 (50.7)
Practice	Inadequate (≤ 16)	65 (47.8)
	Adequate (>16)	71 (52.2)

Table 3 reveals the attitude of hypertensive patients toward each statement. The majority (86; 63.2%) of the patients had a positive attitude towards the positive statement, i.e., hypertension is a major cardiovascular diseases. Sixty-three (34.3%) patients strongly agreed that smoking and alcohol consumption increase the risk of hypertension and no respondent disagreed on that statement. Fifty (36.8%) patients strongly disagreed that a balanced diet and healthy lifestyle play an important role in the causation of hypertension, but also 11 (8.1%) patients agreed to the same statement. Nineteen (14%) patients strongly agreed that only salt intake reduction is sufficient for prevention of hypertension and 41(30.1%) showed strong disagreement in this statement. Similarly, 74 (54.4%) patients strongly agreed that regular physical exercise is essential for maintaining a healthy lifestyle and 11(8.1%) disagreed on it.

Likewise, 59 (43.4%) and 50 (36.8%) patients, respectively, showed strong disagreement and disagreement that regular blood pressure level checking is not necessary for hypertensive patients. Most of the patients (72; 52.9%) strongly agreed and 8(5.9%) strongly disagreed that regular follow-up for hypertensive patients is important. In the negative statement, hypertension usually does not cause any symptoms, only 16(11.8%) patients strongly agreed and 56 (41.2%) disagreed. Fifty-eight (42.6%) disagreed and only 9(6.6%) agreed that regular intake of medication is not necessary for hypertensive patients. In the last positive statement, hypertension may lead to various life-threatening complications if not managed in time, 107 (78.7%) patients showed strong agreement, as shown in Table 3.

Table 3: Attitude of hypertensive patients (n=136)

Statements	Response				
	Strongly agree n (%)	Agree n (%)	Undecided n (%)	Disagree n (%)	Strongly disagree n (%)
Hypertension is common cardiovascular disease in adult.	86 (63.2)	34 (25.0)	8 (5.9)	8 (5.9)	0 (0.0)
Habit of smoking and alcohol consumption increases risk for hypertension.	63 (46.3)	60 (44.1)	1 (0.7)	12 (8.8)	0 (0.0)
Balanced diet and healthy lifestyle plays an important role in causation of hypertension.	18 (13.2)	11 (8.1)	13 (9.6)	44 (32.4)	50 (36.8)
Only salt intake reduction is sufficient for prevention of hypertension.	19 (14.0)	23 (16.9)	18 (13.2)	35 (25.7)	41 (30.1)
Regular exercise is essential for maintaining healthy lifestyle.	74 (54.4)	36 (26.5)	15 (11.0)	11 (8.1)	0 (0.0)
Regular blood pressure level checking is not necessary for hypertensive patient.	6 (4.4)	7 (5.1)	14 (10.3)	50 (36.8)	59 (43.4)
Regular follow up for hypertensive patient is important.	72 (52.9)	34 (25.0)	10 (7.4)	12 (8.8)	8 (5.9)
Hypertension usually does not cause any symptoms.	16 (11.8)	9 (6.6)	14 (10.3)	56 (41.2)	41 (30.1)
Regular intake of medication is not necessary for hypertensive patient.	10 (7.4)	8 (5.9)	5 (3.7)	58 (42.6)	55 (40.4)
Hypertension may lead to various life threatening complication if not managed in time.	107 (78.7)	25 (18.4)	1 (0.7)	0 (0.0)	3 (2.2)

Among 136 patients, the majority (130; 95.6%) were found to follow a healthy dietary pattern. One hundred and six (77.9%) patients did not regularly take fried goods. One hundred and sixteen (85.3%) added extra salt to their diet and only 20 (14.7%) did not add extra salt to their diet. Most of them (97; 71.3%) did not take red or smoked meat. One hundred and thirty (95.6%) and 105 (77.2%) patients reported taking more than 5 servings of green leafy vegetables and fruits, respectively.

Similarly, only 34 (25%) patients were found to have a habit of smoking and drinking alcohol, 94 (69.1%) patients had confessed that they performed physical exercise for more than 30 minutes daily, among which 44 patients exercised six or more days in a week.

Forty-two (44.68%) patients used to exercise with the purpose of reducing hypertension, while 52 (53.32%) used to exercise with the intention of reducing weight. Most of the study patients (112; 82.4%) had reported taking hypertensive medication. About 96 (70.58%) of the patients told that they visited for follow-up as per physician advice, and 124 (91.2%) of them regularly checked their blood pressure, and 54 (43.54%) of them had the habit of checking it weekly, as illustrated in Table 4.

Table 4: Self-care practice of Hypertensive patients (n=136)

Variables	Categories	n (%)
1. Follow a healthy eating pattern	Yes	6 (4.4)
2. Take fried food regularly	Yes	30 (22.1)
3. Adding extra salt in regular diet	Yes	20 (14.7)
4. Take red meat and smoked meat	Yes	39 (28.7)
5. Take ≥ 5 green leafy vegetable servings	Yes	130 (95.6)
6. Take ≥ 5 fruits servings	Yes	105 (77.2)
7. Smoking habit	Yes	34 (25.0)
8. Alcohol consumption habit	Yes	34 (25.0)
8.1. How often (n=34)	Daily	17 (50.0)
	Weekly	4 (11.76)
	Occasionally	13 (38.24)
	As eye opener	0 (0.0)
8.2. Amount of alcohol (n=34)	Quarter(180 ml)	26 (76.48)
	Half bottle (375ml)	3 (8.82)
	Bottle (750ml)	5 (14.7)
9. Physical exercise of at least 30 minutes	Yes	94 (69.1)
10. Specific exercise	Yes	93 (68.4)
11. Days of exercise in a week (n=94)	< 3 days	11 (11.71)
	4 days	13 (13.83)
	5 days	26 (27.66)
	>6 days	44 (46.8)
12. Purpose of exercise (n=94)	Hypertension reduction	42 (44.68)
	Weight management	52 (53.32)
13. Taking any anti-hypertensive medication	Yes	112 (82.4)
13.1. Take blood pressure medicine regularly (n=112)	Yes	111 (99.1)
13.2. Habit of increasing/decreasing doses (n=112)	Yes	2 (1.79)
13.3. Take blood pressure medicine at same time every day (n=112)	Yes	106 (94.65)
13.4. Take recommended number of blood pressure medicine (n=112)	Yes	109 (97.32)
14. Visit for follow up	As physician advice	96 (70.58)
	6 monthly	20 (14.7)
	When symptoms worsen	14 (10.3)
	Rarely	6 (4.42)

15. Habit of checking blood pressure regularly	Yes	124 (91.2)
15.1. If yes, how often(n=124)	Daily	16 (12.90)
	Weekly	54 (43.54)
	Monthly	37 (29.83)
	When symptoms worsen	17 (13.73)

The attitude of hypertensive patients was high with a mean ($\pm$ SD) of 40.78 ($\pm$ 4.37) and a mean percentage of 81.56. Similarly, the self-care practice of the patients was adequate with a mean ($\pm$ SD) value of 14.66 ($\pm$ 3.48) and a mean percentage of 73.3, as shown in Table 5.

Table 5: Attitude and self-care practice of hypertensive patients

Response	Maximum score	Median	Mean $\pm$ SD	Mean percentage (%)
Attitude	50	41	40.78 $\pm$ 4.37	81.56
Self-care practice	20	16	14.66 $\pm$ 3.48	73.3

A significant association was found between selected variables such as educational status, dietary pattern, and attitude of hypertensive patients, whereas there was no significant association with other variables like age, gender, religion, ethnicity, occupation, monthly income, and family history, as illustrated in Table 6. In contrast, no significant association was found between sociodemographic variables and self-care practice, as depicted in Table 7. As a whole, a significant association was found between attitude and self-care practice, as shown in Table 8.

Table 6: Association of attitude of hypertensive patients with demographic variables (n=136)

Variables	Categories	Total scores		p-value
		Median ≤41 n (%)	Median >41 n (%)	
Age	≤ 54	27 (46.6)	31 (53.4)	0.585
	>54	40 (51.3)	38 (48.7)	
Sex	Male	37 (48.1)	40 (51.9)	0.747
	Female	30 (50.8)	29 (49.2)	
Ethnicity	Upper caste	24 (60.0)	16 (40.0)	0.094
	Others	42 (44.2)	53 (55.8)	
Educational status	Illiterate	21 (70.0)	9 (30.0)	0.010*
	Others	46 (43.4)	60 (56.6)	
Monthly family income	Less than 7,000	12 (54.5)	10 (45.5)	0.734
	7,000-14,000	9 (39.1)	14 (60.9)	
	14,000-21,000	13 (52.0)	12 (48.0)	
	21,000 and above	33 (50.0)	33 (50.0)	
Occupation	Home maker	22 (59.5)	15 (40.5)	0.146
	Others	45 (45.5)	54 (54.5)	
Family history of hypertension	Present	33 (47.8)	36 (52.2)	0.733
	Absent	34 (50.7)	33 (49.3)	
Dietary pattern	Vegetarian	22 (75.9)	7 (24.1)	0.001***
	Non vegetarian	45 (42.1)	62 (57.9)	

(***) Highly significant (*) Significant

Table 7: Association of self-care practice of hypertensive patients with demographic variables (n=136)

Variables	Categories	Total scores		p-value
		Median ≤16 n (%)	Median >16 n (%)	
Age	≤ 54	30 (51.7)	28 (48.3)	0.429
	>54	35 (44.9)	43 (55.1)	
Sex	Male	39 (50.6)	38 (49.4)	0.446
	Female	26 (44.1)	33 (55.9)	
Ethnicity	Upper caste	24 (60.0)	16 (40.0)	0.094
	Other caste	42 (44.2)	53 (55.8)	
Educational status	Illiterate	14 (46.7)	16 (53.3)	0.889
	Others	51 (48.1)	55 (51.9)	
Monthly family income	Less than 7,000	12 (54.5)	10 (45.5)	0.734
	7,000-14,000	9 (39.1)	14 (60.9)	
	14,000-21,000	13 (52.0)	12 (48.0)	
	21,000 and above	33 (50.0)	33 (50.0)	
Occupation	Home maker	18 (48.6)	19 (51.4)	0.903
	Others	47 (47.5)	52 (52.5)	
Family history of hypertension	Present	32 (46.4)	37 (53.6)	0.737
	Absent	33 (49.3)	34 (50.7)	
Dietary pattern	Vegetarian	15 (51.7)	14 (48.3)	0.633
	Non vegetarian	50 (46.7)	57 (53.3)	

Table 8: Association between attitude and self-care practice of hypertensive clients (n=136)

Attitude	Total scores of practice		p-value
	Median ≤16 n (%)	Median >16 n (%)	
Negative	41 (61.2)	26 (38.8)	0.002***
Positive	24 (34.8)	45 (65.2)	

(***) Highly significant (*) Significant

Discussion

Our study evaluated the attitude and self-care practices of hypertension among hypertensive patients taking antihypertensive medication attending the medical outpatient department, medical ward, and geriatric ward of a tertiary hospital in Nepal. The majority of our study patients were male, aged (SD) 56.74 (12.58) years, Hindu, upper caste, married, illiterate, home maker, with monthly family income of 21000 and above, and non-vegetarian. Most of them had a family history of hypertension and health services nearby their residence.

Our study revealed that the majority of hypertensive patients had a positive attitude towards hypertension and had implemented adequate self-care practices to control hypertension than a study conducted in

western Nepal and Nigeria [14, 15]. Most of our study patients have a great understanding and positive attitude towards the risk of excessive salt intake to prevent hypertension. Many studies have studied about this issue and found that a reduction in salt intake in the diet is beneficial for hypertensive patients who are under medication [16]. A healthy diet burns up more calories, preventing obesity and in the long run prevents hypertension. Poor dietary habits like salty and fatty foods, smoking and drinking alcohol increase the prevalence of hypertension. The practice of salt reduction was found adequately practiced by our study population. Similarly, many patients quitted smoking, drinking alcohol habit, consuming green vegetables and fruits, to avoid fried food and red and smoked meat in their diet. These findings suggest that patients were aware of their illness, risk factors and protective measures for maintaining normal health. Our findings are supported by a previous study from Japan, which reported that most of its patients believed that salty and fatty diet and smoking are important factors in hypertension [17]. One-fourth of the patients confessed their smoking and alcohol drinking habits, which is comparably higher than that illustrated by the STEPS survey 2013 by NHRC [18]. Drinking alcohol is more associated with smoking and both of them are strong risk factors for cardiovascular diseases [19, 20]. Evidence suggests that the consumption of alcohol beyond 20 g for men and up to 10 g for women is risky [20]. Likewise, the attitude and practice of physical exercise were found good enough in our study. Mostly physical exercise is used in conjugation with weight reduction for the management of hypertension. The majority of patients reported involving themselves in specific exercise for more than 6 days a week for at least 30 minutes. In contrast, a study from Malaysia reported poor attitude and practice of exercise in its hypertensive patients with a low score for exercising at least 3 times per week [21]. Increased physical activity along with health dietary patterns, low salt and fatty foods, and reduction in alcohol intake reduces the relative risk of hypertension [22]. The attitude and practice of patients regarding medication adherence in our study was found superior to that found in studies from central Nepal and Pakistan [23, 24]. The majority of our patients reported taking their antihypertensive medication regularly, the same number and the same recommended dose without alteration at the same time everyday. They confessed that they are more conscious of their health, check their blood pressure mostly once weekly, and visit the hospital for follow-up, as advised by their physician.

Our study showed a significant association between attitude and educational status ($p=0.010$) and dietary pattern ($p=0.001$) in hypertensive patients. This means that patients those with higher education and who prefer vegetarian diets have better attitudes than those with lower education levels and non-vegetarians. Additionally, no sociodemographic variables were found to be significantly associated with self-care practice regarding hypertension among the study population. In contrast, a study from central Nepal showed a significant association of knowledge, attitude and practice with age, level of education, and duration of hypertension [23]. Moreover, a strong relationship was found between attitude and self-care practice in hypertensive patients ($p=0.002$), which was consistent with the finding of a study from Iran [2].

There are some limitations to our study. One of the important limitations is knowledge level of patients with hypertension is not assessed. The other is the low sample size, which limits generalization. In addition, the self-reported practice by participants is prone to bias is the other important limitation.

Despite these limitations, this study provides an insight into the attitude and self-care practices among hypertensive patients in the medical outpatient department, medical ward and geriatric ward and the association between them. The findings abstained might be useful as a source of reference for conducting such research on a large scale in the future.

Conclusion

Our study found a better level of attitude and adequate self-care practice among the study patients. There was a significant association between education status and dietary pattern and attitude of hypertensive patients, whereas no association was found between sociodemographic variables and self-care practices of patients. Moreover, attitude and self-care practice of hypertensive patients were found to be strongly associated ($p=0.002$). Our study highlights the need to focus on educational interventions and awareness programs on the aspect of educational and dietary habits that could improve the attitude and practices of all the patient groups, and hence prevent further complications of the disease.

Declarations

Acknowledgments

The authors are equally thankful to the doctors, nurses, and staff of Western Regional Hospital for their help during data collection and also to the patients for their warm support and coordination throughout this study.

Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) received no financial support for the research, authorship, and/or publication of this article.

References

1. Zhou B, Bentham J, Di Cesare M, Bixby H, Danaei G, Cowan MJ, Paciorek CJ, Singh G, Hajifathalian K, Bennett JE, Taddei C. Worldwide trends in blood pressure from 1975 to 2015: a pooled analysis of 1479 population-based measurement studies with 19.1 million participants. *The Lancet*. 2017;389(10064):37-55.[Google Scholar](#)[PubMed](#)
2. Rashidi Y, Manaflouyan H, Azar FP, Nikniaz Z, Nikniaz L, Ghaffari S. Knowledge, attitude and practice of Iranian hypertensive patients regarding hypertension. *Journal of cardiovascular and thoracic research*. 2018;10(1):14[PubMed](#)

3. World Health Organization. A global brief on hypertension: silent killer, global public health crisis: World Health Day 2013. World Health Organization; 2013.[Web link](#)
4. Peberdy V. Hypertension: Putting the pressure on the silent killer. IFPMA; 2016.[Full text](#)
5. Dhimal M, Bista B, Bhattarai S, Dixit LP, Hyder MKA, Agrawal N, Rani M, Jha AK. Report of Non Communicable Disease Risk Factors: STEPS survey Nepal 2019. Kathmandu: Nepal Health Research Council. 2020.[Full text](#)
6. Kilic M, Uzunçakmak T, Ede H. The effect of knowledge about hypertension on the control of high blood pressure. International Journal of the Cardiovascular Academy. 2016 ; 2(1): 27-32.[PubMed](#)[Google Scholar](#)
7. Ademe S, Aga F, Gela D. Hypertension self-care practice and associated factors among patients in public health facilities of Dessie town, Ethiopia. BMC health services research. 2019 ;19(1):51.[Pubmed](#)
8. Aung MN, Lorga T, Srikrajang J, Promtingkran N, Kreuangchai S, Tonpanya W, Vivarakanon P, Jaiin P, Praipaksin N, Payaprom A. Assessing awareness and knowledge of hypertension in an at-risk population in the Karen ethnic rural community, Thasongyang, Thailand. International Journal of General Medicine. 2012;5:553.[Pubmed](#)
9. Appel LJ, Moore TJ, Obarzanek E, Vollmer WM, Svetkey LP, Sacks FM, Bray GA, Vogt TM, Cutler JA, Windhauser MM, Lin PH. A clinical trial of the effects of dietary patterns on blood pressure. New England journal of medicine. 1997; 336(16):1117-24.[Pubmed](#)
10. Sacks FM, Svetkey LP, Vollmer WM, Appel LJ, Bray GA, Harsha D, Obarzanek E, Conlin PR, Miller ER, Simons-Morton DG, Karanja N. Effects on blood pressure of reduced dietary sodium and the Dietary Approaches to Stop Hypertension (DASH) diet. New England journal of medicine. 2001; 344(1):3-10.[Pubmed](#)
11. Whelton SP, Chin A, Xin X, He J. Effect of aerobic exercise on blood pressure: a meta-analysis of randomized, controlled trials. Annals of internal medicine. 2002 Apr 2; 136(7):493-503.[Pubmed](#)[Google Scholar](#)
12. Roerecke M, Kaczorowski J, Tobe SW, Gmel G, Hasan OS, Rehm J. The effect of a reduction in alcohol consumption on blood pressure: a systematic review and meta-analysis. The Lancet Public Health. 2017; 2(2):e108-20.[Pubmed](#)
13. Stevens VJ, Obarzanek E, Cook NR, Lee IM, Appel LJ, Smith West D, Milas NC, Mattfeldt-Beman M, Belden L, Bragg C, Millstone M. Long-term weight loss and changes in blood pressure: results of the Trials of Hypertension Prevention, phase II. Annals of internal medicine. 2001; 134(1):1-1.[Google Scholar](#)
14. Sharma S, Bhuvan KC, Alrasheedy AA, Kaundinnyayana A, Khanal A. Impact of community pharmacy-based educational intervention on patients with hypertension in Western Nepal. The Australasian medical journal. 2014;7(7):304.[Google Scholar](#)
15. Awotidebe TO, Adedoyin RA, Rasaq WA, Adeyeye VO, Mbada CE, Akinola OT, Otwombe KN. Knowledge, attitude and Practice of Exercise for blood pressure control: A cross-sectional survey.

- Journal of exercise science and physiotherapy. 2014; 10(1):1.[Google Scholar](#)
16. Dennison CR, Peer N, Lombard CJ, Kepe L, Levitt NS, Steyn K, Hill MN. Cardiovascular risk and comorbid conditions among Black South Africans with hypertension in public and private primary care settings: the HiHi study. *Ethnicity & disease*. 2007;17(3):477-83.[Pubmed](#)
 17. Kokubo Y. Prevention of hypertension and cardiovascular diseases: a comparison of lifestyle factors in Westerners and East Asians. *Hypertension*. 2014;63(4):655-60.[Pubmed](#)
 18. Nepal Health Research Council. Non communicable disease risk factors: STEPS Survey Nepal 2013: Nepal Health Research Council. Available from: [Full text](#)
 19. Viridis A, Giannarelli C, Fritsch Neves M, Taddei S, Ghiadoni L. Cigarette smoking and hypertension. *Current pharmaceutical design*. 2010;16(23):2518-25.[Google Scholar](#)
 20. Reed MB, Wang R, Shillington AM, Clapp JD, Lange JE. The relationship between alcohol use and cigarette smoking in a sample of undergraduate college students. *Addictive behaviors*. 2007;32(3):449-64.[Google Scholar](#)
 21. Buang NF, Rahman NA, Haque M. Knowledge, attitude and practice regarding hypertension among residents in a housing area in Selangor, Malaysia. *Medicine and pharmacy reports*. 2019;92(2):145.[Google Scholar](#)
 22. Stefanov T, Vekova A, Bonova I, Tzvetkov S, Kurktschiev D, Blüher M, Temelkova-Kurktschiev T. Effects of supervised vs non-supervised combined aerobic and resistance exercise programme on cardiometabolic risk factors. *Central European journal of public health*. 2013;21(1):8-16.[Google Scholar](#)
 23. Shrestha S, Adhikari B, Poudel RS, Thapaliya K, Kharal T, Bastakoti M, Bhatta NK. Knowledge, attitude and practice on hypertension among antihypertensive medication users. *Journal of the Nepal Medical Association*. 2016;55(204):86.[Pubmed](#)
 24. Naseem S, Sarwar MH, Afzal M, Gilani SA. Knowledge attitude and practice towards hypertension among adult population in a rural area of Lahore, Pakistan. *Int J Sci Eng Res*. 2018;9(5):1674-9.[google scholar](#)