

Comparison of vocal cord view between neutral and sniffing position during orotracheal intubation using fiberoptic bronchoscope: A prospective, randomized cross over study

Sanghee Park

Department of Anesthesiology and Pain Medicine, Chonnam National University Medical School, Chonnam National University Hospital, Gwangju, Korea

Hyung Gon Lee

Department of Anesthesiology and Pain Medicine, Chonnam National University Medical School, Chonnam National University Hospital, Gwangju, Korea

Jeong Il Choi

Department of Anesthesiology and Pain Medicine, Chonnam National University Medical School, Chonnam National University Hospital, Gwangju, Korea

Seongheon Lee

Department of Anesthesiology and Pain Medicine, Chonnam National University Medical School, Chonnam National University Hospital, Gwangju, Korea

Eun-A Jang

Department of Anesthesiology and Pain Medicine, Chonnam National University Medical School, Chonnam National University Hospital, Gwangju, Korea

Hong-Beom Bae

Department of Anesthesiology and Pain Medicine, Chonnam National University Medical School, Chonnam National University Hospital, Gwangju, Korea

Jeeyun Rhee

Department of Anesthesiology and Pain Medicine, Chonnam National University Medical School, Chonnam National University Hospital, Gwangju, Korea

Hyung Chae Yang

Department of Otolaryngology-Head and Neck Surgery, Chonnam National University Medical School, Chonnam National University Hospital, Gwangju, Korea

Seongtae Jeong

Department of Anesthesiology and Pain Medicine, Chonnam National University Medical School, Chonnam National University Hospital, Gwangju, Korea

Keywords:

Posted Date: November 12th, 2018

DOI: <https://doi.org/10.21203/rs.1.13/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published on January 5th, 2019. See the published version at <https://bmcanesthesiol.biomedcentral.com/articles/10.1186/s12871-018-0671-9>.

Abstract

Background: Limitation of sight during endotracheal intubation is a major complicating factor using fiberoptic bronchoscope. Thus, maneuvers to relieve such obstructions are recommended. There have been no studies to determine whether the sniffing or neutral position is superior for this purpose. Therefore, this study was performed to examine the effects of these two positions including vocal core view. **Methods:** Fifty-four patients scheduled to receive general anesthesia by orotracheal intubation were eligible for inclusion in the study with informed consent. After confirmation of proper head positioning depending on the group, the view of the vocal cord was acquired in each position. Images were reviewed using the percentage of glottic opening (POGO) score. **Results:** A total of 106 images of vocal cords from 53 patients were obtained. The mean POGO scores were 26.65 ± 30.18 for the sniffing position and 37.29 ± 32.56 for the neutral position ($P = 0.002$). Neutral position increased POGO score in 31 patients and decreased POGO score in 13 patients compare to sniffing position ($p = 0.017$). There were no significant differences between the two head positions with regard to intubation time or degree of convenience during intubation. **Conclusions:** Neutral position improved the view of glottic opening than sniffing position during oral fiberoptic intubation. However, there was no difference in the difficulty of tube insertion between the two positions. Trial registration: Clinical Trials.gov identifier: NCT02931019, registered on October 12, 2016

Background

Limitation of sight is a major complicating factor in endotracheal intubation using fiberoptic bronchoscope (FOB), and many methods to relieve the obstruction have been reported. As FOB has become a strategic tool for endotracheal intubation, efficient positions for fiberoptic endotracheal intubation including patient head position, have been studied [1, 2].

In the sedated state, the soft palate, tongue base, and epiglottis move posteriorly, thereby obstructing the airway patency [3, 4]. A number of positions including head tilt, jaw thrust, or lingual traction have been recommended for fiberoptic intubation to relieve the obstruction at these three points [5-7]. These positions move the oropharyngeal structures anteriorly with the mouth open and an empty oropharyngeal airspace, thus enabling advancement of the FOB. However, there have been no studies to determine whether the sniffing or neutral position is superior for this purpose. Although the sniffing position has been recommended for laryngoscopic endotracheal intubation, its clinical benefit is controversial, and it is unclear whether this position is suitable for fiberoptic intubation [8-10].

In intubation using FOB, a previous endoscopic study indicated that the most common point where endotracheal tube advancement is blocked is the right arytenoid process [11]. Furthermore, the head position with neck flexion using a 7 cm high pillow aggravates the epiglottic lift and can induce the bumps between the endotracheal tube and the arytenoid process [12]. From these consequences, we deduced that the neck flexion obstructing the advancement of endotracheal tube could limit the view of

FOB during orotracheal intubation. Therefore, we hypothesized the neutral position give better vocal cord view and ease of intubation than the sniffing position during fiberoptic orotracheal intubation.

Methods

This prospective randomized crossover trial was approved by the Institutional Review Board of Chonnam National University Hwasun Hospital (CNUHH-2016-120) and was registered in the ClinicalTrials.gov public registry (NCT02931019). The study was conducted at the Department of Anesthesiology of Chonnam National University Hwasun Hospital, Republic of Korea, between 13 October 2016 and 2 December 2016.

Patient selection and enrolment

Patients aged 19 to 70 years, with an American Society of Anesthesiologists classification of I–II, who were scheduled to receive general anesthesia for surgery of the thyroid, breast, stomach, colon, and uterus with orotracheal intubation, were eligible for the study. To minimize the effects of airway variation, a preoperative airway evaluation was conducted the day before surgery using the total airway score (TAS) including the Mallampati classification, thyromental distance, head and neck movement, body mass index, buck teeth, inter-incisor gap, and upper lip bite test [13]. Patients with a total airway score > 6 points or determined to be class III in the upper lip bite test were defined as being difficult to intubate and were thus excluded. Patients with a history of difficult intubation, cervical spine defect, previous head and neck surgery, or risk of pulmonary aspiration were also excluded. After the examination of 65 patients, 54 qualified for study participation and provided written informed consent.

Study design and data collection

The patients were divided into two groups according to the order of head position: S-N group, the sniffing (S) position followed by the neutral (N) position; and the N-S group, the neutral position followed by the sniffing position. The sequence of positions was randomized by a computer-generated list of random numbers. For the neutral position, patients were placed on the operating table in the supine position without a pillow, and for the sniffing position they were kept in the supine position with a 7 cm high pillow placed under the occiput. Upon arrival in the operating room, all patients were monitored with an electrocardiogram, pulse oximeter, non-invasive blood pressure unit (Aisys; GE Healthcare, Waukesha, WI, USA) and bispectral index score (BIS; Medtronic-Covidien, Dublin, Ireland). Patients were given 0.1 mg glycopyrrolate intravenously to suppress salivation, and anesthesia was induced with 2 mg kg⁻¹ propofol followed first by remifentanyl infusion of up to 3 ng mL⁻¹ using a target-controlled infusion device (Orchestra Base Primea; Fresenius Kabi, Brezins, France), and then by 0.8 mg kg⁻¹ rocuronium. Manual mask ventilation was provided until the patient's oxygen saturation reached 100% as determined by pulse oximetry. Neuromuscular monitoring was applied with train-of-four (TOF) monitoring (Aisys; GE Healthcare). At a TOF count of 0, an Ovassapian airway (Ovassapian Fiber Optic Intubating Airway; Teleflex, Wayne, PA, USA) was inserted to guide the scope insertion through the midline of the tongue base [14, 15]. Then a specific trained anesthesia nurse conducted the jaw thrust with mouth opening,

grasping patient's both mandible angles and protruding the lower teeth until the FOB took a picture of tracheal inlet. All patients were intubated by experienced anesthesiologists using a FOB (MAF-TM Portable Bronchoscope; Olympus, Tokyo, Japan). Under the guidance of the Ovassapian airway, the FOB was inserted until just passing the tongue base and adjusted to get the best view of vocal cord with the epiglottis was acquired. At this point, the glottic opening formed by the epiglottis, vocal cord, and arytenoid cartilage was imaged (vocal cord view). Second images were taken after changing the position according to the patient group, and the FOB was passed into the trachea until it was located just above the carina. With the FOB position maintained, the Ovassapian airway was removed and the endotracheal tube (ETT) advanced smoothly into the trachea. If resistance was detected during passage into the airway, the ETT was withdrawn, rotated 90° anticlockwise, and re-inserted. The time from acquisition of the second image until completion of endotracheal intubation and the number of re-insertion trials were recorded. After proper placement of the ETT above the carina was confirmed by fiberoptic bronchoscopy view, the FOB was removed, and the patient was administered routine anesthetic care. Hypoxia, defined as oxygen saturation (SpO₂) < 93%, and the lowest SpO₂ value during the study were documented. The procedure was terminated if any adverse events occurred. The epiglottic view of each position was assessed by the percentage of glottic opening (POGO) score to evaluate the practitioner's first vision of glottic opening during fiberoptic intubation [16].

Sample size justification and statistical analyses

The POGO score was used to compare the visual range of tracheal inlet during fiberoptic intubation. The effect size to identify a meaningful difference in the respective POGO scores was obtained in a pilot study of 12 patients (effect size = 0.3644069). Thus, 48 patients yielded a power of 80% and a type I error of 0.05. Assuming a dropout rate, 54 patients (27 per group) were needed for the study. The patients were allocated to the S-N or N-S group using a randomization table. POGO scores were assigned by three other anesthesiologists who were blinded to the head position group. Numerical data, including the POGO score of vocal cord images, were analyzed using the paired *t*-test because it follows the normal distribution by the Kolmogorov-Smirnov test. Changes in the POGO score between positions were analyzed using McNemar test. Time to completion of intubation and the difficulty of tube insertion were analyzed using the Mann-Whitney U test. The relationship between difficulty of tube insertion and time to intubation was assessed by Pearson's correlation coefficient. Data are expressed as mean ± standard deviation or as numerical values. In all of the analyses, *P* < 0.05 was taken to indicate statistical significance.

Results

Sixty-five patients were initially enrolled in this study, and 11 were excluded. Detailed information on patients including those who were excluded is provided in Figure 1. The characteristics of these patients are listed in Table 1. The lowest SpO₂ reading by pulse oximetry was 94%, recorded in one patient in the N-S group. No adverse events were observed in either group, nor were there significant differences in the hemodynamic parameters between the two groups during the study period.

As this study had a crossover design and images of each head position from 53 patients were taken, 106 images (53 images for sniffing position and 53 images for neutral position) of the vocal cord view were obtained. The mean POGO score was 26.9 ± 30.4 in the sniffing position and 38.0 ± 32.5 in the neutral position ($p = 0.002$) (Fig. 2). The POGO score was significantly higher for the neutral position. Neutral position increased POGO score in 31 patients and decreased POGO score in 13 patients compare to sniffing position ($p = 0.017$) (Fig. 2). There were no significant differences in blockade during endotracheal tube insertion based on the number of repositioning events or handling of the FOB (Table 2) [17] between the two groups ($P = 0.355$). The times from obtaining images to completion of intubation were 23.3 ± 6.5 s in the sniffing position and 23.4 ± 11.5 s in the neutral position ($P = 0.626$). POGO score and time to completion of intubation showed a correlation of -11.8% ($P = 0.403$).

Discussion

In orotracheal intubation, manipulation of the patient's head and neck position contributes to the success of the procedure. This also applies to endotracheal intubation using FOB, and various positions for this purpose have been studied [5, 6, 18-20]. The advantage of the sniffing position over the neutral position in laryngoscopic endotracheal intubation is controversial [8-10], and it is unclear whether this posture affects intubation using FOB. Therefore, we examined whether there were differences in the effects of these two patient head positions on fiberoptic orotracheal intubation.

The sniffing position, achieved using a pillow, was recommended by Magill in 1936 as being better than the neutral position during endotracheal intubation using laryngoscopy [21]. The average head height needed to obtain an optimal laryngeal view was 55 mm (range: 31–71 mm) [22]. The anatomical basis of this recommendation is the three axes (oral, pharyngeal and laryngeal) alignment theory. Bannister and Macbeth suggested that the alignment of these three axes provides a better view of the vocal cords for endotracheal intubation using a laryngoscope [23].

However, the oral axes or line of view can be ignored during endotracheal intubation using FOB, because the view inside the oral cavity can be secured and the fiberoptic endoscope passed through the patient's mouth. Thus, only the angle between the pharyngeal and laryngeal axes is important during intubation using FOB. According to a previous MRI-based study, this angle is wider in the sniffing position as the pharyngeal axes lean posteriorly with atlanto-occipital extension [24], and poorer visualization of the vocal cord during fiberoptic endotracheal intubation is assumed.

Indeed, in the present study, the POGO score indicating the non-obstructing area of glottic opening was significantly higher in the neutral position than in the sniffing position. This difference could be attributed to the anterior aspect of the vocal cord, which is covered by the epiglottis (Fig. 3). In the sniffing position, because the epiglottis is closer to the posterior pharyngeal wall [25] due to the stretched and collapsed retropharyngeal muscles at the neck flexion point, the area of vocal cord obscured by the epiglottis is wider than in the neutral position. In sedated patients, this tendency is stronger because of the reduced muscle tone of the retropharyngeal structure and soft palate [7, 26].

In the previous studies about fiberoptic intubation, there was no consistent evaluating tool for assessment of glottis view [7, 12]. Therefore, to assess the visualization of vocal cord we used the POGO score which is invented to estimate the glottic inlet in conventional laryngoscopic intubation. For applying the POGO score to the fiberoptic view which has wide angle view, we paid attention to maintain the consistency along the pictures using ovassapian airway. Also, to compensate the inconsistency of the location taking pictures between two head positions, we selected the images viewing the widest glottic opening for scoring the POGO.

In the present study, despite the apparent difference in vocal cord view between the two positions, there were no differences in the ease of intubation and the time for the endotracheal tube insertion. These observations suggested that two different positions do not affect time or ease of orotracheal intubation using FOB. Also, while visualization of the vocal cord affects the difficulty of laryngoscopic endotracheal intubation [27], the effects of vocal cord view on duration or ease of fiberoptic intubation are unclear. The time for tube insertion and the difficulty of tube insertion showed no relationship with better vocal cord view. These observations suggested that visualization of the vocal cord does not affect time or ease of orotracheal intubation using FOB.

The present study has several limitations. First, the intubation time was not measured during whole procedure but only measured from acquiring images to completion of tube insertion. There could be the difference in the time getting the best view of glottis opening between two head position. Measurement of whole intubation time may reveal the relationship between head position and intubation time. Second, we included patients with low TAS score for reducing the airway variation of subjects, excluding patients expected difficult intubation. This could limit the application of this study's result to various clinical situations. To validate the superiority of different head positions during fiberoptic intubation, further studies are required with larger patient populations including various airway difficulty including measurement of the entire procedure time.

Conclusions

Neutral position improved the view of glottic opening than sniffing position during oral fiberoptic intubation. However, there was no difference in the difficulty of tube insertion between the two positions.

The English in this document has been checked by at least two professional editors, both native speakers of English. For a certificate, please see:

<http://www.textcheck.com/certificate/7JKRf9>

Abbreviations

BIS: Bispectral index score; ETT: Endotracheal tube; FOB: Fiberoptic bronchoscope; POGO: Percentage of glottic opening; SpO₂: Oxygen saturation; TAS: total airway score TOF: Train-of-four;

Declarations

Ethics approval and consent to participate

This study was approved by the Institutional Review Board of Chonnam National University Hwasoon Hospital (CNUHH-2016-120), and written informed consent was obtained from all patients.

Consent for publication: All authors have given final approval of the version to be published.

Availability of data and materials: The datasets used and analysed during the current study are available from the corresponding author on reasonable request.

Competing interests: The authors declare that they have no competing interest

Funding: This study was supported by a grant (HCRI17920-1) Chonnam National University Hwasun Hospital Institute for Biomedical Science..

Author's contributions: Park and Jeong designed and conducted the study, collected the data, analyzed the data, and wrote the manuscript; HG Lee, Choi, and Lee helped design the study and collect the data; Jang, Bae, and Rhee conducted the study; Dr. Yang helped design the study and data analysis.

Acknowledgements: None.

Authors information: Dr. Park, HG Lee, Choi, S Lee, Jang, Bae, Rhee and Jeong are at Department of Anesthesiology and Pain Medicine, Dr. Yang is at Department of Otolaryngology-Head and Neck Surgery, Chonnam National University Medical School and Hospital, 42 Jebong-ro, Dong-gu, Gwangju 61469, South Korea.

References

1. Murphy P. A fibre-optic endoscope used for nasal intubation. *Anaesthesia* 1967; 22: 489-91.
2. Apfelbaum JL, Hagberg CA, Caplan RA, Blitt CD, Connis RT, Nickinovich DG, et al. American Society of Anesthesiologists Task Force on Management of the Difficult Airway. Practice guidelines for management of the difficult airway: an updated report by the American Society of Anesthesiologists Task Force on Management of the Difficult Airway. *Anesthesiology* 2013; 118: 251-70.
3. Nandi P, Charlesworth C, Taylor S, Nunn J, Dore C. Effect of general anaesthesia on the pharynx. *Br J Anaesth* 1991; 66: 157-62.
4. Sutthiprapaporn P, Tanimoto K, Ohtsuka M, Nagasaki T, Iida Y, Katsumata A. Positional changes of oropharyngeal structures due to gravity in the upright and supine positions. *Dentomaxillofac Radiol* 2008; 37: 130-35.

5. Iqbal R, Gardner-Thorpe C, Thompson J, Popat MT, Yentis SM, Pandit JJ. A comparison of an anterior jaw lift manoeuvre with the Berman airway for assisting fibreoptic orotracheal intubation. *Anaesthesia* 2006; 61: 1048-52.
6. Durga VK, Millns JP, Smith JE. Manoeuvres used to clear the airway during fibreoptic intubation. *Br J Anaesth* 2001; 87: 207-11.
7. Uzun L, Ugur MB, Altunkaya H, Ozer Y, Ozkocak I, Demirel CB. Effectiveness of the jaw-thrust maneuver in opening the airway: a flexible fiberoptic endoscopic study. *ORL J Otorhinolaryngol Relat Spec* 2005; 67: 39-44.
8. Adnet F, Baillard C, Borron SW, Denantes C, Lefebvre L, Galinski M, et al. Randomized Study Comparing the "Sniffing Position" with Simple Head Extension for Laryngoscopic View in Elective Surgery Patients. *Anesthesiology* 2001; 95: 836-41.
9. Akihisa Y, Hoshijima H, Maruyama K, Koyama Y, Andoh T. Effects of sniffing position for tracheal intubation: a meta-analysis of randomized controlled trials. *Am J Emerg Med* 2015; 33: 1606-11.
10. Isono S. Common Practice and Concepts in Anesthesia: Time for Reassessment Is the Sniffing Position a "Gold Standard" for Laryngoscopy? *Anesthesiology* 2001; 95: 825-27.
11. Johnson DM, From AM, Smith RB, From RP, Maktabi MA. Endoscopic study of mechanisms of failure of endotracheal tube advancement into the trachea during awake fiberoptic orotracheal intubation. *Anesthesiology* 2005; 102: 910-4.
12. Cheng KI, Yun MK, Chang MC, Lee KW, Huang SC, Tang CS, et al. Fiberoptic bronchoscopic view change of laryngopharyngeal tissues by different airway supporting techniques: comparison of patients with and without open mouth limitation. *J Clin Anesth* 2008; 20: 573-9.
13. Seo SH, Lee JG, Yu SB, Kim DS, Ryu SJ, Kim KH. Predictors of difficult intubation defined by the intubation difficulty scale (IDS): predictive value of 7 airway assessment factors. *Korean J Anesthesiol* 2012; 63: 491-7.
14. Aoyama K, Seto A, Takenaka I. Simple Modification of the Ovassapian Fiberoptic Intubating Airway. *Anesthesiology* 1999; 91: 897-97.
15. Ravindran RS. Another Advantage of Marking Ovassapian Fiber-optic Intubating Airway. *Anesthesiology* 2000; 92: 1843-43.
16. Ochroch EA, Hollander JE, Kush S, Shofer FS, Levitan RM. Assessment of laryngeal view: percentage of glottic opening score vs Cormack and Lehane grading. *Can J Anaesth* 1999; 46: 987-90.
17. Greer JR, Smith SP, Strang T. A comparison of tracheal tube tip designs on the passage of an endotracheal tube during oral fiberoptic intubation. *Anesthesiology* 2001; 94: 729-31.

18. Stacey M, Rassam S, Sivasankar R, Hall J, Latto I. A comparison of direct laryngoscopy and jaw thrust to aid fiberoptic intubation. *Anaesthesia* 2005; 60: 445-48.
19. Aoyama K, Takenaka I, Nagaoka E, Kadoya T. Jaw thrust maneuver for endotracheal intubation using a fiberoptic stylet. *Anesth & Analg* 2000; 90: 1457-58.
20. Han S, Oh A, Jung C, Park S, Kim J, Nahm F. The effect of the jaw-thrust manoeuvre on the ability to advance a tracheal tube over a bronchoscope during oral fiberoptic intubation. *Anaesthesia* 2013; 68: 472-77.
21. Magill IW. Endotracheal anesthesia. *Am J Surg* 1936; 34: 450-55.
22. Horton WA, Fahy L, Charters P. Defining a standard intubating position using "angle finder". *Br J Anaesth* 1989; 62: 6-12.
23. Bannister F, Macbeth R. DIRECT LARYNGOSCOPY AND TRACHEAL INTUBATION. *Lancet* 1944; 244: 651-54.
24. Adnet F, Borron SW, Dumas JL, Lapostolle F, Cupa M, Lapandry C. Study of the "sniffing position" by magnetic resonance imaging. *Anesthesiology* 2001; 94: 83-6.
25. Hagberg CA. *Benumof's Airway Management*: Elsevier Health Sciences, 2007.
26. Mathru M, Esch O, Lang J, Herbert ME, Chaljub G, Goodacre B, et al. Magnetic resonance imaging of the upper airway. Effects of propofol anesthesia and nasal continuous positive airway pressure in humans. *Anesthesiology* 1996; 84: 273-9.
27. Adnet F, Borron SW, Racine SX, Clemessy J-L, Fournier J-L, Plaisance P, et al. The Intubation Difficulty Scale (IDS): Proposal and Evaluation of a New Score Characterizing the Complexity of Endotracheal Intubation. *Anesthesiology* 1997; 87: 1290-97.

Tables

Table 1 Characteristics of the subjects and results of preoperative airway evaluation

Characteristic	Group NS	Group SN
Subjects (n=53)	26	27
Age [yr: mean (range)]	53 (32-69)	52 (38-70)
Height [cm: mean (SD)]	161 (7.8)	159 (8.1)
Weight [kg: mean (SD)]	65 (8.9)	63 (13.1)
BMI [kg m ⁻² : mean (SD)]	25 (2.8)	25 (3.6)
Preoperative airway evaluation		
Mallampati class		
Class I	20	19
Class II	6	7
Class III-IV	0	1
ULBT		
Class I	21	23
Class II	5	3
Class III	0	1
Total airway score [mean (SD)]	1.15 (1.02)	1.22 (1.53)

Group NS: Neutral position followed by sniffing position, Group SN: Sniffing position followed by neutral position

Table 2. Grading for difficulty of tube insertion (DTI) and the result

Grade	Difficulty	
0	No hold-up encountered	
1	Hold-up on initial attempt, relieved by withdrawal and rotation of tube 90° anticlockwise.	
2	Hold-up on initial attempt requiring more than one manipulation of the tube, alteration in head or neck position or external manipulation	
Position	DTI Grading	Number of patients
Sniffing	0	22
	1	3
	2	1
Neutral	0	25
	1	1
	2	1

Figures




Figure 1

CONSORT flowchart showing the flow of patients through the trial. Group NS: Neutral position followed by sniffing position, Group SN: Sniffing position followed by neutral position, FOB = fiberoptic bronchoscope.




Figure 2

Comparison of the POGO (percentage of glottic opening) scores. A Solid line (—): Neutral position increased POGO score compare to Sniffing position (n = 31); a dot line (-): decreased POGO score (n = 14); a grey line: no changes in POGO score (n = 8). Neutral position showed higher POGO score than sniffing position (37.29 ± 32.56 vs. 26.65 ± 30.18 , respectively, $p = 0.002$). Neutral position significantly increased the number of patients with increased POGO score compare to sniffing position ($p = 0.017$).




Figure 3

Fiberoptic bronchoscopy views of vocal cord during (A) sniffing position and (B) neutral position.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [supplement1.docx](#)
- [supplement2.doc](#)