

Assessing of the Therapeutic Efficacy of These 2 Treatment Regimens in the Management of Helicobacter pylori Infection in Kinshasa, Democratic Republic of the Congo: A Prospective cohort study

Antoine W. Tshipi

University of Kinshasa

Patrick Ngoma

University of Kinshasa

Aliocha N. Nkodila (✉ nkodilaaliocha@gmail.com)

Protestant University in Congo

Charles Mbendi

University of Kinshasa

Aurore Beia

University of Kinshasa

Youyou Paka

University of Kinshasa

Jean Paul Kipiatu

University of Kinshasa

Trésor Monsere

University of Kinshasa

Pitshou Kengibe

University of Kinshasa

Jean Jacques Matimbo

University of Kinshasa

Mireille N. Nganga

University of Kinshasa

Benjamin Longo-Mbenza

University of Kinshasa

Article

Keywords: Helicobacter pylori, Concomitant and Sequential Quadruple Therapy, Eradication, Kinshasa

Posted Date: April 6th, 2022

DOI: <https://doi.org/10.21203/rs.3.rs-1506758/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background

Helicobacter pylori (Hp) infection is a worldwide public health problem. Unfortunately, its management poses a problem because of resistance to antibiotics. However, there are codified treatment protocols covering sequential and concomitant quadritherapy with regard to first-line probabilistic treatment. The aim of this study was to assess the therapeutic efficacy of these 2 treatment regimens in the management of Hp infection at Kinshasa.

Methods

This was a mixed study, with documentary, descriptive and interventional approaches, carried out between September 1, 2018 and April 30, 2020.

Results

Sixty-four patients were collected, including 36 men against 28 women with a sex ratio of 1H:1F; the mean age was 54 ± 16.5 years. There was an over-representation of senior patients ($n = 29$); an intermediate number of adult patients ($n = 22$) and a lower number of young patients ($n = 13$). 34 and 30 were respectively treated according to the concomitant and sequential regimens. Concomitant quadruple therapy offered an eradication rate of 91.2% compared to 56.7% for sequential quadruple therapy; concomitant treatment, advancing age and absence of risky behavior more quickly predicted the occurrence of eradication success.

Conclusion

The present study showed superiority of concomitant quadruple therapy over sequential quadruple therapy in first-line treatment. Alcohol with active smoking had a negative influence, while concomitant quadruple therapy, advancement in age had a positive influence on the success of the eradication of Hp infection.

Background

Helicobacter pylori (Hp) is a gram-negative microaerophilic bacterium that infects the epithelial mucosa of the stomach¹. It is a universally prevalent condition and is currently recognized as a public health problem, as prevalence of up to 90% has been reported in several regions of the world [2]. The areas with the highest prevalence are Africa, South America, China and India. This situation is explained by the poor socio-economic and hygienic conditions in these regions². The modes of transmission of *H. pylori* remain unclear. However, given the differences in infection rates between developed and developing countries and between urban and rural areas, socioeconomic factors appear to contribute to the transmission of this infection. In most regions, the main mechanism of spread is interfamilial transmission^{1,3-7}. In 2015,

approximately 4.4 billion people worldwide were estimated to be positive for Hp⁴. The highest prevalence of H. pylori was found in Africa (79.1%), Latin America and the Caribbean (63.4%), and Asia (54.7%), and the lowest prevalence in North America (37.1%) and Oceania (24.4%)⁸. However, the prevalence of Helicobacter pylori infection has decreased in recent decades due to improved sanitation, socio-economic development and better living conditions. Unfortunately, the prevalence remains high in developing countries. The high prevalence of Hp infection is compounded by the problem of its management⁹⁻¹¹. Indeed, eradication of Helicobacter pylori infection has become more difficult over the last decade due to the increase in antimicrobial resistance, particularly to clarithromycin and levofloxacin¹¹⁻¹³. In a recent study conducted in Kinshasa, disturbing rates of in vitro resistance were reported in the study population¹⁴. While in the Western world, studies of bacterial ecology have led to the establishment of well codified therapeutic regimens, these are lacking in developing countries, forcing them to align themselves with Western protocols¹⁵.

The 2017 CAG guidelines describe first-line treatment strategies that include clarithromycin triple therapy, bismuth quadruple therapy, concurrent quadruple therapy, sequential quadruple therapy, hybrid therapy, levofloxacin triple therapy, and fluoroquinolone quadruple therapy¹⁶. Italian researchers have described eradication rates of more than 80% with sequential probabilistic quadruple therapy and close to 73% with Clarithromycin resistance¹⁷. The efficacy of concurrent quadruple therapy has been demonstrated in several randomized trials and meta-analyses with eradication rates ranging from 85–94% [18]. The October 2015 European Consensus Conference or Maastricht V recommends abandoning sequential therapy in favor of concomitant and bismuth quadrithery, with bismuth offering an eradication rate of over 90%¹⁸⁻²⁰.

In the DRC, the extent of Hp has been demonstrated through prevalence found in various studies conducted in the provincial city of Kinshasa²¹⁻²⁴. Unfortunately, there is no common protocol for the management of this infection. Some still use triple therapy as first line, others use quadruple therapy over ten days, and others use fourteen days, either sequentially or concurrently. This disparity offers a good number of resistances and does not increase the eradication rate of Hp infection. However, the study conducted in Kinshasa by Lamarque et al.²⁵ clearly showed the superiority of fourteen-day therapy over other therapeutic modalities, without distinguishing between concurrent and sequential four-day therapy. The unanswered question in this study was which of the four-therapy regimens, sequential or concurrent, provided a better eradication rate. In fact, there is already a well-established first line management protocol in the West based on the recommendations of the Maastricht V consensus¹⁷⁻²⁰. It would be interesting to see how effective one or the other therapeutic regimen is in the Kinshasa population.

Methods

Study design : This was a mixed-methods study, involving literature (secondary analysis, retrospective), analytic, and interventional approaches. It included all 64 patients with helicobacter pylori infection followed from September 2018 to April 2020 total duration of 18 months. The study setting was the

Cinquantenaire Hospital, Médical Diamant, Médical Candeur, Spécialisé de Kinshasa, Médical de Kinshasa, and Biomédical de Kinshasa. All patients were dyspeptic with Hp infection after the rapid urease test, treated according to the concomitant and sequential therapeutic modalities (respectively 34 and 30) and submitted to the eradication control 4 weeks after treatment by detection of Hp antigen in the stool.

Data Collection

Data were collected using the OCIP mobile application (Aurore Foundation©, DRC, 2018) developed for the study by the H. pylori infection observatory of the Aurore Foundation, which is a nonprofit association that fights digestive cancers in the DRC.

Variables of Interest

The parameters of interest were age, sex, alcohol excess associated with active smoking (yes or no), results of rapid urease test (positive or negative), treatment (concurrent or sequential quadrithrapy), results of stool antigen test (positive or negative). 2.3. Operational Definitions Some operational definitions were considered in this study. H. pylori infection was having a positive rapid urease test. An adult patient was considered to be between 40 and 50 years of age and a young patient was considered to be under 40 years of age. The senior citizen was any patient at least 60 years old. Senility was the fact of being over 40 years old. For the endpoint, eradication was defined as a negative repeat stool test for H. pylori at least 4 weeks after the end of treatment.

Statistical Analysis

Statistical analyses included evaluation, validation, comparison, and inference of the data using SPSS version 25 software. File cleaning ensured quality control of univariate, bivariate, and multivariate statistical analyses. For the univariate analyses, the categorical data (qualitative and nominal) were presented in the form of frequency (number) and proportions (percentages). To do this, Pearson's Chi-square test was used to compare the percentages between two groups while Chi-square with trend was used to compare the percentages between > to 2 groups to highlight the biological gradient. The quantitative or continuous variables were summarized as mean \pm standard deviation according to their normal (symmetric) and abnormal distribution. Kaplan Meier survival curves were generated to identify the most important, independent and significant predictors according to successful eradication for Hp. The Log Rank test was compared the curves of survival time means. Obtaining the value of probability $p < 0.05$ as a threshold of statistical significance.

Results

Predictors of Eradication Success

Gender

The impact of gender was neutral on the eradication of Hp infection: 77.8% (n = 28/36) success rate in men, similar (P = 0.561) to the 71.4% success rate in women (n = 20/28) (Table 1).

Table 1. Shows sociodemographic and clinical characteristic of the study population.

	Frequency	Percentage
Male	36	56.3
Female	28	43.7
Binges drinking and active smoking	41	64.0
No risk behavior	23	36.0
Therapeutic modalities		
Sequential	30	46.9
Concomitant	34	53.1
Eradication of Hp infection	Success n=48 (75.0%)	Failure n=16 (25.0%)
Male	28(77.8)	8(22.2)
Female	20(71.4)	8(28.6)

Age

There was a highly significant quadratic relationship (P = 0.002) in letter “J” between Hp eradication success and age advancement: the proportions of success being respectively highest at the extremes of age (senior-senior age and young age versus the nadir (hole) of Hp eradication success (Figure 1).

Risk Behaviour

smoking (95.5%; n = 22/23) than in the presence of alcohol-active smoking (63, 4%; n = 26/41), with the statistical difference being highly significant (p = 0.004) (Figure 2).

Therapeutic Modalities

Successful eradication of Hp infection was more frequent in the concomitant type of therapy (91.2%; n = 31/34) than in the sequential type of therapy (56, 7%; n = 17/30), the statistical difference being highly significant (P < 0.001) (Figure 3).

Eradication Success Survival Function

Survival Function and Therapeutic Modalities

Considering survival function time as a dependent variable, concomitant treatment predicted the occurrence of eradication success faster (mean time = 11.3 ± 1.245 ; 95% CI: 8.9 - 13.7 days) than did sequential treatment (mean time = 26.1 ± 0.956 ; 95% CI: 24.2 - 27.9 days), the statistical difference being highly significant (Log Rank (Mantel-cox) Chi-square = 29.530; degree of freedom = 1; $P < 0.001$) according to Kaplan Meir curves (Figure 4).

Risk Behaviour

Considering the time of survival function as a dependent variable, the absence of risky behavior predicted the occurrence of eradication success faster (mean duration = 12.6 ± 1.646 ; 95% CI: 9.3 - 15.8 days) than did the risky behavior (mean duration = 21.4 ± 1.445 ; 95% CI: 18.6 - 24.2 days), the statistical difference being highly significant (Log Rank (Mantel-cox) Chi-square = 14.100; degree of freedom = 1; $P < 0.001$) according to the Kaplan Meier curves (Figure 5)

Age

Considering the survival function time as a dependent variable, the mean durations were summarized around 17 days for senior-senior and young age versus the mean duration around 21 days in adults, the statistical difference being significant (Log Rank (Mantel-cox) Chi-square = 7.294; degree of freedom = 2; $P < 0.026$) according to Kaplan Meir curves (Figure 6).

Discussion

This chapter has commented on the most important results according to the objectives achieved and with regard to certain data in the literature. Indeed, this study examined the socio-demographic characteristics, risk behaviors, therapeutic modalities, and predictors of success in eradicating *Helicobacter pylori* infection in patients followed in different hospital institutions in Kinshasa and subjected to sequential quadruple therapy and concomitant quadruple therapy.

Hp Infection and Sociodemographic Characteristics

Gender

Although some authors^{10, 26-27} support a small contribution of sex differences in the prevalence of Hp infection, there is still controversy as to the predominance of one or the other sex on this prevalence. Indeed, for some data in the literature, the prevalence of *H. pylori* is equal in both sexes, which is in agreement with the prevalence of Hp(+) observed in the 2009 Janulaityte-Gunther study [28] and by other authors^{27, 29}. Other authors support a small influence in the prevalence of Hp infection in favor of male sex^{10, 27}. This is the case of the study conducted by Eisdorfer, where the influence of male sex was demonstrated in the results of the urea breath test (UBT), a test commonly used for the diagnosis of *H. pylori* infection³⁰. Indeed, male sex is more associated with the risk factors for acquiring Hp infection (alcohol and active smoking) often seen in adult males³¹. However, male gender predominance was

associated with a higher prevalence of *H. pylori* infection in both children and adults. This observation reinforces the contention that the factors that might explain these differences, or affect UBT values, are not fully understood³².

In the present study, despite the 1:1 sex ratio, there was an overrepresentation of males in the study population. This may be explained by the lifestyle of men (alcohol, smoking).

Age

The mean age of patients with *Hp* infection (54.2 ± 15.1 years) in the present work reflects the preferred age of onset of symptoms of *Hp* infection. Indeed, the prevalence of infection in developing countries is greater than 80% in adults older than 50 years. Infected individuals usually acquire *H. pylori* before the age of 10 years and the infection peaks in adulthood, between 35 and 44 years of age³². It has also been shown that the seroprevalence of *H. pylori* increases with age, representing the combined effect of a decrease in the rate of exposure during childhood (associated with an improvement in the standard of living) and the acquisition of the infection with age³³, with prevalence increasing at ages above 50 years. A study comparing the infection rates of children aged 1 to 19 years with those of adults aged 20 to 75 years found a lower prevalence in the former group (22% vs. 75%)³⁴. Similarly, Mabeku et al. in a study conducted in Cameroon, showed that 62% of women over 68 years of age were infected with *Hp*³². Factors such as sexual intercourse, particularly fellatio, which is a very common sexual activity, can explain the high prevalence of *Hp*, especially in older age groups³⁵⁻³⁶. Although *Hp* is primarily a gastric organism, studies have reported that infected individuals may carry *H. pylori* permanently or transiently in their mouth and saliva.

Risk Behaviors

The present study highlighted the harmful effect of excess alcohol with active smoking in the occurrence of *Hp* infection. Mehmet in a study that investigated the relationship between blood type, sex, age, smoking and *Hp*, found that there was a susceptibility of smokers to *Hp* compared to non-smokers^{33, 37}. In previous studies, the authors have pointed out that smoking may be another factor in the sharp increase in *H. pylori* infection in persons 15 years and older [38]. In addition, it is possible that alcohol consumption, which in some studies is related to HIV status, is an indirect measure of socioeconomic and cultural variables³⁷. In contrast to the present study and others conducted around the world, Olga Sjomina et al.⁸ in a systemic review showed that there was no association between alcohols, coffee and other factors. Similarly, Wu et al., also found no association between alcohol intake and *Hp* infection^{8,38}.

Treatment Modalities and *Hp* Infection

In general, and after several treatment regimens have been evaluated, few consistently achieve high eradication rates in first-line treatment. In addition, there are also limited data on *Hp* antibiotic resistance rates to guide first-line treatment. However, the treatment regimen chosen should take into account local

patterns of antibiotic resistance (if known), prior exposure, allergies to specific antibiotics, cost, side effects, and ease of administration [39] [40] [41]. According to some recommendations, reasonable eradication targets would be $\geq 90\%$ cure rate on per-protocol analysis and $\geq 80\%$ cure rate on intention-to-treat analysis^{40,42}. In the present study, successful eradication of Hp infection was more frequent in the 1-line concomitant therapy modality (91.2%; n = 31/34) than in the 1-line sequential therapy modality (56, 7%; n = 17/30). These results are in line with much data in the literature, especially for concurrent quadruple therapy⁴³. However, the eradication rate in this study is much lower than the rates found in the West and in some SSA countries for four-line sequential therapy⁴⁴⁻⁴⁶. Although initially the 1-line sequential therapy achieved high eradication rates and was widely used in Italy⁴⁴, its efficacy has decreased over time influenced by resistance to clarithromycin and metronidazole^{21,46}. A systematic review and meta-analysis reported that rates of primary and secondary resistance to clarithromycin, metronidazole, and levofloxacin exceeded 15% (alarming levels) in all World Health Organization (WHO) regions^{21,47-49}. As a result, in 2017, clarithromycin-resistant H. pylori were defined as a high-priority bacterium in the WHO priority list of antibiotic-resistant bacteria⁵⁰. The modality of sequential treatment, which does not allow maximizing the synergy of the different antibiotics used, may explain this low eradication rate. However, Zullo et al.⁴⁵ reported that sequential treatments of 10 and 14 days achieved treatment success (above 90% according to PP analysis) for first-line H. pylori eradication in clinical practice in Italy. In Korea, where antibiotic resistance rates are relatively high, the efficacy of 10-day sequential therapy was reported at 76.3% and 85.0% on an intention-to-treat (ITT) and per-protocol (PP) analysis, respectively. These results were described as unsatisfactory⁵¹. Concomitant therapy contains three antibiotics and a PPI; its eradication rate is not compromised by resistance to clarithromycin or metronidazole, unless there is dual resistance¹³. A prospective study in Greece showed that eradication rates of concomitant therapies were above 90% by PP analysis and well tolerated. However, 31.3% of treated patients had side effects⁵².

Predictors of Successful Eradication of Helicobacter Pylori

Infection Several factors influence the outcome of eradication therapy for Hp infection. A study in South Korea showed that resistance to clarithromycin alone or in combination with metronidazole resistance (dual resistance) significantly reduced the eradication success rate after sequential treatment. However, double resistance did not affect the eradication rate after concurrent treatment⁵³. Other factors may also explain the propensity for low success with sequential therapy.

In particular, major adverse events were defined as predictors of eradication in the sequential group. Adherence to treatment had a significant influence on the efficacy outcome in the concurrent treatment group⁵³. As described in some studies^{32, 54-55}, excess alcohol with active smoking had a deleterious effect in the occurrence of Hp infection. Indeed, it has been reported in the literature that metronidazole produces a disulfiram-like reaction (vomiting, nausea, dizziness) when taken in combination with alcohol with the possibility of negatively influencing the success rate of sequential treatment, especially since the adverse effects are no longer observed in this type of therapeutic scheme. A meta-analysis showed that

smoking increased the treatment failure rate for *H. pylori* eradication^{54,55}. Similarly, a multivariate logistic regression analysis showed that smokers had a 2-fold higher probability of failure in *Helicobacter pylori* eradication than non smokers (OR: 2.0; 95% CI: 1.01 - 3.95)⁵⁵. However, lifestyle factors, including smoking and alcohol consumption, did not worsen *H. pylori* eradication failure in the study by Suzuki et al.⁵⁶. Several studies have investigated the impact of age on the success of eradication of Hp.⁵⁷⁻⁶⁰. In a Chinese study, age was found to be associated with *H. pylori* eradication. Indeed, the eradication rate was higher in patients ≥ 40 years of age than in patients < 40 years of age (85.7% vs. 54.7%, $p = 0.002$). In addition, the eradication rate was 100% in patients older than 60 years⁵⁷. Japanese authors also found a relationship between eradication failure and being in the under-50 age group⁵⁹⁻⁶⁰. These results are in line with the present study which showed that advancing age was a predictor of successful eradication of *H. pylori* infection. This is thought to be related to the fact that the gastric mucosa of advanced age is more atrophic than that of younger patients and therefore secretes less acid than normal mucosa⁵⁷⁻⁵⁹. The ability of gastric acidity to decrease the effectiveness of antibiotics is, therefore, less important in older people than in younger subjects⁵⁹.

Strengths and Limitations of the Study

The present study presented strengths balanced by some degree of limitations. The strength of this study lies in its originality. Indeed, it is the first study in Kinshasa to have investigated the eradication rate of sequential and concomitant quadritherapies in patients with Hp infection. However, the weaknesses of this study are inherent in its secondary and documentary nature, which is often characterized by biases in the completeness of information, as well as in its sample size, which did not allow certain conclusions to be drawn. Thus, an analytical study with a larger sample size would be needed to investigate other variables and to look for determinants and/or predictors of Hp eradication rates not analyzed in this study. It would be interesting to subject patients who have failed first-line therapy to second-line therapy and to evaluate its efficacy; as well as to study the efficacy of other therapeutic modalities such as hybrid triple therapy and hybrid reverse therapy which give eradication rates in ITT and PP of 97.5% and 100% respectively⁶⁰.

Implications

The current results will have implications for routine practice, pharmacovigilance, capacity building, and public health perspectives related to Hp infection in DRC. In addition, they will serve as a database for future large-scale studies.

Conclusion

The present study showed the superiority of the concomitant quadruple therapy compared to the sequential quadruple therapy. It also underlined the negative influence of alcohol with active smoking and the positive influence of quadruple therapy and advancement in age in the success of eradication of *Helicobacter pylori* infection, a prevalent problem in the DRC in general and in Kinshasa in particular.

Declarations

Acknowledgements

The authors warmly thank all the investigators who participated in the study, the mothers who agreed to take a sample from their child who was not sick, as well as the hospitals who accepted the collection of field data.

Authors' contribution

AWT and ANN initiated the study; PKN participated in the study design; AWT, ANN, PKN and BLM wrote the manuscript; CNM supervised data collection; BLM and ANN validated and analyzed the data; AB, YP, JPMK, TM, PK, JJM, MNN interpreted the data; all authors made intellectual contributions to the draft manuscript and approved the final manuscript for submission.

Funding

This research received no external funding.

Ethics approval and consent to participate

The study protocol was approved by the ethics committee of the Protestant University in Congo (n°CEUPC0037) and the study was conducted in accordance with the Helsinki principles. All participants signed written informed consent forms before enrollment.

Consent for publication

Not applicable.

Competing interests

The authors declare no conflict of interest.

Availability of data and materials

The datasets generated and analyzed during the present study are available from the corresponding author upon request : nkodilaaliocha@gmail.com

References

1. Hooi, J.K.Y., Lai, W.Y., Ng, W.K., Suen, M.M.Y., Underwood, F.E., Tanyingoh, D., et al. Global Prevalence of *Helicobacter pylori* Infection: Systematic Review and Meta-Analysis. *Gastroenterology*, 153, 420-429. (2017) <https://doi.org/10.1053/j.gastro.2017.04.022>
2. Vincent, P., Gottrand, F. and Leclerc, H. Epidemiology of *Helicobacter pylori*: Disparities in the Distribution of Infection. *Gastroentérologie Clinique et Biologique*, 20, S27-S33. (1996)

3. Perez-Perez, G.I., Rothenbacher, D. and Brenner, H. Epidemiology of *Helicobacter pylori* Infection. *Helicobacter*, 9, 1-6. (2004)<https://doi.org/10.1111/j.1083-4389.2004.00248.x>
4. Parkinson, A.J., Gold, B.D., Bulkow, L., Wainwright, R.B., Swaminathan, B., Khanna, B., et al. High Prevalence of *Helicobacter pylori* in the Alaska Native Population and Association with Low Serum Ferritin Levels in Young Adults. *Clinical and Diagnostic Laboratory Immunology*, 7, 885-888. (2000)<https://doi.org/10.1128/CDLI.7.6.885-888.2000>
5. Zabala Torres, B., Lucero, Y., Lagomarcino, A.J., Orellana-Manzano, A., George, S., Torres, J.P., et al. Review: Prevalence and Dynamics of *Helicobacter pylori* infection during Childhood. *Helicobacter*, 22, Article No. e12399. (2017)<https://doi.org/10.1111/hel.12399>
6. Lim, S.H., Kwon, J.W., Kim, N., Kim, G.H., Kang, J.M., Park, M.J., et al. Prevalence and Risk Factors of *Helicobacter pylori* Infection in Korea: Nationwide Multicenter Study over 13 Years. *BMC Gastroenterology*, 13, Article No. 104. (2013)<https://doi.org/10.1186/1471-230X-13-104>
7. Jang, K.M., Choe, B.H., Choe, J.Y., Hong, S.J., Park, H.J., Chu, M.A., et al. Changing Prevalence of *Helicobacter pylori* Infections in Korean Children with Recurrent Abdominal Pain. *Pediatric Gastroenterology, Hepatology & Nutrition*, 18, 10-16. (2015)<https://doi.org/10.5223/pghn.2015.18.1.10>
8. Sjomina, O., Pavlova, J., Niv, Y. and Leja, M. Epidemiology of *Helicobacter pylori* Infection. *Helicobacter*, 23, Article No. e12514. (2018)<https://doi.org/10.1111/hel.12514>
9. Eusebi, L.H., Zagari, R.M. and Bazzoli, F. Epidemiology of *Helicobacter pylori* Infection. *Helicobacter*, 19, 1-5. (2014)<https://doi.org/10.1111/hel.12165>
10. Hong, W., Tang, H.-L., Dong, X.-L., Hu, S.-K., Yan, Y., Basharat, Z., et al. Prevalence of *Helicobacter pylori* Infection in a Third-Tier Chinese City: Relationship with Gender, Age, Birth-Year and Survey Years. *Microbiota in Health and Disease*, 1, Article No. e150. (2019).
11. Argueta, E.A. and Moss, S.F. Treatment of *Helicobacter pylori*. *Current Opinion in Gastroenterology*, 35, 544-550. (2019)<https://doi.org/10.1097/MOG.0000000000000578>
12. Matsumoto, H., Shiotani, A. and Graham, D.Y. Current and Future Treatment of *Helicobacter pylori* Infections. In: Kamiya, S. and Backert, S., Eds., *Helicobacter pylori* in Human Diseases, Vol. 1149, Springer, Cham, 211-225. (2019)https://doi.org/10.1007/5584_2019_367
13. Hu, Y., Zhu, Y. and Lu, N.-H. Recent Progress in *Helicobacter pylori* Treatment. *Chinese Medical Journal*, 133, 335-343. (2020)<https://doi.org/10.1097/CM9.0000000000000618>
14. Ngoma, P., Longo-Mbenza, B., Tshibangu, K.E., Matungo, B., Lupande, D., Christian Malu, C. et al. Seroprevalence and Risk Factors of the *Helicobacter pylori* Infection in Bukavu City in the Democratic Republic of Congo. *Open Access Library Journal* 8(10):1-14 (2021)[doi:10.4236/oalib.1108032](https://doi.org/10.4236/oalib.1108032).
15. A Q&A and Howden, C.W. *Helicobacter pylori*: Current Treatment Options and Challenges. *The American Journal of Managed Care*®, 19-20. (2021)
16. Li, B.Z., Threapleton, D.E., Wang, J.Y., Xu, J.M., Yuan, J.Q., Zhang, C., et al. Comparative Effectiveness and Tolerance of Treatments for *Helicobacter pylori*: Systematic Review and Network Meta-Analysis. *BMJ*, 351, Article No. h4052. (2015)<https://doi.org/10.1136/bmj.h4052>

17. Gatta, L., Vakil, N., Vaira, D. and Scarpignato, C. Global Eradication Rates for *Helicobacter pylori* Infection: Systematic Review and Meta-Analysis of Sequential Therapy. *BMJ*, 347, Article No. f4587. (2013)<https://doi.org/10.1136/bmj.f4587>
18. Cuadrado-Lavín, A., Salcines-Caviedes, J.R., Diaz-Perez, A., Carrascosa, M.F., Ochagavía, M., Fernandez-Forcelledo, J.L., et al. First-Line Eradication Rates Comparing Two Shortened Non-Bismuth Quadruple Regimens against *Helicobacter pylori*: An Open-Label, Randomized, Multicentre Clinical Trial. *Journal of Antimicrobial Chemotherapy*, 70, 2376-2381. (2015)<https://doi.org/10.1093/jac/dkv089>
19. Mc Nicholl, A.G., Marin, A.C., Molina-Infante, J., Castro, M., Barrio, J., Ducons, J., et al. Randomised Clinical Trial Comparing Sequential and Concomitant Therapies for *Helicobacter pylori* Eradication in Routine Clinical Practice. *Gut*, 63, 244-249. (2014)<https://doi.org/10.1136/gutjnl-2013-304820>
20. Nkondi, S.J., Longo-Mbenza, B., Kabongo Mpolesha, J.M., Mabwa, L., Mbungu, S. and Vangu, D. Seropositivity for *Helicobacter pylori*, *Helicobacter pylori* Gastritis, Blood Pressure Components: Reality of Cardiovascular Risk. *Journal Africain d'Hépatogastroentérologie*, 1, 87-92. (2007)<https://doi.org/10.1007/s12157-007-0021-x>
21. Malfertheiner, P., Megraud, F., O'morain, C., Gisbert, J., Kuipers, E., Axon, A., et al. Management of *Helicobacter pylori* Infection—The Maastricht V/Florence Consensus Report. *Gut*, 66, 6-30. (2017)<https://doi.org/10.1136/gutjnl-2016-312288>
22. Monteiro, L. and Mégraud, F. By What Means Should *Helicobacter pylori* Be Tested before and after Eradication? *Clinical and Biological Gastroenterology*, 23, 3C-19C. (1999)
23. Correa, P. A Human Model of Gastric Carcinogenesis. *Cancer Research*, 48, 3554-3560. (1988)
24. Crowe, S.E. *Helicobacter pylori* Infection. *New England Journal of Medicine*, 380, 1158-1165. (2019)<https://doi.org/10.1056/NEJMcp1710945>.
25. Lamarque, D., Burucoa, C., Courillon-Mallet, A., de Korwin, J.-D., Delchier, J.-C., Heluwaert, F., et al. Recommendations for Treatment of *Helicobacter pylori* Infection in Adults. *Hepato-Gastro & Digestive Oncology*, 24, 157-170. (2017)
26. Ibrahim, A., Morais, S., Ferro, A., Lunet, N. and Peleteiro, B. Sex-Differences in the Prevalence of *Helicobacter pylori* Infection in Pediatric and Adult Populations: Systematic Review and Meta-Analysis of 244 Studies. *Digestive and Liver Disease*, 49, 742-749. (2017)
<https://doi.org/10.1016/j.dld.2017.03.019>
27. The Eurogast Study Group Epidemiology of, and Risk Factors for, *Helicobacter pylori* Infection among 3194 Asymptomatic Subjects in 17 Populations. *Gut*, 34, 1672-1676. (1993)
<https://doi.org/10.1136/gut.34.12.1672>
28. Janulaityte-Gunther, D., Kupcinskas, L., Pavidonis, A., Valuckas, K., Andersen, L.P. and Wadstrom, T. *Helicobacter pylori* Antibodies and Gastric Cancer: A Gender-Related Difference. *FEMS Immun Med Microbiol*, 44, 191-195. (2009) <https://doi.org/10.1016/j.femsim.2004.11.011>
29. Nijevitch, A., Khamidullina, S. and Khamidullina, F. Childhood Duodenal Ulcer Associated with *Helicobacter pylori* and ABO Blood Groups. *American Journal of Gastroenterology*, 94, 1424-1425.

- (1999)<https://doi.org/10.1111/j.1572-0241.1999.01424.x>
30. Ozaydin, N., Turkyilmaz, S.A. and Cali, S. Prevalence and Risk Factors of Helicobacter pylori in Turkey: A Nationally-Representative, Cross-Sectional, Screening with the 13C-Urea Breath Test. *BMC Public Health*, 13, Article No. 1215. (2013) <https://doi.org/10.1186/1471-2458-13-1215>
 31. Eisdorfer, I., Shalev, V., Goren, S., Chodick, G. and Muhsen, K. Sex Differences in Urea Breath Test Results for the Diagnosis of Helicobacter pylori Infection: A Large Cross-Sectional Study. *Biology of Sex Differences*, 9, Article No. 1. (2018)<https://doi.org/10.1186/s13293-017-0161-7>
 32. Mabeku, L.B.K., Ngamga, M.L.N. and Leundji, H. Potential Risk Factors and Prevalence of Helicobacter pylori Infection among Adult Patients with Dyspepsia Symptoms in Cameroon. *BMC Infectious Diseases*, 18, Article No. 278. (2018)<https://doi.org/10.1007/s10620-005-2762-y>
 33. Kanbay, M., Gür, G., Arslan, H., Yilmaz, U. and Boyaciog˘ Lu, S. The Relationship of ABO Blood Group, Age, Gender, Smoking, and Helicobacter pylori Infection. *Digestive Diseases and Sciences*, 50, 1214-1217. (2005)<https://doi.org/10.1007/s10620-005-2762-y>
 34. Park, J.S., Jun, J.S., Seo, J.-H., Youn, H.-S. and Rhee, K.-H. Changing Prevalence of Helicobacter pylori Infection in Children and Adolescents. *Clinical and Experimental Pediatrics*, 64, 21-25. (2021)<https://doi.org/10.3345/cep.2019.01543>
 35. Dimitriadi, D. Helicobacter pylori: A Sexually Transmitted Bacterium? *Central European Journal of Urology*, 67, 407-409. (2014)<https://doi.org/10.5173/cej.2014.04.art18>
 36. Sgambato, D., Visciola, G., Ferrante, E., Miranda, A., Romano, L., Tuccillo, C., et al. Prevalence of Helicobacter pylori Infection in Sexual Partners of H. pylori Infected Subjects: Role of Gastroesophageal Reflux. *United European Gastroenterology Journal*, 6, 1470-1476. (2018)<https://doi.org/10.1177/2050640618800628>
 37. Roger, A., Feldman, A., Eccersley, J.P. and Hardie, J.M. Epidemiology of Helicobacter pylori: Acquisition, Transmission, Population Prevalence and Disease To Infection Ratio. *British Medical Bulletin*, 54, 39-53. (1998)<https://doi.org/10.1093/oxfordjournals.bmb.a011678>
 38. Wu, C.T., Chen, L.K. and Hwang, S.J. Seroprevalence of Helicobacter pylori in School-Aged Chinese in Tapei City and Relationship between ABO Blood Groups. *World Journal of Gastroenterology*, 9, 1752-1755. (2003)<https://doi.org/10.3748/wjg.v9.i8.1752>
 39. Amaral, O., Fernandes, I., Veiga, N., Pereira, C., Chaves, C., Nelas, P., et al. Living Conditions and Helicobacter pylori in Adults. *BioMed Research International*, 2017, Article ID: 9082716. (2017)<https://doi.org/10.1155/2017/9082716>
 40. Myran, L. and Zarbock, S.D. Management of Helicobacter pylori Infection. *U.S. Pharmacist*, 43, 27-32. (2018)
 41. Siddalingam, R. and Chidambaram, K. Helicobacter pylori. Current Therapy and Future Therapeutic Strategies. In: Roesler, M., Ed., *Trends in Helicobacter pylori Infection*, IntechOpen London, 200, 279-302. (2014)<https://doi.org/10.5772/58338>
 42. Selecedo, J.A. and Al-Kawas, F. Treatment of Helicobacter pylori. *JAMA Internal Medicine*, 158, 842-851. (1998)<https://doi.org/10.1001/archinte.158.8.842>

43. Mégraud, F., O'Morain, C., Malfertheiner, P., on behalf of the Working Party of the European Helicobacter pylori Study Group Guidelines for Clinical Trials in Helicobacter 294 Trends in Helicobacter pylori Infection. Statistical Annex: Statistical Aspects of Clinical Trials in Helicobacter pylori Infection. *Gut*, 41, S19-S23. (1997)
44. Macias-Garcia, F., Baston-Rey, I., de la Iglesia-Garcia, D., Calvino-Suarez, C., Nieto-Garcia, L., Dominguez-Munoz, J.E. Bismuth-Containing Quadruple Therapy versus Concomitant Quadruple Therapy as First-Line Treatment for Helicobacter Pylori Infection in an Area of High Resistance to Clarithromycin: A Prospective, Cross-Sectional, Comparative, Open Trial. *Helicobacter*, 24, Article ID: e12546. (2019)<https://doi.org/10.1111/hel.12546>
45. Zullo, A., Vaira, D., Vakil, N., Hassan, C., Gatta, L., Ricci, C., et al. High Eradication Rates of Helicobacter pylori with a New Sequential Treatment. *Alimentary Pharmacology & Therapeutics*, 17, 719-726. (2003)<https://doi.org/10.1046/j.1365-2036.2003.01461.x>
46. Zullo, A., Fiorini, G., Scaccianoce, G., Portincasa, P., De Francesco, V., Vassallo, R., et al. Sequential Therapy for First-Line Helicobacter pylori Eradication: 10- or 14-Day Regimen? *Journal of Gastrointestinal and Liver Diseases*, 28, 11-14. (2019)<https://doi.org/10.15403/jgld.2014.1121.281.hpy>
47. Laving, A., Kamenwa, R., Sayed, S., Kimang'a, A.N. and Revathi, G. Effectiveness of Sequential v. Standard Triple Therapy for Treatment of Helicobacter pylori Infection in Children in Nairobi, Kenya. *South African Medical Journal*, 103, 921-924. (2013)<https://doi.org/10.7196/SAMJ.7012>
48. Nyssen, O.P., Mc Nicholl, A.G., Megraud, F., Savarino, V., Oderda, G., Fallone, C.A., et al. Sequential versus Standard Triple First-Line Therapy for Helicobacter pylori Eradication. *Cochrane Database of Systematic Reviews*, No. 6, Article No. CD009034. (2016)
<https://doi.org/10.1002/14651858.CD009034.pub2>
49. Sugano, K., Tack, J., Kuipers, E.J., Graham, D.Y., El-Omar, E.M., Miura, S., et al. Kyoto Global Consensus Report on Helicobacter pylori Gastritis. *Gut*, 64, 1353-1367. (2015)<https://doi.org/10.1136/gutjnl-2015-309252>
50. El-Serag, H.B., Kao, J.Y., Kanwal, F., Gilger, M., LoVecchio, F., Moss, S.F., et al. Houston consensus Conference on Testing for Helicobacter pylori Infection in the United States. *Clinical Gastroenterology and Hepatology*, 16, 992-1002.E6. (2018)<https://doi.org/10.1016/j.cgh.2018.03.013>
51. Liu, W.Z., Xie, Y., Lu, H., Cheng, H., Zeng, Z.R., Zhou, L.Y., et al. Fifth Chinese National Consensus Report on the Management of Helicobacter pylori infection. *Helicobacter*, 23, Article ID: e12475. (2018)<https://doi.org/10.1111/hel.12475>
52. Tacconelli, E., Carrara, E., Savoldi, A., Harbarth, S., Mendelson, M., Monnet, D.L., et al. Discovery, Research, and Development of New Antibiotics: The WHO Priority List of Antibiotic-Resistant Bacteria and Tuberculosis. *The Lancet Infectious Diseases*, 18, 318-327. (2018)[https://doi.org/10.1016/S1473-3099\(17\)30753-3](https://doi.org/10.1016/S1473-3099(17)30753-3)
53. Kim, B.J., Lee, H., Lee, Y.C., Jeon, S.W., Kim, G.H., Kim, H.S., et al. Ten-Day Concomitant, 10-Day Sequential, and 7-Day Triple Therapy as First-Line Treatment for Pylori Infection: A Nationwide

- Randomized Trial in Korea. *Gut and Liver*, 13, 531-540. (2019)<https://doi.org/10.5009/gnl19136>
54. Kapizioni, C., Koutoufaris, G., Ntouli, V., Makris, K., Milioni, K., Kourkoulis, P., et al. Optimal Duration of Concomitant Non-Bismuth Quadruple Therapy as First-Line Therapy for *Helicobacter pylori*: A Prospective, Open Label, Comparative Study. *European Journal of Gastroenterology & Hepatology*, 31, 1206-1210. (2019)<https://doi.org/10.1097/MEG.0000000000001517>
55. Kuo, C.-H., Kuo, F.-C., Hu, H.-M., Liu, C.-J., Wang, S.S.W., Chen, Y.-H., et al. Sequential and Concomitant Therapy with Four Drugs Is Equally Effective for Eradication of *H pylori* Infection. *Clinical Gastroenterology and Hepatology*, 8, 36-41.e1. (2010)<https://doi.org/10.1016/j.cgh.2009.09.030>
56. Suzuki, T., Matsuo, K., Ito, H., Sawaki, A. and Hirose, K. Smoking Increases the Treatment Failure for *Helicobacter pylori* Eradication. *The American Journal of Medicine*, 119, 217-224. (2006)<https://doi.org/10.1016/j.amjmed.2005.10.003>
57. Blanca Piazuolo, M., Mera, R. and Correa, P. Effect of Smoking on Failure of *H. pylori* Therapy and Gastric Histology in a High Gastric Cancer Risk Area of Colombia. *Acta Gastroenterologica Latino Americana*, 37, 238-245. (2008)
58. Takara, Y., Endo, H., Nakano, R., Kawachi, K., Hidaka, H., Matsunaga, T., et al. Smoking and Drinking Did Not Increase the Failure of Therapeutic *Helicobacter pylori* Eradication by Vonoprazan, Clarithromycin, and Amoxicillin. *Digestion*, 99, 172-178. (2019)<https://doi.org/10.1159/000490889>
59. Tang, Y., Tang, G., Pan, L., Zhu, H., Zhou, S. and Wei, Z. Clinical Factors Associated with Initial *Helicobacter pylori* Eradication Therapy: A Retrospective Study in China. *Scientific Reports*, 10, Article No. 15403. (2020)<https://doi.org/10.1038/s41598-020-72400-0>
60. Hassan, M., Nouredine, M., Assi, F. and Houmani, Z. (2018) Eradication Rate of *Helicobacter pylori* by Classic Triple Therapy in Lebanon: Is It Still Effective? *International Journal of Clinical Medicine*, 2, 1-5.<https://doi.org/10.15761/ICM.1000129>

Figures




Figure 1

Success in eradicating Hp infection in relation to advancing age.




Figure 2

Eradication success versus risk behavior.




Figure 3

Successful eradication of Hp infection in relation to therapeutic modalities.




Figure 4

Incidence of eradication for Hp in relation to category of treatment.




Figure 5

Incidence of eradication for Hp in relation to excess alcohol-active smoking.




Figure 6

Incidence of eradication for Hp versus advancement in age.