Spatio-Temporal Variability of Malaria Infection in Chahbahar County, Iran: Association with the ENSO and Rainfall Variability

Seyed Mohammad Jafar Nazemosadat

Shiraz University

Reza Shafiei

North Khorasan University of Medical Sciences

Habib Ghaedamini

Shiraz University

Mohsen Najjari

Shiraz University of Medical Sciences

Zahra Nazemosadat

Shiraz University of Medical Sciences

Gholamreza Hatam (✉ hatamghr@sums.ac.ir)

Shiraz University of Medical Sciences <https://orcid.org/0000-0002-7807-4793>

Research Article

Keywords: Malaria, Plasmodium falciparum, Plasmodium vivax, ENSU, Chahbahar, Iran

Posted Date: September 28th, 2021

DOI: <https://doi.org/10.21203/rs.2.24532/v2>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Environmental Science and Pollution Research on January 31st, 2022. See the published version at <https://doi.org/10.1007/s11356-021-18326-0>.

Abstract

Malaria is one of the most widespread communicable diseases in the southeast of Iran particularly in Chahbahar County. Comprehensive understanding of the influence of climate on this disease is a key element for success in the environmental-based malaria elimination programs. After analyzing the spatio-temporal distribution of the disease, the relationships between the infection statistics and some important climate indices particularly the El Niño-Southern Oscillation (ENSO) and rainfall were investigated. The malaria statistics collected from various health centers were composited with the corresponding data of Southern Oscillation Index (SOI), ground-based meteorological records and satellite-based rainfall data. Non-parametric Mann-whitely test was applied to detect the significant difference between patient number between El Niño and La Niña and between the adopted wet and dry episodes. Patient number from highest to lowest was associated to summer, autumn, spring and winter, respectively. *Plasmodium falciparum*, *Plasmodium vivax* and other species were responsible for 22%, 75% and 3% of the sickness, respectively. While the outbreak of *P. falciparum* is in autumn; *P. vivax* is erupted in summer. When the epidemic statistics were collected from rural rather than urban areas, the effect of climate on the infection statistic was more obvious. Interpretation: For rural / urban areas, the infection statistics exhibited a significant decline / increase during El Niño episodes. In autumn, spring and winter, patient number has significantly increased / decreased during the dry / wet epochs, respectively. These relationships were, however, reversed during summertime of health indicators are rarely available for every population and period of interest, and available data.

Introduction

Parasitic diseases, including incoming blood and tissue protozoa, have always originated from environmental pollution (Omidian et al., 2021; Rayani et al., 2020). Malaria is a leading life-threatening mosquito-borne parasitic disease across the globe that causes debilitating disease and sometimes terrible death in different parts of the world, especially in the tropics and subtropics, where *Anopheles* mosquitoes breed abundantly in mud and swamps (Hay et al., 2002). Researchers estimate that 219 million cases of malaria occurred worldwide in 2017, causing 435,000 deaths (Organization, 2018). In 1924, malaria infection was a serious barrier to the Iran's economic growth since more than one-third of Iran's 13 million population suffered from this infection (Ziony, 1950). Malaria control programs initiating from 1945 resulted in eradication/significant suppression of the disease in the northern/southern parts of the country by 1977, respectively (Edrissian, 2006). In 2010, 90% of the reported cases in Iran belonged to the southeastern parts of the country (Hatam, 2010; Hemami et al., 2013; Kazemi et al., 2005; Mohammadzadeh et al., 2014; Rezanezhad et al., 2011). In spite of the fact that the climate-malaria connection is not yet completely understood, scientific communities know that there is an important balance of temperature, rainfall, and humidity that creates ideal conditions for mosquitoes to breed and transmit malaria parasites (Abiodun et al., 2016; Krefis et al., 2011; Parham et al., 2012; Sena et al., 2015; Soverow et al., 2009).

The El Niño-Southern Oscillation (ENSO) is a natural phenomenon described by periodic warming of the ocean surface between the eastern and western parts of the equatorial Pacific Ocean. The warmer water stream essentially oscillates back and forth across this part of the Pacific, much like water in a large container. The El Niño (or warm phase) and La Niña (or cold phase) are the extreme phases of ENSO, which contribute to the predictability of climate anomalies in the global scale. Comprehensive definition of ENSO and its effects on global climate are presented in the web pages of and the Australian Bureau of Meteorology (BOM) and National Oceanic and atmospheric Administration (NOAA) (<http://oceanservice.noaa.gov/facts/ninonina.html>; <http://www.bom.gov.au/climate/enso/history/In-2010-12/ENSO-what.shtml>). Recent studies have shown that ENSO in conjunction with variations in the sea surface temperatures over tropical and extra-tropical water bodies induces a significant impact on precipitation variability and climate change in Iran and West Asia (Nazemosadat and Cordery, 2000; Nazemosadat and Ghaedamini, 2010; Nazemosadat and Ghasemi, 2004; Nazemosadat et al., 2006; Nazemosadat and Shahgholian, 2017).

The study area, Chahbahar, is a county located in Sistan- Baluchistan province, in the southeast of Iran with an area of 9739 Km² and population of 291910 (Zubair et al., 2008). Southern and eastern borders of the county are confined by the Indian Ocean (115 Km) and the Iran-Pakistan border (131 Km), respectively (Fig. 1). This specific geographical position is the main reason that makes this area susceptible to malaria (Fekri et al., 2014; Halimi, 2016; Hanafi-Bojd et al., 2010).

In this study, we utilized our own methodology for understanding the malaria-climate associations. In contrast to most of the previous studies, we analyzed this association in the seasonal scale and for each of the individual health center. We composited malaria statistics with the Southern Oscillation index (SOI) data that is represented as the ENSO indicator. For the first time, we utilized the satellite-based rainfall data for analyzing the malaria-rainfall relationships. To avoid misleading conclusion, we took the effects of the malaria-elimination policies on these examinations into account.

Data

Monthly values of temperature, rainfall and relative humidity of Chahbahar station (Fig. 2) were obtained from the Iranian meteorological organization for the period 1986–2016 to evaluate climatological behaviors of this region. To distinguish the El Niño, and La Niña and phases of ENSO, we extracted the corresponding monthly SOI data from the BOM's website (<http://www.bom.gov.au/climate/enso/history/In-2010-12/ENSO-what.shtml>). Monthly statistics of the number of patients collected in various health centers of Chahbahar were obtained from the regional office of the Ministry of Health and Medical Education in Iran for the period 2001 to 2011.

While the infection statistics were gathered from fourteen health centers, the sole meteorological station was located near the city center on the coast of the Indian Ocean (Fig. 2). The difference between the number of health centers and meteorological stations was a big barrier for spatial assessment of the malaria-rainfall relationships. The satellite-based Tropical Rainfall Measuring Mission (TRMM) 3B43 (V7)

rainfall data were hence utilized on a calendar month temporal resolution and a 0.25° by 0.25° spatial resolution for the period 2001 to 2012(<https://pmm.nasa.gov/data-access/downloads/trmm>; <https://trmm.gsfc.nasa.gov/>). The capabilities of the Geographical Information System (GIS) were utilized to convert the TRMM gridded data into the point data for the specified latitudes and longitudes of the health centers. Although the Goddard Earth Sciences Data and Information Services Center (GES-DISC) has recommended the application of TRMM data for the Malaria-Modeling Approaches (<https://pmm.nasa.gov/data-access/downloads/trmm>), the authors could not find any references for the application of this recommendation, as conducted in the present study.

Methods

Our preliminary examination revealed that the effects of climate indices on malaria are more obvious if their relationships are investigated in seasonal rather than monthly timescale. The collected monthly data were, hence, averaged into seasonal records so that the January-March, April-June, July-September and October-December periods were adopted as winter, spring, summer and autumn, respectively. The non-parametric Mann-Whitney and Fisher Exact tests were used to investigate the significance of the difference in the weather data or number of patients between various phases of ENSO or between the adopted wet and dry epochs(Gagnon et al., 2002; Parham et al., 2012; Wardrop et al., 2013; Zubair et al., 2008).

Construction of the SOI-climate composites

Time-series of seasonal SOI (1986–2016) were firstly sorted in ascending order. All years with seasonal SOI equal or less than -5 or equal or greater than $+5$ were assigned as the El Niño or La Niña events, respectively. Seasonal values of rainfall, temperature and relative humidity in Chahbahar station were matched with the El Niño and La Niña years to construct the ENSO composites for each variable. The significance of the difference in the mean values of these variables between El Niño and La Niña episodes was then investigated.

Construction of the SOI-malaria composites

Since malaria data were available for the period 2002–2011, seasonal values of SOI data for this period were firstly sorted. Three years with lowest or highest values of SOI were then considered as the El Niño or La Niña epoch, respectively. The other four years in the middle were assigned as the neutral epochs. The Number of Malaria Infected patients (NMI) during the El Niño and La Niña events denoted by NMI_{EI} and NMI_{La} were then statistically compared for each individual season and health center. Furthermore, the values of NMI_{EI} / NMI_{La} were used for assessing the measure of the effects of ENSO extreme phases on the malaria statistics. When this ratio is greater/less than unity, the occurrence of the El Niño generally harmonizes with the increase /decrease of the patient number. We also re-examined all the mentioned mathematical procedures when the El Niño, neutral and La Niña epochs were respectively chosen as 4, 2 and 4 years, but the results were generally similar to what is shown here.

Rainfall-malaria composites

For each individual season or health center, the TRMM rainfall data between 2002 and 2011 were sorted in ascending order. The sorted dataset were then divided into three parts including three years with highest or lowest rainfall and four years in the middle. The significance of the difference in the number of patients was then assessed between the years with highest and lowest rainfall (hereinafter wet and dry episodes). The ratio of P_w/P_d was also examined for each individual health center to assess the departure of the ratio from unity which demonstrates the effects of rainfall variability on malaria statistics. The mentioned procedures were also re-examined when the wet or dry episodes contain four years, but the results were mostly similar to those depicted in this study.

Results And Discussion

Climatic features and associations with ENSO

The mean values of annual rainfall at Chahbahar meteorological station that emphasizes the severity of dryness over the study area is 120 mm for the period 1986–2016 (Table 1). Winter, spring, summer and autumn rainfalls constitute 53.5%, 8%, 4.5% and 34% of this rainfall, respectively. As indicated, the corresponding seasonal values of temperature or relative humidity were 22. °C, 29.3°C, 29.2°C and 25.1°C or 69%, 79%, 82% and 70 %, respectively. These statistics suggests that, in contrast to rainfall, relative humidity is generally greater for warm rather than cool months of the year.

Table 1

Mean values of seasonal rainfall, temperature and relative humidity in Chahbahar for the period 1986–2016. The corresponding values of these variables during the El Niño and La Niña are also presented.

Seasons	Rainfall mm			Temperature °C			Relative Humidity %		
	Mean	El Niño	La Niña	Mean	El Niño	La Niña	Mean	El Niño	La Niña
Winter	63.9	69.7	54.3	22.0	22.1	22.0	68.8	72.1	64.8
Spring	9.6	1.1	19.5	29.3	29.1	29.3	79.4	78.4	78.0
Summer	5.4	2.4	4.2	29.2	29.1	29.5	82.0	84.0	80.6
Autumn	41.4	58.8	21.0	25.1	25.0	25.3	70.0	70.7	67.0
Annual	120.3	132.0	99.0						
Mean				26.4	26.3	26.5	75.1	76.3	72.6
* Difference in mean values is statistically significant between El Niño and La Niña (Mann-Whitney test)									
** Difference in the frequency is statistically significant El Niño and La Niña (Fisher Exact test)									

Autumn/winter rainfall is significantly/remarkably greater for the El Niño as compared with La Niña years. This relationship, however, is reversed for spring and summer, when the mean seasonal rainfall is

negligible and deviates from 9.6 to 5.4 mm, respectively. Since rainfall time-series in Chahbahar are affected by frequent near zero and sporadic torrential rain, the presented ENSO-rainfall relationships could change over time for winter, spring and summer, when these relationships are not strongly significant. Compared to El Niño, air temperature tends to be warmer during La Niña episodes from 0.2°C to 0.4°C. Relative humidity, however, is consistently greater during El Niño episodes from about 0.4% in spring to 7% in winter.

The impact of ENSO on malaria potentially increases when the difference in the climatic features is significant between El Niño and La Niña, as shown in Table 1. This influence is, hence, more remarkable during autumn, winter, summer and spring, respectively.(Zubair et al., 2008) It is concluded that long-term records of malaria data which are less affected by the disease control programs are needed to establish the ENSO-malaria relationship.

Seasonal distribution of malaria

Table 2 illustrates the historical records of patient's statistics in twelve health centers during summertime, when total infection is in its highest status. Since the dataset of the other two stations contained some missing data, their statistics are not presented. Similar statistics were also obtained for the other seasons, but due to result briefing, these data are presented in appendix 1. As indicated, mean values of the number of patients for winter, spring, summer and autumn are correspondingly 267, 1501, 5159, and 2885. While the highest records are related to the hot and dry months, the lowest values are linked to winter when rainfall is remarkably higher than the other seasons. These suggest the significant impact of temperature rather than rainfall on the number of patients.

Table 2

; Historical records of the patient number during summertime; when the presented statistics are greater than the other seasons. The statistics of twelve health centers with lowest values of missed data are presented. The name of these centers is presented as the footnote of the Table.

Year	Station Number											
	1*	2	3	4	5	6	7	8	9	10	11	12
Summer												
2002	4	25	25	6	28	37	3	13	-	185	92	8
2003	88	215	420	19	54	171	254	127	-	254	64	161
2004	15	14	44	8	6	44	40	17	2	104	61	20
2005	22	55	65	12	14	78	59	20	7	70	33	49
2006	2	15	53	3	5	23	9	3	3	36	17	2
2007	20	108	132	11	4	108	153	66	0	47	11	30
2008	14	49	312	5	1	83	50	19	6	26	4	21
2009	6	18	210	3	3	17	2	5	1	12	7	1
2010	1	26	105	2	2	9	1	7	1	34	15	7
2011	3	8	68	2	0	5	6	1	0	-	-	8
Average	17	53	143	7	12	58	58	28	3	85	34	31
1.Arabzahi, 2.Bahookalat, 3.Darges, 4.Kambal Soleyman, 5.Nagoor, 6.Nobandian, 7.Pir Sohrab, 8. Plan, 9.Sangan, 10.Shahri1, 11.Shahri2, 12.Talang												

Infection Types

P. falciparum, *P. vivax* and the other types of malaria accounted for 22%, 75%, and 3% of infections, respectively. Figure 3 illustrates the spatial distribution of autumnal *P. falciparum*-infection when the frequency of this type of infection is greater than the other seasons. Similar illustrations related to other seasons are presented in Appendix 2. In spite of the fact that the infection statistics are different from season to season, the spatial patterns of the epidemic are in general agreement, suggesting that the frequency of seasonal infection is firstly the highest over the northeastern and secondly over the southwestern parts of the county (Fig. 3 and Appendix 2). According to Fig. 4, after autumn, the infection statistics are in the second to fourth ranks during summer, spring and winter, respectively.

Figure 5 is similar to Fig. 3 except for *P. vivax* and for summer; when this infection is more acute than the other seasons. Comparing Figs. 3 and 5 suggests that, although seasons and infection types are different, the geographical patterns of the infections are almost identical for both Figs. In spite of this coherence, patient number is more than three times for *P. vivax* in Fig. 5 as compared with *P. falciparum* in Fig. 3. After summer, the risk of *P. vivax* is, respectively, more severe during autumn, spring and winter

(Fig. 6). The highest frequencies of *P. falciparum* and *P. vivax* are related to the year 2003 one of the driest year during the study period (Figs. 4 and 6). Our examination, however, did not exhibit any exceptional condition in climatic indices for this year. It is noteworthy to mention that the unusually high incidence of malaria is not necessarily driven by unusual weather patterns, and rather, is mediated by other factors such as human behavior, the mosquitos movement patterns, migration and agricultural practices (Wardrop et al., 2013).

ENSO phase and malaria

Table 3 illustrates the autumnal values of the number of patients for the warm and cold phases of the ENSO at Dagres health center as an example. The data are presented for two choices including the options that each of the ENSO phases contains 3 or 4 years. For instance, if three years with lowest or highest values of SOI are adopted as the El Niño or La Niña episodes, the mean values of the number of patients and their associated NMIEI/NMILa are 78.6, 101 and 0.78, respectively. For four years option, the statistics of the years 2004 or 2007 are also included as the El Niño or La Niña year, respectively.

Table 3

The magnitudes of autumnal SOI and their corresponding patient numbers and the ratio of NMI_{EI} / NMI_{La} for the El Niño and La Niña episodes. Patient numbers are related to Dagres health center.

Year	SOI	Patients' number	ENSO Status	Mean infection	NMI_{EI}/NMI_{La}
2009	-9.2	66	El Niño	78.6	78.6/101 = 0.78
2002	-8.0	42	El Niño		
2006	-7.1	128	El Niño		
2004	-6.80	56	El Niño If 4 years considered	73	73/96 = 0.76 If 4 years considered
2007	9.7*	80	La Niña if 4 years considered	101	
2008	14.6	191	La Niña		
2011	14.7	44	La Niña		
2010	20.6	68	La Niña If four	96	
* The statistics related to the La Nina years are shown by bold and italic.					

Figure 7 depicts the spatial distribution of the NMIEI/NMILa for autumn, when the difference between NMIEI and NMILa is significant for most of the health centers. Similar Figures were also obtained for the other seasons, but they are presented in appendix 3. (Gagnon et al., 2002) We found a statistically significant relationship between El Niño and malaria epidemics in Colombia, Guyana, Peru, and Venezuela (Hanafi-Bojd et al., 2010).

As indicated in Fig. 7, in contrast to the northern and particularly northeastern parts of the county, for the urban areas in the southwestern corner of the study area, the ratio of NMIEI/NMILa is greater than unity and deviates from 1.5 to 2.36. This suggests about 50–136% increase in the infection statistics of urban areas during El Niño as compared with the corresponding statistics during La Niña events. According to this Fig, the effects of the ENSO phases on the number of patients are reversed between eastern and western sides of the county.

Impacts of the malaria elimination policies

Since ENSO is a global climate phenomenon, it is difficult to accept that the prevalence of the El Niño or La Niña induces an opposite effect on the malaria statistics of the study area, as presented in the previous section. Our close examination revealed that the implementation of malaria eradication policies by Iranian government is probably the main cause of the observed contradiction. While the number of infected people in urban regions has been consistently reduced from 2002 to 2011, such decline in the number of patients is not found for rural regions

Table 4 delineates the ratio of the number of patients during five years period of 2002–2006 to the corresponding statistics during 2007–2011. As indicated, except for one case (summer in Bahookalat), the ratio is consistently greater than unity. This indicates a considerable reduction in the infection frequency during the 2007–2011, as compared to its preceding five years. The reduction rate is, however, different between health centers and from one season to another. The biggest or smallest ratio values are generally associated to the urban (i.e. Shahri 2 and Shahri 3) or rural areas (i.e. Darges and Bahookalat), respectively. Statistics in Table 4 prove that the malaria elimination policies are more effective in urban areas as compared to the rural parts of the county. The given results by (Fekri et al., 2014) and (Hanafi-Bojd et al., 2010) generally support these differences.

Table 4

The ratio of the infected people during the period 2002–2006 to the next five years period of 2007 to 2011. The number 1 to 12 refer to the station name as indicated in the footnote of this Table.

Season	Station Number											
	1*	2	3	4	5	6	7	8	9	10	11	12
$N_{(2002-2006)}/N_{(2006-2011)}$												
Spring	-	1.19	1.45	-	3.70	1.75	2.00	1.90	-	4.20	2.10	2.30
Summer	-	1.40	0.80	-	3.80	1.85	1.75	1.6	-	4.10	2.40	3.15
Autumn	-	1.10	1.05	-	2.80	2.50	-	2.05	-	5.05	2.45	-
Winter	-	1.05	1.50	-	4.00	1.55	2.95	1.20	-	2.80	1.25	-
*The station name are as follow:												
1.Arabzehi, 2.Bahookalat, 3.Darges, 4.Kambal-Soleyman, 5.Nagoor, 6.Nobandian, 7.PirSohrab, 8. Plan, 9.Sangan, 10.Shahri1, 11.Shahri2, 12.Talang												

According to the SOI data, in the 10 years of the study period the autumnal SOI was negative for 4 years and positive for the remaining 6 years. As indicated in Table 3, while the El Niño events had mostly occurred during the first half of the study period, the La Niña years are frequently associated to the second half of this period when the effect of technology for immunity improvement is more obvious. The considered El Niño or La Niña events, therefore, coincide with the periods with low or high effects of technology on the infection statistics, respectively. This means that if the effects of technology on the malaria statistics were removed, El Niño or La Niña events would be generally associated to the decrease or increase in the malaria epidemic as indicated for Sangan, Arabzahi, Dagres and Bahookalat in Fig. 7. In other words, the ENSO-malaria relationships are more accurate if these relations are assessed for the rural rather than urban regions.

The ratio of Pw/pd

Table 5 depicts the mean values of autumnal rainfall for the assigned dry and wet periods at Dagres health center (Fig. 2), that is presented as an example. The corresponding numbers of patients (Pw and Pd) and the ratio of pw/pd are also presented. The three years with lowest or highest rainfall are 2005, 2007 and 2003 or 2011, 2006 and 2004, respectively. Mean values of rainfall for these dry and wet periods, their corresponding number of patients and the ratio of Pw/Pd are presented. These statistics suggest that the autumnal upsurge or decline in the number of patients in Dagres health center is statistically associated to the adopted dry or wet events, respectively. The results of the statistical tests remained stationary static when Pw and Pd were four years.

Table 5

The measure of rainfall, patient number, mean rainfall, mean infection and the ratio of Pw/Pd. All data are related to the Degres health center during autumn.

	Year	Rainfall mm	Patients' number		Mean Rainfall mm/month	Mean patient	The ratios of Pw/Pd
Low Rainfall	2005	0.74	84	151	2.6 (3 years	71	120
	2007	2.72	53	80	4.8 (4 years)	65	106
	2003	4.23	76	128			51/71 = 0.72
	2010	11.71	47	68			52/65 = 0.78
	2009	37.92	49	66			76/120=
High Rainfall	2011	55.99	21	44			74/104=
	2006	71.61	94	128	67.8 (3 years	51	76
	2004	75.97	39	56	60.4(4 years)	52	74
Mean		32.61	58				

Table 6 is the seasonal values of Pw/Pd for all twelve stations. As indicated, ratios are consistently and significantly less than unity for autumn, winter and spring and greater than 1.0 for summer. This means that the occurrence of wet or dry episodes harmonize is in accordance with the reduction or increase in the number of patients during autumn, winter and spring, respectively. This relationship, however, is reversed for summer. Some direct or indirect malaria-rainfall associations were also reported by other investigators (Odongo-Aginya et al., 2005; Wardrop et al., 2013). Although precipitation provides essential habitat for larvae during the aquatic stages of some infectious diseases, drought can indirectly expand the vector's range while other researchers have also reported some negative relationships between the amount of rainfall and malaria during persistent regional drought.

Table 6
Seasonal values of the ratio of Pw/pd for all stations and various stations.

Station name	Autumn	winter	Spring	Summer
Darges	0.72	0.75	0.33	0.41
Bahokalal	0.62	0.95	0.43	1.97
sangan	0.44		0.19	
Nobandian	0.47	0.91	0.34	1.85
Nagoor	0.45	0.27	0.09	8.33
Kambal solyman	0.24	Almost no patient	Almost no patient	1.73
Talang	0.35	0.25	0.43	3.20
Pir sohrab	0.08	0.50	0.11	1.62
Plan	0.16	1.67	0.72	1.76
Arabzehi	0.22	1.00	0.36	2.89
Shahri 2	0.33	0.42	0.20	2.02
Shari 3	0.40	0.30	0.43	1.08

Conclusion

In spite of the fact that the patient number for *P. vivax* was more than three times as much as the *P. falciparum* patients, spatial patterns of these diseases were almost identical. The outbreaks of *P. falciparum* or *p. vivax* were observed in autumn and summer, respectively. Relative humidity consistently increased / decreased during El Nino/La Nina. Increase or decrease in the epidemic frequency in rural areas was associated to the La Niña or El Niño episodes. Since this relationship was reversed for urban regions, the study justified that, due to the malaria control programs, the given results for rural areas are more reliable for assessing the malaria-climate connections. The infection rate lower or higher than usual

was in general agreement with the proposed wet or dry periods in autumn, spring and winter, respectively. This association was, however, reversed for summer.

Declarations

Ethics approval and consent to participate: All data can be accessed by vice chancellor of research, Baluchistan University of Medical Sciences and is available. The corresponding author on behalf of all author confirm that the manuscript has not been and will not be published or submitted elsewhere for publication.

Consent for publication: Not applicable

Availability of data and materials: The datasets used and analyzed during the current study are available from the corresponding author on reasonable request.

Competing interests: The authors declare that they have no competing interests

Funding: No fund received

Authors' contributions: Conceptualization: [G. R. Hatam and Nazemosadat, S. M. J; Methodology: [Shafiei, R. Ghaedamini, H; Nazemosadat, Z; Najjar2,M]; Formal analysis and investigation: [G. R. Hatam and Nazemosadat, S. M. J], Writing - original draft preparation: [Shafiei, R. Ghaedamini]; Writing - review and editing: [G. R. Hatam and Nazemosadat, S. M. J]; Funding acquisition: [G. R. Hatam]; Resources: [G. R. Hatam,]; Supervision: [G. R. Hatam and Nazemosadat, S. M. J].

Acknowledgment:

The study was supported by the office of Vice-chancellor for research of Shiraz University of Medical Sciences and also technically supported by Baluchistan University of Medical Sciences.

References

1. Abiodun GJ, Maharaj R, Witbooi P, Okosun KO (2016) Modelling the influence of temperature and rainfall on the population dynamics of *Anopheles arabiensis*. *Malar J* 15:364
2. Edrissian G (2006) Malaria in Iran: Past and present situation. *Iranian Journal of Parasitology* 1:1–14
3. Fekri S, Vatandoost H, Daryanavard A, Shahi M, Safari R, Raeisi A, Omar AS, Sharif M, Azizi A, Ali AA (2014) Malaria situation in an endemic area, southeastern iran. *Journal of arthropod-borne diseases* 8:82
4. Gagnon AS, Smoyer-Tomic KE, Bush AB (2002) The El Nino southern oscillation and malaria epidemics in South America. *Int J Biometeorol* 46:81–89

5. Halimi MZ-C, Z and Jafari-Modrek M (2016) Impact of El Niño Southern Oscillation (ENSO) on Annual Malaria Occurrence in Iran. *Iranian Journal of Health Environment* 9:369–383
6. Hanafi-Bojd A, Vatandoost H, Philip E, Stepanova E, Abdi A, Safari R, Mohseni G, Bruhi M, Peter A, Abdulrazag S (2010) Malaria situation analysis and stratification in bandar abbas county, southern Iran, 2004–2008. *Journal of Arthropod-Borne Diseases* 4:31
7. Hatam G (2010) Malaria, control and diagnosis. *Iranian Red Crescent Medical Journal* 2010, 239–241
8. Hay SI, Cox J, Rogers DJ, Randolph SE, Stern DI, Shanks GD, Myers MF, Snow RW (2002) Climate change and the resurgence of malaria in the East African highlands. *Nature* 415:905–909
9. Hemami MR, Sari AA, Raisi A, Vatandoost H, Majdzadeh R (2013) Malaria elimination in iran, importance and challenges. *International journal of preventive medicine* 4
10. <http://oceanservice.noaa.gov/facts/ninonina.html>.
<http://oceanservice.noaa.gov/facts/ninonina.html>
11. <http://www.bom.gov.au/climate/enso/history/In-2010-12/ENSO-what.shtml>.
<http://www.bom.gov.au/climate/enso/history/In-2010-12/ENSO-what.shtml>
12. <https://pmm.nasa.gov/data-access/downloads/trmm>. <https://pmm.nasa.gov/data-access/downloads/trmm>
13. <https://trmm.gsfc.nasa.gov/>. <https://trmm.gsfc.nasa.gov/>
14. Kazemi B, Najari M, Saneimoghaddam E, Bandehpour M, Seyed N (2005) Detection of Plasmodium parasites in healthy blood donors using polymerase chain reaction. *Archives of Iranian Medicine* 8:135–138
15. Krefis AC, Schwarz NG, Krüger A, Fobil J, Nkrumah B, Acquah S, Loag W, Sarpong N, Adu-Sarkodie Y, Ranft U (2011) Modeling the relationship between precipitation and malaria incidence in children from a holoendemic area in Ghana. *Am J Trop Med Hyg* 84:285–291
16. Mohammadzadeh T, Hatam G, Kalantari M, Sarkari B, Motazedian MH, Sadjjadi SM, Safari R (2014) Molecular and microscopic-based characterization of Plasmodium spp. in fars and Hormozgan Provinces, South of Iran. *Journal of tropical medicine* 2014
17. Nazemosadat M, Cordery I (2000) On the relationships between ENSO and autumn rainfall in Iran. *Int J Climatol* 20:47–61
18. Nazemosadat M, Ghaedamini H (2010) On the relationships between the Madden–Julian oscillation and precipitation variability in southern Iran and the Arabian Peninsula: Atmospheric circulation analysis. *J Clim* 23:887–904
19. Nazemosadat M, Ghasemi A (2004) Quantifying the ENSO-related shifts in the intensity and probability of drought and wet periods in Iran. *J Clim* 17:4005–4018
20. Nazemosadat M, Samani N, Barry D (2006) ENSO forcing on climate change in Iran: Precipitation analysis. *Iranian Journal of Science Technology Transaction B: Engineering* 30:555–565

21. Nazemosadat MJ, Shahgholian K (2017) Heavy precipitation in the southwest of Iran: association with the Madden–Julian Oscillation and synoptic scale analysis. *Climate Dynamics*, 1–19
22. Odongo-Aginya E, Ssegwanyi G, Kategere P, Vuzi P (2005) Relationship between malaria infection intensity and rainfall pattern in Entebbe peninsula, Uganda. *Afr Health Sci* 5:238–245
23. Omidian M, Ganjkarimi AH, Asgari Q, Hatam G (2021) Molecular and serological study on congenital toxoplasmosis in newborn of Shiraz, Southern Iran. *Environ Sci Pollut Res* 28:16122–16128
24. Organization WH (2018) WHO expert consultation on rabies: third report, vol 1012. World Health Organization
25. Parham PE, Pople D, Christiansen-Jucht C, Lindsay S, Hinsley W, Michael E (2012) Modeling the role of environmental variables on the population dynamics of the malaria vector *Anopheles gambiae sensu stricto*. *Malaria Journal* 11:271
26. Rayani M, Unyah NZ, Vafafar A, Hatam GR (2020) Isoenzyme profiles and phylogenetic analysis of *Giardia duodenalis* isolates from Iranian patients. *Environ Sci Pollut Res* 27:40652–40663
27. Rezanezhad H, Menegon M, Sarkari B, Hatam G, Severini C (2011) Characterization of the metacaspase 1 gene in *Plasmodium vivax* field isolates from southern Iran and Italian imported cases. *Acta Trop* 119:57–60
28. Sena L, Deressa W, Ali A (2015) Correlation of climate variability and malaria: a retrospective comparative study, Southwest Ethiopia. *Ethiopian journal of health sciences* 25:129–138
29. Soverow JE, Wellenius GA, Fisman DN, Mittleman MA (2009) Infectious disease in a warming world: how weather influenced West Nile virus in the United States (2001–2005). *Environmental health perspectives* 117, 1049
30. Wardrop NA, Barnett AG, Atkinson J-A, Clements AC (2013) *Plasmodium vivax* malaria incidence over time and its association with temperature and rainfall in four counties of Yunnan Province, China. *Malar J* 12:452
31. Ziony M (1950) Malaria control in Iran. *Public Health Rep* 65:351–367
32. Zubair L, Galappaththy GN, Yang H, Chandimala J, Yahiya Z, Amerasinghe P, Ward N, Connor SJ (2008) Epochal changes in the association between malaria epidemics and El Niño in Sri Lanka. *Malar J* 7:140

Figures

Figure 1

Malaria risk map in Iran according to the districts classification. Chahbahar is located in the southeast of the country on the coast of Oman Sea denoted by solid black line. (Red) with Malaria transmission; (Orange) without transmission, but high risk; (Yellow) without transmission and low risk; (Green) free malaria districts. This Fig is the enhanced version of the Fig obtained in Hemami et al. (2013).

Figure 2

The spatial distribution of the health centers around Chahbahar. The position of the meteorological station is denoted by "chahbahar" in red color.

Figure 3

Spatial distribution of *P. falciparum* during autumn; when the frequency of the infected people is higher than the other seasons.

Figure 4

Seasonal variations of *P. falciparum* for for the study period.

Figure 5

Spatial distribution of *P. vivax* during summer; when the frequency of this infection is greater than the other seasons.

Figure 6

Seasonal variations of *P. vivax* for the study period.

Figure 7

Spatial distribution of the NMIEI / NMILa for autumn.