

Association between Birth Circumstances and Prenatal Maternal Mental Health: A Retrospective Study of Intergenerational Transmission

Vahideh MoghaddamHosseini

University of Pécs

Alexandra Makai (✉ alexandra.makai@etk.pte.hu)

University of Pécs

Katalin Varga

Eötvös Loránd University

Annick Bogaerts

Women & Child KU Leuven

Ákos Várnagy

University of Pécs

Research Article

Keywords: Anxiety, Breast feeding, Depression, Fear of childbirth/ W-DEQ A, Oxytocin

Posted Date: April 15th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-150994/v2>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background

The evidence on long-lasting effects of certain specific aspects of birth circumstances on the maternal mental health later during pregnancy in offspring is scarce.

Aim

This study aimed to investigate the association between birth circumstances and prenatal maternal mental health in offspring.

Method

In this retrospective survey, 380 pregnant women completed the Hungarian translation of Wijma Delivery Expectancy/Experience Questionnaire A, Beck Depression Inventory-Short Form, Beck Anxiety Inventory to measure prenatal fear of childbirth (FOC), depression, and anxiety, respectively. Information on peri and postnatal events were obtained from participants' biological mothers through Mother's Birth Circumstances Questionnaire. This included questions regarding mode of birth, administered medical interventions during labor, early life care, and breastfeeding during infancy. Multiple Linear regression was performed for statistical analysis.

Results

After adjusting for potential confounders, administration of Oxytocin (OT) induction during labor was significantly associated with higher levels of prenatal FOC ($B = 7.64$; 95% CI, 0.55, 14.73), depression ($B = 1.60$; 95% CI, 0.47, 2.73), and anxiety ($B = 3.80$; 95% CI, 0.55, 6.99). Moreover, having been breastfed for more than 12 months was significantly associated with lower level of prenatal FOC in offspring ($B = -10.45$; 95% CI, -18.79, -2.11).

Conclusion

Poor prenatal mental health might be rooted in administered OT induction as a common medical practice during labor while long duration of breast feeding can have a positive effect on improving prenatal maternal mental health in offspring. Further studies of prospective design are on demand to explore the biological trajectories of these findings in humans.

Introduction

Recently there is an increasing research interest in exploring the long-term effects of birth circumstances including peri and postnatal events on offspring's mental health across the lifespan. Allen et al (1998) defined perinatal events as surgical delivery and obstetrical complications, and categorized postnatal events under two groups of early neonatal environment such as prematurity, low birth weight and anoxia,

and late neonatal environment like breast feeding and infant health during the first year of life (1). In this regard, several studies have documented a significant association of low birth weight with anxiety disorders and depressive symptoms in adulthood (2,3). Also, breastfeeding during infancy, as a factor in late neonatal environment, was found to be a significant predictor for lower levels of depression, anxiety, and other mental problems in young adults (4,5). In contrast, some others reported no any significant link between low birth weight and breastfeeding with mental health problems later in adolescents (6).

Despite of these informative findings, there is no any evidence on the long-term effects of other aspects of birth circumstances such as medical interventions during labor and also early life care on the maternal mental health during pregnancy in offspring. Maternal fear of childbirth (FOC), depression, and anxiety are relatively common mental health problems during pregnancy with the global estimated prevalence of 11.1% (7), 11.9% (8), and 15.2% (9), respectively. Few attention has been paid to the prenatal mental health in Eastern-European countries so far. Based on one study conducted in Western-Hungary, the prevalence of prenatal mild to severe depression was estimated to be 19.9% and trait anxiety was reported to be 14.2% (10).

Prenatal maternal depression and anxiety have been frequently reported to be strong predictors for a wide range of deleterious maternal and neonatal outcomes such as lack of receiving adequate prenatal care, poor nutrition, sleep disorders, increased risk of alcohol and drugs use, suicide and infanticide, increased rate of low birth weight and preterm birth, poor maternal bonding and attachment, and certain psychological disorders in childhood and adulthood (11,12). Because of the high importance of prenatal maternal mental health, it has been recently considered as one of the essential patient safety factors by the Council on Patient Safety in Women's Health Care (13).

Although a wide range of socio-demographic, obstetric, and psychological factors have been listed as contributors to prenatal FOC (14), depression (15), and anxiety (16), there is a gap in the in-depth understanding of the root of these prenatal maternal psychological problems with respect to the intergenerational transmission. The evidence on intergenerational transmission of maternal behaviors to the rearing environment that probably via epigenetic processes (such as DNA methylation) has been displayed by several studies in rats (17–19). Consistent results from primate and rodent studies confirm strongly effectiveness of the maternal environment on offspring phenotype and it's mediating by means of changes in gene expression (17,20). Nevertheless, the evidence on humans is scares. Therefore, we aimed to investigate the association of birth circumstances including peri and postnatal events with maternal FOC, depression and anxiety, as the most common prenatal mental health problems, in offspring.

Our aim was to contribute to the already existing knowledge about the long-lasting or even transgenerational effects of maternity care received by mothers on their offspring to ameliorate prenatal maternal mental health as one of the patient safety issues.

Method

Recruitment and procedure

A population-based cross-sectional study of healthy pregnant women was carried out from April 2016 to February 2017. Recruitment of a convenience sample of women in the third trimester of pregnancy took place at the Non-Stress Test laboratory of the Department of Obstetrics and Gynecology, Hungary.

Inclusion criteria were to be able to speak and read Hungarian, and to have their mother within reach. Women who had a history of drug abuse and/or were suffering from any major psychiatric disorder were excluded from the study.

Of the 400 pregnant women invited, 380 agreed to participate after having given informed and written consent. They were then given a booklet of questionnaires and asked to return it at the subsequent check-up one week later. Information on the birth circumstances was obtained from participants' biological mothers.

Measurements

The socio-demographic and obstetric checklist included questions about participants' age, educational level, occupational status, any economic hardship, marital status, place of residence, number of pregnancies, gestational age, history of miscarriage(s), whether this pregnancy was a result of previous planning or not.

Peri and postnatal events

Mother's Birth Circumstances Questioner (MBCQ) was mapped by questions about the date of birth (due date, preterm or postterm), mode of birth (vaginal or cesarean section), interventions during labor (administration of Oxytocin (OT), amniotomy, epidural analgesia, instrumental delivery, and episiotomy), the early life care of infants (the presence of their father during their birth, immediate skin-to-skin contact with their mother, having been breastfed during the first few hours, early rooming-in), and late life care of infants (having been breastfed over the first two years).

Prenatal Mental Health

- *The Hungarian Version of Wijma Delivery Expectancy/Experience Questionnaire (W-DEQ A)* that consists of 30 items was used to measure prenatal fear of childbirth. All items of this instrument are assessed on a six-point Likert scale with a range score of 0 for 'not at all' to 5 for 'extremely', according to the degree of agreement, ranging a total score from 0 to 150 (21). The Cronbach alpha in the present study accounted for 0.88.
- *The Hungarian Version of Beck Anxiety Inventory (BAI)* consists of 21 items was used to measure prenatal anxiety. Each item is measured on a four-point Likert scale ranging from 0 ("not at all") to 3 ("severely"). The total possible score for this instrument ranges between 0 (indicating low level of anxiety) and 63 (an indication of a high level of anxiety) (22). The Cronbach alpha in the present study accounted for 0.89.

- *The Hungarian Version of Beck Depression Inventory-Short Form (BDI-SF)* consists of 9 items was used to measure prenatal depression.. *Each item* is assessed on a four-point likert scale (from 0 to 3), with a total possible score ranging from 0 to 27 (23). The Cronbach alpha for this study tool was 0.71.

Potential confounding variable

- *Multidimensional Scale of Perceived Social Support (MSPSS)* consists of 12 items was used to measure social support as a potential confounder. This instrument measures both the perceived availability and adequacy of emotional and instrumental social support from three sources of support, including family, friends, and significant others. Each item is assessed on a seven-point scale, ranging from 1 (very strongly disagree) to 7 (very strongly agree) (24). In this study, the Cronbach alpha of .86 was found for total score.

Ethical statements:

The present study was the secondary analysis of participants data recruited for the Ph.D. thesis project of the corresponding author approved by The Regional and Institutional Committee of Science and Research Ethics of the University of Pécs, Hungary (No. 5923). The research was performed in accordance with the [Declaration of Helsinki](#).

Statistical analysis

Descriptive statistics was performed to describe socio-demographic and obstetric features of participants, and psychological outcomes. In order to do the main analysis, firstly, we employed the univariate linear regression model to define the association of each section of birth circumstances questionnaire with each outcome namely prenatal FOC, depression, and anxiety, independently. After that, multivariate regression models were separately performed to adjust obtained models for maternal age, maternal education, parity, economic status, and social support as potential confounders using stepwise method.

All estimates were reported with 95% confidence intervals (95% CI) and statistical significance was considered at the level of $p < 0.05$. IBM SPSS® version 22.0 was used for the analyses.

Results

Of the 380 women who agreed to participate in the study, 371 completed and returned the questionnaires. Ten participants were excluded as they did not respond to more than half of the main items. Therefore, data of 361 participants were subject to analysis.

Basic socio-demographic and obstetric characteristics

Basic socio-demographic and obstetric characteristics of the respondents are shown in Table 1. mean age of the participants was slightly above 32 years. Over half of the women had an academic educational level and the majority of them (80.7%) were full-time employees. Married participants accounted for almost two-thirds of the sample.

In terms of obstetric characteristics, mean gestational age of the respondents was almost 37 weeks. Almost 60% of the women were multiparous and four-fifths of these women were expecting their second or third child (data not shown in Table 1). The pregnancy was unplanned in 11.5% of the cases. Almost one-third of the women reported any history of pregnancy loss until the 24th gestational week.

Descriptive analysis of the Mother's Birth Circumstances

Two-thirds of the respondents were born due date, and the majority of them (85.8%) were born vaginally, according to their mothers' recollections. At least 59% of the participants' mothers had an episiotomy and a minimum of 5,7% of them were born by means of instrumental delivery (forceps/vacuum). The incidence of the administration of EDA and OT during their birth was at least 20% and 12%, respectively.

Immediate skin-to-skin contact, early breastfeeding, and rooming-in took place in a minimum of 15%, 24%, and 13% of the cases, respectively. In addition, slightly more than half of women had been breastfed for less than six months. Table 2 illustrates all details.

Descriptive analysis of Prenatal Fear of Childbirth, Depression, Anxiety

Mean scores (SD) on scales W-DEQ A, BDI-SF, and BAI was obtained 52.43(20.58), 10.91(9.30), and 4.25(3.61), respectively.

The Association between Mother's birth Circumstances and Prenatal Fear of childbirth

After adjusting the regression models for potential confounders, the findings revealed a significant association between administration of OT induction during labor and higher levels of prenatal FOC in offspring (B= 7.64; 95% CI, 0.55, 14.73). Also, confounder parity remained in the adjusted model and was significantly linked with the lower levels of prenatal FOC (B= -3.94; 95% CI, -6.38, -1.50) (Table 3).

Moreover, in another adjusted model related to the late life care of infants, having been breastfed for more than 12 months was significantly associated with the lower level of prenatal FOC in offspring (B= -10.45; 95% CI, -18.79, -2.11). Confounders including parity, social support and having the college and

university education remained significantly in the adjusted model (Table 4). Any significant relationship was observed between other peri and postnatal events and prenatal FOC in offspring.

The Association between Mother's birth Circumstances and Prenatal Depression and Anxiety

Fully adjusted model showed the significant association of administration of OT induction during labor and higher levels of prenatal depression (B= 1.60; 95% CI, 0.47, 2.73), and anxiety (B= 3.80; 95% CI, 0.55, 6.99). Also, confounders including social support and having high economic status remained in adjusted models and significantly was related with lower levels of depression (B= -0.04; 95% CI, -0.08, -0.007) and anxiety (B= -3.02; 95% CI, -5.53, -0.51), respectively (Table 3).

Neither all the other medical interventions during labor, nor the early nor the late life care factors, was significantly linked with the prenatal depression and anxiety before and after controlling for confounders.

Discussion

This study aims to be the first one to map the link between peri and postnatal events and offspring's prenatal FOC, depression, and anxiety as the most common maternal psychological problems during pregnancy with regard to the intergenerational transmission. This issue makes it difficult to compare our results with those of previous studies.

Our results revealed that the administration of OT induction is significantly associated with higher levels of fear of childbirth, depression, and anxiety later during pregnancy in offspring. Moreover, while, there was a significant relationship between being breast fed for more than 12 months and lower level of fear of childbirth, this result was not obtained for prenatal depression and anxiety.

Hormonal plausibility

The trajectory that could justify our findings is related to an anxiolytic feature of endogenous OT in the reduction of maternal fear and anxiety. Increasing number of human and animal studies (25,26) support the role of endogenous OT in attenuating fear-related anxiety through inhibition of the stress-induced activation of the HPA axis and the reduction of the amygdala activity (27). Infusing OT receptor antagonists into the central amygdala resulted in an anxiogenic behavior in pregnant rats (28). Certain studies called the attention to the possible fetal effects of exogenous OT administered in the perinatal period. A decreased amount of face-to-face contact between mother and newborn in the first hour postpartum (29), weaker neonatal sucking reflexes (30), and interrupted pre-feeding (31) were observed in connection with labor induction by administration of exogenous OT. Syntactic OT administered during labor is able to alter physiological and emotional regulation in the mother that can influence the newborn indirectly via epigenetically transmission from mother to offspring (32). Given that in our study, the mother's recollections of OT administration during labor was significantly associated with poor prenatal

mental health in their offspring, a tentative long-term effect of the insufficient level of endogenous OT and/or OT receptor on the offspring's psychological and emotional development could be concluded.

The association of breast feeding with lower levels of fear of childbirth could be interpreted through the level of OT in lactating women and also developing greater maternal bonding and attachment style in offspring. The higher levels of OT and consequently, more positive mood and less aggression and depression in lactating women in comparison with non-lactating ones have been well documented across studies(33). Furthermore, the longer duration of breast feeding plays a known role in forming bonding and secure attachment style. On the other hand, a correlation between self-report of attachment style and the levels of plasma OT has been observed, demonstrating the higher levels of OT in women with secure attachment with their parents (34,35). The increased level of OT during the postnatal period can be transmitted to the following generation and have positive impact on them. Recently, Porta et al. (2016) pointed to the intergenerational transmission of breastfeeding behavior. It possibly happens through epigenetic mechanisms that can affect the offspring's OT receptor signaling(36). In consistent with our findings, a longitudinal prospective study showed a significant prediction of disruptive behavior in adulthood at age 30 by breastfeeding during infancy. However, family support was a significant mediator for this relationship(37). Similarly, social support remained significantly in our final model, as well.

In relation to early life care, we did not gain significant results. In contrast, the positive impact of early contact between mother and newborn (such as skin-to-skin, eye contact, touching and verbal cues) in the activation of the OT system has been documented (38). Stolzenberg and Champagne (2016) pointed to the experience-induced basis of maternal behavior which similarly to hormone-induced pathways might be able to lead to epigenetic changes via chromatin remodeling. This can play a significant role in the initiation and continuation of the mother–infant interactions (39).

A good deal of attention should be given to the role of other hormones (such as cortisol, estrogen, progesterone, endogenous opioids) and their receptor systems in maternal neuropsychological health and their interaction with OT (40). Furthermore, basic individual differences, and previous and current maternal experiences should be taken into account.

Strength and limitation

Considering the mother's birth circumstances in the examination of intergenerational transmission of an effective component of reproductive behavior in human, likely through an epigenetic mechanism brings a principle novelty in this area, presenting a new scientific approach for future studies.

One limitation of the study is that certain obstetric practices such as EDA, skin-to-skin contact with mother, rooming-in, and father's presence at birth were not common at the time when participants were born. Therefore, the rate of negative answers to these items was relatively high, resulting in a non-significant relationship. Also, fading memories of childbirth circumstances might have led to recall bias. Nevertheless, certain studies have noted that maternal recall of some peri and postnatal events, especially regarding initiation and duration of breast feeding, have acceptable accuracy(41).

Conclusion

This study, after adjusting for potential confounders, revealed that administration of OT induction can significantly predict poor prenatal maternal mental health while, breast feeding for more than 12 months was a significant predictor for the lower level of fear of childbirth, resulting in better maternal mental health later during pregnancy in offspring. Our findings support this theory that poor prenatal maternal mental health might be rooted in perinatal events and late neonatal care, showing the high importance of attention from maternity care givers to long-lasting effects of perinatal and postnatal events. This effect might be mediated through the tentative role of plasma OT and/or OT receptors in mothers and its capability of being transmitted to the next generation and having a negative or positive effect on them.

Further studies of prospective design are on demand to explore the precise biological trajectories and molecular background of these findings in humans.

Declarations

Ethics approval and consent to participate

The present study was the secondary analysis of participants data recruited for the Ph.D. thesis project of the corresponding author approved by The Regional and Institutional Committee of Science and Research Ethics of the University of Pécs, Hungary (No. 5923). The research was performed in accordance with the [Declaration of Helsinki](#).

Consent for publication: Not applicable for this paper.

Availability of data and materials: The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

Competing interests: The authors declare that they have no any conflicts of interest.

Funding: This work was supported by the Doctoral School of Health Sciences, Faculty of Health Sciences, University of Pécs, Pécs Hungary.

Authors' contributions: Vahideh MoghaddamHosseini designed and performed study, and wrote the manuscript. Katalin Varga and Ákos Várnagy supervised the study and revised the manuscript. Alexandra Makai conducted statistical analysis. Annick Bogaerts revised the manuscript. All authors contributed to and have approved the final manuscript.

Acknowledgment: We appreciate all pregnant women who agreed to participate in this study and especially midwives working at the NST laboratory for their great cooperation.

References

1. Allen NB, Lewinsohn PM, Seeley JR. Prenatal and perinatal influences on risk for psychopathology in childhood and adolescence. *Dev Psychopathol.* 1998;10(3):513–529.
2. Alati R, Najman JM, O’Callaghan M, Bor W, Williams GM, Clavarino A. Fetal growth and behaviour problems in early adolescence: findings from the Mater University Study of Pregnancy. *Int J Epidemiol.* 2009;38(5):1390–1400.
3. Nomura Y, Wickramaratne PJ, Pilowsky DJ, Newcorn JH, Bruder-Costello B, Davey C, et al. Low birth weight and risk of affective disorders and selected medical illness in offspring at high and low risk for depression. *Compr Psychiatry.* 2007;48(5):470–478.
4. de Mola CL, Horta BL, Gonçalves H, de Avila Quevedo L, Pinheiro R, Gigante DP, et al. Breastfeeding and mental health in adulthood: A birth cohort study in Brazil. *J Affect Disord.* 2016;202:115–119.
5. Hayatbakhsh MR, O’Callaghan MJ, Bor W, Williams GM, Najman JM. Association of breastfeeding and adolescents’ psychopathology: a large prospective study. *Breastfeed Med.* 2012;7(6):480–486.
6. Wiles NJ, Peters TJ, Heron J, Gunnell D, Emond A, Lewis G. Fetal growth and childhood behavioral problems: results from the ALSPAC cohort. *Am J Epidemiol.* 2006;163(9):829–837.
7. Nilsson C, Hessman E, Sjöblom H, Dencker A, Jangsten E, Mollberg M, et al. Definitions, measurements and prevalence of fear of childbirth: a systematic review. *BMC Pregnancy Childbirth.* 2018;18(1):28.
8. Woody CA, Ferrari AJ, Siskind DJ, Whiteford HA, Harris MG. A systematic review and meta-regression of the prevalence and incidence of perinatal depression. *J Affect Disord.* 2017;219:86–92.
9. Dennis C-L, Falah-Hassani K, Shiri R. Prevalence of antenatal and postnatal anxiety: systematic review and meta-analysis. *Br J Psychiatry.* 2017;210(5):315–323.
10. Bödecs T, Szilágyi E, Cholnoky P, Sándor J, Gonda X, Rihmer Z, et al. Prevalence and psychosocial background of anxiety and depression emerging during the first trimester of pregnancy: data from a Hungarian population-based sample. *Psychiatr Danub.* 2013;25(4):0–358.
11. Leight KL, Fitelson EM, Weston CA, Wisner KL. Childbirth and mental disorders. *Int Rev Psychiatry.* 2010;22(5):453–471.
12. Schetter CD, Tanner L. Anxiety, depression and stress in pregnancy: implications for mothers, children, research, and practice. *Curr Opin Psychiatry.* 2012;25(2):141.
13. Kendig S, Keats JP, Hoffman MC, Kay LB, Miller ES, Simas TAM, et al. Consensus bundle on maternal mental health: perinatal depression and anxiety. *J Obstet Gynecol Neonatal Nurs.* 2017;46(2):272–281.
14. Dencker A, Nilsson C, Begley C, Jangsten E, Mollberg M, Patel H, et al. Causes and outcomes in studies of fear of childbirth: a systematic review. *Women Birth.* 2019;32(2):99–111.
15. Gelaye B, Rondon MB, Araya R, Williams MA. Epidemiology of maternal depression, risk factors, and child outcomes in low-income and middle-income countries. *Lancet Psychiatry.* 2016;3(10):973–982.
16. Leach LS, Poyser C, Fairweather-Schmidt K. Maternal perinatal anxiety: A review of prevalence and correlates. *Clin Psychol.* 2017;21(1):4–19.

17. Champagne FA. Epigenetic mechanisms and the transgenerational effects of maternal care. *Front Neuroendocrinol.* 2008;29(3):386–397.
18. Champagne F, Diorio J, Sharma S, Meaney MJ. Naturally occurring variations in maternal behavior in the rat are associated with differences in estrogen-inducible central oxytocin receptors. *Proc Natl Acad Sci.* 2001;98(22):12736–12741.
19. Francis DD, Young LJ, Meaney MJ, Insel TR. Naturally occurring differences in maternal care are associated with the expression of oxytocin and vasopressin (V1a) receptors: gender differences. *J Neuroendocrinol.* 2002;14(5):349–353.
20. Lovic V, Gonzalez A, Fleming AS. Maternally separated rats show deficits in maternal care in adulthood. *Dev Psychobiol.* 2001;39(1):19–33.
21. MoghaddamHosseini V, Makai A, Dweik D, Várnagy Á. Factor analysis study of the Hungarian translation of Wijma Delivery Expectancy/Experience Questionnaire (version A). *Curr Psychol.* 2018;1–8.
22. Beek AT, Epstein N, Brown G, Steer RA. An inventory for measuring clinical anxiety: Psychometric properties. *J OfConsulting Clin Psychol.* 1988;56(6):893–897.
23. Rózsa S, Szádóczy E, Furedi J. Psychometric properties of the Hungarian version of the shortened Beck Depression Inventory. *Psychiatr Hung.* 2001;16(4):384–402.
24. Zimet GD, Powell SS, Farley GK, Werkman S, Berkoff KA. Psychometric characteristics of the multidimensional scale of perceived social support. *J Pers Assess.* 1990;55(3–4):610–617.
25. Mezzacappa ES, Katkin ES. Breast-feeding is associated with reduced perceived stress and negative mood in mothers. *Health Psychol.* 2002;21(2):187.
26. Lahoud N, Maroun M. Oxytocinergic manipulations in corticolimbic circuit differentially affect fear acquisition and extinction. *Psychoneuroendocrinology.* 2013;38(10):2184–2195.
27. Lee H-J, Macbeth AH, Pagani JH, Young WS. Oxytocin: the great facilitator of life. *Prog Neurobiol.* 2009;88(2):127–151.
28. Neumann ID, Krömer SA, Toschi N, Ebner K. Brain oxytocin inhibits the (re) activity of the hypothalamo–pituitary–adrenal axis in male rats: involvement of hypothalamic and limbic brain regions. *Regul Pept.* 2000;96(1):31–38.
29. Trevathan WR. Maternal “en face” orientation during the first hour after birth. *Am J Orthopsychiatry.* 1983;53(1):92.
30. Olza Fernández I, Marín Gabriel M, Malalana Martínez A, Fernández-Cañadas Morillo A, López Sánchez F, Costarelli V. Newborn feeding behaviour depressed by intrapartum oxytocin: a pilot study. *Acta Paediatr.* 2012;101(7):749–754.
31. Bell AF, White-Traut R, Rankin K. Fetal exposure to synthetic oxytocin and the relationship with prefeeding cues within one hour postbirth. *Early Hum Dev.* 2013;89(3):137–143.
32. Kenkel WM, Yee JR, Carter CS. Is oxytocin a maternal–foetal signalling molecule at birth? Implications for development. *J Neuroendocrinol.* 2014;26(10):739–749.

33. Uvnäs-Moberg K. Role of efferent and afferent vagal nerve activity during reproduction: integrating function of oxytocin on metabolism and behaviour. *Psychoneuroendocrinology*. 1994;19(5–7):687–695.
34. Strathearn L, Fonagy P, Amico J, Montague PR. Adult attachment predicts maternal brain and oxytocin response to infant cues. *Neuropsychopharmacology*. 2009;34(13):2655–2666.
35. Feldman R, Zagoory-Sharon O, Weisman O, Schneiderman I, Gordon I, Maoz R, et al. Sensitive parenting is associated with plasma oxytocin and polymorphisms in the OXTR and CD38 genes. *Biol Psychiatry*. 2012;72(3):175–181.
36. Porta F, Mussa A, Baldassarre G, Perduca V, Farina D, Spada M, et al. Genealogy of breastfeeding. *Eur J Pediatr*. 2016;175(1):105–112.
37. Essau CA, Sasagawa S, Lewinsohn PM, Rohde P. The impact of pre-and perinatal factors on psychopathology in adulthood. *J Affect Disord*. 2018;236:52–59.
38. Matthiesen A-S, Ransjö-Arvidson A-B, Nissen E, Uvnäs-Moberg K. Postpartum maternal oxytocin release by newborns: effects of infant hand massage and sucking. *Birth*. 2001;28(1):13–19.
39. Stolzenberg DS, Champagne FA. Hormonal and non-hormonal bases of maternal behavior: the role of experience and epigenetic mechanisms. *Horm Behav*. 2016;77:204–210.
40. Swain JE, Kim P, Ho SS. Neuroendocrinology of parental response to baby-cry. *J Neuroendocrinol*. 2011;23(11):1036–1041.
41. Li R, Scanlon KS, Serdula MK. The validity and reliability of maternal recall of breastfeeding practice. *Nutr Rev*. 2005;63(4):103–110.

Author Contribution

Vahideh MoghaddamHosseini designed and performed study, and wrote the manuscript. Katalin Varga and Ákos Várnagy supervised the study and revised the manuscript. Alexandra Makai conducted the statistical analysis. Annick Bogaerts revised the manuscript.

All authors contributed to and have approved the final manuscript.

Tables

Table 1: Participants' socio-demographic and obstetric characteristics

Socio-demographic characteristics	
Age (Mean± SD)	32.41±5.32
Education <i>n</i> (%)	
Less than high school	18(5.0)
High school	154(42.9)
College and university	187(52.1)
Employment status <i>n</i> (%)	
Full time employee	298(80.7)
Part time employee	7(2.0)
Self-employed	28(7.8)
Housewife	34(9.5)
Marital status <i>n</i> (%)	
Married	222(62.0)
Cohabiting	128(35.8)
Single/divorced/widow	8(2.5)
Place of residence <i>n</i> (%)	
Capital/county seat	134(37.5)
City	132(36.7)
Village	92(25.8)
Economic hardship <i>n</i> (%)	
Yes	75(21.5)
No	281(78.9)
Obstetrics characteristics	
Gestational age (Mean± SD)	36.98±1.25
Parity <i>n</i> (%)	
Nulliparous	144(40.1)
Multiparous	215(59.9)
Pregnancy Status <i>n</i> (%)	
Planned	316(88.5)
Unplanned	41(11.5)
History of miscarriage(s) and/or termination(s) of pregnancy <i>n</i> (%)	
Yes	109(30.4)
No	249(69.6)

Table 2: Descriptive statistics on mother's birth circumstances (based on the recollections of their mothers)

Date of Birth		n(%)		
Full term		231(64.5)		
Preterm		42(11.7)		
Post term		57(15.9)		
I do not know		28(7.8)		
Mode of Birth				
Vaginal delivery		308 (85.8)		
Elective Cesarean section		36 (10.0)		
Emergency Cesarean section		13 (3.6)		
I do not know		2 (0.6)		
Medical Interventions		Yes n(%)	No n(%)	I do not know n(%)
Episiotomy		182 (59.1)	47(15.3)	79 (25.6)
Amniotomy		71 (20.7)	113 (32.9)	159 (46.4)
Instrumental delivery		19 (5.7)	204 (61.3)	110(33.0)
Epidural anesthesia		70(20.6)	200(58.9)	69(20.5)
Oxytocin induction		40(11.8)	186(54.7)	114(33.5)
Early Life Care				
Father present at labor and delivery		22(6.3)	279(80.4)	46(13.3)
Immediate skin-to-skin contact with mother after birth		54(15.5)	129(37.1)	165(47.4)
Having been breastfed in the first few hours after birth		83(24.1)	85(24.6)	177(51.3)
Rooming-in with mother during the hospital stay		46(13.3)	167(48.4)	132(38.3)
late Life Care (breast feeding over the first 2 years)				
Not having been breastfed		3(1.1)		
Having been breastfed for	less than 6 months	138(52.3)		
Having been breastfed for	6-12 months	95(27.0)		
Having been breastfed for	more than 12 months	28(10.6)		

Table 3: Fully adjusted¹ model of association between Administrated OT Induction and prenatal mental health. (Only remained variables in the final model have been presented).

Variables	Prenatal Fear of Childbirth	Prenatal Depression	Prenatal Anxiety
	B (95%CI)	B (95%CI)	B (95%CI)
Administrated OT Induction	7.64(0.55, 14.73) *	1.60(0.47, 2.73) **	3.80(0.55, 6.99) *
Parity	-3.94(-6.38, -1.50) **		
Social Support		-0.04(-0.08, -0.007) *	
High economic status			-3.02(-5.53, -0.51) *
Constants	56.52	7.30	10.40
R ²	0.07	0.06	0.05
F	7.63**	6.73**	4.76**

* $p < 0.05$ ** $p < 0.01$

1- Adjusted for maternal age, maternal education, parity, economic status, and social support.

Table 4: Fully adjusted¹ model of association between having been breastfed and prenatal mental health. (Only remained variables in the final model have been presented).

Variables	Prenatal Fear of Childbirth	Prenatal Depression	Prenatal Anxiety
	B (95%CI)	B (95%CI)	B (95%CI)
Having been breast fed for more than 12 months	-10.45(-18.79, -2.11) *		
Parity	-4.44(-6.41, -2.48) **		
Social Support	-0.18(-0.36, 0.00) *	-0.07(-0.10, -0.04) **	-0.10(-0.18, -0.02) *
College and university Education	-2.57(-4.70, -0.44) *		
High economic status			-3.16(-5.28, -1.04) *
Constants	82.40	10.48	19.90
R ²	0.10	0.08	0.05
F	8.44**	7.63**	5.74**

* $p < 0.05$ ** $p < 0.01$

1- Adjusted for maternal age, maternal education, parity, economic status, and social support.