

Efficacy of Mobile App-Based Training on Health Literacy among Pregnant Women: A Randomized Controlled Trial study

Parisa Farzi Karamolahi

Guilan University of Medical Sciences

Zahra Bostani Khalesi (✉ z_bostani@yahoo.com)

Guilan University of Medical Sciences

Maryam Niknami

Guilan University of Medical Sciences

Research Article

Keywords: Training, health literacy, pregnancy, mobile application, women

Posted Date: February 15th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-151311/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background: Health literacy of mothers during pregnancy shows social and cognitive skills indicating the motivation and ability of mothers to receive and use useful knowledge to maintain and promote the health of themselves and their children. The present study was conducted aimed to determine the effect of mobile app-based training on health literacy among pregnant women.

Methods: This randomized controlled trial (RCT) study Among 500 eligible pregnant women who attended al-Zahra hospital (Rasht, Iran) between 2019 to 2020 years, was conducted on 140 pregnant women by random sampling method. They were divided into experimental and control groups. Data collection tools included personal information and maternal health literacy questionnaires. The questionnaires were completed and then a mobile app-based training intervention was performed for the experimental group. The participants of the experimental group were asked to read the contents of the software once a week for 8 weeks. The questionnaires were completed again after 8 weeks of the training intervention. The data were analyzed using SPSS software version 21. Independent t-test, Mann-Whitney U, Wilcoxon, Fisher, Chi-square, Kruskal-Wallis, and one-way analysis of variance was used for this purpose.

Results: The subjects showed no statistically significant difference in terms of demographic-social information ($p > 0.05$). The mean change in health literacy scores after the intervention was statistically significant between the experimental and control groups ($p < 0.001$). Also, the mean change in health literacy scores before and after the intervention in the experimental group was statistically significant ($p < 0.001$). But this change in the control group was not statistically significant ($p = 0.609$).

Conclusions: For the first time we used mobile app-based training and results showed that it is effective in the health literacy of pregnant women especially in situations like the Corona Virus pandemic. Therefore, it is suggested that health care providers, especially midwives, use this training method to promote the health literacy of pregnant women.

Trial registration

This paper is taken from the master thesis student of midwifery training with ethics code IR.GUMS.REC.1397.490. This is a RCT study (clinical trial code IRCT20180707040364N1: <https://www.irct.ir/search/result?query=IRCT20180707040364N1-2019-03-27>).

Background

Health literacy is one of the biggest determinants of health that is essential for improving social services and the economic development of individuals [1]. Health literacy is an emerging concept based on the two beliefs of literacy and health, which are vital sources for daily life [2]. Today, health literacy is recognized as a vital and important index of health costs and its absence increases the hospitalization rate and the use of medical services [3]. Health literacy in women is also very important because of the

important periods of life and their key role in the family [4]. According to a study by Dadipour et al., half of the pregnant mothers have inadequate health literacy and inadequate health literacy prevents proper understanding of health messages and information [5].

The health literacy of pregnant women is effective in receiving correct and timely care services from health centers [6]. Maternal health literacy is specific knowledge and special social skills to diagnose the risk factors of pregnancy, a healthy lifestyle, and proper nutrition during pregnancy and is effective on pregnancy outcomes by improving the quality of pregnancy care [7]. Prenatal care is one of the maternal and child health programs that, if adequately provided, can be an effective intervention to improve pregnancy, including reducing infant mortality, maternal mortality due to pregnancy and childbirth, and especially perinatal mortality [8].

The World Health Organization (WHO) has also emphasized the promotion of health literacy for pregnant women as a primary population [9]. Maternal health literacy is very important in pregnant mothers' understanding of prenatal risks because the mothers' understanding of risks may affect their desire to follow pregnancy advice [10]. In a study by Cho et al., mothers with lower health literacy began prenatal care later and their infants were more prone to low birth weight [11]. The promotion of health literacy requires effective training, planning, and intervention [3]. With the spread of information technology, training methods have changed significantly [12]. One of these changes was the use of new communication tools to facilitate the transfer of information and improve the quality of training and learning [13]. One of the new training methods that are effective in reducing costs and eliminating barriers to access to services such as spatial and temporal factors was training through mobile apps [14]. Learning through mobile can be limitless due to the flexibility and high computing power of mobile technologies [15]. This means, of communication, has been able to change the traditional method of in-person training and provide a new definition of training, as well as pave the way for learners to learn at home or at work in terms of time and place [16].

Regarding the high importance of health literacy of pregnant women and the ability of apps with data transfer as well as ease of access to training materials at any time of the day and night, this study was conducted aimed to determine the effect of mobile app training on health literacy in pregnant women.

Methods

This is an RCT study aimed to determine the effect of mobile app training on health literacy in pregnant women. Among 500 eligible pregnant women who attended al-Zahra hospital (Rasht, Iran) between 2019 to 2020 years), 140 passed inclusion criteria (Iranian citizenship, minimum age 18 years, maximum age 45 years, pregnant for the first time, gestational age 6–12 weeks, reading and writing literacy, no underlying and chronic disease, no medical training, a mobile phone with system Android and the ability to install the app, the ability to use and work with the mobile app, lack of experience in participating in a training program of pregnancy care, and interest in participating in the study). Exclusion criteria included: unwillingness to continue participating in the study or not using the training app installed the desired

number of times. The number of samples was determined $n = 53$ in each group, taking into account a 95% confidence interval and 90% test power. Taking into account a 25% drop, the sample size increased to 70 in each group. The sampling method of this study was the random method so that the subjects were assigned numbers 1 to 140, respectively. A Table with 35 rows was designed as a Block and each part of it was named with the letters A and B. At the next stage, the numbers were placed in each cell in order, then all the numbers were placed in blocks. 35 quadruple blocks were randomly drawn. The blocks were selected daily after the start of the study by a person who did not know the type of blocks. Those in house A were placed in the experimental group and those in house B were placed in the control group.

The random blocks were placed in a sealed envelope and kept in the nursing center of Al-Zahra Hospital, and no one knew their sequence. The blocks were selected daily after the start of the study by a person who did not know the type of blocks. People in house A were placed in the experimental group and people in house B were placed in the control group.

The data collection tool in this study was a questionnaire consisting of two parts. The first part was the demographic information questionnaire (women and spouse age, women and spouse education level, women and spouse's occupation, women's income, and place of residence) and the second part was the Maternal Health Literacy and Pregnancy (MHLAP). The MHLAP has 14 items. The range of maternal health literacy scores is from 14 to 70. The items of this questionnaire are based on a 5-option Likert scale of Totally Agree (Score 5), Agree (Score 4), No idea (Score 3), Disagree (Score 2), and Totally Disagree (Score 1). A higher score shows more health literacy. The Maternal Health Literacy Questionnaire was developed by Mojuyinola (2011) in Nigeria[17], and localized in Iran by Kharazi et al[18]. The Cronbach's alpha coefficient was 0.89 for health literacy and 0.67 for pregnancy outcome.

The mobile app, based on the items of the MHLAP questionnaire (changes and adaptations of the body during pregnancy, the fetus development, pregnancy care, tests, vaccinations, nutrition, taking supplements, personal hygiene, employment, travel, rest, exercise, sex, addiction, taking medication, pregnancy risk factors, neonatal care, bathing and breastfeeding, neonatal risk factors, vitamin taking, baby position, and coverage) was installed on smartphones in the experimental group and the mother was taught how to use the training app. One week after installing the app on the mobile phone of the mothers of the experimental group, possible questions and problems in using the app were answered during a telephone call. Then, every week, participants were reminded about using the app via SMS, and the researcher's phone number was provided to the participants so that they could contact the researcher if she had any questions about the app. Eight weeks later, MHLAP was completed by the questioner in both groups. The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Statistical analysis

Statistical analysis was performed using descriptive and inferential statistical methods. In order to compare the experimental and control groups before and after the training intervention based on the

mobile app, we used statistical tests, including independent t-test, U Mann-Whitney test, and Wilcoxon test by SPSS software version 16. The difference was considered significant at a significant level of $P < 0.05$.

Results

At the end of the study, among 140 pregnant women (70 pregnant women in the experimental group and 70 pregnant women in the control group), 10 women in the experimental group (1

participant due to unwillingness to continue participating in the study, 6 participants due to app deletion by themselves, 1 participant due to abortion and 2 participants due to not reading the app) and 17 participants in the control group (7 participants due to unwillingness to continue participating in the study, 9 participants due to abortion, and 1 participant due to ectopic pregnancy) were excluded from the study (Fig. 1).

The mean age of participants was 30.1 ± 5.66 years, the majority of participants had a diploma (38.05%) and were housewives (84.96%).

The mean age of the spouses was 33.6 ± 5.7 years. The majority of spouses had a high school education and diplomas (40.71%), and the majority of spouses were self-employed (76.11%). The monthly household income of the majority (46.90%) of the subjects had incomes between one to one and a half million.

According to the information in Table 1, before and after the intervention, the changes in the health literacy score are statistically significant. The mean score of health literacy before the intervention in the experimental group was 45.35 ± 6.75 and 42.02 ± 7.41 in the control group. Also, the mean score of health literacy after the intervention in the experimental group was 58.03 ± 6.57 and 42.23 ± 8.47 in the control group. The change in health literacy score before and after the intervention was 12.68 ± 6.31 in the experimental group and 0.21 ± 4.25 in the control group. In the study of intragroup changes, the changes were significant only in the experimental group based on the Wilcoxon test ($p < 0.001$).

Table 1
Comparison of health literacy of the two groups: before and after intervention

Health literacy		Study group		Significance level	Statistical test type
		Experimental	Control		
Before the intervention	M	45.35	42.02	* <0.014	Independent t-test
	SD	6.75	7.41		
	Median	44	42		
After the intervention	M	58.03	42.23	* <0.001	Mann-Whitney U
	SD	6.57	8.47		
	Median	59	42		
Health literacy score changes before and after the intervention	M	12.68	0.21	* <0.001	
	SD	6.31	4.25		
	Median	13	0.001		
Significance level		* <0.001	0.609		
* Significance					

According to the inter-group comparison of the health literacy score before and after the intervention according to the age of the spouse using the Wilcoxon test, a statistically significant difference was observed in the age groups of 30 years and younger and 31 years and older ($p < 0.001$). According to the intragroup comparison of the health literacy score before the intervention in terms of the level of education of the spouse using a one-way analysis of variance test, a statistically significant difference was between the subjects ($p = 0.004$). According to the intergroup comparison, health literacy score before and after the intervention in terms of the spouse's education using Wilcoxon test, the score changes before and after the training intervention in the primary school ($p = 0.014$), elementary school ($p = 0.001$), high school and diploma groups ($p < 0.001$), Associate degree ($p = 0.036$) and bachelor and higher ($p = 0.006$) were significantly different.

According to the intragroup comparison of the health literacy score before the intervention according to the occupation of the spouse using a one-way analysis of variance test, a statistically significant difference was observed between the subjects ($p < 0.001$). According to the intergroup comparison of the health literacy score before and after the intervention in terms of the level of occupation of the spouse using the Wilcoxon test, a significant difference was between the subjects in the group of employee ($p = 0.009$), worker ($p = 0.018$) and self-employed ($p < 0.001$).

According to the intragroup comparison of the health literacy score before the intervention in terms of monthly household income using a one-way analysis of variance test, a statistically significant difference

was observed between the subjects ($p = 0.012$).

According to the intergroup comparison of health literacy score before and after the intervention in terms of monthly household income using Wilcoxon signed-rank test, a significant difference was between the subjects in the group with income less than one million ($p = 0.009$), one to one and a half million ($p < 0.001$), one and a half to two million ($p = 0.003$), and two million and more ($p = 0.005$).

As shown in Table 2, a multiple regression model has been implemented using the step-by-step method, and the effect of training intervention after controlling health literacy score before training intervention, age of women and spouse, education level of women and spouse, occupation of women and spouse, monthly income and place of residence was still significant in the final linear regression model as a predictor variable related to health literacy ($p < 0.001$), so that the training intervention increased health literacy score by 13.4% ($\beta \pm SE = 13.4 \pm 1$). In addition to the effect of the training intervention, the health literacy score before the intervention also had a significant relationship with changes in health literacy score ($p = 0.015$). With increasing health literacy scores before the intervention, the rate of change in health literacy score reduces ($\beta \pm SE = -0.181 \pm 0.073$).

Table 2
Intervention effect on changes in health literacy score after controlling the effects of confounding variables

Variable	Unstandardized Coefficients		Standard error	Significance level	Beta 95% confidence interval	
	Beta Regression coefficient	Regression coefficient standard error			Beta	Lower bound
(Constant)	-12.268	1.652		0.000	-15.542	-8.995
Intervention	12.476	1.026	0.756	0.000	10.443	14.509
(Constant)	-3.358	3.141		0.287	-9.582	2.865
Intervention	13.243	1.011	0.802	0.000	11.24	15.246
Health literacy scores before the intervention	-0.230	-0.070	-0.201	0.001	-0.369	-0.091
(Constant)	-3.066	3.099		0.325	-9.208	3.077
Intervention	13.421	1	0.813	0.000	11.438	15.403
Health literacy scores before the intervention	-0.181	0.073	-0.158	0.015	-0.326	-0.036
Education of spouse	-0.931	0.455	-0.129	0.043	-1.832	-0.030
a. Dependent Variable: Health literacy score changes before and after the intervention R Square =) 0.624)						

Also, spouse's education was another factor related to changes in health literacy score in this study, with increasing spouse education level, health literacy score reduced ($\beta \pm SE = -0.931 \pm 0.455$).

Discussion

According to the results of the present study, the mean score of health literacy after the intervention was significantly different ($p < 0.001$). The results of the present study were consistent with the study results of Kharazi et al., [19]. In a study by Kharazi, the mean score of health literacy after the intervention in the experimental group was 45.76 ± 5.29 and 39.21 ± 7.03 in the control group. A significant difference was between the two groups in terms of health literacy after the intervention using the statistical analysis ($p < 0.001$).

The study results showed that the mean change in health literacy score before and after the intervention was significantly different in the experimental group ($p < 0.001$) while no significant difference was found

in the control group ($p = 0.609$). The significant changes indicate that the implemented training program has been effective in promoting maternal health literacy in the intervention group. The results of the present study were consistent with the study results of Kamali et al. In a study by Kamali in terms of changes in health literacy score before and after the intervention a significant difference was found in the experimental group ($p < 0.001$) while no significant difference was found in the control group ($p = 0.909$) [20].

According to the intergroup comparison of health literacy scores before and after the intervention in terms of age of the women, a statistically significant difference was observed between the subjects ($p < 0.001$). According to the intragroup comparison of health literacy scores before the intervention in terms of education of the mother, a statistically significant difference was observed between the subjects ($p < 0.001$). Also, according to the intragroup comparison of the health literacy score after the intervention in terms of education of the participants, a statistically significant difference was observed between the subjects ($p < 0.001$). This finding is consistent with the study results of Kohan et al., [21]. Ghanbari et al., [22], and Safari, et al., [5]. The high level of health literacy in those with higher education confirms the role of education in the level of health literacy. Because general literacy is the basis for the level of health literacy [19].

According to the intergroup comparison of health literacy scores before and after the intervention in terms of age, the level of education of the spouse ($p < 0.001$), the spouse's occupation ($p < 0.001$), monthly income ($p = 0.012$) showed a statistically significant difference between the participants.

The results of the present study showed that the effect of training intervention after controlling health literacy score before training intervention, age of women and spouse, education level of women and spouse, occupation of women and spouse, monthly household income, and residence is still significant in the final model of linear regression as a predictor variable related to health literacy ($p < 0.001$) so that on average, training intervention increases the score of health literacy level by 13.4%.

In addition to the effect of the training intervention, the health literacy score before the intervention had a significant relationship with changes in health literacy score ($p = 0.015$) with increasing health literacy score before the intervention, the rate of score changes reduces. Also, the education of the spouse was another factor related to the changes in health literacy score in this study, with increasing the level of education of the spouse, the changes in the score reduce.

According to a study by Safari, et al., [5] the most effective factor in predicting health training was education $OR = 7.345$ and the lowest effect was related to occupation $OR = 2.542$, which is inconsistent with the results of the present study which is due to the type of study. The present is a clinical trial and a study by Safari was a descriptive-analytical study.

Conclusion

In general, a mobile app was tested in this study and was found to be efficacious in promoting the health literacy of pregnant women. Also, this study showed that mobile-based interventions can be an option for mass dissemination for improving health in pregnant women. Health providers should be aware of this technology and familiarize themselves with the various features of each app because this method will enable them to instruct women pregnant remotely. On the other hand, health literacy is key for self-care during pregnancy and mobile health offers pregnant women a new option for self-care during pregnancy.

Limitations of the study

The major limitation of this study is related to the difficulty to interpret or generalize the results because the studied population is very different from the population treated in normal life.

Abbreviations

WHO: World Health Organization

RCT: Randomized Controlled Trial

MHLAP: Maternal Health Literacy and Pregnancy

Declarations

All methods were performed in accordance with the relevant guidelines and regulations (Declaration of Helsinki).

Ethics approval and consent to participate

This paper is taken from the master thesis student of midwifery training with ethics code IR.GUMS.REC.1397.490. The researchers sincerely appreciate the cooperation and support of the Department of Research and Technology of Guilan University of Medical Sciences. Informed consent to participate in the study obtained from participants.

We send an email to the editor and request for waiver. He confirms that we will be able to provide a full waiver for the APC for our manuscript after it is accepted for publication. (RE: Request to free of publishing charges Ticket ID [#5193100]).

Consent for publication:

Not applicable.

Availability of data and materials:

The data that support the findings of this study are available from the corresponding author upon a reasonable request.

Competing interests:

No conflict of interest

Funding:

Not applicable

Authors' contributions:

Z.B. and P.F participated in the Conceptualization, design, and implementation of the intervention, analysis of the findings, and drafting of the manuscript. M.N. participated in the design of the study and writing—review and editing of the manuscript. All authors read and approved the final manuscript.

Acknowledgments:

We appreciate the support and assistance of all the pregnant women who participated in this study.

Authors' information (optional):

Parisa Farzi Karamolahi: Social Determinants of Health Research Center, School of Nursing and Midwifery, Guilan University of Medical Sciences, Rasht, Iran

Zahra Bostani Khaledi: Social Determinants of Health Research Center, School of Nursing and Midwifery, Guilan University of Medical Sciences, Rasht, Iran

Maryam Niknami: Social Determinants of Health Research Center, School of Nursing and Midwifery, Guilan University of Medical Sciences, Rasht, Iran

References

1. Kandula NR, Nsiah-Kumi PA, Makoul G, Sager J, Zei CP, Glass S, et al. The relationship between health literacy and knowledge improvement after a multimedia type 2 diabetes education program. *PEC*. 2009;75(3):321-7.

2. Kanj M, Mitic W, editors. Health literacy and health promotion: Definitions, concepts and examples in the Eastern Mediterranean region. Kenya: WHO 7th Global Conference on Health Promotion Promoting Health and Development: Closing the Implementation Gap; 2009.
3. Tol A, Pourreza A, Tavasoli E, Rahimi Foroshani A. Determination of knowledge and health literacy among women with type 2 diabetes in teaching hospitals of TUMS. *HOSPITAL*. 2012;11(3):45_52.
4. Ferguson B. Health literacy and health disparities: The role they play in maternal and child health. *NWH*. 2008;12(4):286-98.
5. Safari -Moradabadi A, Aghamolaei T, Ramezankhani A, Dadipoor S. The Health Literacy of Pregnant Women in Bandar Abbas, Iran. *J Sch Publ Health Inst Publ Health Res*. 2017;15(2):121-32.
6. Tabandeh A, Kashani E. Effects of maternal body mass index and weight gain during pregnancy on the outcome of delivery. *J Gorgan Univ Med Sci*. 2007;9(1):20_4.
7. Ohnishi M, Nakamura K, Takano T. Improvement in maternal health literacy among pregnant women who did not complete compulsory education: policy implications for community care services. *Health Policy*. 2005 May;72(2):157-64.
8. Gashtasbi A, Fereshte M, Rahimi Froshani A, Nadim A. Evaluation the quality of prenatal care in Kohgiluyeh va Boyerahmad. *PAYESH*. 2002;1(1):7_12.
9. Sedaghati P, Ardjmand A, Sedaghati N. Does Regular Ergometric Training Have Any Effect On The Pregnancy Outcome?. *IJP*. 2006;16(3):325-31.
10. Tol A, Pourreza A, Rahimi Foroshani A, Tavassoli E. Assessing the effect of educational program based on small group on promoting knowledge and health literacy among women with type2 diabetes referring to selected hospitals affiliated to Tehran University of Medical Sciences. *RJMS*. 2013;19(104): 10-9.
11. Cho YI, Lee S-YD, Arozullah AM, Crittenden KS. Effects of health literacy on health status and health service utilization amongst the elderly. *SOC SCI MED*. 2008;66(8):1809-16.
12. Soufi Siavash S. Assessing e_learning readiness at Gilaun University by using Fuzzy Logic Method. A thesis submitted to the Graduate Studies Office in partial fulfillment of the requirements for The degree of Master Of Art in Information Technology Management. Guilan: Mehr Alborz virtual University; 2012.
13. Alzaza NS, Yaakub AR. Students' awareness and requirements of mobile learning services in the higher education environment. *American Journal of Economics and Business Administration*. 2011;3(1):95-100.
14. Farzi Karamolahi P, Bostani Kholesi Z, Niknami M. Pregnant Women's Health Literacy Promotion Strategies: A Review. *Journal of Pharmaceutical Research International*. 2019;28(2): 1-8.
15. Zakerimoghadam M, Bassampour S, Rjab A, Faghihzadeh S, Nesari M. Effect of nurse-led telephone follow ups (tele-nursing) on diet adherence among type 2 diabetic patients. *H* 2008;14(2):63-71.
16. Zamani BE, Babri H, Ghorbani S. Strategies for the development of mobile learning through teaching-learning activities in medical education: perspectives of medical students and IT professionals in Isfahan university of medical sciences. *Iranian Journal of Medical Education*. 2013;13(2):87-97.

17. Mojinyinola J. Influence of maternal health literacy on healthy pregnancy and pregnancy outcomes of women attending public hospitals in Ibadan, Oyo State, Nigeria. *African Research Review*. 2011;5(3):28_39.
18. Kharazi S, Peyman N, HA E. An Evaluation of the Validity and Reliability of the Maternal Health Literacy and Pregnancy Outcome Questionnaire *Journal of Health System Research*.12 (4): 512-9.
19. Kharazi SS, Peyman N, Esmaily H. Effect of an Educational Intervention Based on Self-efficacy Theory and Health Literacy Strategies on Pregnancy Outcomes: A Randomized Clinical Trial. *The Iranian Journal of Obstetrics, Gynecology and Infertility*. 2018;21(5):33-46.
20. Kamali Z, Abedian Z, Mohammad A, Mohebbi-Dehnavi Z. The Effect of Small-Group Teaching on Health Literacy in Pregnant Females with Nausea and Vomiting: A Clinical Trial. *Journal of Nursing Education* 2018;6 (6): 25-32.
21. Kohan S, Ghasemi S, Dodangeh M. Associations between maternal health literacy and prenatal care and pregnancy outcome. 2008;12(4):146-52.
22. Ghanbari S, Majlessi F, Ghaffari M, Mahmoodi Majdabadi M. Evaluation of health literacy of pregnant women in urban health centers of Shahid Beheshti Medical University. *Daneshvar*. 2012;19(97):1-12.
23. Kandula NR, Nsiah-Kumi PA, Makoul G, Sager J, Zei CP, Glass S, et al. The relationship between health literacy and knowledge improvement after a multimedia type 2 diabetes education program. *PEC*. 2009;75(3):321-7.
24. Kanj M, Mitic W, editors. Health literacy and health promotion: Definitions, concepts and examples in the Eastern Mediterranean region. Kenya: WHO 7th Global Conference on Health Promotion Promoting Health and Development: Closing the Implementation Gap; 2009.
25. Tol A, Pourreza A, Tavasoli E, Rahimi Foroshani A. Determination of knowledge and health literacy among women with type 2 diabetes in teaching hospitals of TUMS. *HOSPITAL*. 2012;11(3):45_52.
26. Ferguson B. Health literacy and health disparities: The role they play in maternal and child health. *NWH*. 2008;12(4):286-98.
27. Safari -Moradabadi A, Aghamolaei T, Ramezankhani A, Dadipoor S. The Health Literacy of Pregnant Women in Bandar Abbas, Iran. *J Sch Publ Health Inst Publ Health Res*. 2017;15(2):121-32.
28. Tabandeh A, Kashani E. Effects of maternal body mass index and weight gain during pregnancy on the outcome of delivery. **J Gorgan Univ Med Sci**. 2007;9(1):20_4.
29. Ohnishi M, Nakamura K, Takano T. Improvement in maternal health literacy among pregnant women who did not complete compulsory education: policy implications for community care services. *Health Policy*. 2005 May;72(2):157-64.
30. Gashtasbi A, Fereshte M, Rahimi Froshani A, Nadim A. Evaluation the quality of prenatal care in Kohgiluyeh va Boyerahmad. *PAYESH*. 2002;1(1):7_12.
31. Sedaghati P, Ardjmand A, Sedaghati N. Does Regular Ergometric Training Have Any Effect On The Pregnancy Outcome?. *IJP*. 2006;16(3):325-31.

32. Tol A, Pourreza A, Rahimi Foroshani A, Tavassoli E. Assessing the effect of educational program based on small group on promoting knowledge and health literacy among women with type2 diabetes referring to selected hospitals affiliated to Tehran University of Medical Sciences. *RJMS*. 2013;19(104): 10-9.
33. Cho YI, Lee S-YD, Arozullah AM, Crittenden KS. Effects of health literacy on health status and health service utilization amongst the elderly. *SOC SCI MED*. 2008;66(8):1809-16.
34. Soufi Siavash S. Assessing e_learning readiness at Gilaun University by using Fuzzy Logic Method. A thesis submitted to the Graduate Studies Office in partial fulfillment of the requirements for The degree of Master Of Art in Information Technology Management. Guilan: Mehr Alborz virtual University; 2012.
35. Alzaza NS, Yaakub AR. Students' awareness and requirements of mobile learning services in the higher education environment. *American Journal of Economics and Business Administration*. 2011;3(1):95-100.
36. Farzi Karamolahi P, Bostani Khalesi Z, Niknami M. Pregnant Women's Health Literacy Promotion Strategies: A Review. *Journal of Pharmaceutical Research International*. 2019;28 (2): 1-8.
37. Zakerimoghadam M, Bassampour S, Rjab A, Faghihzadeh S, Nesari M. Effect of nurse-led telephone follow ups (tele-nursing) on diet adherence among type 2 diabetic patients. *H* 2008;14(2):63-71.
38. Zamani BE, Babri H, Ghorbani S. Strategies for the development of mobile learning through teaching-learning activities in medical education: perspectives of medical students and IT professionals in Isfahan university of medical sciences. *Iranian Journal of Medical Education*. 2013;13(2):87-97.
39. Mojinyinola J. Influence of maternal health literacy on healthy pregnancy and pregnancy outcomes of women attending public hospitals in Ibadan, Oyo State, Nigeria. *African Research Review*. 2011;5(3):28_39.
40. Kharazi S, Peyman N, HA E. An Evaluation of the Validity and Reliability of the Maternal Health Literacy and Pregnancy Outcome Questionnaire *Journal of Health System Research*. 12 (4): 512-9.
41. Kharazi SS, Peyman N, Esmaily H. Effect of an Educational Intervention Based on Self-efficacy Theory and Health Literacy Strategies on Pregnancy Outcomes: A Randomized Clinical Trial. *The Iranian Journal of Obstetrics, Gynecology and Infertility*. 2018;21(5):33-46.
42. Kamali Z, Abedian Z, Mohammad A, Mohebbi-Dehnavi Z. The Effect of Small-Group Teaching on Health Literacy in Pregnant Females with Nausea and Vomiting: A Clinical Trial. *Journal of Nursing Education* 2018;6 (6): 25-32.
43. Kohan S, Ghasemi S, Dodangeh M. Associations between maternal health literacy and prenatal care and pregnancy outcome. 2008;12(4):146-52.
44. Ghanbari S, Majlessi F, Ghaffari M, Mahmoodi Majdabadi M. Evaluation of health literacy of pregnant women in urban health centers of Shahid Beheshti Medical University. *Daneshvar*. 2012;19(97):1-12.

Figures




Figure 1

Consort flow diagram