

Globulin, albumin to globulin ratio, and plasma fibrinogen may be potential diagnostic biomarkers for tibial infected nonunion

Guangyue Xu (✉ xugy0813@163.com)

Nanjing Drum Tower Hospital, The Affiliated Hospital of Nanjing University Medical School

Haijun Mao

Nanjing Drum Tower Hospital, The Affiliated Hospital of Nanjing University Medical School

Zhen Wang

Nanjing Drum Tower Hospital, The Affiliated Hospital of Nanjing University Medical School

Research Article

Keywords: globulin, albumin to globulin ratio, plasma fibrinogen, tibial infected nonunion, diagnostic test

Posted Date: April 13th, 2022

DOI: <https://doi.org/10.21203/rs.3.rs-1514230/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Aims: The accurate preoperative diagnosis of infected nonunion remains challenging. Here, we evaluated the diagnostic potential of novel biomarkers for tibial infected nonunion.

Methods: A single-center retrospective study was conducted on 155 patients divided into two groups: 59 with tibial infected nonunion (Group A) and 96 with tibial aseptic nonunion (Group B). Preoperatively analyzed clinical parameters included plasma D-dimer, plasma fibrinogen, albumin (ALB), globulin (GLB), albumin-to-globulin ratio (AGR), erythrocyte sedimentation rate (ESR), C-reactive protein (CRP), and white blood cell (WBC) count. Receiver operating characteristic (ROC) curves, sensitivity, and specificity were utilized to compare the diagnostic potential of these biomarkers.

Results: The WBC, ESR, CRP, plasma D-dimer, plasma fibrinogen, and GLB levels of Group A were significantly higher ($P < 0.05$) than in Group B. The ALB and AGR levels of Group A were significantly lower ($P < 0.05$) than in Group B. The area under the curve (AUC) of ESR, plasma fibrinogen, and AGR were greater than 0.8. An analysis of a combination of ESR, plasma fibrinogen, GLB, and AGR had the highest AUC. In patients with comorbidities, the diagnostic accuracy of ESR, plasma fibrinogen, GLB, and AGR also performed well. Higher levels of WBC, CRP, and ESR were detected in patients who had recently used antibiotics ($P < 0.05$).

Conclusion: GLB, AGR, and plasma fibrinogen are promising biomarkers for improving the diagnosis of tibial infected nonunion. The integrated analysis of ESR, plasma fibrinogen, GLB, and AGR provided more accurate and more specific diagnosis than the four biomarkers individually.

Introduction

Postoperative tibial infected nonunion is a chronic and debilitating disorder associated with that high hospitalization costs, lengthy treatment, and increased risk of patient morbidity and mortality [1–4]. It is often difficult to discriminate between infected nonunion and aseptic nonunion [5, 6]. If infected nonunion is not promptly diagnosed or managed, it can result in devastating consequences, including permanent loss of function, amputation, or death [7–9]. Therefore, the timely and accurate diagnosis of infected nonunion is critical to facilitate any necessary reoperation, which can substantially affect treatment plans [10, 11].

Blood-based assays provide a potential solution for diagnosing infected nonunion, because they can be rapidly performed in advance of intraoperative culture [12]. Traditional biomarkers, like white blood cell count (WBC), erythrocyte sedimentation rate (ESR), and C-reactive protein (CRP) are important biomarkers in the diagnostic evaluation of inflammation, and can reduce the rate of missed infections [8, 13]. These biomarkers can be widely used to diagnose acute fracture-related infections, but sometimes remain normal in late or chronic infections caused by low-virulence pathogens [14]. Therefore, new biomarkers need to be evaluated for the diagnosis of tibial infected nonunion.

Albumin (ALB) is one of the main components of serum proteins, and is associated with surgical site infection in orthopedics [15, 16]. Globulin (GLB), which is a component of complements and ceruloplasmin, generally increases during the inflammatory process [17]. Hence, the albumin-to-globulin ratio (AGR), which considers both ALB and GLB levels, is also a promising biomarker for inflammation [18]. Furthermore, several studies reported that GLB and AGR were informative biomarkers in the diagnosis of periprosthetic joint infection (PJI) [19, 20]. In addition, serum D-dimer and plasma fibrinogen may be novel biomarkers to detect infected nonunion, and they have shown good performance in past studies [21, 22]. The tibia is the most commonly involved bone in infected nonunion of open fractures [23, 24]. However, the accuracy of these blood biomarkers in diagnosing infected nonunion of tibia fracture after surgery is unknown.

In the present study, we evaluated the diagnostic accuracy of ALB, GLB, AGR, plasma D-dimer, and plasma fibrinogen in tibial infected nonunion. We hypothesize that these new blood biomarkers will be associated with tibial infected nonunion fractures after surgery, and will therefore help guide orthopedic surgeons in the assessment of patients with tibial infected nonunion.

Methods

Study design and participants.

This single-center retrospective study was conducted in a tertiary hospital providing specialist treatment to patients with musculoskeletal infections. Over a period of approximately 3 years (June 2018 to July 2021), a total of 211 patients with tibial fracture who underwent surgery for nonunion after failed fracture operation were eligible for inclusion in this study. Ethical approval was obtained from the Clinical Research Ethics Committee of The Affiliated Drum Tower Hospital of Nanjing University Medical School. The methods were carried out in accordance with the relevant guidelines and regulations. Patients were excluded based on the following exclusion criteria: 1) age < 18 years (n = 3); 2) without complete blood workup or medical records (n = 7); 3) unavailability of complete examinations (n = 4). Patients were also divided into subgroups for analysis to avoid bias: concurrent infections in other organs (n = 8), venous thrombosis (n = 13), malignancies (n = 2), autoimmune diseases (n = 4), and recent use of antibiotics 2 weeks before operation (n = 15). A nonunion was determined as radiographic evidence of nonprogression of healing for at least 3 months, or lack of healing by 9 months since the initial injury [25]. According to the fracture-related infections definition criteria, multiple gross tissue specimens (≥ 5 samples) were cultured, when the same organism was grown in at least 2 cultures of the intraoperative sample, a positive diagnosis of infection was made [8]. Ultimately, the patients with tibial nonunion were separated into two groups: 59 in Group A (operation for infected tibial nonunion) and 96 in Group B (operation for aseptic tibial nonunion) (Fig. 1).

Demographic features and blood biomarkers.

The clinical features of patients with tibial nonunion were comprehensively interpreted by the attending physician after admission. Baseline demographic features, including age, gender, BMI, smoking, and fracture position were collected. Fasting venous blood samples were obtained from patients on admission and were sent to the clinical laboratory within 2 h. These blood samples were used to test WBC, CRP, ESR, liver function tests (including ALB, GLB, AGR), and plasma coagulation examinations (including plasma D-dimer, and plasma fibrinogen). Antimicrobial treatment was stopped for at least 14 days until intraoperative specimens were collected, unless the patients needed anti-infective therapy urgently; those patients were evaluated separately.

Statistical analysis.

Descriptive statistics are presented as mean \pm SD (standard deviation) for normally distributed continuous variables or median (IQR) for non-normally distributed data, and frequency (percentage) for categorical variables. Kolmogorov-Smirnov (K-S) test was used to identify data as normally distributed variables or non-normally distributed data. Student's t-tests were used to analyze normal distributed numerical variables, and Mann-Whitney U test was used to analyze numerical variables with non-normal distribution or unequal variance. Pearson χ^2 test or Fisher exact test were used to analyze qualitative variables. Differences with a p-value of $P < 0.05$ are considered to be significant. Receiver operating characteristic (ROC) curves were plotted to determine the diagnostic value of each biomarker for assessing tibial infected nonunion, and the area under the curve (AUC) and 95% confidence interval (CI) were calculated to compare different biomarkers. The Youden index ($J = [\text{sensitivity} + \text{specificity}] - 1$) was used to determine the optimal predictive cut-offs for calculating AUC. The sensitivity, specificity, positive predictive value (PPV), and negative predictive value (NPV) of each test were calculated. The AUC values were determined to be excellent (0.900 to 1.000), good (0.800 to 0.899), fair (0.700 to 0.799), poor (0.600 to 0.699), and having no discriminatory capacity (< 0.599). All statistical analyses were performed using STATA version 18.0 (SPSS Inc., Chicago, IL, USA).

Results

The basic demographic data of the two groups are shown in Table I. There were no significant differences between the two groups in terms of age, gender, BMI, smoking history, or fracture position. ($P > 0.05$).

Group A had significantly higher WBC, CRP, ESR, plasma D-dimer, plasma fibrinogen, and GLB levels than Group B ($P < 0.05$; Table II). Nonetheless, the ALB and AGR levels of patients in Group A were significantly lower than those of Group B ($P < 0.05$). Table III shows the AUCs, Youden index values, optimal cutoff values, and predictive values (sensitivity, specificity, PPV, NPV) of the various biomarkers. Based on ROC curve analysis (Fig. 2), the diagnostic performance of ESR (AUC = 0.883), plasma fibrinogen (AUC = 0.819), AGR (AUC = 0.815), and GLB (AUC = 0.805) were good. Moreover, the AUC of a composite biomarker combining ESR with plasma fibrinogen, GLB, and AGR (0.926) had the highest AUC. However,

other biomarkers' diagnostic accuracy were fair or poor, including WBC, CRP, plasma D-dimer, and ALB. At the optimal cutoff value, the sensitivity, specificity, PPV, and NPV were 38.98, 84.38, 60.53, and 69.23% for WBC; 69.49, 77.08, 65.08, and 80.43% for CRP; 71.19, 90.63, 82.35, and 83.65% for ESR; 64.41, 83.33, 70.37, and 79.21% for plasma D-dimer; 72.88, 75.00, 64.18, and 81.82% for plasma fibrinogen; 88.14, 9.38, 37.41, and 56.25% for ALB; 55.93, 96.88, 91.67, and 78.15% for GLB; and 57.63, 92.71, 82.93, and 78.07% for AGR; and 76.27, 97.92, 95.75, and 87.04% for the combination of ESR with plasma fibrinogen, GLB, and AGR.

Based on the optimal threshold, table IV shows that the diagnostic performance of tested markers for patients with comorbidities, including concurrent infections in other organs, venous thrombosis, autoimmune diseases, and malignancies. The range of the diagnostic accuracy of ESR, plasma fibrinogen, and AGR was 60–100% in patients with these comorbidities. The overall diagnostic accuracies were good.

Table V shows that there were significant differences between the cases of infected tibial nonunion patients who had recently used and had not used antibiotics in terms of WBC, CRP, and ESR ($P < 0.05$). However, the levels of plasma D-dimer, plasma fibrinogen, ALB, GLB, and AGR were not significantly different between the two groups ($P > 0.05$).

Discussion

This study was carried to compare the accuracy of plasma D-dimer, plasma fibrinogen, ALB, GLB, and AGR with traditional inflammatory biomarkers (WBC, CRP, and ESR) in diagnosing tibial infected nonunion. Our data demonstrated that the blood-based biomarkers plasma fibrinogen and GLB were associated with infected tibial nonunion, and could be used as new biomarkers to diagnose infected tibial nonunion. Furthermore, the integrated analysis of four promising biomarkers (ESR, plasma fibrinogen, GLB, and AGR) provided more accurate and more specific evaluation of tibial infected nonunion than the four biomarkers individually.

ALB and GLB are two easily accessible and reliable biomarkers evaluated in liver function tests that are routinely performed before surgery, and are able to be performed rapidly and economically. ALB is a negative acute phase protein, and is considered to be a biomarker of inflammation and nutritional status [26–28]. A meta-analysis conducted by Yuwen *et al.*[15] reported that ALB levels < 3.5 g/dL was associated with an almost 2.5 fold increased risk of surgical site infection (SSI) in orthopaedics. However, Wang *et al.*[20] reported that ALB was not useful for diagnosis of patients suspected of having PJI. Similarly, Ye *et al.*[19] found that the levels of ALB were not associated with PJI. In our study, the diagnostic accuracy of ALB in tibial infected nonunion was poor, with an AUC of 0.671, sensitivity of 88.14%, specificity of 9.38%, PPV of 37.41%, and NPV of 56.25%. GLB is another major serum protein component that consists of antibodies and inflammatory cytokines, including complements, interleukin-6, and immunoglobulins [29, 30]. There is an inverse relationship between GLB and ALB in response to inflammation and infection, and the AGR could indicate an inflammatory state more accurately. GLB and

AGR have been validated to have potential roles in the pathogenesis of inflammatory and infectious diseases in various studies [31, 32]. Moreover, recent studies have demonstrated that GLB and AGR are associated with PJI and may serve as potential adjuvant biomarkers in the diagnosis of PJI [19, 20]. In accordance with these results, we also observed that high globulin levels and low AGR were independently associated with the risk of tibial infected nonunion. ROC curve analysis revealed that globulin and AGR showed acceptable predictive value for the diagnosis of tibial infected nonunion. The diagnostic value of GLB (AUC = 0.805) was slightly lower than that of AGR (0.815), which were higher than 0.8. Both biomarkers demonstrated fair sensitivity (GLB 55.93%, AGR 57.63%), but high specificity (GLB 96.88%, AGR 92.71%), reducing the misdiagnosis rate of tibial infected nonunion. Moreover, the PPV of GLB (91.67%) was the highest; suggesting that GLB may be superior as a single biomarker in predicting the diagnosis of tibial infected nonunion.

Recently, it has been demonstrated that systemic and local infections result in fibrinolytic activity [33, 34]. Moreover, coagulation-related indicators, including D-dimer and fibrinogen, are useful diagnostic markers for PJI [35–39]. Serum D-dimer and plasma fibrinogen have performed well in the diagnosis of infected nonunion [21, 22]. In our study, plasma fibrinogen (AUC = 0.819) demonstrated good diagnostic performance in infected nonunion, similar to the results of another recent study [21]. These data suggest that plasma fibrinogen may be a novel biomarker to diagnose tibial infected nonunion. However, the diagnostic value of plasma D-dimer was limited in our study, with a fair AUC of 0.776. Conversely, the D-dimer, assayed in serum rather than plasma, has proven to be useful for preoperative prediction of infected nonunion after ORIF [22]. Some studies demonstrated that serum D-dimer is superior to plasma D-dimer in the diagnosis of PJI [40, 41]. Consequently, high-quality prospective studies that address these research gaps are needed to validate the use of D-dimer as a biomarker for tibial infected nonunion.

WBC, CRP, and ESR are the most commonly used biomarkers of tibial infected nonunion. Unfortunately, they are usually affected by other factors such as physiological stress, treatment, and other diseases [5, 42]. Peripheral WBC is frequently normal in low-grade infections and affords little diagnostic help. The diagnostic value of WBC was limited in our study, with a fair AUC of 0.599. The standard WBC biomarker may not provide enough information to clearly distinguish between infected nonunion and aseptic nonunion. On the contrary, ESR had good performance in diagnosis of suspicious infection with higher sensitivity, which is verified by past studies [43, 44].

In subgroup analysis, we found that the diagnostic accuracy of these novel biomarkers was good in patients with tibial infected nonunion who also had other comorbidities. Considering the small number of cases, a larger number of cases should be enrolled into evaluate the diagnostic accuracies of these biomarkers and reduce potential bias. For example, recent use of antibiotics might influence inflammatory biomarkers. Our subgroup analysis showed that there were significant differences between patients with infected nonunion who recently used antibiotics vs. those who had not in a number of biomarkers, including WBC, CRP, and ESR levels ($P < 0.05$); other biomarkers were not significantly different, including plasma D-dimer, plasma fibrinogen, ALB, GLB, and AGR ($P > 0.05$). Therefore, our data

suggest that plasma fibrinogen, GLB, and AGR may be better biomarkers to diagnose patients with suspected infected nonunion who have recently used antibiotics.

There are several limitations to our study. Firstly, this study was retrospective, with inherent biases. Secondly, the sample size of this study is fairly small. Finally, the different comorbidities and application of antibiotics among the patients may have influenced the observed results of the diagnostic accuracy of the biomarkers in patients with infected nonunion. Therefore, multicenter, prospective, comparative studies with larger samples are required to more thoroughly determine the accuracy of these biomarkers for predicting tibial infected nonunion.

It was a secondary analysis that recursive partitioning analysis of the inflammatory marker predictive values for infected nonunion, and is limited by the relatively small sample size of the cohort included. In order to avoid external validity, we will calibrate and externally validate this model with a future large observational cohort of patients with suspected nonunion to inform clinical utility.

In conclusion, our results show that GLB, AGR, and plasma fibrinogen may be reliable biomarkers to screen for tibial infected nonunion. The integrated analysis of four biomarkers (ESR, plasma fibrinogen, GLB, and AGR) may provide more accurate and more specific evaluation of tibial infected nonunion. Therefore, we recommend that surgeons begin to evaluate GLB, AGR, plasma fibrinogen, and the combination of ESR, plasma fibrinogen, GLB, and AGR as a potential screening tool for patients who might be at risk for tibial infected nonunion.

Abbreviations

AGR: albumin-to-globulin ratio; AUC: area under the curve; BMI: body mass index; CI: confidence interval; CRP: C-reactive protein; ESR: erythrocyte sedimentation rate; IQR: interquartile range; NPV: negative predictive value; ORIF: open reduction and internal fixation; PJI: periprosthetic joint infection; PPV: positive predictive value; ROC: receiver operating characteristic; TKA: total knee arthroplasty; WBC: white blood cell.

Declarations

Acknowledgements

We highly acknowledge the contribution by all the doctors and nurses in our department.

Author contributions

Conception and design were the responsibility of H.M, Z.W and G.X. Data collection was the responsibility of H.M and Z.W. Statistical analysis was the responsibility of H.M and Z.W. Analysis and interpretation were the responsibility of H.M and Z.W. Composition of the manuscript was the responsibility of H.M and Z.W. Revision of the manuscript for important intellectual content and approval of the final draft were

done by all authors. G.X took responsibility for the paper as a whole. All authors read and approved the final manuscript.

Funding

This study received financial support from Key Project supported by Medical Science and Technology Development Foundation, Nanjing Department of Health, Jiangsu, China (ZKX16034).

Availability of data and materials

The final dataset will be available from the corresponding author.

Declarations

Ethics approval and consent to participate

This study was approved by the ethics committee of Nanjing Drum Tower Hospital, The Affiliated Hospital of Nanjing University Medical School, China (approval no.2021–135). All patients signed an informed consent form approved by the Institutional Review Board.

Consent for publication

Not applicable

Competing Interests

All the authors declare that there is no conflict of interest

References

1. Chaudhary MM: **Infected nonunion of tibia.** *Indian journal of orthopaedics*2017, **51**(3):256-268.
2. Motsitsi NS: **Management of infected nonunion of long bones: the last decade (1996-2006).** *Injury*2008, **39**(2):155-160.
3. Berkes M, Obrebsky WT, Scannell B, Ellington JK, Hymes RA, Bosse M: **Maintenance of hardware after early postoperative infection following fracture internal fixation.** *The Journal of bone and joint surgery American volume*2010, **92**(4):823-828.
4. Shen J, Sun D, Fu J, Wang S, Wang X, Xie Z: **Management of surgical site infection post-open reduction and internal fixation for tibial plateau fractures.** *Bone & joint research*2021, **10**(7):380-387.
5. Wang S, Yin P, Quan C, Khan K, Wang G, Wang L, Cui L, Zhang L, Zhang L, Tang P: **Evaluating the Use of Serum Inflammatory Markers for Preoperative Diagnosis of Infection in Patients with Nonunions.** *BioMed research international*2017, **2017**:9146317.
6. Olszewski D, Streubel PN, Stucken C, Ricci WM, Hoffmann MF, Jones CB, Sietsema DL, Tornetta P, 3rd: **Fate of Patients With a "Surprise" Positive Culture After Nonunion Surgery.** *Journal of*

*orthopaedic trauma*2016, **30**(1):e19-23.

7. Bonneville P: **Operative treatment of early infection after internal fixation of limb fractures (exclusive of severe open fractures)**. *Orthopaedics & traumatology, surgery & research : OTSR*2017, **103**(1s):S67-s73.
8. Metsemakers WJ, Morgenstern M, McNally MA, Moriarty TF, McFadyen I, Scarborough M, Athanasou NA, Ochsner PE, Kuehl R, Raschke Met al: **Fracture-related infection: A consensus on definition from an international expert group**. *Injury*2018, **49**(3):505-510.
9. Walter N, Rupp M, Hierl K, Pfeifer C, Kerschbaum M, Hinterberger T, Alt V: **Long-term patient-related quality of life after fracture-related infections of the long bones**. *Bone & joint research*2021, **10**(5):321-327.
10. McNally MA: **Decision-making in infected nonunion: is the surgery more important than the condition?** *The bone & joint journal*2016, **98-b**(4):435-436.
11. Patzakis MJ, Zalavras CG: **Chronic posttraumatic osteomyelitis and infected nonunion of the tibia: current management concepts**. *The Journal of the American Academy of Orthopaedic Surgeons*2005, **13**(6):417-427.
12. Morgenstern M, Kuehl R, Eckardt H, Acklin Y, Stanic B, Garcia M, Baumhoer D, Metsemakers WJ: **Diagnostic challenges and future perspectives in fracture-related infection**. *Injury*2018, **49** Suppl 1:S83-s90.
13. Bishop JA, Palanca AA, Bellino MJ, Lowenberg DW: **Assessment of compromised fracture healing**. *The Journal of the American Academy of Orthopaedic Surgeons*2012, **20**(5):273-282.
14. van den Kieboom J, Bosch P, Plate JDJ, FFA IJ, Kuehl R, McNally MA, Metsemakers WJ, Govaert GAM: **Diagnostic accuracy of serum inflammatory markers in late fracture-related infection: a systematic review and meta-analysis**. *The bone & joint journal*2018, **100-b**(12):1542-1550.
15. Yuwen P, Chen W, Lv H, Feng C, Li Y, Zhang T, Hu P, Guo J, Tian Y, Liu Let al: **Albumin and surgical site infection risk in orthopaedics: a meta-analysis**. *BMC surgery*2017, **17**(1):7.
16. Yang G, Zhu Y, Zhang Y: **Prognostic risk factors of surgical site infection after primary joint arthroplasty: A retrospective cohort study**. *Medicine*2020, **99**(8):e19283.
17. Li K, Fu W, Bo Y, Zhu Y: **Effect of albumin-globulin score and albumin to globulin ratio on survival in patients with heart failure: a retrospective cohort study in China**. *BMJ open*2018, **8**(7):e022960.
18. Ukibe NR, Ndiuwem CK, Ogbu, II, Ukibe SN, Ehiaghe FA, Ikimi CG: **Prognostic value of some serum protein fractions as Early Index of Clinical Recovery in Pulmonary Tuberculosis subjects**. *The Indian journal of tuberculosis*2020, **67**(2):167-171.
19. Ye Y, Chen W, Gu M, Xian G, Pan B, Zheng L, Zhang Z, Sheng P: **Serum globulin and albumin to globulin ratio as potential diagnostic biomarkers for periprosthetic joint infection: a retrospective review**. *Journal of orthopaedic surgery and research*2020, **15**(1):459.
20. Wang H, Zhou H, Jiang R, Qian Z, Wang F, Cao L: **Globulin, the albumin-to-globulin ratio, and fibrinogen perform well in the diagnosis of Periprosthetic joint infection**. *BMC musculoskeletal disorders*2021, **22**(1):583.

21. Wang XJ, Wang Z, Zhang ZT, Qiu XS, Chen M, Chen YX: **Plasma Fibrinogen as a Diagnostic Marker of Infection in Patients with Nonunions.** *Infection and drug resistance*2020, **13**:4003-4008.
22. Wang Z, Zheng C, Wen S, Wang J, Zhang Z, Qiu X, Chen Y: **Usefulness of serum D-dimer for preoperative diagnosis of infected nonunion after open reduction and internal fixation.** *Infection and drug resistance*2019, **12**:1827-1831.
23. Bhowmick K, Boopalan P, Gunasekeran C, Livingston A, Jepegnanam TS: **Management of Chronic Infected Intra-Articular Fractures of the Proximal Tibia with Ilizarov Ring Fixation.** *The journal of knee surgery*2020, **33**(2):213-222.
24. Thakore RV, Francois EL, Nwosu SK, Attum B, Whiting PS, Siuta MA, Benvenuti MA, Smith AK, Shen MS, Mousavi *et al.*: **The Gustilo-Anderson classification system as predictor of nonunion and infection in open tibia fractures.** *European journal of trauma and emergency surgery : official publication of the European Trauma Society*2017, **43**(5):651-656.
25. Bell A, Templeman D, Weinlein JC: **Nonunion of the Femur and Tibia: An Update.** *The Orthopedic clinics of North America*2016, **47**(2):365-375.
26. Don BR, Kaysen G: **Serum albumin: relationship to inflammation and nutrition.** *Seminars in dialysis*2004, **17**(6):432-437.
27. Eckart A, Struja T, Kutz A, Baumgartner A, Baumgartner T, Zurfluh S, Neeser O, Huber A, Stanga Z, Mueller *Bet et al.*: **Relationship of Nutritional Status, Inflammation, and Serum Albumin Levels During Acute Illness: A Prospective Study.** *The American journal of medicine*2020, **133**(6):713-722.e717.
28. Deng S, Gao J, Zhao Z, Tian M, Li Y, Gong Y: **Albumin/Procalcitonin Ratio Is a Sensitive Early Marker of Nosocomial Blood Stream Infection in Patients with Intra-Cerebral Hemorrhage.** *Surgical infections*2019, **20**(8):643-649.
29. Zhou T, Yu ST, Chen WZ, Xie R, Yu JC: **Pretreatment albumin globulin ratio has a superior prognostic value in laryngeal squamous cell carcinoma patients: a comparison study.** *Journal of Cancer*2019, **10**(3):594-601.
30. Wu PP, Hsieh YP, Kor CT, Chiu PF: **Association between Albumin-Globulin Ratio and Mortality in Patients with Chronic Kidney Disease.** *Journal of clinical medicine*2019, **8**(11).
31. Hoffman LK, Ghias MH, Cohen SR, Lowes MA: **Polyclonal hyperglobulinaemia and elevated acute-phase reactants in hidradenitis suppurativa.** *The British journal of dermatology*2018, **178**(2):e134-e135.
32. Schmilovitz-Weiss H, Cohen M, Pappo O, Sulkes J, Braun M, Tur-Kaspa R, Ben-Ari Z: **Serum globulin levels in predicting the extent of hepatic fibrosis in patients with recurrent post-transplant hepatitis C infection.** *Clinical transplantation*2007, **21**(3):391-397.
33. Stark K, Massberg S: **Interplay between inflammation and thrombosis in cardiovascular pathology.** *Nature reviews Cardiology*2021, **18**(9):666-682.
34. Han H, Yang L, Liu R, Liu F, Wu KL, Li J, Liu XH, Zhu CL: **Prominent changes in blood coagulation of patients with SARS-CoV-2 infection.** *Clinical chemistry and laboratory medicine*2020, **58**(7):1116-1120.

35. Xu H, Shang G, Wang Y, Xiang S: **Plasma fibrinogen is a reliable marker for diagnosing periprosthetic joint infection and determining the timing of second-stage revision.** *International journal of infectious diseases : IJID : official publication of the International Society for Infectious Diseases*2021, **108**:220-225.
36. Yang F, Zhao C, Huang R, Ma H, Wang X, Wang G, Zhao X: **Plasma fibrinogen in the diagnosis of periprosthetic joint infection.** *Scientific reports*2021, **11**(1):677.
37. Wu H, Meng Z, Pan L, Liu H, Yang X, Yongping C: **Plasma Fibrinogen Performs Better Than Plasma d-Dimer and Fibrin Degradation Product in the Diagnosis of Periprosthetic Joint Infection and Determination of Reimplantation Timing.** *The Journal of arthroplasty*2020, **35**(8):2230-2236.
38. Hu Q, Fu Y, Tang L: **Serum D-dimer as a diagnostic index of PJI and retrospective analysis of etiology in patients with PJI.** *Clinica chimica acta; international journal of clinical chemistry*2020, **506**:67-71.
39. Grzelecki D, Walczak P, Grajek A, Szostek M, Dudek P, Bartosz P, Olewnik Ł, Czubak-Wrzosek M, Marczak D, Tyrakowski M: **Elevated plasma D-dimer concentration has higher efficacy for the diagnosis of periprosthetic joint infection of the knee than of the hip-A single-center, retrospective study.** *Journal of orthopaedic research : official publication of the Orthopaedic Research Society*2021, **39**(2):291-298.
40. Li C, Margaryan D, Ojeda-Thies C, Perka C, Trampuz A: **Meta-analysis of serum and/or plasma D-dimer in the diagnosis of periprosthetic joint infection.** *Journal of orthopaedic surgery and research*2020, **15**(1):298.
41. Xu H, Xie JW, Yang JL, Huang ZY, Pei FX: **Role of D-dimer and Fibrinogen in the Diagnosis of Periprosthetic Joint Infection: A Systematic Review and Meta-Analysis.** *Orthopaedic surgery*2021, **13**(3):692-700.
42. Metsemakers WJ, Kuehl R, Moriarty TF, Richards RG, Verhofstad MHJ, Borens O, Kates S, Morgenstern M: **Infection after fracture fixation: Current surgical and microbiological concepts.** *Injury*2018, **49**(3):511-522.
43. Vasso M, Schiavone Panni A: **Low-grade periprosthetic knee infection: diagnosis and management.** *Journal of orthopaedics and traumatology : official journal of the Italian Society of Orthopaedics and Traumatology*2015, **16**(1):1-7.
44. Garvin KL, Konigsberg BS: **Infection following total knee arthroplasty: prevention and management.** *The Journal of bone and joint surgery American volume*2011, **93**(12):1167-1175.

Tables

Table I

Patient characteristics for the two groups

	Group A (n = 59)	Group B (n = 96)	P-value
Number of women	7/52 (11.9%)	20/76 (20.8%)	0.193†
Age (year, mean ± SD)	47.8 ± 15.2	45.7 ± 13.0	0.381#
BMI (kg/m ² , mean ± SD)	25.1 ± 3.6	24.8 ± 3.2	0.487#
Smoking history	12/47 (21.1%)	18/78 (17.3%)	0.836†
Position (left)	30/29 (50.8%)	53/43 (55.2%)	0.662†
Fracture position (actual sites)			0.476†
Proximal	8/51 (13.6%)	19/77 (19.8%)	
Mid-diaphysis	30/29 (50.8%)	50/46 (52.1%)	
Distal	21/38 (35.6%)	27/69 (28.1%)	
<p>Notes: Group A = tibial infected nonunion; Group B = tibial aseptic nonunion; # Independent-samples <i>t</i>-test; † Chi-squared test (linear by linear); *P < 0.05 indicates statistical significance.</p> <p>Abbreviation: BMI = body mass index.</p>			

Table II

Comparison of the tested markers in the two groups

	Group A (n = 59)	Group B (n = 96)	P-value
WBC (10 ⁹ /μL)			
Median	6.6	6.2	0.038*
P25, P75	5.5 ~ 8.2	5.4 ~ 7.2	
CRP (mg/L)			<0.001*
Median	7.0	3.5	
P25, P75	3.8~ 17.5	2.5 ~ 4.9	
ESR (mm/hr)			<0.001*
Median	25.0	6.5	
P25, P75	13.0 ~ 39.0	5.0 ~ 11.0	
Plasma D-dimer (mg/L)			<0.001*
Median	1.19	0.34	
P25, P75	0.53 ~ 2.9	0.17 ~ 0.75	
Plasma fibrinogen (mg/L)			<0.001*
Median	3.4	2.4	
P25, P75	2.8 ~ 5.1	2.1 ~ 2.9	
ALB (g/L)			<0.001*
Median	38.9	41.2	
P25, P75	37.7 ~ 41.1	39.2 ~ 43.1	
GLB (g/L)			<0.001*
Median	32.2	24.2	
P25, P75	26.4 ~ 39.2	21.1 ~ 27.2	
AGR			<0.001*
Median	1.30	1.71	
P25, P75	0.98 ~ 1.56	1.48 ~ 1.91	

Notes: Group A = tibial infected nonunion; Group B = tibial aseptic nonunion; All P-values calculated using Mann-Whitney U test; *P < 0.05 indicates statistical significance.

Abbreviation: WBC = white blood cell; CRP = C-reactive protein; ESR = erythrocyte sedimentation rate; ALB = albumin; GLB = globulin; AGR = albumin-to-globulin ratio

Table III

The diagnostic value of tested markers in patients with tibial infected nonunion.

Variables	AUC (95%CI)	Optimal Cutoff Value	Sensitivity (%)	Specificity(%)	PPV (%)	NPV (%)
WBC	0.599 (0.504-0.695)	7.65×10 ⁹ /L	38.98	84.37	60.53	69.23
CRP	0.748 (0.665-0.832)	4.95 mg/L	69.49	77.08	65.08	80.43
ESR	0.883 (0.830-0.937)	17.5 mm/h	71.19	90.63	82.35	83.65
Plasma D-dimer	0.776 (0.700-0.852)	0.95 mg/L	64.41	83.33	70.37	79.21
Plasma fibrinogen	0.819 (0.752-0.887)	2.85 g/L	72.88	75.00	64.18	81.82
ALB	0.671 (0.580-0.762)	44.45 g/L	88.14	9.38	37.41	56.25
GLB	0.805 (0.726-0.883)	31.5 g/L	55.93	96.88	91.67	78.15
AGR	0.815 (0.742-0.889)	1.35	57.63	92.71	82.93	78.07
ESR+ Plasma Fibrinogen +GLB+AGR	0.926 (0.880-0.972)	0.41	76.27	97.92	95.75	87.04

Abbreviation: WBC = white blood cell; CRP = C-reactive protein; ESR = erythrocyte sedimentation rate; ALB = albumin; GLB = globulin; AGR = albumin-to-globulin ratio; AUC = areas under the curve; CI = confidence interval; PPV = positive predictive value; NPV = negative predictive value.

Table IV

Diagnostic performance of tested markers for patients with comorbidities.

Comorbidities and Numbers		Concurrent infections in other organs (n = 8)	Venous thrombosis (n = 13)	Autoimmune diseases (n = 4)	Malignancies (n = 2)
ESR	Group A	7(6)	5(4)	1(1)	2(2)
	Group B	1(0)	8(8)	3(2)	0
Plasma fibrinogen	Group A	7(6)	5(2)	1(0)	2(2)
	Group B	1(0)	8(7)	3(3)	0
GLB	Group A	7(6)	5(4)	1(1)	2(2)
	Group B	1(1)	8(7)	3(3)	0
AGR	Group A	7(7)	5(3)	1(1)	2(2)
	Group B	1(1)	8(8)	3(3)	0
<p>Notes: Group A = tibial infected nonunion; Group B = tibial aseptic nonunion. Abbreviation: ESR = erythrocyte sedimentation rate; ALB = albumin; GLB = globulin; AGR = albumin-to-globulin ratio</p>					

Table V

Comparison of tested markers between patients with tibial infected nonunion who used antibiotics two weeks before and patients with tibial infected nonunion who had not used antibiotics recently

Variables	Normal range	Tibial Infected Nonunion Recent Use of Antibiotics (n = 15) (median, P25–75)	Tibial Infected Nonunion Recent No Use of Antibiotics (n = 59) (median, P25–75)	<i>P</i> -value
WBC($10^9/\mu\text{L}$)	3.5-9.5	5.6(5.1-6.4)	6.6 (5.5-8.2)	0.029*
CRP(mg/L)	0-8	4.4(2.8-6.1)	7.0(3.8-17.5)	0.035*
ESR(mm/hr)	0-15	15.0(12.0-18.0)	25.0(13.0-39.0)	0.006*
Plasma D-dimer (mg/L)	0-0.5	0.9(0.7-1.5)	1.2(0.5-2.9)	0.273
Plasma Fibrinogen(mg/L)	2-4	3.6(2.9-4.5)	3.4(2.8-5.1)	0.777
ALB (g/L)	40-55	39.4(37.9-41.8)	38.9(37.7-41.1)	0.727
GLB (g/L)	20-40	38.5(27.3-39.5)	32.2(26.4-39.2)	0.224
AGR	1.2-2.4	1.1(1.0-1.4)	1.3(1.0-1.6)	0.378
<p>Notes: All <i>P</i>-values calculated using the Mann-Whitney U test; *<i>P</i> < 0.05 indicates statistical significance.</p> <p>Abbreviation: WBC = white blood cell; CRP = C-reactive protein; ESR = erythrocyte sedimentation rate; ALB = albumin; GLB = globulin; AGR = albumin-to-globulin ratio.</p>				

Figures




Figure 1

Flow diagram of the study design.




Figure 2

The ROC curves of biomarkers in the diagnosis of tibial infected nonunion.

Notes: WBC white blood cell; CRP C-reactive protein; ESR erythrocyte sedimentation rate; ALB albumin; GLB globulin; AGR albumin-to-globulin ratio; ROC, receiver operating characteristic; AUC, areas under the curve.