

# Streamlining sporozoite isolation from mosquitoes by leveraging the dynamics of migration to the salivary glands

Ashutosh K. Pathak ([✉ ash1@uga.edu](mailto:ash1@uga.edu))

University of Georgia

Justine C. Shiao

University of Georgia

Blandine Franke-Fayard

Leiden University Medical Center

Lisa M. Shollenberger

Old Dominion University

Donald A. Harn

University of Georgia

Dennis E. Kyle

University of Georgia

Courtney C. Murdock

Cornell University

---

## Research Article

**Keywords:** Plasmodium, mosquitoes, oocysts, sporozoites, salivary glands, dissections, density dependence, EIP

**Posted Date:** April 4th, 2022

**DOI:** <https://doi.org/10.21203/rs.3.rs-1514945/v1>

**License:**  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---

**Version of Record:** A version of this preprint was published at Malaria Journal on September 13th, 2022.  
See the published version at <https://doi.org/10.1186/s12936-022-04270-y>.

## Abstract

## Background

Sporozoites isolated from the salivary glands of *Plasmodium*-infected mosquitoes are a prerequisite for several basic and pre-clinical applications. Although salivary glands are pooled to maximize sporozoite recovery, insufficient yields pose logistical and analytical hurdles; thus, predicting yields prior to isolation would be valuable. Preceding oocyst densities in the midgut is an obvious candidate, although it is unclear whether our current understanding of its relationship with sporozoite densities can be used to maximize yields.

## Methods

This study presents a retrospective analysis of mosquitoes infected with two strains of the rodent-specific *P. berghei*. Mean oocyst densities were estimated in the midguts earlier in the infection (11–15 days post-bloodmeal), with sporozoites pooled from the salivary glands later in the infection (17–29 days). Generalized linear mixed effects models were used to determine if 1) mean oocyst densities can predict sporozoite yields from pooled salivary glands, 2) whether these densities can capture differences in rates of sporozoite invasion of salivary glands, and 3) if the interaction between oocyst densities and time could be leveraged to boost overall yields.

## Results

The non-linear effect of mean oocyst densities confirmed the role of density-dependent constraints in limiting yields beyond certain oocyst densities. Irrespective of oocyst densities however, the continued invasion of salivary glands by the sporozoites boosted recoveries over time (17–29 days post-bloodmeal) for either parasite strain.

## Conclusions

Sporozoite invasion of the salivary glands over time can be leveraged to maximize yields for *P. berghei*. In general, however, invasion of the salivary glands over time is a critical fitness determinant for all *Plasmodium* species (extrinsic incubation period, EIP). Thus, delaying sporozoite collection could, in principle, substantially reduce dissection effort for any parasite within the genus, although our results also recommend closer examination of how changes in sporozoites densities over time modify parasite infectivity for the next host.

## Background

Preventing sporozoite establishment in the human liver has been a cornerstone of anti-malarial therapies for more than a century [1]. While the RTS/S vaccine underscored the potential for targeting the most devastating of human malaria, *Plasmodium falciparum*, its limited efficacy served as a reminder of the need for continual improvements to the vaccine design pipeline. Sporozoites isolated from the salivary glands of mosquitoes infected with human or rodent *Plasmodium* species offer several benefits that are either not possible, or difficult to assess with mosquito challenge models. Indeed, isolated sporozoites have served as starting points for identifying novel mechanisms of host-parasite interactions and developing pre-clinical assays for evaluating drugs and vaccines [2–12]. Developing approaches to maximize sporozoite yields from the salivary glands would help streamline existing workflows [13].

Despite the benefits of working with purified sporozoites, isolating sporozoites is a labor- and technique-intensive endeavor [2, 14]. The limited carrying capacity of salivary glands requires isolating sporozoites from large pools of salivary glands dissected from multiple mosquitoes [2–12]. However, low sporozoite yields present a hurdle to the design, execution, and interpretation of most assays. For instance, low yields may compel a researcher to reduce the number of study groups and/or replicates on an ad hoc basis at the cost of assay throughput and statistical power. Alternatively, yields could be fortified by dissecting more salivary glands, although low infection status and/or sporozoite densities in individuals may counter any net benefits gained from increasing dissection efforts. The ability to predict sporozoite yields *a priori* would be useful in overcoming these limitations. Following ingestion, parasites penetrate the mosquito midgut before establishing as oocysts, lining the epithelium under the basal lamina. Sporozoite replication is initiated in the oocysts from where they are released into the hemocoel before migrating to the salivary glands, ready for injection into another host. One commonly suggested predictor of sporozoite densities in the salivary glands is the density of oocysts in the midgut, where sporozoite replication occurs. Identifying a relationship between oocyst density and sporozoite yield would be beneficial as quantifying oocyst requires relatively little investment in labor and expertise [15, 16].

Previous studies investigating relationships between oocyst and sporozoite densities suggest the relationship is non-linear, with increasing oocyst densities generally associated with diminishing returns in sporozoite yields, likely due to density-dependent constraints on sporozoite production. These constraints could involve bottom-up and top-down processes limiting sporozoite production, such as increased competition for host resources, immune activation, or vector mortality [17–22]. Density-dependent effects on sporozoite replication could also have implications for the extrinsic incubation period (EIP) or the average time it takes for sporozoites to establish in the salivary glands. While differences in EIP are critical to *Plasmodium* fitness in general [23, 24], the potential for density-dependent asynchronicity in the dynamics of sporozoite migration is also supported by ecological theory, which documents similar density-dependent effects on the population growth rates of other organisms [25, 26]. In fact, changes in EIP with oocyst density might explain the lack of consensus in the literature regarding when (day post-bloodmeal) to recover *P. berghei* sporozoites, with times cited ranging from as early as 17 days to 29 days post-bloodmeal [2, 5, 17–19, 21, 27–38]. Thus, by accounting for shifts in the temporal dynamics of sporozoite invasion due to variation in oocyst burdens, we assumed oocyst densities would predict optimal sporozoite yields.

Since its initial demonstration [39], mosquitoes infected with the rodent-specific *P. berghei* constitute an important component of the vaccine and drug discovery pipeline, with sporozoite yields from the salivary glands a critical bottleneck for several downstream applications [2]. In a retrospective, statistical analysis of a dataset consisting of 46 groups of mosquitoes infected with either one of the two strains of *P. berghei*, we asked the following questions: 1) How does oocyst density affect overall sporozoite yields, 2) can oocyst density predict rates of sporozoite migration, and 3) can we use this interaction to optimize sporozoite yields? Oocyst densities were estimated once between 11–15 days post-bloodmeal from a subgroup of individuals' midguts, with sporozoites pooled from the salivary glands of another subgroup at one to four time points between 17 to 29 days post-bloodmeal (supplementary table 1). Sporozoite yields were then modeled with the (arithmetic) mean oocyst densities estimated earlier in the infection to determine whether yields were also dependent on time and/or parasite strain. Specifically, we hypothesized that as oocyst densities increase, delayed replication and/or migration of sporozoites will enhance sporozoite yields over time, before eventually saturating and/or declining for both parasite strains. We show that irrespective of parasite strain, sporozoite yields from pooled salivary glands were dependent on a non-linear relationship with the mean oocyst densities. In addition, sporozoite migration over the 17-29-day sampling period facilitated a linear increase in yields, albeit independent of the mean oocyst densities. Taken together, we show how sporozoite yields could be streamlined further by leveraging the dynamics of sporozoite migration. Finally, at the oocyst densities tested here, density-dependent migration may yet be the primary reason for increased sporozoite yields, although pooling salivary glands may only provide a qualitative overview of how sporozoite yields change over time; we suggest the rates of migration are likely to vary quantitatively between individuals, especially at oocyst densities where yields could be maximized.

## Methods

### Chemical and consumables

Unless stated otherwise, all chemicals and consumables were purchased from Thermo Scientific Inc. (Hampton, NY).

### Parasite strains

Mice were infected with one of the two parasite strains, *P. berghei* WT (clone ANKA, referred to hereon as PbANKA, kind gift from Dr. Evlina Angov, Walter Reed Army Institute of Research, Silver Spring, MD), or a transgenic strain of *P. berghei*. Transgenic parasites were generated in female OF1 mice (6–7 weeks; Charles River, NL) in accordance with the European Guideline 86/609/EEC and follow the FELASA (Federation of European Laboratory Animal Science Associations) guidelines and recommendations concerning laboratory animal welfare. Animal experiments performed in Leiden University Medical Center (LUMC, Leiden, The Netherlands) were approved by the Animal Experiments Committee of the Leiden University Medical Center (DEC 10099).

To generate the reporter line, GFP-Luc<sub>ama1-eef1a</sub>, the reference clone cl15cy1 of the *P. berghei* ANKA strain was used [40]. Two GFP-Luciferase expression cassettes were inserted into the neutral *p230p* gene locus (PBANKA\_0306000) using standard transfection technologies (Supplementary Fig. 1a) [40]. We generated DNA construct pL1308 that is based on two standard DNA constructs, pL1063 [41, 42] and pL1156 [43]. The *eef1a*-GFP-Luc expression cassette of pL1063 (Sacl fragment) was cloned into the Sacl site of pL1165, generating plasmid pL1308 that contains the two *gfp-luc* fusion genes under the control of the *eef1a* (PBANKA\_1133300) promoter or the *ama1* (PBANKA\_0915000) promoter, respectively, and contains the two 5' and 3' *p230p* target regions (TR). The construct was linearized using SacII restriction sites outside of the 5' and 3' *p230p* TR before transfection. After transfection (exp. 1052), transfected parasites were obtained by flow-sorting based on GFP expression as described [41, 42], followed by cloning using the method of limiting dilution [40], resulting in line GFP-Luc<sub>ama1-eef1a</sub> (line 1052cl1). Correct integration of DNA construct into the genome of GFP-Luc<sub>ama1-eef1a</sub> was confirmed by Southern analyses of Pulsed Field Gel (PFG)-separated chromosomes (Supplementary Fig. 1b) [40]. PFG-separated chromosomes were hybridized with the 3'utr *Pbdhfr/ts* probe [40].

## Mice infections

All procedures described herein were approved by the University of Georgia's Institutional Animal Care and Use Committee GA under Animal Use Protocol number A2016 06-010-Y1-A0 and A2020 01-013-Y2-A3. Three to four days before mosquito feeds, female C57BL/6 mice or Hsd:ICR(CD-1) mice (Envigo, Indianapolis, IN), aged 6–8 weeks, were injected intraperitoneally with 500 µl parasite suspended in sterile PBS with a density of  $5 \times 10^6$  or  $10^7$  parasites per mouse with either PbANKA or the GFP-luciferase expressing strain of *P. berghei* (GFP-Luc<sub>ama1-eef1a</sub>, referred to herein as PbGFP-LUC<sub>CON</sub>). Total parasitemia (trophozoites, schizonts, and gametocytes) was estimated starting two days post-infection with a Giemsa-stained smear of 1–2 µl of blood collected from the tail vein of each mouse [44]. Once total parasitemia had reached 2–6% (usually four days), each mouse was anesthetized with ~ 0.5 ml of 1.25% 2,2,2-Tribromoethanol (v/v, Avertin, Sigma-Aldrich) and placed atop mosquito cages.

### Mosquito infections with *P. berghei* infected mice

*An. stephensi* colonies were maintained as described previously, with adult mosquitoes maintained on a diet of 5% dextrose (w/v) and 0.05% para-aminobenzoic acid (w/v), a constant temperature of 27°C and relative humidity of 75–85%, on a 12h day/night cycle [44, 45]. Infections were performed with 3- to 7-day old, host-seeking female *An. stephensi* sorted into cages (17.5 cm L x 17.5 cm W x 17.5 cm H, BugDorm Inc., Taiwan). Approximately 24 hours prior to infections, mosquitoes were transferred to a 20 °C chamber (Percival Scientific Inc., Dallas, IA) and starved before being fed on anesthetized mice for 15–20 minutes.

## Measuring oocyst densities in the midguts and sporozoite isolations

For quantifying oocyst densities in the midguts, mosquitoes from each group were vacuum aspirated directly into 70% ethanol, washed twice with ice-cold PBS, and oocysts enumerated as described previously [44, 45]. Salivary glands from mosquitoes in each group were used to isolate sporozoites, as described previously, with modifications [46]. Mosquitoes were first transferred to 16–ounce paper cups and cold anesthetized at -20 °C for two to three minutes. Anesthetized mosquitoes were then transferred to a sterile petri dish (10 cm) maintained at -2 to -4 °C on a portable chill table (BioQuip Products, Rancho Dominguez, CA). Mosquitoes were surface sterilized with 70% ethanol before being dissected in chilled, sterile Schneider 2 (S2) insect cell culture media supplemented with 0.1% (v/v) fetal bovine serum. Using a tungsten micro-dissecting needle (0.25 mm diameter, 1 µm tip, Roboz Surgical Instrument Co., Gaithersburg, MD), salivary glands were pooled in a 1.5 ml microcentrifuge tube containing 100–500 µl of the dissection media; care was taken to ensure all six lobes were accounted for. Pooled glands were then disrupted with 5–10 passages through a sterile, 30G needle (Becton–Dickinson, Franklin Lakes, NJ) to release sporozoites [5]. Sporozoites were counted in a 10µL aliquot with a hemocytometer and total yields were expressed as follows.

$$Y = (m * 10) * v \quad (1)$$

Where  $Y$  denotes sporozoite yields,  $m$  indicates mean sporozoites counted from the four corner grids in a volume of 0.1µL, with a constant of '10' to further express 'm' per µL (i.e.,  $0.1 * 10 = 1\mu\text{L}$ ), and  $v$  indicates original volume of sporozoite suspension that the aliquot was drawn from (e.g., 500µL). To express yields per mosquito, the following equation was used.

$$y = Y/n \quad (2)$$

where  $y$  denotes sporozoite yields per mosquito,  $Y$  indicates total sporozoite yields (from Eq. 1 above), with  $n$  representing the number of salivary glands pooled initially.

## Data analyses and statistical modeling

The dataset used in the current study represents 46 groups of *P. berghei*-infected mosquitoes, of which 36 were infected with PbANKA and the remaining 10 with PbGFP-LUC<sub>CON</sub>. Each group comprised 70–200 mosquitoes, distinguished based on being exposed to the same mouse (i.e., the same source of infection). We assumed time of sampling to be replicated over the 46 independent groups (i.e., a cross-sectional/partially nested sampling schedule where, for instance, sporozoite yields at 26 days post-bloodmeal were replicated across groups 4, 5, 17, 20, 27, 38, and 45) (Supplementary table 1) [47, 48]. To accommodate the imbalanced nature of data collection, we tested our hypothesis using generalized linear mixed-effects models (GLMMs) [47–50], with slopes of sporozoite yields dependent on the fixed effects/predictors (averaged oocyst densities, time, and/or parasite strain), but with intercepts allowed to vary randomly among the 46 groups ("fixed slopes, random intercepts"). Specifying groups as a random effect allows for the possibility that the relationship between oocyst and sporozoite densities measured in different individuals (since a mosquito can either be sampled for oocysts or sporozoites [2]) may show

a greater correlation between mosquitoes that were exposed to the same source of infection (i.e., the same mouse) [18, 19, 21].

All data analyses and modeling were performed in RStudio [51], an integrated development environment for the open-source R package (version 4.1.0) [52], and associated packages. Graphical analyses were performed with the “ggplot2” package (version 3.3.5) [53] with the number of sporozoites recovered displayed as yields per mosquito (i.e.,  $y$  from Eq. 2 above, rounded to the nearest integer). Because pooling prevents the estimation of each individual’s contribution of sporozoites, this method of expression does not represent a “true” average [54]. GLMMs were specified as suggested by the “glmmTMB” package [50]. Dispersion characteristics of residuals were tested in the “DHARMA” package (version 0.4.3). Tables were prepared for presentation with the “sjPlot” package (version 2.8.9) [55].

To maintain consistency with studies that rely on pooled sporozoites [2, 5, 17–19, 21, 27–38], statistical modeling was performed with sporozoite yields per mosquito (i.e.,  $y$  from Eq. 2 above) specified as the dependent variable (model referred to herein as the ‘default’ model). However, to determine if differences in the number of salivary glands can alter model fit, we also built a model with total sporozoite yields expressed as the dependent variable (i.e.,  $Y$  from Eq. 1 above), but with the number of salivary glands pooled added as an offset ( $n$  from Eq. 2 above) (model referred to herein as the ‘offset’ model). For instance, if 243,000 sporozoites were recovered from 31 mosquitoes, the dependent variable ( $y$ ) in the ‘default’ model would be 7,839 (or the quotient of 243000 / 31, rounded to the nearest integer); the dependent variable in the ‘offset’ model would be 243,000, with the number of salivary glands dissected (i.e., 31) included as an ‘offset’ to account for differences in the number of salivary glands pooled between groups and/or timepoints [50]. Note that neither method of expression is based on true averages and thus, the ‘mean’ predicted by the statistical analysis should also not be considered a ‘true’ mean [55]. In addition, because of the distinct dependent variables, it was not possible to compare their fits with likelihood-based criteria [50].

Of the three fixed effects, parasite strain was coded as a categorical variable, while averaged oocyst densities and days post-bloodmeal were specified as continuous variables and centered and scaled with the mean and standard deviations respectively [45]. Averaged oocyst densities were estimated by deriving the arithmetic mean of oocyst densities in all midguts, irrespective of infection status (i.e., midguts with oocysts  $\geq 0$ ). Expressing oocyst densities as the arithmetic mean was in keeping with how sporozoite yields were expressed, which, despite not being a true average, is more similar to how the arithmetic means ( $y$  in Eq. 2 above) are expressed, compared to other representations of central tendency [54]. Although only oocyst-positive mosquitoes will contribute to the sporozoite pool, our rationale for including all midguts irrespective of infection status is because salivary glands are generally dissected from all available mosquitoes irrespective of infection status (e.g., [2]).

In general, both ‘default’ and ‘offset’ models included sporozoite yields as a linear function of averaged oocyst densities, days post-bloodmeal (i.e., time), and parasite strain as main effects, and up to a three-way interaction. While a clear interaction between the three predictors would indicate differences between

strains, a clear two-way interaction between oocyst densities and time on sporozoite density in the salivary glands would be indicative of density-dependence in sporozoite migration. To test the potential for saturation and decline in sporozoite yields due to mean oocyst densities and/or time (i.e., to model the hump-shaped effect on yields [45]), we also considered quadratic terms (or second-order polynomial) of the two predictors as additive (main) effects. However, in keeping with our objective of describing a simple predictive framework, automated stepwise elimination (based on likelihood-ratio tests and Akaike's information criteria, corrected for small sample sizes ( $\Delta\text{AICc}$ )) was used to identify the most parsimonious combination of predictors (predictors with  $p < 0.05$ ) from either the 'default' or 'offset' models described above (referred to herein as 'minimal' models) [50]; note that distinct dependent variables meant direct statistical comparisons (using likelihood-based criteria) could not be performed 'default' and 'offset' models (either full or minimal) [50]. However, to enable indirect comparison, all visual displays of model fits were based on the minimal versions of 'default' and 'offset' models, with predicted means and 95% confidence intervals (95%CI) expressed per mosquito; predictions were derived via *post-hoc* simulations, as specified in the "ggeffects" package (version 1.1.1) [55].

## Results

### Summary of data.

The results are based on data from 2,557 mosquitoes. Between 11 to 15 days post-bloodmeal, oocyst densities were quantified once from mosquitoes infected with PbANKA or PbGFP-LUC<sub>CON</sub> (Supplementary table 2). Starting 17 days post-bloodmeal, salivary glands from either a subsample or all remaining individuals were pooled to isolate sporozoites at one to four time points until 29 days post-bloodmeal (Fig. 1 and Supplementary table 1). In general, heterogeneity (or differences) in oocyst densities between individuals, represented by the standard deviations, increased linearly with the corresponding mean for the 46 groups of mosquitoes (Pearson's  $r$  (95%CI) = 0.88 (0.79–0.93),  $p < 0.001$ ) (Fig. 2).

### Sporozoite yields increase non-linearly with mean oocyst densities, and linearly over time.

The full model (Table 1) suggested sporozoite yields ( $y$  in Eq. 2) were driven by a non-linear relationship with the (arithmetic) mean of oocyst densities estimated earlier in the infection: the initial increase in yields ('linear',  $Z$ -value = 5.52,  $p < 0.001$ , Table 1) was eventually followed by a decline ('quadratic',  $Z$ -value = -2.22,  $p = 0.027$ , Table 1) (Fig. 1a). Although the model was unable to identify a clear interaction between mean oocyst densities and time ( $Z$ -value = -1.01,  $p = 0.312$ , Table 1), it suggested a relatively weaker (vs. oocyst densities), yet positive, linear effect of time on yields ( $Z$ -value = 2.62,  $p = 0.009$ , Table 1); in other words, at all mean oocyst densities tested here, the sporozoite migration dynamic resulted in higher sporozoite yields over time (Fig. 1b). Unlike the non-linear effect of mean oocyst densities ('quadratic'), the model suggested yields did not decline with time post-infection ('quadratic',  $Z$ -value = 0.49,  $p = 0.623$ , Table 1). Additionally, the influence of mean oocyst densities and time on sporozoite yields did not differ between parasite strains (PbANKA or PbGFP-LUC<sub>CON</sub>) (rows 6, 8, 9, and 10, Table 1).

While mean oocyst densities and time were able to explain 47.6% of the variation in the dataset (Marginal  $R^2$ , Table 1), allowing random variation among groups in the contribution of these predictors improved the model's account of the overall variation in the dataset (Conditional  $R^2 = 77.7\%$ , Table 1). Excluding group as a random effect significantly worsened model fit ( $\Delta AICc = + 13.2$ ). Finally, stepwise elimination of the least significant predictors from the full model ('Minimal model', Table 1) confirmed the non-linear effects of mean oocyst densities ('linear', Z-value = 6.17,  $p < 0.001$  and 'quadratic', Z-value = -2.83,  $p = 0.005$ ) and time of sporozoite collection (i.e., days post-bloodmeal) were sufficient to model the overall trends ('linear', Z-value = 2.73,  $p = 0.006$ ).

### Sporozoite yields increase with time, irrespective of how many salivary glands were pooled.

In general, both the 'default' (Table 1) and 'offset' models (Supplementary table 3) were consistent in highlighting the contribution of mean oocyst densities and time (rows 2, 3, and 4, Table 1 and Supplementary table 3, differences in model structure are explained in 'Data analyses and statistical modeling' section). Sporozoite yields simulated over the same parameters for both models were also similar (Fig. 3); specifically, sporozoite yields increased with mean oocyst densities and time (colored lines), albeit with a corresponding increase in uncertainty (shaded areas, Fig. 3).

## Discussion

While mean oocyst densities estimated earlier in the infection were able to capture the non-linear trends in sporozoite recovery, the continued migration of sporozoites to the salivary glands also resulted in higher sporozoite yields over 17–29 days post-bloodmeal. The effect of mean oocyst densities and time were independent of parasite strain, and the number of salivary glands pooled, further suggesting that the effects described here should be broadly applicable. Taken together, we describe a framework where mean oocyst densities estimated earlier in the infection could be used to approximate yields a priori, with the potential for yields to be maximized by leveraging the relationship between sporozoite yields and time. For example, at all mean oocyst densities, our model suggests that compared to 17 days post-bloodmeal, yields may be 1.3-, 1.7-, and 2.2-fold higher at 21, 25, and 29 days post-bloodmeal, respectively (Fig. 3). While yields will be dependent on mean oocyst densities, in general, by accounting for the effect of time, our results offer several benefits such as dissecting fewer mosquitoes, increasing the efficiency of otherwise cumbersome technical procedures, and providing greater reliability in the results of the downstream assays.

The range of sporozoites reported here generally corroborates previous findings from individual or pooled mosquitoes infected with *P. berghei* [17–19, 21, 33–38, 56–60]. The quadratic effect of mean oocyst densities on sporozoite yields increased with mean oocyst densities in a non-linear manner, wherein the initial increase in yields until ~ 50 oocysts were followed by a decline. In general, this non-linearity has been noted for sporozoites produced by both rodent and human *Plasmodium* species; current evidence suggests sporozoite densities may be dependent on complex interactions between nutritional availability, vector mortality, and immune responses to the oocysts and/or sporozoites [22, 61–63]. For *P. berghei*, the

decline in sporozoite densities was noted at mean oocyst densities  $\geq 50$  per midgut [19, 21], with at least one study suggesting vector mortality during sporozoite egress as the most likely explanation for the reduced recovery [21]. A major benefit of this analysis is that we can now leverage our knowledge of the relationship between sporozoite yields and time post-infection to reduce the risk of losing sporozoites at high mean oocyst densities (e.g.,  $> 50$ ) [19, 21], while also reducing dissection effort by boosting yields at lower oocyst densities. Although the model was unable to detect any evidence of a density-dependent delay (lack of clear association between mean oocyst densities and time), the clear effect of time suggested that the continued invasion of salivary glands should still be beneficial to sporozoite yields. Taken together, the effects of oocyst densities and time carry two implications for maximizing sporozoite yields. First, the non-linear effect of mean oocyst densities suggests that as oocyst densities increase ( $> 50$ ), more mosquitoes would need to be dissected to obtain yields comparable to groups with lower densities. Second, although yields may increase linearly over time for all oocyst densities, groups with mean oocyst densities over 50 may still yield more sporozoites than groups with oocyst densities under 50, even if, for instance, sporozoite collection was delayed to 29 days post-bloodmeal (Fig. 3).

In addition to testing the potential for yields to decline with increasing mean oocyst densities ( $\geq 50$ ), we also tested whether yields would decline over time by including a quadratic ("humped") effect of days post-bloodmeal on sporozoite yields (supplementary tables 2 and 3). While the model was unable to identify a time point when yields started declining, simulations based on the model fit suggested that irrespective of mean oocyst density, yields became increasingly uncertain starting  $\sim 25$  days post-bloodmeal (Fig. 1b and Fig. 2). Whether this was due to a lack of data, nutritional resources, heightened immune responses, a sign of deteriorating host health and increased mortality, or sporozoite senescence [19–21], is not possible to ascertain here. As such, until more data is available, we suggest sampling mosquitoes at  $\sim 25$  days post-bloodmeal, but no later than 29 days when sporozoite infectivity has been suggested to decline [33]. Based on our results, the model predicts 1.7-fold higher yields at 25 days (vs 17 days), irrespective of mean oocyst densities.

Together, mean oocyst densities and time were able to account for just under 50% of the overall variation in sporozoite yields from our dataset (e.g., Table 1). Allowing the contributions of the two predictors to vary randomly among groups significantly improved the model's account of the total variation ( $R^2 > 70\%$ , Table 1), indicating the presence of other confounders specific to each group contributed significantly to the variation around the predicted yields (Fig. 1 and Fig. 3). For instance, of the groups where more information was available, one potential source of variation was a group of PbANKA-infected mosquitoes, which, despite only carrying a mean of 10.1 oocysts/midgut provided unexpectedly high sporozoite yields over time. However, only seven of the 17 individuals dissected (41%) showed evidence of having fed on the mouse (presence of eggs in ovaries during oocyst quantifications, mean feeding rates for all groups  $\pm$  se =  $72.4\% \pm 2.7$ ), suggesting that  $< 50\%$  of the group could contribute to the sporozoite pool (albeit with the caveat that estimation of feeding rates may be masked by eggs being reabsorbed by the mother [64–66]). These seven blood-fed individuals were carrying a mean of 25 oocysts per midgut, which was higher than the 10.1 calculated earlier, but more likely to explain the higher

sporozoite yields observed for the group. In contrast, a group of PbGFP-LUC<sub>CON</sub>-infected mosquitoes with a mean oocyst density of 46.2 should have yielded, according to the model, significantly more sporozoites than the 5,250 recovered from 15 salivary glands (or 350 per mosquito) at 17 days post-bloodmeal. To determine why the yields were low, we quantified oocysts in the midguts of these 15 individuals at the time of sporozoite collections (i.e., 17 days) and found that they were carrying 17 mean oocysts per midgut, which was lower than the 46.2 mean oocysts measured earlier from the same group. Although we are uncertain whether this was due to excess mortality in the cage or overestimation of mean oocyst densities earlier, the low oocyst densities were the likely reason for the low yields observed. While the models identified these groups as sources of variation, in general, they were consistent with the conclusion that yields were driven by mean oocyst densities and time.

While the instances above highlight how ‘accurate’ estimation of mean oocyst densities is also critical to predicting sporozoite yields over time, despite the consistency, the effect of time on sporozoite yields was less clear (Table 1). Future experiments with sporozoites collected over a more balanced sampling schedule, and a fixed number of individuals could address some of the study limitations and offer clearer insight into whether rates of sporozoite migration do indeed vary with oocyst density. As oocyst densities increase, sporozoite migration over time may be the primary reason for increased yields (Fig. 1a), however, a clear estimation of its contribution may be confounded by pooling sporozoites from individuals with increasingly heterogenous oocyst densities (shaded areas, Supplementary Fig. 2). Further, this heterogeneity between individuals in their contribution to the sporozoite pool could explain why adjusting for differences in the number of salivary glands did not identify a clearer effect of time (Supplementary table 3). Thus, estimating the relationship between oocyst density and sporozoite yields from pooled salivary glands likely only provides qualitative insight into a process that may vary quantitatively among individuals. Considering this, we suggest that isolating sporozoites from all mosquitoes at a single day post-bloodmeal (e.g., 25–26 days), instead of at fixed intervals, may be a more efficient use of mosquitoes and time, while still achieving maximal sporozoite yields.

There are two additional caveats to our study and analysis. First, although the main effects of mean oocyst densities and time were common to both parasite strains, our objective was not to compare strains. Thus, we cannot disregard the possibility that strains may differ in overall sporozoite output [21]. Second, differences between mosquito colonies due to genetic (e.g., *An. stephensi* “Indian” vs “SDA-500”) [67, 68] and/or epigenetic effects (for instance, larval culture conditions and its carry-over effect on adults) [69], may manifest as differences in parasite infection, replication, and sporozoite yields. While these differences suggest that more careful consideration may be necessary before settling on specific (or range of) parameters, the fundamental nature of the relationship between oocyst densities, time, and sporozoite yields should be consistent.

## Conclusions

Taken together, our results suggest that as with other organismal systems, density-dependent effects in *Plasmodium* parasites resulted in a non-linear relationship between the density of parasites successfully

establishing (oocysts) and the transmission stages (sporozoites) [23, 70]. In general, the temporal dynamics of sporozoite migration are critical to *Plasmodium* fitness. Thus, in addition to indicating how prior oocyst densities could be leveraged to streamline sporozoite isolations for other *Plasmodium* species [2], our results point to the potential for oocyst densities to predict when a mosquito becomes infectious (EIP), but also whether the changes in sporozoite densities over time can alter parasite infectivity to the next host [33, 71]. Irrespective of the downstream application, the versatility offered by rodent malaria models underscores their continued utility in evaluating strategies to manage parasite transmission, but also how these efforts can be aided by an increased understanding of fundamental aspects of vector-parasite interactions.

## Declarations

**Ethics approval and consent to participate:** Not applicable.

**Consent for publication:** Not applicable.

**Availability of data and materials:** Upon acceptance, all data and statistical code from the study will be posted to a public repository.

**Competing interests:** "The authors declare that they have no competing interests".

**Funding:** This work was funded by the University of Georgia and the NIH (5R01AI110793-04). JCS was funded by NIH T32 AI1060546. The funding bodies did not have any say in the design of the study and collection, analysis, interpretation of data or in writing the manuscript.

### Author contributions:

AKP: conception, design, acquisition, analysis, interpretation, drafting the manuscript.

JCS: acquisition, design, analysis, interpretation, revisions to the manuscript.

BFF: acquisition, design.

LS: acquisition.

DAH: acquisition.

DEK: acquisition.

CCM: acquisition, analysis, interpretation, substantive revisions to the manuscript.

All authors approved the submitted version.

**Acknowledgments:** The authors would like to thank Courtnie Vickery, Tyrone Williams, Anne Elliott, Adam McHugh, and Jared Cooper for help with mosquito husbandry, and the UGA animal resources staff for

mouse husbandry.

## References

1. Goh YS, McGuire D, Renia L: **Vaccination With Sporozoites: Models and Correlates of Protection.** Front Immunol 2019, **10**:1227.
2. Various: Methods In Molecular Biology: **Malaria** In *Methods and Protocols* (Menard R ed., vol. 923, 2nd edition. pp. 626: Springer; 2013:626.
3. Cowell AN, Winzeler EA: **Advances in omics-based methods to identify novel targets for malaria and other parasitic protozoan infections.** Genome Med 2019, **11**:63.
4. Rosenberg R, Rungswongse J: **The number of sporozoites produced by individual malaria oocysts.** Am J Trop Med Hyg 1991, **45**:574–577.
5. Leitner WW, Bergmann-Leitner ES, Angov E: **Comparison of *Plasmodium berghei* challenge models for the evaluation of pre-erythrocytic malaria vaccines and their effect on perceived vaccine efficacy.** Malar J 2010, **9**:145.
6. Kreutzfeld O, Muller K, Matuschewski K: **Engineering of Genetically Arrested Parasites (GAPs) For a Precision Malaria Vaccine.** Front Cell Infect Microbiol 2017, **7**:198.
7. Kori LD, Valecha N, Anvikar AR: **Insights into the early liver stage biology of *Plasmodium*.** J Vector Borne Dis 2018, **55**:9–13.
8. Parmar R, Patel H, Yadav N, Patidar M, Tyagi RK, Dalai SK: **Route of administration of attenuated sporozoites is instrumental in rendering immunity against Plasmodia infection.** Vaccine 2016, **34**:3229–3234.
9. Sauerwein RW, Bousema T: **Transmission blocking malaria vaccines: Assays and candidates in clinical development.** Vaccine 2015, **33**:7476–7482.
10. Mordmuller B, Supan C, Sim KL, Gomez-Perez GP, Ospina Salazar CL, Held J, Bolte S, Esen M, Tschan S, Joanny F, et al: **Direct venous inoculation of *Plasmodium falciparum* sporozoites for controlled human malaria infection: a dose-finding trial in two centres.** Malar J 2015, **14**:117.
11. Singh N, Barnes SJ, Kennedy S, Adams JH: **Experimental evaluation of cryopreservative solutions to maintain in vitro and in vivo infectivity of *P. berghei* sporozoites.** PLoS One 2017, **12**:e0177304.
12. Marques-da-Silva C, Peissig K, Kurup SP: **Pre-Erythrocytic Vaccines against Malaria.** Vaccines (Basel) 2020, **8**.
13. WHO: **World malaria report 2021** In *World malaria report 2021* (Organization WH ed. Geneva, Switzerland: World Health Organization; 2021.
14. Schrum M., Canezin A., Chakravarty S., Laskowski M., Comert S., Sevimli Y., Chirikjian G. S., Hoffman S.L., R.H. T: **An Efficient Production Process for Extracting Salivary Glands from Mosquitoes.** arXiv 2019.
15. Stone WJ, Eldering M, van Gemert GJ, Lanke KH, Grignard L, van de Vegte-Bolmer MG, Siebelink-Stoter R, Graumans W, Roeffen WF, Drakeley CJ, et al: **The relevance and applicability of oocyst**

- prevalence as a read-out for mosquito feeding assays. *Sci Rep* 2013, **3**:3418.
- 16. Miura K, Swihart BJ, Deng B, Zhou L, Pham TP, Diouf A, Fay MP, Long CA: **Strong concordance between percent inhibition in oocyst and sporozoite intensities in a *Plasmodium falciparum* standard membrane-feeding assay.** *Parasit Vectors* 2019, **12**:206.
  - 17. Vanderberg JP, Yoeli M: **Effects of temperature on sporogonic development of *Plasmodium berghei*.** *J Parasitol* 1966, **52**:559–564.
  - 18. Sinden RE, Dawes EJ, Alavi Y, Waldock J, Finney O, Mendoza J, Butcher GA, Andrews L, Hill AV, Gilbert SC, Basanez MG: **Progression of *Plasmodium berghei* through *Anopheles stephensi* is density-dependent.** *PLoS Pathog* 2007, **3**:e195.
  - 19. Dawes EJ, Churcher TS, Zhuang S, Sinden RE, Basanez MG: **Anopheles mortality is both age- and *Plasmodium*-density dependent: implications for malaria transmission.** *Malar J* 2009, **8**:228.
  - 20. Miazgowicz KL, Shocket MS, Ryan SJ, Villena OC, Hall RJ, Owen J, Adanlawo T, Balaji K, Johnson LR, Mordecai EA, Murdock CC: **Age influences the thermal suitability of *Plasmodium falciparum* transmission in the Asian malaria vector *Anopheles stephensi*.** *Proc Biol Sci* 2020, **287**:20201093.
  - 21. Pollitt LC, Churcher TS, Dawes EJ, Khan SM, Sajid M, Basanez MG, Colegrave N, Reece SE: **Costs of crowding for the transmission of malaria parasites.** *Evol Appl* 2013, **6**:617–629.
  - 22. Vaughan JA: **Population dynamics of *Plasmodium* sporogony.** *Trends Parasitol* 2007, **23**:63–70.
  - 23. Childs LM, Prosper OF: **The impact of within-vector parasite development on the extrinsic incubation period.** *R Soc Open Sci* 2020, **7**:192173.
  - 24. Ohm JR, Baldini F, Barreaux P, Lefevre T, Lynch PA, Suh E, Whitehead SA, Thomas MB: **Rethinking the extrinsic incubation period of malaria parasites.** *Parasit Vectors* 2018, **11**:178.
  - 25. Wilson K, Bjørnstad ON, Dobson AP, Merler S, Poglayen G, Randolph SE, Read AF, Skorping A: **Heterogeneities in macroparasite infections: patterns and processes.** In *The Ecology of Wildlife Diseases*. Edited by Hudson P.J., Rizzoli A., Grenfell B.T., Heesterbeek H., A.P. D. Oxford, UK: Oxford University Press; 2002: 1–48
  - 26. Beckerman AP, Benton TG, Ranta E, Kaitala V, Lundberg P: **Population dynamic consequences of delayed life-history effects.** *Trends in Ecology & Evolution* 2002, **17**:263–269.
  - 27. Tartz S, Deschermeier C, Retzlaff S, Heussler V, Sebo P, Fleischer B, Jacobs T: ***Plasmodium berghei* sporozoite challenge of vaccinated BALB/c mice leads to the induction of humoral immunity and improved function of CD8(+) memory T cells.** *Eur J Immunol* 2013, **43**:693–704.
  - 28. Gilbert SC, Schneider J, Hannan CM, Hu JT, Plebanski M, Sinden R, Hill AVS: **Enhanced CD8 T cell immunogenicity and protective efficacy in a mouse malaria model using a recombinant adenoviral vaccine in heterologous prime-boost immunisation regimes.** *Vaccine* 2002, **20**:1039–1045.
  - 29. Sato Y, Montagna GN, Matuschewski K: ***Plasmodium berghei* sporozoites acquire virulence and immunogenicity during mosquito hemocoel transit.** *Infect Immun* 2014, **82**:1164–1172.
  - 30. Vaughan JA, Scheller LF, Wirtz RA, Azad AF: **Infectivity of *Plasmodium berghei* sporozoites delivered by intravenous inoculation versus mosquito bite: implications for sporozoite vaccine trials.** *Infect*

Immun 1999, **67**:4285–4289.

31. Nganou-Makamdop K, van Gemert GJ, Arens T, Hermsen CC, Sauerwein RW: **Long term protection after immunization with *P. berghei* sporozoites correlates with sustained IFNgamma responses of hepatic CD8 + memory T cells.** PLoS One 2012, **7**:e36508.
32. Manzoni G, Marinach C, Topcu S, Briquet S, Grand M, Tolle M, Gransagne M, Lescar J, Andolina C, Franetich JF, et al: ***Plasmodium* P36 determines host cell receptor usage during sporozoite invasion.** Elife 2017, **6**.
33. Coosemans M, Wery M, Van Marck E, Timperman G: **Studies on the infectivity of *Plasmodium berghei* sporozoites in experimental hosts.** Ann Soc Belg Med Trop 1981, **61**:349–368.
34. Costa G, Gildenhard M, Eldering M, Lindquist RL, Hauser AE, Sauerwein R, Goosmann C, Brinkmann V, Carrillo-Bustamante P, Levashina EA: **Non-competitive resource exploitation within mosquito shapes within-host malaria infectivity and virulence.** Nat Commun 2018, **9**:3474.
35. Choudhary HH, Gupta R, Mishra S: **PKAc is not required for the preerythrocytic stages of *Plasmodium berghei*.** Life Sci Alliance 2019, **2**.
36. Wang Q, Fujioka H, Nussenzweig V: **Exit of *Plasmodium* sporozoites from oocysts is an active process that involves the circumsporozoite protein.** PLoS Pathog 2005, **1**:e9.
37. Porter MD, Nicki J, Pool CD, DeBot M, Illam RM, Brando C, Bozick B, De La Vega P, Angra D, Spaccapelo R, et al: **Transgenic parasites stably expressing full-length *Plasmodium falciparum* circumsporozoite protein as a model for vaccine down-selection in mice using sterile protection as an endpoint.** Clin Vaccine Immunol 2013, **20**:803–810.
38. Yang J, Schleicher TR, Dong Y, Park HB, Lan J, Cresswell P, Crawford J, Dimopoulos G, Fikrig E: **Disruption of mosGILT in *Anopheles gambiae* impairs ovarian development and *Plasmodium* infection.** J Exp Med 2020, **217**.
39. Nussenzweig RS, Vanderberg J, Most H, Orton C: **Protective immunity produced by the injection of x-irradiated sporozoites of *Plasmodium berghei*.** Nature 1967, **216**:160–162.
40. Janse CJ, Ramesar J, Waters AP: **High-efficiency transfection and drug selection of genetically transformed blood stages of the rodent malaria parasite *Plasmodium berghei*.** Nat Protoc 2006, **1**:346–356.
41. Janse CJ, Franke-Fayard B, Mair GR, Ramesar J, Thiel C, Engelmann S, Matuschewski K, van Gemert GJ, Sauerwein RW, Waters AP: **High efficiency transfection of *Plasmodium berghei* facilitates novel selection procedures.** Mol Biochem Parasitol 2006, **145**:60–70.
42. Janse CJ, Franke-Fayard B, Waters AP: **Selection by flow-sorting of genetically transformed, GFP-expressing blood stages of the rodent malaria parasite, *Plasmodium berghei*.** Nat Protoc 2006, **1**:614–623.
43. Fonager J, Pasini EM, Braks JA, Klop O, Ramesar J, Remarque EJ, Vroegrijk IO, van Duinen SG, Thomas AW, Khan SM, et al: **Reduced CD36-dependent tissue sequestration of *Plasmodium*-infected erythrocytes is detrimental to malaria parasite growth in vivo.** J Exp Med 2012, **209**:93–107.



44. Pathak AK, Shiao JC, Thomas MB, Murdock CC: **Cryogenically preserved RBCs support gametocytogenesis of *Plasmodium falciparum* in vitro and gametogenesis in mosquitoes.** Malar J 2018, **17**:457.
45. Pathak AK, Shiao JC, Thomas MB, Murdock CC: **Field Relevant Variation in Ambient Temperature Modifies Density-Dependent Establishment of *Plasmodium falciparum* Gametocytes in Mosquitoes.** Front Microbiol 2019, **10**:2651.
46. Schmid MA, Kauffman E, Payne A, Harris E, Kramer LD: **Preparation of Mosquito Salivary Gland Extract and Intradermal Inoculation of Mice.** Bio Protoc 2017, **7**.
47. Schielzeth H, Nakagawa S: **Nested by design: model fitting and interpretation in a mixed model era.** Methods in Ecology and Evolution 2013, **4**:14–24.
48. Harrison XA, Donaldson L, Correa-Cano ME, Evans J, Fisher DN, Goodwin CED, Robinson BS, Hodgson DJ, Inger R: **A brief introduction to mixed effects modelling and multi-model inference in ecology.** PeerJ 2018, **6**:e4794.
49. Silk MJ, Harrison XA, Hodgson DJ: **Perils and pitfalls of mixed-effects regression models in biology.** PeerJ 2020, **8**.
50. Brooks ME, Kristensen K, van Benthem KJ, Magnusson A, Berg CW, Nielsen A, Skaug HJ, Machler M, Bolker BM: **glmmTMB Balances Speed and Flexibility Among Packages for Zero-inflated Generalized Linear Mixed Modeling.** R Journal 2017, **9**:378–400.
51. RStudio Team: **RStudio: Integrated Development Environment for R.** Boston, MA: RStudio, Inc.; 2016.
52. R Core Team: **R: A Language and Environment for Statistical Computing.** Vienna, Austria: R Foundation for Statistical Computing; 2018.
53. Wickham H: *ggplot2: Elegant Graphics for Data Analysis.* New York: Springer-Verlag; 2016.
54. Manikandan S: **Measures of central tendency: The mean.** J Pharmacol Pharmacother 2011, **2**:140–142.
55. Lüdecke D: **strengejacke: Load Packages Associated with Strenge Jacke!** R package version 0.3.0 edition2018.
56. Armistead JS, Jennison C, O'Neill MT, Lopaticki S, Liehl P, Hanson KK, Annoura T, Rajasekaran P, Erickson SM, Tonkin CJ, et al: ***Plasmodium falciparum* subtilisin-like ookinete protein SOPT plays an important and conserved role during ookinete infection of the *Anopheles stephensi* midgut.** Mol Microbiol 2018, **109**:458–473.
57. Siden-Kiamos I, Pace T, Klonizakis A, Nardini M, Garcia CRS, Curra C: **Identification of *Plasmodium berghei* Oocyst Rupture Protein 2 (ORP2) domains involved in sporozoite egress from the oocyst.** Int J Parasitol 2018, **48**:1127–1136.
58. Saeed S, Tremp AZ, Dessens JT: **The *Plasmodium* LAP complex affects crystalloid biogenesis and oocyst cell division.** Int J Parasitol 2018, **48**:1073–1078.
59. Ukegbu CV, Giorgalli M, Tapanelli S, Rona LDP, Jaye A, Wyer C, Angrisano F, Blagborough AM, Christophides GK, Vlachou D: **PIMMS43 is required for malaria parasite immune evasion and**

- sporogonic development in the mosquito vector.** Proc Natl Acad Sci U S A 2020, **117**:7363–7373.
60. Hollin T, De Witte C, Freville A, Guerrera IC, Chhuon C, Saliou JM, Herbert F, Pierrot C, Khalife J: **Essential role of GEXP15, a specific Protein Phosphatase type 1 partner, in *Plasmodium berghei* in asexual erythrocytic proliferation and transmission.** PLoS Pathog 2019, **15**:e1007973.
61. Ponnudurai T, Lensen AH, van Gemert GJ, Bensink MP, Bolmer M, Meuwissen JH: **Sporozoite load of mosquitoes infected with *Plasmodium falciparum*.** Trans R Soc Trop Med Hyg 1989, **83**:67–70.
62. Hillyer JF, Strand MR: **Mosquito hemocyte-mediated immune responses.** Curr Opin Insect Sci 2014, **3**:14–21.
63. Murdock CC, Moller-Jacobs LL, Thomas MB: **Complex environmental drivers of immunity and resistance in malaria mosquitoes.** Proc Biol Sci 2013, **280**:20132030.
64. Hopwood JA, Ahmed AM, Polwart A, Williams GT, Hurd H: **Malaria-induced apoptosis in mosquito ovaries: a mechanism to control vector egg production.** J Exp Biol 2001, **204**:2773–2780.
65. Ahmed AM, Hurd H: **Immune stimulation and malaria infection impose reproductive costs in *Anopheles gambiae* via follicular apoptosis.** Microbes Infect 2006, **8**:308–315.
66. Pigeault R, Villa M: **Long-term pathogenic response to *Plasmodium relictum* infection in *Culex pipiens* mosquito.** PLoS One 2018, **13**:e0192315.
67. Jiang X, Peery A, Hall AB, Sharma A, Chen XG, Waterhouse RM, Komissarov A, Riehle MM, Shouche Y, Sharakhova MV, et al: **Genome analysis of a major urban malaria vector mosquito, *Anopheles stephensi*.** Genome Biol 2014, **15**:459.
68. Sinka ME, Pironon S, Massey NC, Longbottom J, Hemingway J, Moyes CL, Willis KJ: **A new malaria vector in Africa: Predicting the expansion range of *Anopheles stephensi* and identifying the urban populations at risk.** Proc Natl Acad Sci U S A 2020, **117**:24900–24908.
69. Moller-Jacobs LL, Murdock CC, Thomas MB: **Capacity of mosquitoes to transmit malaria depends on larval environment.** Parasit Vectors 2014, **7**:593.
70. Bompard A, Da DF, Yerbanga SR, Morlais I, Awono-Ambene PH, Dabire RK, Ouedraogo JB, Lefevre T, Churcher TS, Cohuet A: **High *Plasmodium* infection intensity in naturally infected malaria vectors in Africa.** Int J Parasitol 2020, **50**:985–996.
71. Graumans W, Jacobs E, Bousema T, Sinnis P: **When Is a *Plasmodium*-Infected Mosquito an Infectious Mosquito?** Trends Parasitol 2020, **36**:705–716.

## Tables

Tables 1 is available in the Supplementary Files section.

## Figures

**Figure 1****a****b**


*P. berghei* strain • PbANKA ▲ PbGFP-LUC<sub>CON</sub>

**Figure 1**

Sporozoite yields increase (a) non-linearly with (arithmetic) mean oocyst densities earlier in the infection (11-15 days post-bloodmeal), and (b) linearly over time (17 to 29 days post-bloodmeal) for mosquitoes infected with *P. berghei* strains PbANKA (circles) or PbGFP-LUC<sub>CON</sub> (triangles). Lines and shaded areas indicate the mean and uncertainty (95% CI), respectively, predicted by the minimal model (Table 1). Mean oocyst densities were derived from counts of individual midguts, irrespective of infection status (oocysts

$\geq 0$ ). Note that although sporozoite yields are expressed per mosquito, values do not represent a true mean as counts are from pooled salivary glands, with no information of densities contributed by individuals.


**Figure 2**



**Figure 2**

Mean oocyst densities for the 46 groups of mosquitoes increase linearly with the respective standard deviations (orange crosses), indicating the potential for collecting sporozoites from individuals with increasingly heterogeneous oocyst densities. Mosquitoes were infected with *P. berghei* strains PbANKA (circles) or PbGFP-LUC<sub>CON</sub> (triangles).

### Figure 3



### Figure 3

Predicted sporozoite yields over time (17, 24, and 29 days post-bloodmeal) at the respective mean oocyst densities (x-axis). Predictions after modeling sporozoite yields as (a) the 'mean' ('Minimal model', Table 1) or (b) after considering differences in the number of salivary glands ('Minimal model', Supplementary table 3). Lines and shaded areas indicate the mean and uncertainty (95%CI), respectively.

## Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Table1.pdf](#)
- [Supplementaryfigure1.pdf](#)
- [Supplementaryfigure2.pdf](#)
- [Supplementarytable1.pdf](#)
- [Supplementarytable2.pdf](#)
- [Supplementarytable3.pdf](#)