

A National Hiv Clinical Mentorship Program: Enabling Zambia to Accelerate Towards the Controlling the Hiv Epidemic

Mary Adetunke Boyd (✉ moq6@cdc.gov)

CDC: Centers for Disease Control and Prevention <https://orcid.org/0000-0001-6565-680X>

Sombo Fwoloshi

Zambia Ministry of Health

Peter A. Minchella

CDC: Centers for Disease Control and Prevention

James Simpungwe

CDC: Centers for Disease Control and Prevention

Terence Siansalama

CDC: Centers for Disease Control and Prevention

Danielle T. Barradas

CDC: Centers for Disease Control and Prevention

Minesh Shah

CDC: Centers for Disease Control and Prevention

Lloyd Mulenga

Zambia Ministry of Health

Simon Agolory

CDC: Centers for Disease Control and Prevention

Research

Keywords: Zambia, HIV, mentorship

Posted Date: January 22nd, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-151655/v1>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at PLOS Global Public Health on February 23rd, 2022. See the published version at <https://doi.org/10.1371/journal.pgph.0000074>.

Abstract

Background

Although Zambia has increased the proportion of people living with HIV (PLHIV) who are on antiretroviral therapy (ART) in recent years, progress toward HIV epidemic control remains inconsistent. Some districts are still failing to meet the UNAIDS 90/90/90 targets where 90% of PLHIV should know their status, 90% of those who know their status should be receiving sustained ART, and 90% of those on ART should have documented viral load suppression (VLS) by 2020. Providing consistently excellent HIV services at all ART health facilities is critical for achieving the UNAIDS 90/90/90 targets and controlling the HIV epidemic in Zambia. Zambia Ministry of Health (MoH), in collaboration with the U.S. Centers for Disease Control and Prevention (CDC), aimed to achieve these targets through establishing a national HIV clinical mentorship program in which government-employed mentors were assigned to specific facilities with a mandate to identify and ameliorate programmatic challenges.

Methods

Mentors were hired, trained and deployed to individual facilities in four provinces to mentor staff on quality HIV clinical and program management. The pre-mentorship period was July 2018–September 2018 and the post-mentorship period was July 2019–September 2019.

Results

Review of key programmatic indicators from the pre and post-deployment periods revealed HIV testing yield improved from 4.2–6.8% ($P < 0.001$) as fewer HIV tests were needed despite the number of PLHIV being identified and placed on ART increasing from 492,613 to 521,775, and VLS increased from 84.8–90.1% ($p < 0.001$).

Conclusions

Key considerations in the establishment of an HIV clinical mentorship program include having a government-led process of regular site level data review and continuous clinical mentorship underpinned by quality improvement methodology.

Background

With an estimated 1.2 million people living with HIV (PLHIV) and 12% prevalence, Zambia is severely affected by the HIV epidemic¹. By the end of September 2018, 892,362 (72%) PLHIV in Zambia were on life saving antiretroviral therapy (ART).² Despite improvements in ART coverage, overall progress toward UNAIDS 90-90-90 targets for HIV epidemic control (90% of PLHIV know their HIV sero-status, 90% of those

who know their status sustain ART, and 90% of those on ART achieve HIV viral load suppression by 2020) is slow and inconsistent. While the factors limiting progress are diverse, an effort to evaluate the provision of services within Zambia's HIV program revealed that a lack of continuous capacity building and mentorship for healthcare workers from experienced HIV clinicians was a key gap³. Due to a strained health workforce, clinical services in Zambia are generally provided by mid-level practitioners such as clinical officers and nurses who often have limited access to in-service training, ongoing continuous medical education or clinical mentorship in the delivery of quality client services³. This is compounded by the reality that specialist care is more often concentrated in urban areas, far out of the reach of much of Zambia's population. In the United States and Sub-Saharan Africa, HIV clinical mentorship has been shown to improve clinical practices and outcomes related to linking and retaining patients in HIV care and across varying disciplines^{4,5,6,7}. Furthermore, in resource limited settings where distance is a barrier to providing ongoing clinical training and mentorship, Project ECHO (Extension for Community Healthcare Outcomes), which uses video-conferencing technology to bridge knowledge gaps between experts and providers, has demonstrated an impact on patient outcomes^{8,9,10}. Thus, the Zambian Ministry of Health (MoH), with support from U.S. Centers for Disease Control and Prevention (CDC), established a national HIV clinical mentorship program with the aim of improving performance across key HIV program indicators. Improved case finding, increasing the number of HIV infected individuals on treatment and viral load suppression were particular areas of focus.

Methods

Study design and setting

We conducted a retrospective pre-post analysis of routine program data collected during two periods: July – September 2018 and July – September 2019. The HIV clinical mentorship program was implemented across four of Zambia's ten provinces, Eastern, Western, Southern, and Lusaka, which enabled the program to reach a wide array of clinical care settings (e.g., urban, rural, semi-urban) and geographies with varying HIV prevalence (Fig. 1).

Program

Starting in October 2018, 27 mentors were hired, trained, and deployed to 45 districts with high HIV burdens across the four selected provinces. Mentors had a variety of clinical backgrounds (nurses, clinical officers, and medical officers), were hired through a competitive process, and participated in a two-week training focused on the fundamentals of HIV clinical and program management. Following training, each mentor was assigned 2–10 health facilities, each of which employed more than 15 health workers and managed patient volumes ranging from 500 to 12,000 clients on ART. In general, each mentor was assigned to one district, though there were exceptions in which mentors were tasked with covering multiple districts or multiple mentors were assigned to facilities within the same district. Deployment of mentors and assignment of health facilities was done strategically in order to leverage interconnectedness. Specifically, more senior mentors were assigned to cover all districts that did not

have a dedicated mentor and to perform visits supplemented with telephone calls with the same frequency as districts with a dedicated mentor. Mentor deployment was also done in collaboration with district health offices to ensure local support for the program. Overall, there were 28 mentor-supported districts that accounted for approximately 83% of the patients on ART within the four selected provinces at the time the program was initiated.

Following their training and subsequent deployment, mentors continued to interact with experienced public health professionals and HIV clinical care providers from CDC, local and international implementing partners and academic institutions, and the Zambia Ministry of Health chief mentor via weekly video conferences, on-site mentoring, and Tele ECHO™ sessions¹². Weekly Tele ECHO sessions aimed to both build knowledge through presentation and discussion of common or complex HIV cases, and to facilitate interaction among healthcare workers from across different health facilities to help create a community of practice with substantial cross-learning across the network. Additionally, weekly check-ins and on-site mentoring focused on data review using provided reporting templates and implementation of continuous quality improvement (CQI) methodology. Overall, ongoing interactions with mentors aimed to improve clinical practice by addressing key programmatic barriers to epidemic control and by sharing best practices. Specific areas of focus included mentored use of HIV risk screening tools in outpatient departments and safe and ethical index client elicitation to improve HIV case finding. Mentors implemented the national risk screening tool (RST) at facility level and provided facility-based group training and 1:1 mentoring to HIV testers and other facility staff on the proper use of RST, routinely observing and providing on the spot feedback. To address retention in care, mentors implemented scheduled facility appointments in ART clinics, monitored missed appointments and provided training in patient satisfaction and customer care. Mentors also guided facility staff in the implementation of routine HIV viral load testing and established high viral load clinic days for the management of patients with high HIV viral load. Notably, mentors institutionalized facility mid-week CQI meetings where the facility nurse-in-charge leads facility staff in the review and accompanying process improvements of key HIV indicators.

Outcomes

The outcomes of interest in our analysis were related to HIV testing services (HTS) and HIV viral load (VL) monitoring. Those related to HTS were (1) overall positive yield, defined as the number of positive HIV tests divided by the total number of individuals who received HIV testing per national guidelines; and (2) overall positive yield from index case testing (ICT), defined as the number of index testing contacts who were identified HIV-positive divided by the total number of index contacts who received HTS and received their test results. Index testing contacts were defined as sexual contacts or biological children of PLHIV who were elicited and offered HTS¹³. Outcomes for HIV viral load testing were (3) viral load coverage, defined as the number of PLHIV with documented HIV viral load test results during the last 12 months divided by the number of PLHIV eligible for viral load testing, and (4) HIV viral load suppression, defined as the number of PLHIV with viral load test results < 1,000 cp/ml during the previous 12 months divided by the number with a HIV viral load test result. For all outcomes we used definitions and guidance from

the U.S President's Emergency Plan For AIDS Relief (PEPFAR) Monitoring, Evaluation and Reporting (MER) Indicator Reference Sheet version 2.3¹³.

Data source and analysis

Aggregate program data were abstracted from PEPFAR's Data for Accountability Transparency and Impact (DATIM) system. The pre-mentorship period was July 2018–September 2018 and the post-mentorship period was July 2019–September 2019. Proportions were calculated for outcomes of interest and analyzed using descriptive statistics. Chi-square tests were used to compare positivity yield, viral load coverage, and viral load suppression. All analyses were performed using R (version 4.0.2).

Results

Given the high proportion of PLHIV represented by mentor-supported districts and the intended impacts of mentors beyond their districts, results from all districts in the four selected provinces were included in the analysis. Between the pre- and post-mentorship periods the HIV testing yield increased from 4.2–6.8% ($p < 0.001$) and this resulted in total of 38,555 PLHIV diagnosed in the post mentorship period compared to 37,206 in the pre-mentorship period. The number of cases identified in the post-mentorship period included 9,091 cases identified through ICT up from 4,115 and index-case testing yields from 11.6–16.9% ($p < 0.001$) as well as increased contribution of index-case testing to overall case finding from 12% (4,006/33,574) to 26% (9,039/34,944) in one year. Number of PLHIV on ART increased from 492,613 to 521,775 and documented viral load coverage increased from 40.8–74.1% ($p < 0.001$) with an increasing rate of viral suppression from 84.8–90.1% ($p < 0.001$). Also, number of HIV tests performed decreased from 878,948 to 565,226. Table 1.

Table 1

Aggregate data for the pre-mentorship (July 1, 2018 - Sept 30, 2018) and post-mentorship (July 1, 2019 - Sept 30, 2019) periods reported by PEPFAR-supported health facilities in four Zambian provinces (Eastern, Western, Southern, Lusaka)

	Pre-mentorship period (July 1, 2018 - Sept 30, 2018)	Post-mentorship period (July 1, 2019 - Sept 30, 2019)	Absolute Difference	<i>p</i>-value
Number of HIV tests performed	878,948	565,226	(313,722)	
Number of positive HIV tests	37,206	38,555	1,349	
Positive yield from HIV testing	4.2%	6.8%		p < 0.001
Number of index contact HIV tests	35,386	53,859	18,473	
Number of positive index contact HIV tests	4,115	9,091	4,976	
Positive yield from index testing	11.6%	16.9%		p < 0.001
Number of PLHIV currently on ART	492,613	521,775	29,162	
Number of PLHIV eligible for VL testing	452,239	497,560	45,321	
Number of PLHIV receiving VL test	184,488	368,801	184,313	
VL coverage among PLHIV	40.8%	74.1%		p < 0.001
Number of VL tests with suppressed viral load	156,383	334,453	178,070	
Viral load suppression	84.8%	90.1%		p < 0.001
¹ Chi-square tests were used compare frequencies				

Discussion

Implementation of Zambia's national HIV clinical mentorship program aligned with progress toward the UNAIDS 90/90/90 targets in the four selected provinces. Program data indicated that rates of case-finding, index-case testing, HIV testing efficiency (i.e. reduction of unnecessary testing), ART initiation, viral load testing coverage, and viral load suppression significantly improved during the implementation period. In particular, the decrease in the number of HIV tests pre and post program, coupled with an

increase in the number of total positive cases identified suggest that RST implementation contributed to testing efficiency implemented by the mentors in facilities. These results justify continued support for the mentorship program and make a case for implementation of similar programs in other settings.

To achieve these results, a “team of teams” approach, with solutions derived not from a hierarchical dictatorship, but rather from more complex multi-directional relationships, were critical to program success. More experienced public health managers at CDC and MOH provided onsite and remote sessions on evidence-based innovations to mentors who in turn provided context to the strategies and mentorship to facility staff ^{4,14,15}. Key aspects of Zambia’s national HIV clinical mentorship program were that it was government led, which ensured authority and sustainability; incorporated video-conferencing technology, which enabled mentors to interact with mentees on a regular basis, and that there was a commitment to a continuous quality improvement methodology, which led to innovation and improvement, rather than just performance monitoring. We also recognized that accountability of mentors was critical, such that poor performance as indicated by lack of visitation to assigned sites and lack of persistence in applying continuous quality improvement methodology to achieve results was addressed through performance management improvement plans. Conversely, exceptional performance was recognized by awarding mentor’s mid-year certificates of achievements.

The PEPFAR model of working through international and non-governmental implementing partners enabled countries to rapidly scale up ART in the initial phase of the HIV response. In the current phase of the response, there is a greater focus on ensuring country ownership in order to promote sustainability while maintaining epidemic control ¹⁶. Zambia’s national HIV clinical mentorship program is an example of an effort to develop local ownership as it helps to not only establish a capable workforce that can manage both quality for the care individuals, but also quality of programs at sites, and ultimately a successful national HIV program that utilizes a public health approach. However, as additional evidence in the optimal management of HIV and new public health approaches to epidemic control continues to emerge, even though mentors when hired were clinicians in their own right and may even have public health degrees, a key lesson learned is that mentors need ongoing mentoring, which requires significant resources. Zambia’s national HIV clinical mentorship program was the recipient of significant investment from the national ministry of health and the U.S. government, which is consistent with training and mentorship programs in epidemiology, clinical services and non-clinical health systems in the region ^{17,18,19}. Specifically, for the 27 mentors, 10 CDC and 8 MOH staff were involved in regular mentorship and supervision. Additionally, annual trainings and quarterly retreats were held to review current practice guidelines and reflect on its application in their assigned districts and sites. While the up-front investment is substantial, the mentors graduate and provide the same training and mentorship to upcoming mentors thus contributing to the building of national capacity poised to take-over management of the HIV program in Zambia from international and non-governmental organizations funded by the U.S and other donors. The success of this program will indicate that Zambia’s HIV clinical mentorship program leveraged donor funding to contribute towards building a sustainable system for delivery of HIV services going forward.

There are some limitations to our analysis; the improvements seen in the post-mentorship period could be due to improved data quality at the site level from implementation of a CQI methodology by mentors who systematically reviewed and queried site data on a weekly and continuous basis, and cannot be attributed fully to improvement in service provision. Additionally, it may not be possible to attribute all the improvements in the monitored indicators seen in the four provinces to the mentorship program, it is plausible that other factors at the individual, societal, structural and policy levels by the MOH and other implementing partners working on HIV in the provinces and not necessarily an intervention implemented contributed significantly to these achievements. However, to our knowledge, no new initiatives or interventions were implemented in the four provinces during the evaluation period that can fully explain the significant improvements in case finding, treatment coverage and viral load testing and suppression.

Conclusions

Sound clinical knowledge and practice combined with public health practices in real-time, with continuous quality improvement as the backbone can enable program impact at facility and provincial level. To our knowledge, this is the first published report of a government-led national HIV clinical mentorship program in sub-Saharan Africa that aligned with significant improvement in key HIV programmatic indicators. Implementation of this approach should be considered in other resource limited settings.

Abbreviations

PLHIV

People living with HIV

ART

Antiretroviral therapy

UNAIDS

Joint United Nations Programme on HIV/AIDS

VLS

Viral Load Suppression

VL

Viral Load

MOH

Ministry of Health

CDC

U.S. Centers for Disease Control and Prevention

CQI

Continuous Quality Improvement

HTS

HIV Testing Services

PEPFAR

U.S President's Emergency Plan For AIDS Relief

MER

Monitoring, Evaluation and Reporting

DATIM

Data for Accountability Transparency and Impact

ECHO

Extension for Community Healthcare Outcomes

RST

Risk Screening/Stratification Tool

Declarations

Ethics approval and consent to participate

Ethical approval for this retrospective analysis of routine program data was approved by the ERES Converge Zambian Institutional Review Board (IRB) and U.S. Centers for Disease Control and Prevention.

Consent for publication

Not applicable

Availability of data and material

The data is publicly available and openly accessible at <https://data.pepfar.gov/dashboards>

Competing interests

The authors have no conflicts of interest to disclose.

Funding

This work has been supported by the President's Emergency Plan for AIDS Relief (PEPFAR) through the Centers for Disease Control and Prevention (CDC). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Authors' contributions

MAB, SF, MS, LM, and SA designed the project; MAB, SF, MS, LM, JS, TS and SA directed the project, MAB, PAM, and DTB conducted the analysis; MAB and PAM drafted the manuscript, and all authors read and approved the final version.

Acknowledgements

We would like to thank each of the mentors and site staff for their dedication to excellence and to public health.

Authorship Disclaimer

The findings and conclusions in this report are those of the author(s) and do not necessarily represent the official position of the funding agencies.

References

1. Barradas DT, Gupta S, Moyo C, Sachathep K, Dzekedzeke K, Nkumbula T, et al. Findings from the 2016 Zambia Population-based HIV Impact Assessment (ZAMPHIA): HIV prevalence, incidence and progress towards the 90-90-90 goals. Abstract TUAC0301. 9th International AIDS Society Meeting; 2017 Jul 23–26; Paris, France.
2. PEPFAR Data for Accountability, Transparency and Impact Monitoring (DATIM)
3. Mwamba, C., Sharma, A., Mukamba, N., Beres, L., Geng, E., Holmes, C. B., ... Topp, S. M. (2018). 'They care rudely!': resourcing and relational health system factors that influence retention in care for people living with HIV in Zambia. *BMJ global health*, 3(5), e001007. doi:10.1136/bmjgh-2018-001007
4. Wagner CMJ, Carlberg-Racich S, Linsk NL, et al. Impacts of Longitudinal Mentorship to Strengthen the HIV Workforce: Qualitative Evidence of Changes in Clinicians' Knowledge, Attitudes, and Practice. *The Journal of the Association of Nurses in AIDS Care: JANAC*. 2017;28(6):938–952.
5. Naikoba S, Senjovu KD, Mugabe P, et al. Improved HIV and TB Knowledge and Competence Among Mid-level Providers in a Cluster-Randomized Trial of One-on-One Mentorship for Task Shifting. *Journal of Acquired Immune Deficiency Syndromes (1999)*. 2017 Aug;75(5):e120-e127. DOI: 10.1097/qai.0000000000001378.
6. Caine V, Mill J, O'Brien K, et al. Implementation Process of a Canadian Community-based Nurse Mentorship Intervention in HIV Care. *J Assoc Nurses AIDS Care*. 2016;27(3):274–284. doi:10.1016/j.jana.2015.11.001
7. Burgess A, van Diggele C, Mellis C. Mentorship in the health professions: a review. *Clin Teach*. 2018;15:197–202
8. Arora S, Thornton K, Komaromy M, Kalishman S, Katzman J, Duhigg D. Demonopolizing medical knowledge. *Acad Med*. 2014;89:30–32.
9. Arora S, Thornton K, Murata G, et al. Outcomes of treatment for hepatitis C virus infection by primary care providers. *N Engl J Med*. 2011;364:2199–2207.
10. L. Bikinesi¹, N. Hamunime et al. Adaptation, implementation and evaluation of a tele-mentoring program for the Namibian HIV health workforce. <http://programme.aids2018.org/Abstract/Abstract/6455>

11. Zambia Statistics Agency, Ministry of Health Zambia, ICF. Zambia demographic and health survey 2018. Lusaka, Zambia, and Rockville, MD: Zambia Statistics Agency, Ministry of Health, and ICF; 2019.
12. Arora S, Kalishman S, Thornton K, et al. Expanding access to hepatitis C virus treatment—Extension for Community Health Outcomes (ECHO) project: disruptive innovation in specialty care. *Hepatology* 2010; 52(3):1124–33.
13. PEPFAR. Monitoring, evaluation and reporting indicator reference sheet version 2.3. 2018.
14. McChrystal, S. A., Collins, T., Silverman, D., & Fussell, C. (2015). Team of teams: New rules of engagement for a complex world.
15. Mbuagbaw L, Thabane L. How to set-up a long-distance mentoring program: a framework and case description of mentorship in HIV clinical trials. *J Multidiscip Healthc*. 2013;6:17–23. doi:10.2147/JMDH.S39731
16. Goosby E, Dybul M, Fauci AS, et al. The United States President's Emergency Plan for AIDS Relief: a story of partnerships and smart investments to turn the tide of the global AIDS pandemic [published correction appears in *J Acquir Immune Defic Syndr*. 2012 Oct 1;61(2):e24. Fauci, Anthony A [corrected to Fauci, Anthony S]]. *J Acquir Immune Defic Syndr*. 2012;60 Suppl 3:S51-S56. doi:10.1097/QAI.0b013e31825ca721
17. Reddy et al. South Africa field epidemiology training program: developing and building applied epidemiology capacity, 2007–2016 *MC Public Health* 2019, 19(Suppl 3):469
18. Matovu JK, Wanyenze RK, Mawemuko S, et al. Building capacity for HIV/AIDS program leadership and management in Uganda through mentored Fellowships. *Glob Health Action*. 2011;4:5815
19. Edwards LJ, Moises A, Nzaramba M. et al. Implementation of a health management mentoring program: year-1 evaluation of its impact on health system strengthening in Zambézia province, Mozambique. *Int J Health Policy Manag*. 2015;4(6):353–361. doi: 10.15171/ijhpm.2015.58.

Figures




Figure 1

Map of Zambia with HIV prevalence among adults aged 15-49 by province. Labels indicate the provinces in which the HIV clinical mentorship program was implemented 11.