

Is Age of Self-Harm Onset Associated With Increased Frequency of Non-Suicidal Self-Injury and Suicide Attempts in Adolescent Outpatients?

Anne Brager-Larsen (✉ a.t.brager-larsen@studmed.uio.no)

Oslo University Hospital

Pål Zeiner

Oslo University Hospital

Ole Klungsøyr

Oslo University Hospital

Lars Mehlum

University of Oslo

Research Article

Keywords: Self-harm, Suicide attempt, Non-suicidal self-injury (NSSI), Adolescents, Age of Onset, Borderline symptoms

Posted Date: February 5th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-153519/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at BMC Psychiatry on January 26th, 2022. See the published version at <https://doi.org/10.1186/s12888-022-03712-w>.

Abstract

Background

Self-harm in adolescents is an increasing public health concern and an important risk factor for suicide. We aimed to examine how age of self-harm onset in adolescents was associated with frequency of subsequent suicidal and non-suicidal self-harm (NSSI) episodes, and how age of onset and duration of self-harm may influence the frequency of self-harm.

Methods

Data from 103 adolescents with recurrent self-harm recruited from child and adolescent psychiatric outpatient clinics were collected through clinical interviews and self-reports, and analysed with negative binomial and hurdle models.

Results

A lower age of self-harm onset and a longer duration of self-harm were both significantly associated with increased frequency of subsequent episodes of NSSI and risk of a first suicide attempt. There was an increase in repeated suicide attempts when the age of onset of self-harm decreased and the duration increased, and dramatically more for long duration of NSSI before first suicide attempt.

Conclusion

Initiating self-harm behaviour at the youngest age had the highest risk of increased frequency of NSSI and suicide attempts. Longer duration of self-harm behaviour increased this risk. This underlines the importance of early identification of self-harm behaviour in adolescents, and the use of effective interventions.

Background

Suicide attempts and non-suicidal self-injury (NSSI) (collectively labelled “self-harm”) in adolescents are public health concerns (1, 2). Both suicide attempts and NSSI in adolescence are found to be robust risk factors for future suicidal behaviour (3–5). Self-harm behaviour increases up to age 12 and peaks between ages 14–16, with decreasing rates after 18 years of age (6, 7). Age of self-harm onset among adolescents is decreasing, while rates of self-harm are increasing (8, 9). The age with highest probability of NSSI onset is between 14 and 15 years, suggesting that adolescence is the most critical period for such behaviour (7). Determining whether an early age of self-harm onset is associated with a clinical course characterized by increasing self-harm behaviour, merits considerable interest. Understanding how different ages of self-harm onset may be associated with a later course of NSSI and suicide attempts,

could help us more specifically target risk groups in clinical settings. So far, only a limited number of studies have examined age of onset as a specific risk factor of NSSI and suicide attempts. Ammerman et al. (10), found that onset before the age of 12 was associated with an increased lifetime frequency of NSSI episodes and more severe self-harm behaviour. Whether the age of onset was totally or partially a function of duration, however, was not investigated. Muehlenkamp et al. (11), similarly found that self-harm onset before the age of 12 was associated with a significantly higher lifetime frequency of NSSI episodes even when controlled for duration. These studies were both based on general population samples and are not readily generalizable to adolescent clinical populations.

Studies focusing on duration of self-harm behaviour have found that longer duration is associated with increased frequency and severity of self-harm behaviour (12, 13). Although these studies did not address how the different ages of onset in children and adolescents may affect the frequency of self-harm behaviour, the results illustrate that the duration of self-harm is an important factor to consider when assessing age of self-harm onset in relation to frequency of NSSI and suicide attempts.

Adolescence is an important developmental time-period and the prevalence of many psychiatric illnesses increases sharply from childhood to adolescence (14), as is also the case with self-harm behaviour, making adolescence a very important time-period for prevention. However, we still lack knowledge on how initiating self-harm behaviour at earlier or later ages may give rise to different self-harm trajectories. Thus, there is a need for studies to investigate if the age of self-harm onset and duration of self-harm, affect the following clinical course of self-harming behaviour.

To investigate if age of onset and duration, uniquely and in interaction, are associated with frequency of NSSI and suicide attempts in an adolescent psychiatric population, it is essential to also consider other relevant factors which may influence these associations. We know that factors such as history of sexual or physical abuse, (15), relationship with parents, (16, 17), and borderline symptoms, (18, 19), are strongly associated with both NSSI and suicide attempts, and age of onset. We also know that these factors are commonly observed in clinical adolescent populations (20–22). What we do not know is whether these factors affect how age of self-harm onset and duration of self-harm are associated with frequency of NSSI and risk of suicide attempts.

Given the current state of knowledge, we wanted to study if age of self-harm onset is associated with increased lifetime frequency of NSSI episodes, uniquely and in interaction with duration of self-harm, and adjusted for covariates. We also wanted to study if age of self-harm onset, uniquely and in interaction with duration of self-harm, is associated with an increased risk of having made a suicide attempt and of having repeated suicide attempts, adjusted for covariates.

Method

Participants and procedures

Participants were 103 adolescents (age 12–18) recruited from a child and adolescent psychiatric outpatient clinic at Oslo University Hospital, Norway, serving a catchment area in the city of Oslo. Adolescents were screened for self-harm behaviour. Self-harm was defined as “intentional poisoning or self-injury, regardless of intention to die” (23). Suicide attempt was defined as “a potentially self-injurious act committed with at least some wish to die, as a result of the act” (24), while non-suicidal self-injury (NSSI) was defined as “the deliberate, self-inflicted destruction of body tissue without suicidal intent, and for purposes not socially sanctioned” (25). Inclusion criteria were recurrent self-harm behaviour (two or more episodes) with the last episode having occurred within the past 6 months. Exclusion criteria were mental retardation or insufficient Norwegian language skills to understand or answer the interviews or questionnaires. Adolescents and their parents provided written informed consent prior to inclusion. The study was approved by the Regional Committee for Medical Research Ethics, South-East Norway. The interviewers were experienced clinicians, having received training and supervision in the use of the study instruments.

Measures

The *Suicide Attempt Self-Injury Interview (SASII)* (26), was used to collect data on age of self-harm onset, frequency of non-suicidal self-injury episodes (NSSI), first and lifetime suicide attempts and methods in use. The *Schedule for Affective Disorders-Present and Lifetime version, 2013 (K-SADS-PL)* (27), was used to obtain socio-demographic data and DSM-5 Axis I diagnoses. *Previous experience of sexual and physical abuse* was assessed using the CARE instrument (28). *Childhood interview for Borderline Personality Disorder (CI-BPD)*, (29) was used to assess BPD criteria based on *DSM-5*. The self-report *Borderline Symptom List (BSL-23)* (30) was used to assess borderline-specific symptoms. *Attachment to Parents and Peers* was measured through a shortened version of the self-report *Inventory of Parental and Peer Attachment (IPPA)* (31).

Statistics

Means and standard deviations or median and interquartile ranges were computed for normally and non-normally distributed variables respectively (Table 2). As is commonly observed in clinical studies of self-harm, number of NSSI episodes and suicide attempts showed over dispersion leading to more zeros and heavier tails than in a Poisson-model (32).

Therefore, for NSSI as outcome, a negative binomial regression model was fitted and the results are presented in both table and plot of expected frequency of NSSI episodes. For first and repeated suicide attempts as outcome, a two-component “hurdle” model was fitted (32–34). One component models zero versus positive counts, where a logistic regression model (logistic link) was employed. The other component is a truncated count component for positive counts, where a negative binomial regression (log link) model was employed. Regression results from each component are presented in tables. Expected frequency of suicide attempts was calculated based on the results from the total model (both first and repeated suicide attempts), and represented in a plot diagram. Significant covariates in either component will be significant in the total hurdle model. Maximum likelihood estimation was achieved

with the function `glm.nb()` from the R-package MASS and the function `hurdle()` in the R-package pscl (35).

The variation in age at time of interview made it possible to model both the effect of age of onset and duration of self-harm (Fig. 1a). For illustrative purposes, we assessed group differences in time to first suicide attempt and median NSSI frequency (Fig. 1b and 1c).

Age of onset was defined as the age (years) at the time of first self-harm episode. Total duration of self-harm was defined as the time-span between the age of onset and the time of interview. When analysing risk of suicide attempts, duration was defined as “duration of NSSI before first suicide attempt” (if any). Age at time of interview was considered to reflect total frequency of years under risk, and was entered as an offset-term in all regression models.

Adjusting for covariates

Age of onset and duration could be analysed with respect to “mediation” (indirect/direct effects) and “effect-modification” (interaction), but would rely on a complete set of measured confounders (36, 37). With duration as a mediator between age of onset and suicide attempts, confounders for duration and suicide attempt association should be adjusted for (37). In the present study, no attempt was made to do causal inference, but rather to indicate mechanisms, and the associations found must be interpreted with this in mind. Selected covariates that may influence the association between the age of onset, separately and in interaction with duration, as well as the outcome variables, were sexual and physical abuse, borderline symptoms (BSL) and relationship with parents. These covariates are thought to precede age of self-harm onset and duration of self-harm. All demographic and clinical characteristics were analysed using the IBM SPSS statistics version 25, on descriptives and frequencies. All regression analyses and figures were conducted with the R package R Core Team (38), and the significance level was set to .05.

Results

Sample characteristics

A total of 103 adolescents with a mean age of 15.9 years (SD = 1.47) were enrolled in the study between January 2017 and June 2019. Sociodemographic characteristics are shown in Table 1. Mean number of DSM-5 diagnoses was 3.2 (range = 0–8) with mood disorders (84.5%) and anxiety disorders (73.8%) as the most frequent. Nearly one third (28.2%) of the adolescents fulfilled diagnostic criteria for Borderline personality disorder.

Table 1
Sociodemographic and clinical characteristics of adolescents with repetitive self-harm (N = 103)

	n	%
Gender (females)	89	86.4
Age, mean (years) (SD = 1.47) 15.9		
Born in Norway	87	84.5
One or both parents born in foreign countries	43	41.7
Living with	39	37.9
- both parents		
- alternating between parents	15	14.6
- single parent	34	33.0
Current DSM-5 diagnosis¹		
Mood Disorder, any	87	84.5
Psychotic Disorder, any	13	12.6
Anxiety Disorder, any	76	73.8
Eating Disorder, any	18	17.5
Substance-Use Disorder, any	10	9.7
ADHD, any	24	23.3
Borderline Personality Disorder	29	28.2
Other diagnosis ²	27	26.2
¹ Not mutually exclusive categories		
² Conduct disorder, Oppositional defiant disorder, Unspecified disruptive behaviour disorder, Chronic motor or Vocal tic disorder, Autism spectrum disorder, Adjustment disorder		
Note: ADHD = Attention-deficit/hyperactivity disorder.		

Characteristics of the self-harm behaviour

Distribution of self-harm characteristics and selected covariates are presented in Table 2. The median age of self-harm onset was 13.2 years (range 7.1–17.4), and the median duration of self-harm was 2.5 years (range 0.1–10.3) (Fig. 1a). The lifetime frequency of NSSI episodes ranged from 0 to 990 (median = 50; IQR = 128; 75 percentile = 142). Only one adolescent reported having had no NSSI episodes, but this participant reported repeated suicide attempts. The lifetime number of methods used for self-harm varied from 1 to 10. Cutting with a sharp object was the most commonly reported NSSI method (93.2%),

followed by self-battery (41.7%), and stabbing (16.3%). Fifty adolescents (48.5%) reported at least one lifetime suicide attempt, of which twenty-seven (26.2% of the total sample) repeated the attempt at least once, with cutting (21.4%), drug overdose (21.4%) and asphyxia (14.6%) as the most frequently reported methods.

Age of self-harm onset and duration of self-harm by frequency of NSSI

A 3D-barplot of the median lifetime NSSI episodes in three categories of age of onset against four categories of duration of self-harm behaviour is shown in Fig. 1c. For each category of age of onset, an increasing median lifetime frequency of NSSI episodes was observed for increasing duration of self-harm, except for high values of age of onset and duration. The adjusted regression analysis, (Table 3), showed that age of onset, interaction between age of onset and duration, and their squared terms all remained significantly associated with the frequency of NSSI episodes.

Table 2 Self-harm characteristics and selected covariates in adolescents with repetitive self-harm (N=103)

	Mean	Std.Dev	Median	Interquartile range
Self-harm characteristics				
Age of onset (years)	13.1	1.8	13.2	1.9
Duration (years)	2.8	2.1	2.5	2.9
Suicide attempts, lifetime	1.8	4.4	0.0	2.0
NSSI, lifetime	119.7	174.3	50.0	128.0
Duration before first SA	2.2	1.9	1.6	6.5
NSSI before first SA	89.7	134.9	34.5	90.0
Number of methods (first SH episode)	1.1	0.2	1.0	0.0
Number of methods (lifetime)	2.7	1.6	2.0	2.0
Selected covariates ¹				
Borderline Symptoms (BSL-23)	35.15	23.07	31.0	39.0
Sexual abuse	0.31	0.46	0	1.0
Physical abuse	0.33	0.47	0	1.0
Relationship with mother	32.83	10.15	35.0	17.0
Relationship with father	31.50	10.20	33.0	12.5
¹ Potential confounders SA=Suicide attempt SH=Self harm				

Table 3 Negative binomial regression model for NSSI lifetime episodes with age of self-harm onset, duration of self-harm and selected covariates, in adolescents with repetitive self-harm (N=103).

	Estimate	St. Error	z value	p-value	95% CI	
(Intercept)	2.95702	0.4334	6.823	<0.001 ***	2.06727	3.91199
Relationship with mother	-0.02548	0.01106	-2.304	0.021222 *	-0.04977	-0.00193
Physical abuse	-0.11589	0.24865	-0.466	0.64116	-0.67141	0.45547
Sexual abuse	0.43374	0.25855	1.678	0.093426 .	-0.09431	0.98325
Age of onset	-0.19883	0.09587	-2.074	0.038082 *	-0.39068	-0.01305
Duration of self-harm	0.15894	0.08739	1.819	0.068954 .	-0.00692	0.32822
Age of onset - squared	-0.19423	0.05803	-3.347	0.000816 ***	-0.30633	-0.07348
Duration - squared	-0.19337	0.0556	-3.478	0.000505 ***	-0.31399	-0.06109
Age of onset x Duration (Interaction)	-0.36343	0.10241	-3.549	0.000387 ***	-0.57832	-0.12634

*p < .05; **p < .01; ***p < .001.

The expected frequency of mean NSSI episodes were plotted against age of self-harm onset and duration of self-harm (Fig. 2a). The figure shows that there was an increase in frequency of NSSI episodes for decreasing age of onset and increasing duration, except for short duration, where the frequency of NSSI increased and then decreased for lower values of age of onset. The drop in the front of the plot for small values of age of onset and duration of self-harm in Fig. 2a corresponds with the upper and inner left corner in Fig. 1c.

Age of self-harm onset and duration of NSSI on risk by suicide attempts

A Kaplan-Meier plot of age at first suicide attempt, stratified by three categories of age of onset of self-harm (Fig. 1b), illustrates that an early age of self-harm onset was associated with an earlier first suicide attempt (unadjusted). Adjusted regression results (Table 4) show that both age of onset and duration of NSSI before first suicide attempt, separately and in interaction with each other, were significantly associated with probability of a first suicide attempt. With regard to repeated suicide attempts, duration of NSSI before the first suicide attempt was strongly associated with increased frequency of suicide attempts (p = 0.059). Age of onset was not significantly associated with repeated suicide attempts in this analysis.

Table 4

Fitted two-component hurdle model for first suicide attempt (logit link) and repeated suicide attempts (SA) (log link) with age of self-harm onset, duration of self-harm and selected covariates, in adolescents with repetitive self-harm (N = 103)

First suicide attempt						
	Estimate	St. Error	z value	p-value	95% CI	
(Intercept)	-2.04006	0.54001	-3.778	0.000158 ***	-3.09847	-0.98164
Borderline Symptoms (BSL-23)	0.04569	0.01213	3.767	0.000165 ***	0.02191	0.06946
Age of onset	-0.67436	0.21059	-3.202	0.001364 **	-1.08711	-0.26160
Duration before first SA	-0.63442	0.19544	-3.246	0.001170 **	-1.01748	-0.25135
Age of onset x Duration before first SA (Interaction)	-0.1877	0.07931	-2.367	0.017951 *	-0.34314	-0.03225
Repeated suicide attempts						
	Estimate	Std. Error	z value	p-value	95% CI	
(Intercept)	-5.64526	0.94526	-5.972	2.34e-09 ***	-7.49796	-3.79255
Borderline Symptoms (BSL-23)	0.04552	0.0113	4.028	5.62e-05 ***	0.02337	0.06766
Sexual abuse	1.13342	0.5182	2.187	0.0287 *	0.11774	2.14909
Duration before first SA	-0.08972	0.12136	-0.739	0.4597	-0.32758	0.14814
Duration before first SA - squared (interaction)	0.14685	0.07784	1.887	0.0592 .	-0.00571	0.29941
*p < .05; **p < .01; ***p < .001.						

Both age of onset and duration of self-harm were significantly associated with the expected frequency of suicide attempts, in the combined hurdle model (Fig. 2b). The figure shows that the expected frequency of suicide attempts increased with increased duration of NSSI before first suicide attempt, for all low and moderate values of age of onset, but dramatically more for lower age of onset. In addition, the frequency of suicide attempts increased with decreasing age of onset when the duration of NSSI before first suicide attempt increased, and dramatically more for longer duration before first suicide attempt.

The selected covariates were considered potential confounding factors, and were adjusted for in all analysis. With respect to frequency of NSSI episodes, a positive relationship with mother was

significantly associated with a decrease in the frequency of NSSI and with increasing age of onset (Table 3). Higher scores on borderline symptom list (BSL) were significantly associated with earlier age of onset, longer duration of self-harm and higher risk of a first suicide attempt, as well as repeated suicide attempts (Table 4), while, sexual abuse was significantly associated with lower age of onset and increased frequency of suicide attempts.

Discussion

Our study is the first to examine whether age of onset, uniquely and in interaction with duration of self-harm is associated with frequency of NSSI and risk of suicide attempts in a clinical adolescent population. The main findings were that both earlier age of onset and longer duration were associated with increased frequency of NSSI and suicide attempts and this co-varies in such a way that those who started self-harming at the youngest age, had the highest risk of further self-harm, and that an increasing duration of self-harm behaviour amplifies this risk.

Age of onset, duration and NSSI

Our findings are compatible with findings in adult populations (10, 11) indicating that an earlier age of onset is a potential risk factor for increased lifetime frequency of NSSI, even when controlled for duration and selected covariates. It would seem self-evident that a longer duration of self-harm leads to more NSSI episodes irrespective of the age of onset. However, we found that among adolescents with similar duration but different age of onset, the highest frequency of NSSI episodes was found in those who started at an earlier age.

We also found that a positive relationship with mother was significantly associated with a decrease in the frequency of NSSI episodes and increasing age of onset. This finding corresponds with previous studies indicating that perceived family support predicted termination of NSSI, and that negative perceptions of parents were related to a higher frequency of NSSI (17, 39).

Age of onset, duration and suicide attempts

About half of our participating adolescents reported having had at least one suicide attempt. Suicide attempts were more likely in participants with a lower age of self-harm onset and a longer duration. This is in line with recent research on adults (11, 40) indicating that the risk of suicide attempts increases with earlier age of self-harm onset. These studies are, however, based on adult samples and are not readily comparable with a clinical sample of adolescents.

Mental disorders are associated with an increased risk of suicidal behaviour in adolescents (20). In addition, mental illness combined with ongoing self-harm, further increases risk of suicide attempts (3). In our study, the average number of DSM-5 diagnoses was 3.2, (range 0–8), and 84.5% reported symptoms consistent with a depressive disorder. However, we did not find significant associations

between early age of onset and number of diagnoses, nor between age of first depressive episode and suicide attempts (preliminary analyses).

Age of onset, duration and repeated suicide attempts

About a fourth of participating adolescents reported repeating the suicide attempt at least once. When we examined the expected number of suicide attempts, we found an increase in suicide attempts when the age of onset decreased and the duration increased, and dramatically more for long duration of NSSI before first suicide attempt. This is consistent with a possible direct effect of age of onset. This would mean that if you compare two adolescents with self-harm behaviour and equal duration of NSSI (before the first suicide attempt), it would be the adolescent with earlier age of onset that has the highest expected number of suicide attempts. Furthermore, the number of suicide attempts increased with longer duration of NSSI (before the first suicide attempt) for early and moderate onset age, and dramatically more for early onset age (possible effect modification from age of onset on the effect of duration). Likewise, if you compare two adolescents, with self-harm behaviour and equal age of self-harm onset, then based on our findings, the adolescent with the longer duration of NSSI (before the first suicide attempt) would have the highest expected number of suicide attempts.

These results are in line with previous research suggesting that longer duration of NSSI is associated with increased risk of suicide attempts (41, 42), that suicide attempts are one of the most important risk factors for re-attempts (43), and that the highest risk of re-attempts is relatively shortly after the first suicide attempt (43). A possibility as to why NSSI elevates the risk of suicide attempts was made by Joiner (2005). He hypothesised that the transition from NSSI to suicide attempts may be explained by an acquired capability for suicide attempts through repetition of self-harm over time. Studies that confirm this theory find that increased risk of suicide attempts is associated with reduced pain and fear (41), and an increased belief in own capabilities to do what is needed (44).

In addition, we found that a higher level of borderline symptoms among these adolescents was associated with an increased risk of suicide attempts (first and re-attempts). This finding corresponds with previous studies indicating that borderline symptoms is associated with greater severity of self-harm (45) and suggests that it would be important to assess borderline symptoms in adolescents with recurrent self-harm behaviour as a part of suicide risk assessment and management (46–48). Certainly, it could assist clinicians in selecting treatments specifically designed to address these symptoms and challenges in adolescents (49–52). Finally, we found that having been sexually abused was significantly associated with an increased risk of repeated suicide attempts. Due to small numbers, this finding must be interpreted with caution, but it corresponds well with numerous other studies having made the same finding (3).

Strengths and limitations

A strength of this study is that it is based on a clinical sample of adolescents with recurrent self-harm. Among limitations are the study's cross-sectional design precluding causal inference, the retrospective retrieval of information entailing risks of memory and information biases, and the limited number of

males included in the sample preventing us from generalizing these findings to males. Due to the nature of the age inclusion criteria, we lack data beyond older age of onset for the oldest adolescents. This is natural given the time horizon of the current study, and is important to take into consideration while interpreting these results. Future studies with more diverse samples and with a longitudinal design would improve confidence in the results, as well as generalizability.

Conclusions

Despite the mentioned limitations, our study suggests that both an early age of self-harm onset and a longer duration of self-harm may entail an increased frequency of further NSSI and suicide attempts in adolescents with recurrent self-harm, and that the youngest are particularly at risk of increased self-harm, especially when the duration increases. Recurrent self-harm over time may lead to an acquired capability for suicide attempts, and emphasizes the importance of early detection and interventions adapted to both age and severity.

Abbreviations

NSSI

Non-Suicidal Self-Injury; SA:Suicide Attempt; SH:Self-Harm; BSL-23:Borderline Symptom List

Declarations

Acknowledgements

The authors wish to thank the patients, clinicians and leaders at the participating clinics. They also thank Grete Sundal, Marit Mytting Kristin F. Moe and Lise M. Hassel for conducting assessment interviews; and Grethe Hestnes, Aina L. Hansen, Christian Lie, Grethe Borgersen and Nazia Ahmad, for administrative support.

Ethics approval and consent to participate

Adolescents and their parents provided written informed consent prior to inclusion. The study was approved by the Regional Committee for Medical Research Ethics, South-East Norway. All methods were carried out in accordance with relevant guidelines and regulations

Consent for publication

Covered by information to patients approved by Regional Committee for Medical Research, South-East Norway.

Availability of data and materials

The datasets analysed during the current study are available from the corresponding author on reasonable request.

Competing interests

The authors declare that they have no competing interests.

Funding

The study received funding from the Oslo University Hospital and University of Oslo and by Olafstiftelsen for Research on suicidality among children and adolescents. These funds helped finance staff for data collection. None of the funders interfered with the design of the study and collection, analysis and interpretation of data and in writing the manuscript.

Authors' contributions

AB-L wrote the main manuscript and participated in the conception and developing the design of the study, acquisition of data, and participated in analysis and interpretation of data. OK served as the statistical expert and a major contributor with statistical analysis, tables, figures and interpretation of data. PZ and LM was major contributors in conception and developing the design of the study, research questions and protocol, interpretation of data, major contributors in writing the manuscript, and substantively revised the work. All authors read and approved the final manuscript.

Authors' information (optional)

¹ Oslo University Hospital, Child and Adolescent Mental Health Research Unit, Department of Research and Innovation, Division of Mental Health and Addiction, Norway.² Oslo University Hospital, Section for treatment research, Department of Research and Innovation, Division of Mental Health and Addiction, Norway. ³University of Oslo, National Centre for Suicide Research and Prevention, Institute of Clinical Medicine, Norway.

Questionnaires and interviews

We have only used standard questionnaires and interviews that have been previously published elsewhere, and in according to guidelines.

References

1. Jacobson CM, Gould M. The epidemiology and phenomenology of non-suicidal self-injurious behavior among adolescents: a critical review of the literature. *Archives of suicide research : official journal of the International Academy for Suicide Research*. 2007;11(2):129-47.
2. Hawton K, Saunders KE, O'Connor RC. Self-harm and suicide in adolescents. *Lancet (London, England)*. 2012;379(9834):2373-82.

3. Victor SE, Klonsky ED. Correlates of suicide attempts among self-injurers: a meta-analysis. *Clinical psychology review*. 2014;34(4):282-97.
4. Tormoen AJ, Rossow I, Larsson B, Mehlum L. Nonsuicidal self-harm and suicide attempts in adolescents: differences in kind or in degree? *Social psychiatry and psychiatric epidemiology*. 2013;48(9):1447-55.
5. Castellvi P, Lucas-Romero E, Miranda-Mendizabal A, Pares-Badell O, Almenara J, Alonso I, et al. Longitudinal association between self-injurious thoughts and behaviors and suicidal behavior in adolescents and young adults: A systematic review with meta-analysis. *Journal of affective disorders*. 2017;215:37-48.
6. Plener PL, Schumacher TS, Munz LM, Groschwitz RC. The longitudinal course of non-suicidal self-injury and deliberate self-harm: a systematic review of the literature. *Borderline personality disorder and emotion dysregulation*. 2015;2:2.
7. Gandhi A, Luyckx K, Baetens I, Kiekens G, Sleuwaegen E, Berens A, et al. Age of onset of non-suicidal self-injury in Dutch-speaking adolescents and emerging adults: An event history analysis of pooled data. *Comprehensive psychiatry*. 2018;80:170-8.
8. Griffin E, McMahon E, McNicholas F, Corcoran P, Perry IJ, Arensman E. Increasing rates of self-harm among children, adolescents and young adults: a 10-year national registry study 2007-2016. *Social psychiatry and psychiatric epidemiology*. 2018;53(7):663-71.
9. Tørmoen AJ, Myhre M, Walby FA, Grøholt B, Rossow I. Change in prevalence of self-harm from 2002 to 2018 among Norwegian adolescents. *Eur J Public Health*. 2020;30(4):688-92.
10. Ammerman BA, Jacobucci R, Kleiman EM, Uyeji LL, McCloskey MS. The Relationship Between Nonsuicidal Self-Injury Age of Onset and Severity of Self-Harm. *Suicide & life-threatening behavior*. 2018;48(1):31-7.
11. Muehlenkamp JJ, Xhunga N, Brausch AM. Self-injury Age of Onset: A Risk Factor for NSSI Severity and Suicidal Behavior. *Archives of suicide research : official journal of the International Academy for Suicide Research*. 2019;23(4):551-63.
12. Victor SE, Styer D, Washburn JJ. Functions of nonsuicidal self-injury (NSSI): Cross-sectional associations with NSSI duration and longitudinal changes over time and following treatment. *Psychiatry research*. 2016;241:83-90.
13. Voss C, Ollmann TM, Miché M, Venz J, Hoyer J, Pieper L, et al. Prevalence, Onset, and Course of Suicidal Behavior Among Adolescents and Young Adults in Germany. *JAMA network open*. 2019;2(10):e1914386.
14. Kessler RC, Chiu WT, Demler O, Merikangas KR, Walters EE. Prevalence, severity, and comorbidity of 12-month DSM-IV disorders in the National Comorbidity Survey Replication. *Archives of general psychiatry*. 2005;62(6):617-27.
15. Ford JD, Gomez JM. The relationship of psychological trauma and dissociative and posttraumatic stress disorders to nonsuicidal self-injury and suicidality: a review. *Journal of trauma & dissociation* :

- the official journal of the International Society for the Study of Dissociation (ISSD). 2015;16(3):232-71.
16. Horváth LO, Győri D, Komáromy D, Mészáros G, Szentiványi D, Balázs J. Nonsuicidal Self-Injury and Suicide: The Role of Life Events in Clinical and Non-Clinical Populations of Adolescents. *Frontiers in psychiatry*. 2020;11:370.
 17. Kaess M, Parzer P, Mattern M, Plener PL, Bifulco A, Resch F, et al. Adverse childhood experiences and their impact on frequency, severity, and the individual function of nonsuicidal self-injury in youth. *Psychiatry research*. 2013;206(2-3):265-72.
 18. Jacobson CM, Muehlenkamp JJ, Miller AL, Turner JB. Psychiatric impairment among adolescents engaging in different types of deliberate self-harm. *Journal of clinical child and adolescent psychology : the official journal for the Society of Clinical Child and Adolescent Psychology, American Psychological Association, Division 53*. 2008;37(2):363-75.
 19. Ramleth RK, Groholt B, Diep LM, Walby FA, Mehlum L. The impact of borderline personality disorder and sub-threshold borderline personality disorder on the course of self-reported and clinician-rated depression in self-harming adolescents. *Borderline personality disorder and emotion dysregulation*. 2017;4:22.
 20. Bjureberg J, Ohlis A, Ljótsson B, D'Onofrio BM, Hedman-Lagerlöf E, Jokinen J, et al. Adolescent self-harm with and without suicidality: cross-sectional and longitudinal analyses of a Swedish regional register. *Journal of child psychology and psychiatry, and allied disciplines*. 2019;60(3):295-304.
 21. Kaess M, Fischer-Waldschmidt G, Resch F, Koenig J. Health related quality of life and psychopathological distress in risk taking and self-harming adolescents with full-syndrome, subthreshold and without borderline personality disorder: rethinking the clinical cut-off? *Borderline personality disorder and emotion dysregulation*. 2017;4:7.
 22. Carballo JJ, Llorente C, Kehrmann L, Flamarique I, Zuddas A, Purper-Ouakil D, et al. Psychosocial risk factors for suicidality in children and adolescents. *European child & adolescent psychiatry*. 2020;29(6):759-76.
 23. Hawton K. Psychiatric assessment and management of deliberate self-poisoning patients. *Medicine (Abingdon 1995, UK ed)*. 2007;35(10):565-9.
 24. Posner K, Melvin GA, Stanley B, Oquendo MA, Gould M. Factors in the assessment of suicidality in youth. *CNS spectrums*. 2007;12(2):156-62.
 25. Klonsky ED. The functions of deliberate self-injury: a review of the evidence. *Clinical psychology review*. 2007;27(2):226-39.
 26. Linehan MM, Comtois KA, Brown MZ, Heard HL, Wagner A. Suicide Attempt Self-Injury Interview (SASII): development, reliability, and validity of a scale to assess suicide attempts and intentional self-injury. *Psychol Assess*. 2006;18(3):303-12.
 27. Kaufman J, Birmaher B, Brent D, Rao U, Flynn C, Moreci P, et al. Schedule for Affective Disorders and Schizophrenia for School-Age Children-Present and Lifetime Version (K-SADS-PL): initial reliability

- and validity data. *Journal of the American Academy of Child and Adolescent Psychiatry*. 1997;36(7):980-8.
28. Chaffin M, Wherry JN, Newlin C, Crutchfield A, Dykman R. The Abuse Dimensions Inventory: Initial Data on a Research Measure of Abuse Severity. *Journal of Interpersonal Violence*. 1997;12(4):569-89.
 29. Sharp C, Ha C, Michonski J, Venta A, Carbone C. Borderline personality disorder in adolescents: evidence in support of the Childhood Interview for DSM-IV Borderline Personality Disorder in a sample of adolescent inpatients. *Comprehensive psychiatry*. 2012;53(6):765-74.
 30. Bohus M, Limberger MF, Frank U, Chapman AL, Kuhler T, Stieglitz RD. Psychometric properties of the Borderline Symptom List (BSL). *Psychopathology*. 2007;40(2):126-32.
 31. Armsden GC, Greenberg MT. The inventory of parent and peer attachment: individual differences and their relationship to psychological well-being in adolescence. *Journal of youth and adolescence*. 1987;16(5):427-54.
 32. Bethell J, Rhodes AE, Bondy SJ, Lou WYW, Guttman A. Repeat self-harm: application of hurdle models. *The British journal of psychiatry : the journal of mental science*. 2010;196(3):243-4.
 33. Mullahy J. Specification and testing of some modified count data models. *Journal of econometrics*. 1986;33(3):341-65.
 34. Glenn CR, Lanzillo EC, Esposito EC, Santee AC, Nock MK, Auerbach RP. Examining the Course of Suicidal and Nonsuicidal Self-Injurious Thoughts and Behaviors in Outpatient and Inpatient Adolescents. *J Abnorm Child Psychol*. 2017;45(5):971-83.
 35. Zeileis A, Kleiber C, Jackman S. Regression Models for Count Data in R. *Journal of statistical software*. 2008;27(8).
 36. VanderWeele TJ. Explanation in causal inference: developments in mediation and interaction. *Int J Epidemiol*. 2016;45(6):dyw277-1908.
 37. Cole SR, Hernán MA. Fallibility in estimating direct effects. *Int J Epidemiol*. 2002;31(1):163-5.
 38. R: A language and environment for statistical computing [Internet]. R Foundation for Statistical Computing. Vienna, Austria. 2020. Available from: <https://.R-project.org/> aug.11,2020.
 39. Tatnell R, Hasking P, Newman L, Taffe J, Martin G. Attachment, Emotion Regulation, Childhood Abuse and Assault: Examining Predictors of NSSI Among Adolescents. *Archives of suicide research : official journal of the International Academy for Suicide Research*. 2017;21(4):610-20.
 40. O'Connor RC, Wetherall K, Cleare S, Eschle S, Drummond J, Ferguson E, et al. Suicide attempts and non-suicidal self-harm: national prevalence study of young adults. *BJPsych open*. 2018;4(3):142-8.
 41. Nock MK, Joiner TE, Jr., Gordon KH, Lloyd-Richardson E, Prinstein MJ. Non-suicidal self-injury among adolescents: diagnostic correlates and relation to suicide attempts. *Psychiatry research*. 2006;144(1):65-72.
 42. Willoughby T, Heffer T, Hamza CA. The link between nonsuicidal self-injury and acquired capability for suicide: A longitudinal study. *Journal of abnormal psychology*. 2015;124(4):1110-5.

43. Irigoyen M, Porrás-Segovia A, Galvan L, Puigdevall M, Giner L, De Leon S, et al. Predictors of re-attempt in a cohort of suicide attempters: A survival analysis. *Journal of affective disorders*. 2019;247:20-8.
44. Klonsky ED, Qiu T, Saffer BY. Recent advances in differentiating suicide attempters from suicide ideators. *Current opinion in psychiatry*. 2017;30(1):15-20.
45. Ayodeji E, Green J, Roberts C, Trainor G, Rothwell J, Woodham A, et al. The influence of personality disorder on outcome in adolescent self-harm. *The British journal of psychiatry : the journal of mental science*. 2015;207(4):313-9.
46. Witt K, Milner A, Spittal MJ, Hetrick S, Robinson J, Pirkis J, et al. Population attributable risk of factors associated with the repetition of self-harm behaviour in young people presenting to clinical services: a systematic review and meta-analysis. *European child & adolescent psychiatry*. 2019;28(1):5-18.
47. Fonagy P, Speranza M, Luyten P, Kaess M, Hessels C, Bohus M. ESCAP Expert Article: borderline personality disorder in adolescence: an expert research review with implications for clinical practice. *European child & adolescent psychiatry*. 2015;24(11):1307-20.
48. Consoli A, Garny de La Riviere S, Gerardin P, Cohen D, Mirkovic B, Aouidad A, et al. Borderline personality disorder and prior suicide attempts define a severity gradient among hospitalized adolescent suicide attempters. *BMC psychiatry*. 2020;20(1):525-.
49. Mehlum L, Tormoen AJ, Ramberg M, Haga E, Diep LM, Laberg S, et al. Dialectical behavior therapy for adolescents with repeated suicidal and self-harming behavior: a randomized trial. *Journal of the American Academy of Child and Adolescent Psychiatry*. 2014;53(10):1082-91.
50. Rossouw TI, Fonagy P. Mentalization-Based Treatment for Self-Harm in Adolescents: A Randomized Controlled Trial. *Journal of the American Academy of Child and Adolescent Psychiatry*. 2012;51(12):1304-13.e3.
51. Sharp C, Fonagy P. Practitioner Review: Borderline personality disorder in adolescence - recent conceptualization, intervention, and implications for clinical practice. *Journal of child psychology and psychiatry, and allied disciplines*. 2015;56(12):1266-88.
52. Mehlum L, Ramleth RK, Tormoen AJ, Haga E, Diep LM, Stanley BH, et al. Long term effectiveness of dialectical behavior therapy versus enhanced usual care for adolescents with self-harming and suicidal behavior. *Journal of child psychology and psychiatry, and allied disciplines*. 2019.

Figures




Figure 1

a) Variation in age of self-harm onset, for each level of duration of self-harm. b) Kaplan-Meyer plot of age at first suicide attempt for three strata of age of self-harm onset, c) 3d bar plot with group median NSSI lifetime episodes for three strata age of self-harm onset and four strata duration of self-harm, in adolescents with repetitive self-harm (N=103)




Figure 2

a) Expected frequency of NSSI lifetime episodes and b) Expected suicide attempts lifetime (prediction) (z-axis) as a function of age of self-harm onset and duration of self-harm, in adolescents with repetitive self-harm (N=103).