

Predictors and histological effects of preoperative chemoradiotherapy for rectal cancer and control of lateral lymph node metastasis

Hiroshi Miyakita (✉ mykt0124@tokai-u.jp)

Tokai University

Lin Fung Chan

Tokai University

Kazutake Okada

Tokai University

Hajime Kayano

Tokai University

Masaki Mori

Tokai University

Sotaro Sadahiro

Tokai University

Seiichiro Yamamoto

Tokai University

Research Article

Keywords: chemoradiotherapy, lateral lymph node, rectal cancer

Posted Date: April 12th, 2022

DOI: <https://doi.org/10.21203/rs.3.rs-1535431/v1>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

[Introduction] Standard treatment strategy for low rectal cancer in Japan is different from Western countries. Total mesorectum excision (TME) + lateral lymph node dissection (LLND) is mainly carried out in Japan, whereas neoadjuvant chemoradiotherapy (nCRT) + TME is selected in Western countries. There is no clear definition of preoperative diagnosis of lateral lymph node metastasis. If we can predict lateral lymph node swelling that can be managed by nCRT from lateral lymph node swelling that require surgical resection, clinical benefit is significant. In the current study we assessed characteristics of the lateral lymph node recurrence (LLNR) and LLND that can be managed by nCRT.

[Patients and Methods] Patients with low rectal cancer (n=168) underwent nCRT between 2009 and 2016. We evaluated CEA, neutrophil / lymphocyte ratio (NLR), platelet / lymphocyte ratio (PLR), and lateral lymph node short axis pre and post nCRT, respectively, and also evaluated tumor shrinkage rate, tumor regression grade (TRG). We evaluated the relationship between each and LLNR.

[Results] LLND was not carried out all patients. Factors associated with LLNR were PLR and lymph node short axis pre and post nCRT. ($p=0.0269$, 0.0278 , $p\leq 0.0001$, $p\leq 0.0001$, respectively). Positive recurrence cut-off values of lateral lymph node short-axis calculated were 11.6 mm pre nCRT and 5.5 mm post nCRT.

[Conclusion] Results suggest that PLR before and after CRT was associated with control of LLNR, and LLND should be performed on lateral lymph nodes with short-axis of 5 mm and 11 mm pre and post nCRT.

Introduction

A standard of care for low rectal cancer is different between Japan and Western countries. While total mesorectum excision (TME) combined with lateral lymph node dissection (LLND) is the golden standard in Japan, (1) neoadjuvant chemoradiotherapy (nCRT) combined with TME is the first choice in Western countries. LLND is usually not carried out in Western countries since lateral lymph node metastasis is considered remote metastasis. (2) Previous studies showed that combined nCRT and LLND for low rectal cancer was effective to reduce local recurrence. (3,4) and it became a therapeutic option to reduce local recurrence. However, there are no established criteria for LLND indication in which thresholds sizes of lateral lymph node were varied such as 5 or 8 mm in short-axis diameter, or 8 mm in long-axis diameter. (5–9)

There are studies comparing the incidence of lateral lymph node metastases between nCRT + TME + LLND and TME + LLND. Patients who underwent LLND had a positive lymph node metastasis rate of 52.4% and a cut-off value for the size of the lymph node was 8 mm. And the study suggests that the resected lateral lymph node microscopic residual cancer cells were probably destined to disappear if left unresected. (10) Another study showed that tumor regression effect of nCRT increases over time. (11) Previous studies have not shown how much the tissue regression effect of nCRT, which increases over time, can be controlled if metastatic-positive lymph nodes are not removed at surgery. Our group has

been carrying out nCRT + TME without LLND as a standard of care, and systemic chemotherapy was applied if the lymph node size increased as postoperatively. It is clinically significant if we can distinguish the lymph nodes which require LLND from the lymph nodes which can avoid LLND by nCRT. Previous studies have not yet assessed factors for LLND indication beside size of the lymph node.

Additionally, in the present study, we evaluated neutrophil/lymphocyte (NLR), platelet/lymphocyte (PLR), and CEA, which have been reported as predictors of the effects of nCRT. (12–15) Other than that, tumor shrinkage rate, tumor regression grade (TRG) (16), and lateral lymph node short axis are used to determine if they are predictors of lateral lymph node recurrence (LLNR).

Subjects And Methods

Patients

Between 2009 and 2016, 168 patients with low rectal cancer (cT3-4, N any) whose anal verge (AV) was 7 cm and less and underwent nCRT (40-45Gy) with oral Uracil/Tegafur or S-1 chemotherapy. TME was carried out at 6 to 8 weeks after nCRT, and LLND was not performed.

Assessment of therapeutic effect

As predictive factors of nCRT outcome, CEA levels, NLR, PLR prior to and after the nCRT were assessed. As factors of therapeutic effect, TRG and lateral lymph node short-axis diameter were assessed. For CEA, we used 5ng/mL as our institutional cut off value (COV). Both NLR and PLR were measured from blood samples prior to and after the CRT. We determined COV from the receiver operating characteristic curve (ROC) in which recurrence was positive. Tumor shrinkage rate was calculated by T2-weighted MRI images prior to and after the CRT according to the formula: Tumor shrinkage rate (%) = $100 \times [(pre-CRT \text{ tumor volume}) - (post-CRT \text{ tumor volume})] / (pre-CRT \text{ tumor volume})$. (17)

Histological assessment was carried out according to the tumor regression grade (TRG) in which grade 1 as complete regression; grade 2 as presence of rare residual cancer cells; grade 3 as increased number of residual cancer cells; grade 4 as residual cancer outgrowing fibrosis, and grade 5 as absence of regression change. (16) Responder was defined as TRGs 1 and 2.

Lateral lymph node short-axis diameter was measured using MRI and CT images. Cancer recurrence was diagnosed by CT, and only initial recurrence lesion was evaluated to rule out lateral lymph node metastases from local recurrence.

Statistical analysis

The COV for each score was calculated by risk evaluation analysis, performed using ROC in which the recurrence of LLND was considered as a positive result. The patients were divided into 2 groups according to whether their score was less than the COV or equal to or greater than the COV, and the incidence of recurrence was compared. Lateral lymph node recurrence rate was assessed by Kaplan-

Meier analysis and multiple logistic regression analysis. We defined $p < 0.05$ as statistical significance. All statistical analyses were carried out using MP®10 software (SAS Institute Inc., Cary, NC, USA).

This study was approved by the institutional review board of Tokai University (21R167).

Results

Patient characteristics

Table 1 shows patient characteristics. Among 168 patients, 124 patients were male and 44 patients were female. Based on ypT staging, 26 patients displayed pCR (15.4%). The TRGs 1 and 2 were 65 patients, and TRGs 3 and 4 were 103 patients. Local recurrence was found in 28 patients (16.6%) and the lateral lymph node recurrence was found in 5 cases (3%). Median observational period was 1,874 days.

Association between predictive factors of CRT effect and lateral lymph node recurrence

Pre- and post-CRT CEA levels were not statistically related with lateral lymph node recurrence ($p = 0.5136$, $p = 0.0787$, respectively, Figs. 1a, 1b). The COVs of NLR were 1.76 at pre-CRT and 6.30 at post-CRT, respectively. There were no statistical relations between COVs at pre- and post-CRT and lateral lymph node recurrence ($p = 0.2988$, $p = 0.0701$, Figs. 2a and 2b). The COVs of PLR were 123.8 at pre-CRT and 157.7 at post-CRT, respectively. There were significant statistical relations between pre- and post-CRT PLR and lateral lymph node recurrence. The PLR at COV and above significantly increased in patients with lateral lymph node recurrence ($p = 0.0269$, $p = 0.0278$, Figs. 3a and 3b).

Association between tumor shrinkage rate, histological effect, lateral lymph node short-axis diameter with CRT, and the lateral lymph node recurrence

Median value of tumor shrinkage rate was 77.0%. There was no statistical relation between tumor regression rate and lateral lymph node recurrence ($p = 0.5613$, Fig. 4a). The TRGs 1 and 2 were 65 patients (%) and the TRGs 3 and 4 were 103 patients. There were no statistical relations between TRGs and lateral lymph node recurrence ($p = 0.5847$, Fig. 4b).

The COV of lateral lymph node short-axis diameter was 11.6 mm at pre-CRT and 5.5 mm at post-CRT, respectively. There were 7 patients whose pre-CRT diameters were 11.6 mm or larger (4.2%). Of those, five patients developed lateral lymph node recurrence, and the recurrence rate was 71.4%. No patients whose pre-CRT lymph node diameters less than 11.6 mm developed the lateral lymph node recurrence. There were 11 patients (6.5%) whose post-CRT lateral lymph node short-axis diameters were 5.5 mm and larger. Of those, five patients developed lateral lymph node recurrence, and the recurrence rate was 45.5%. There was a statistically significant relations between pre- or post-CRT and lateral lymph node short-axis diameter ($p < 0.001$, $p < 0.001$, respectively, Figs. 5a and 5b).

Discussion

Our study suggests that the histological effect of rectal lesions by CRT does not support the omission of LLND. And the minor axis of the lateral lymph node and PLR are risk factors for recurrence of the lateral lymph node.

The American Society of Colon and Rectal Surgeons (ASCRS) strongly recommends not performing LLND in the absence of swollen lateral lymph nodes. However, it does not mention appropriate treatment strategy for patients with swollen lateral lymph nodes. (18)

Fujita et al. reported that 5-year relapse-free survival rate in the TME + LLND group was not improved compared with TME group. Moreover, lateral pelvis recurrence was significantly high in TME group and, recurrence of lateral lymph nodes has been reported to be the leading cause of pelvic recurrence in the absence of LLND after CRT, (19, 20) and the usefulness of LLND has also been recognized in Western countries.

Hazen et al. reviewed the treatment of lateral lymph nodes in rectal cancer and state that they will change towards selective LLND in the future. (21) Therefore, it is important to develop predictors of lateral lymph node metastasis.

Regarding lymph node size and cancer recurrence, previous studies investigated LLND indication by lymph node size, and specific criteria have not yet been established. Kawai et al. reported that lateral lymph node recurrence rate was 1.6% in post-CRT + TME and TME + LLND was performed on patients with 8 mm LLN before surgery, and the positive rate for lateral lymph node metastasis was 52.4%. And the estimated incidence of LLN metastases after nCRT was 9.3%. (10)

They also reported that some of patients who underwent LLND did not develop local cancer recurrence through post-operative observation and assessment of lymph node size had 1-mm interobserver error, and there is a limit to the evaluation based on the lymph node size alone. (10) In the currency study, patients with pre-CRT lymph node minor-axis diameter less than 11.6 mm and post-CRT diameter less than 5.5 mm did not develop lateral lymph node recurrence. While the results in the current study did not show significant difference compared to previous studies, only 5 patients developed lateral lymph node recurrence. Further studies with additional cases are necessary.

Predictive factors of therapeutic effect, histological effect, and lateral lymph node recurrence

Previous studies showed that patients with good histological effect had favorable outcome. (22,23) However, some patients with TRG 1 exhibited lymph node metastasis.

It has been reported that 4.6% of patients with TRG1 were positive for lymph nodes in the mesorectum, and that the lymph nodes in the mesorectum were associated with the TRG of the tumor itself. (24,25)

However, its association with lateral lymph nodes has not been investigated. Although it is unclear why there is a difference in the control of lymph nodes in the irradiation range, our study found no association between lateral lymph node recurrence, TRG, and tumor shrinkage.

Inflammation is associated with tumor formation, and various predictive factors are investigated. Previous studies showed that CEA, NLR, and PLR were related to the histological effect. (12–15) Therefore, in the present study, we assessed the relations between each factor and the lateral lymph node recurrence, and found significant relations between pre- and post-CRT PLRs and lateral lymph node recurrence. Lee et al. reported PLR had a close relation with pathological complete response (pCR). (14) Current results and previous report suggest that PLR is useful not only for assessment of histological effect, but also it can be used as a predictive factor of lateral lymph node recurrence. While PLR can be easily calculated from standard blood sampling, there is no consensus regarding COV.

We are aware of several limitations in the current study, which include a single institutional study and small sample size. In the current study, positive cases were defined as lateral lymph node recurrence without local recurrence, however, true lateral lymph node recurrence might not be captured. Despite these study limitations, the current study is the first study which assessed lateral lymph node recurrence by various factors not limited to the lymph node size. Further studies with additional cases are necessary.

Conclusion

The current study showed that patients whose lymph node short-axis diameters less than 11.6 mm at pre-CRT and 5.5 mm at post-CRT did not develop lateral lymph node recurrence. There was no relation between histological effect and lateral lymph node recurrence. Results suggest that pre- and post-CRT PLRs may be a useful predictive factor for management of lateral lymph node recurrence in patients with rectal cancer.

Declarations

Conflict of interest : None. Conflict of Interest: The authors have no conflicts of interest to declare.

Ethical approval : The institutional review board approved the present study.

Funding sources: There was no funding source for this study.

Authors contributions:

Conception and design: H.Miyakita, S Yamamoto

Provision of study materials or patients: H Miyakita.

Collection and assembly of data: H Miyakita, LF Chan.

Data analysis and interpretation: H Miyakita, K Okada, H Kayano

Manuscript writing and prepared figure: H Miyakita, K Okada

All authors reviewed the manuscript.

References

1. Watanabe T, Muro K, Ajioka Y, Hashiguchi Y, Ito Y, Saito Y, Hamaguchi T, Ishida H, Ishiguro M, Ishihara S, Kanemitsu Y, Kawano H, Kinugasa Y, Kokudo N, Murofushi K, Nakajima T, Oka S, Sakai Y, Tsuji A, Uehara K, Ueno H, Yamazaki K, Yoshida M, Yoshino T, Boku N, Fujimori T, Itabashi M, Koinuma N, Morita T, Nishimura G, Sakata Y, Shimada Y, Takahashi K, Tanaka S, Tsuruta O, Yamaguchi T, Yamaguchi N, Tanaka T, Kotake K, Sugihara K (2018) Japanese Society for Cancer of the Colon and Rectum (JSCCR) guidelines 2016 for the treatment of colorectal cancer. *Int J Clin Oncol.* **23**(1):1–34.
2. Kapiteijn E, Marijnen CA, Nagtegaal ID, Putter H, Steup WH, Wiggers T, Rutten HJ, Pahlman L, Glimelius B, van Krieken JH, Leer JW, van de Velde CJ (2001) Preoperative radiotherapy combined with total mesorectal excision for resectable rectal cancer. *N Engl J Med.* **345**(9) : 638–46.
3. Ogura A, Konishi T, Cunningham C, Gracia-Aguilar J, Iversen H, Toda S, Lee IK, Lee HX, Uehara K, Lee P, Putter H, van de Velde CJH, Beets GL, Rutten HJT, Kusters M. (2018) Neoadjuvant (Chemo) radiotherapy with Total Mesorectal excision only is not sufficient to prevent lateral local recurrence in enlarged nodes : results of the multicenter lateral node study of patients with low cT3/4 rectal cancer. *J Clin Oncol.* **37**: 33-43
4. Yang X, Yang S, Hu T, Gu C, Wei M, Deng X, Wang X, Zhou Z. (2020) What is the role of lateral lymph node dissection in rectal cancer patients with clinically suspected lateral lymph node metastasis after preoperative chemoradiotherapy? A meta-analysis and systematic review. *Cancer med.* **9**: 4477-4489
5. Yamaoka Y, Kinugasa Y, Shiomi A, Yamaguchi T, Kagawa H, Yamakawa Y, Numata M, Furutani A. (2017) Preoperative chemoradiotherapy changes the size criterion for predicting lateral lymph node metastasis in lower rectal cancer. *Int J Colorectal Dis.* **32**: 1631-1637
6. Akiyoshi T, Matsueda K, Hiratsuka M, Unno T, Nagata J, Nagasaki T, Konishi T, Fujimoto Y, Nagayama S, Fukunaga Y, Ueno M. (2015) Indications for lateral pelvic lymph node dissection based on magnetic resonance imaging before and after preoperative chemoradiotherapy in patients with advanced low-rectal cancer. *Ann Surg Oncol.* **22**: S614-S620
7. Kim HJ, Choi GS, Park JS, Park SY, Cho SH, Lee SJ, Kang BW, Kim JG. (2017) Optimal treatment strategies for clinically suspicious lateral pelvic lymph node metastasis in rectal cancer. *Oncotarget.* **8**: 100724-100733
8. Kim MJ, Kim TH, Kim DY, Kim SY, Baek JY, Chang HJ, Park SC, Park JW, Oh JH. (2015) Can chemoradiation allow for omission of lateral pelvic node dissection for locally advanced rectal cancer? *J Surg Oncol.* **111**: 459-464

9. Ishihara S, Kawai K, Tanaka T, Kiyomatsu T, Hata K, Nozawa H, Morikawa T, Watanabe T. (2017) Oncological outcomes of lateral pelvic lymph node metastasis in rectal cancer treated with preoperative chemoradiotherapy. *Dis Colon Rectum*. 60: 469-476
10. Kawai K, Shiratori H, Hata K, Nozawa H, Tanaka T, Nishikawa T, Muroto K, Ishihara S. (2021) Optimal size criteria for lateral lymph node dissection after neoadjuvant chemoradiotherapy for rectal cancer. *Dis Colon Rectum*. 64: 274-283
11. Lefevre JH, Mineur L, Kotti S, Rullier R, Rouanet P, Chaisemartin C, Meunier B, Mehrdad J, Cotte E, Desrame J, Karoui M, Benoist S, Kirzin S, Berger A, Panis Y, Piessen G, Saudemont A, Prudhomme M, Peschaud F, Dubois A, Loriau J, Tuech JJ, Meurette G, Lupinacci R, Goasgen N, Parc Y, Simon T, Tiret E. (2016) Effect of interval (7 or 11 weeks) between Neoadjuvant Radiochemotherapy and Surgery on complete pathologic response in Rectal cancer: Multicenter, Randomized, controlled Trial (GRECCAR-6). *J Clin Oncol*. 34: 3773-80
12. Kim IY, You SH, Kim YW. (2014) Neutrophil-lymphocyte ratio predicts pathologic tumor response and survival after preoperative chemoradiation for rectal cancer. *BMC Surg*. 14 : 94 doi: 10.1186/1471-2482-14-94
13. Shen L, Zhang H, Liang L, Li G, Fan M, Wu Y, Zhu J, Zhang Z. (2014) Baseline neutrophil-lymphocyte ratio (≥ 2.8) as a prognostic factor for patients with locally advanced rectal cancer undergoing neoadjuvant chemoradiation. *Radiat Oncol*. 9: 295
14. Lee JH, Song C, Kang SB, Lee HS, Lee KW, Kim JS. (2018) Predicting pathological complete regression with haematological markers during neoadjuvant chemoradiotherapy for locally advanced rectal cancer. *Anticancer res*. 38: 6905-6910
15. P Wang, Z Wang, Y Liu, J Xie, Y Ren. (2021) Prognostic value of platelet-associated biomarkers in rectal cancer patients received neoadjuvant chemoradiation: A retrospective study. *Cancer/Radioterapie*. 25: 147-154
16. Mandard AM, Dalibard F, Mandard JF, J Marnay, M Henry-Amar, JF Petiot, Roussel A, Jacob JH, Segol P, Samama G, Ollivier JM, Bonvalot S, Gignoux M. (1994) Pathologic assessment of tumor regression after preoperative chemoradiotherapy of esophageal carcinoma. Clinicopathologic correlations. *Cancer*. 73: 2680-2686
17. Suzuki T, Sadahiro S, Tanaka A, Okada K, Saito G, Kamijo A, Akiba T, Kawada S. (2014) Relationship between histologic response and degree of tumor shrinkage after chemoradiotherapy in patients with locally advanced rectal cancer. *J Surg Oncol*. 109: 659-664
18. You YN, Hardiman K, Bafford A, Poylin V, Francone TD, Davis K, Paquette IM, Steele SR, Feingold DL. (2020) The American society of colon and rectal surgeons clinical practice guidelines for the management of rectal cancer. *Dis Colon and Rectum*. 63: 1191-222
19. Fujita S, Mizusawa J, Kanemitsu Y, Ito Masaaki, Kinugasa Y, Komori K, Ohue M, Ota M, Akazai Y, Shiozawa M, Yamaguchi T, Bandou H, Katsumata K, Murata K, Akagi Y, Takiguchi N, Saida Y, Nakamura K, Fukuda H, Akasu T, Moriya Y. (2017) Mesorectal Excision with or without lateral lymph

- node dissection for clinical stage II/III lower rectal cancer (JCOG0212): A multicenter, randomized controlled, noninferiority trial. *Ann Surg.* 266: 201-207
20. Kim TH, Jeong SY, Choi DH, Kim DY, Jung KH, Moon SH, Chang HJ, Lim SB, Choi HS, Park JG. (2008) Lateral lymph node metastasis is major cause of locoregional recurrence in rectal cancer treated with preoperative chemoradiotherapy and curative resection. *Ann Surg Oncol.* 15: 729-737
21. SJA Hazen, TC Sluckin, T Konishi, M Kusters. (2021) Lateral lymph node dissection in rectal cancer: Stage of the art review. *Eur J Surg Oncol.* <http://doi.org/10.1016/j.ejso.2021.11.003>
22. Peeter KC, Marijnen CA, Nagtegaal ID, Kranenbarg EK, Putter H, Wiggers T, Rutten H, Pahlman L, Glimelius B, Leer JW. (2007) The TME trial after a median follow-up of 6 years: increased local control but no survival benefit in irradiated patients with resectable rectal carcinoma. *Ann Surg.* 246: 693-701
23. Sauer R, Becker H, Hohenberger W, Rodel C, Wittekind C, Fietkau R, Martus P, Tschmelitsch J, Hager E, Hess CF, Karstens JH, Liersch T, Schmidberger H, Raab R. (2006) Preoperative versus postoperative chemoradiotherapy for rectal cancer. *351:* 1731-1740
24. Wee IJY, Cao HM, Ngu JCY. (2019) The risk of nodal disease in patients with pathological complete response after neoadjuvant chemoradiation for rectal cancer: a systematic review, meta-analysis, and meta-regression. *Int J Colorectal Dis.* 34: 1349-1357
25. Vecchio FM, Valentini V, Minsky BD, Padula GDA, Venkatraman ES, Baldicci M, Micciché F, Ricci R, Morganti AG, Gambacorta MA, Maurizi F, Coco C. (2005) The relationship of pathologic tumor regression grade (TRG) and outcomes after preoperative therapy in rectal cancer. *Int J Radiation Oncology Biol Phys.* 62: 752-760

Tables

Table.1 Patient's Characteristic

| | | | | | |
|-----------|--------|-------|-------------------------------|----------------|-----|
| Sex | Male | 124 | TRG‡ | 1-2 | 65 |
| | Female | 44 | | 3-4 | 103 |
| age | Range | 33-81 | Adjuvant Chemotherapy | no | 106 |
| | Median | 62 | | 5-FU | 58 |
| cT Stage | 3 | 129 | | 5-FU+OX | 4 |
| | 4a | 30 | Local recurrence | Yes | 28 |
| | 4b | 9 | | No | 140 |
| cN Stage | - | 65 | Lateral lymph node recurrence | Yes | 5 |
| | + | 103 | | No | 163 |
| ypT Stage | 0 | 7 | Distant recurrence | Yes | 43 |
| | 1a/b | 9 | | No | 125 |
| | 2 | 40 | Procedure | LAR§ | 119 |
| | 3 | 85 | | APR¶ | 42 |
| | 4a/b | 1 | | Hartmann | 3 |
| | pCR† | 26 | | Local Excision | 4 |
| ypN Stage | - | 121 | | | |
| | + | 44 | | | |
| | X | 3 | | | |


† : pCR; Pathological complete response

‡: TRG; Tumor regression grade


§: LAR; Low anterior resection

¶: APR; Abdominoperioperineal resection

Figures



A




B

Figure 1


Association between predictive factors of CEA level and lateral lymph node recurrence.

a. Association between predictive factors of Pre-CRT CEA level and lateral lymph node recurrence.

b. Association between predictive factors of Post-CRT CEA level and lateral lymph node recurrence.



A




B

Figure 2


Association between predictive factors of NLR and lateral lymph node recurrence.

a. Association between predictive factors of Pre-CRT NLR and lateral lymph node recurrence.

b. Association between predictive factors of Post-CRT NLR and lateral lymph node recurrence.



A




B

Figure 3


Association between predictive factors of PLR and lateral lymph node recurrence.

a. Association between predictive factors of Pre-CRT PLR and lateral lymph node recurrence.

b. Association between predictive factors of Post-CRT PLR and lateral lymph node recurrence.



A




B

Figure 4


Association between tumor shrinkage rate, histological effect of CRT, and the lateral lymph node recurrence.

a. Association between tumor shrinkage rate and the lateral lymph node recurrence.

b. Association between histological effect of CRT, and the lateral lymph node recurrence.



A



B

Figure 5

Association between lateral lymph node short-axis diameter of CRT, and the lateral lymph node recurrence.

a Association between lateral lymph node short-axis diameter of pre- CRT, and the lateral lymph node recurrence.

b. Association between lateral lymph node short-axis diameter of post- CRT, and the lateral lymph node recurrence.