

Greater hypothalamic microglial activation with aging in women as compared to men: potential relevance to sex and species differences in reproductive and biologic aging.

Tracy Butler (✉ tab2006@med.cornell.edu)

Weill Cornell Medicine

Lidia Glodzik

Weill Cornell Medicine

Xiuyuan Hugh Wang

Weill Cornell Medicine

Ke Xi

Weill Cornell Medicine

Yi Li

Weill Cornell Medicine

Hong Pan

Brigham and Women's Hospital

Liangdong Zhou

Weill Cornell Medicine

Gloria Chiang

Weill Cornell Medicine

Simon Morim

Weill Cornell Medicine

Nimmi Wickramasuriya

Weill Cornell Medicine

Emily Tanzi

Weill Cornell Medicine

Thomas Maloney

Weill Cornell Medicine

Patrick Harvey

Weill Cornell Medicine

Xiangling Mao

Weill Cornell Medicine

Ray Qolamreza Razlighi

Weill Cornell Medicine

Henry Rusinek

New York University Langone Medical Center

Dikoma Shungu

Weill Cornell Medicine

Mony de Leon

Weill Cornell Medicine

Craig S. Atwood

University of Wisconsin–Madison

P. David Mozley

Weill Cornell Medicine

Research Article

Keywords:

Posted Date: May 4th, 2022

DOI: <https://doi.org/10.21203/rs.3.rs-1538708/v1>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

In rodents, hypothalamic inflammation plays a critical role in aging and age-related diseases. We used Positron Emission Tomography with a radiotracer sensitive to the TSPO molecule expressed by activated microglia to assess, for the first time, human hypothalamic inflammation. We found that women but not men showed age-correlated hypothalamic TSPO expression, and suggest this is relevant to understanding a stark sex difference in human aging, namely, that only women undergo menopause.

Full Text

The hypothalamus is best known as the controller of hormone secretion to regulate essential bodily functions such as reproduction and feeding, and is one of the only brain structures to demonstrate a sexually dimorphic cell population in humans.¹ Recent work in animal models has shown that the hypothalamus also serves as a critical pathophysiologic hub that translates systemic inflammation into neural inflammation, mediating and perpetuating age-related brain and body changes such as cognitive decline and metabolic syndrome.^{2,3} Reducing hypothalamic microglial activation by a variety of means including medication, genetic manipulation, and dietary changes, reduces age-related diseases and extends life in animal models, with marked sex differences in this effect.^{4,5} To our knowledge, hypothalamic inflammation has not previously been assessed *in vivo* in humans.

We used Positron Emission Tomography (PET) with ¹¹C-PK11195, a radiotracer sensitive to the translocator protein (TSPO) expressed by activated microglia,⁶ to assess the contribution of age and sex to hypothalamic TSPO expression (quantified as non-displaceable binding potential, BPnd) in 43 healthy subjects ranging in age from 23 to 78 (19 female; mean age 55.) In addition to chronologic age, we used a marker of brain biological age based on the degree of atrophy present on MRI.⁷ We controlled for the potential confound of Body Mass Index (BMI) which differs by age and sex and correlates with whole-brain TSPO expression.⁸ We took two imaging approaches: 1) quantification of average hypothalamic BPnd within subjects' native space using a deep learning technique for accurately segmenting the hypothalamus⁹ and 2) morphing individual brains into template space to localize age and sex effects over the whole brain regardless of traditional neuroanatomic boundaries (Statistical Parametric Mapping; SPM.)

In the combined group of women and men (who did not differ in chronologic age, brain age, difference between brain and chronologic age, nor BMI; subject demographics are presented in Supplementary Table 1) SPM analysis showed greater BPnd in association with greater age (both chronologic and brain) in bilateral thalamus (Fig. 1), in accord with prior TSPO PET studies.¹⁰⁻¹² We did not find age-correlated TSPO expression in a probabilistic atlas-defined hypothalamic region of interest¹³ in this mixed-sex group, as might have been expected based on animal studies.^{2,3}

However, both imaging approaches revealed sex-specific associations of age with TSPO expression in hypothalamus. An optimal regression model including brain age, sex and BMI ($R^2=.457$, $F(7,35)=4.203$,

p=0.002, AIC = -223.72) showed that interaction between all of these factors predicted average hypothalamic BPnd ($\beta = -.002$, p=0.001). Partial correlation (controlling for BMI) between brain age and hypothalamic BPnd was significant only in women (R=0.49, p=0.032), as shown in **Figure 2**. A model including chronologic rather than brain age was slightly less accurate (R²= .404, F(7,35)=3.384, p=0.007, AIC = -219.72) suggesting greater relevance of biologic than chronologic age to hypothalamic TSPO expression. Additional details of regression analysis are presented in Supplementary Tables 3 and 4. Voxelwise analyses over the whole brain provided convergent results, demonstrating that only women had age-correlated TSPO expression within hypothalamus, and further localizing this sex difference to a region of hypothalamus extending to thalamus, as shown in **Figure 3**.

This first assessment of hypothalamic microglial activation in humans demonstrates important age and sex effects. Contrary to findings in rodents,^{2,3} hypothalamic microglial activation, indexed by TSPO expression, does not appear to be a universal feature of aging in humans; we detected age-correlated hypothalamic TSPO expression – stronger for a biologic measure of brain age than for chronologic age – only in women. What appears to be universal in humans is age-correlated thalamic TSPO expression, as demonstrated here and in multiple prior studies.¹⁰⁻¹² It is intriguing that in humans, the main focus of TSPO expression is thalamus, while in rodents it is hypothalamus.^{2,3,14} Our preliminary finding that age-correlated thalamic TSPO expression may extend into hypothalamus in women (Figure 2) potentially mirrors rodent studies showing hypothalamic inflammation may extend into thalamus.¹⁴ This difference between rodents and humans, perhaps reflecting evolutionary changes in the size and function of thalamus versus hypothalamus, may be relevant to the appropriateness of using rodents to model human aging and age-related diseases. Rodents are commonly used experimental animal in large part because of their rapid maturity, high reproductive rate, and short lifespan, and these interrelated characteristics (all potentially mediated by hypothalamus) are very different in humans.¹⁵

Our finding of age-correlated hypothalamic TSPO expression only in women may relate to a stark difference in how women and men age: that women but not men experience loss of fertility – menopause – at mid-life. Menopause is associated with markedly increased risk of age-related disease including Alzheimer's¹⁶ and cardiovascular disease.¹⁷ Unlike in rodents, this increased risk is not clearly ameliorated by estrogen replacement,^{18,19} again highlighting important differences between humans and rodents, and the need for new paradigms for understanding human reproductive and brain aging. While menopause is generally attributed to changes at the level of the ovary, hypothalamic changes occur early in the menopause transition.²⁰⁻²² Our results suggesting a role for hypothalamic inflammation in this change could have implications for understanding and perhaps altering reproductive aging in women.

This study has several limitations, including its reliance upon cross section data to study the longitudinal process of aging, lack of information about menopausal status of participants, and the fact that TSPO is expressed not just by microglia but by several cell types in the brain,²³ and has important functions unrelated to inflammation. In particular, the role of TSPO in steroidogenesis, with estrogen-regulated expression in hypothalamus²⁴ may be highly relevant to our findings and to female reproductive aging.

Additional limitations include the use of PET, with its limited spatial resolution, to assess a small structure such as the hypothalamus, as well as use of a first generation TSPO radiotracer with inferior signal properties to newer tracers.⁶ However, TSPO PET is the only available method to assess microglial activation *in vivo* in humans, and we apply optimal image processing and analysis methods to quantify PET signal²⁵ and accurately segment the hypothalamus.^{9,13} Current results suggest intriguing differences between rodents and humans, and between women and men, in hypothalamic TSPO expression in aging. Understanding such differences is essential to the development of effective therapies for age-related diseases, and perhaps aging itself.

Methods

Subjects

Healthy subjects (n = 43) who underwent 11C-PK11195 Positron Emission Tomography (PK PET) as normal controls for several different studies conducted at Weill Cornell Citigroup Biomedical Imaging Center between 2013 and 2019 (investigating epilepsy, hypertension, chronic fatigue syndrome, Parkinson's Disease and Gulf War Syndrome) were included in this analysis. All subjects were free of significant medical, neurologic and psychiatric disease.

Image Acquisition

PET images were acquired over one hour in list mode, starting at the time of injection of ~ 550 MBq of 11C-PK11195 on a Siemens Biograph PET-CT scanner.

Three-dimensional volumetric T1 weighted BRAVO/MPRAGE MR images (1.2 × 1.2 × 1.2 mm³ isotropic voxels) were acquired using a GE Signa or Siemens Skyra 3T scanner.

Image Processing

PET images were reconstructed into 22 frames and motion corrected using MCFLIRT²⁶ within FSL.²⁷ Binding Potential (BPnd) images reflecting the concentration of translocator protein (TSPO) expressed by activated microglia, irrespective of tracer delivery/blood flow, were generated from dynamic PET using a multilinear reference tissue model (MRTM0)^{28,29} implemented in the freely available software package FireVoxel (<https://firevoxel.org>.) The reference tissue time-activity curve was identified via optimized supervised cluster analysis (SVCA),³⁰ which is considered the optimal method for analyzing PK PET scans.²⁵ Each subject's T1 MRI was linearly co-registered to his/her BPnd image with rigid body transformation in FSL.

Brain Age

An estimate of subjects' biological (as distinct from chronological) age, termed brain age, was derived from each subject's MRI using a cloud-based framework based on cortical thickness.⁷

Whole Brain Statistical Parametric Analyses

MRI scans were warped to MNI template space using Advanced Neuroimaging Tools.³¹ MRI-derived normalization parameters were applied to co-registered BPnd images to bring them into standard space. Using Statistical Parametric Mapping³² (SPM 12), regional BPnd was assessed in the group as a whole and in women and men separately. Within SPM, multiple regression was used to assess brain regions in which BPnd correlated with (chronologic and brain) age. Body Mass Index (BMI) was included as a covariate of no interest because it has been shown to correlate with whole-brain TSPO expression and in association with age and sex.⁸ Maps were brain masked to exclude strong extracranial signal which contaminated peripheral cortical regions. Results were considered significant at a threshold of $p < .05$ corrected for False Discovery Rate (FDR)³³ with a minimum cluster volume of 1cm^3 . Results within the hypothalamic region of *a priori* interest were considered significant at a small volume corrected threshold of $p < .05$. The hypothalamic region of interest (ROI) was defined based on a high resolution probabilistic atlas of bilateral hypothalamus¹³ threshold at 50%. T-maps generated using SPM were converted to Z-maps within SPM to directly compare age-correlated TSPO expression between women and men. Results were considered significant at $Z > 3$. XJview (<https://www.alivelearn.net/xjview>) was used to calculate the center of gravity of clusters and generate figures.

Native Space Analyses

Regional BPnd was quantified within hypothalamus within the subject's native space using Freesurfer³⁴ with a highly-accurate deep learning technique for segmenting the hypothalamus.⁹ Segmented regions were transformed to PET space with the inverse transformation matrix from the co-registration step and eroded 1mm in plane to minimize partial volume effects.

Statistical Analyses

Analyses were performed in R. All results were considered significant at $p < .05$.

T-tests were used to assess differences between men and women in age, brain age, difference between chronologic age and brain age, and BMI.

Stepwise linear regression models were built to assess if sex, age (chronologic and brain), BMI and their interactions significantly predicted average hypothalamic BPnd (from the native space analysis). These predictors were chosen based on prior evidence implicating them in hypothalamic structure/function and whole brain TSPO PET.⁸ The optimal regression model was chosen based on Akaike Information Criterion (AIC.) To further assess significant effects, partial correlations between hypothalamic BPnd and age, controlling for BMI, were assessed separately in women and men using Pearson's test, and compared using Fisher Z-comparison.

Declarations

Funding Acknowledgement: NIA R01 AG057681 and NINDS K23 NS057579

Author Contribution: Conceptualization: TB; Data Acquisition: TB, LG, YL, SM, ET, PH, XM, DS, MdL, DM; Data Analysis: TB, HW, KX, YL, HP, LZ, GC, NW, ET, TM, PH, RR, HR; Data Interpretation: TB, LG, KX, YL, RR, HR, MDL, CSA, DM; Drafting Manuscript: TB; Edit and Review of Manuscript: All authors.

Data Availability: All data, in de-identified format, is available upon written or emailed request to the corresponding author.

Competing Interests: No authors have relevant competing interests.

References

1. Swaab, D.F. & Fliers, E. A sexually dimorphic nucleus in the human brain. *Science* **228**, 1112–1115 (1985).
2. Cai, D. & Khor, S. “Hypothalamic microinflammation” paradigm in aging and metabolic diseases. *Cell metabolism* **30**, 19–35 (2019).
3. Ishii, M. & Iadecola, C. Metabolic and non-cognitive manifestations of Alzheimer’s disease: the hypothalamus as both culprit and target of pathology. *Cell metabolism* **22**, 761–776 (2015).
4. Sadagurski, M., Cady, G. & Miller, R.A.J.A.C. Anti-aging drugs reduce hypothalamic inflammation in a sex-specific manner. **16**, 652–660 (2017).
5. Liu, T., Xu, Y., Yi, C.-X., Tong, Q., Cai, D.J.P. & cell The hypothalamus for whole-body physiology: from metabolism to aging. 1–28 (2021).
6. Kreisl, W.C., Kim, M.-J., Coughlin, J.M., Henter, I.D., Owen, D.R. & Innis, R.B. PET imaging of neuroinflammation in neurological disorders. *The Lancet Neurology* **19**, 940–950 (2020).
7. Pardoe, H.R. & Kuzniecky, R.J.N. NAPR: a cloud-based framework for neuroanatomical age prediction. **16**, 43–49 (2018).
8. Tuisku, J., Plavén-Sigra, P., Gaiser, E.C., Airas, L., Al-Abdulrasul, H., Brück, A., Carson, R.E., Chen, M.-K., Cosgrove, K.P., Ekblad, L.J.E.j.o.n.m. & imaging, m. Effects of age, BMI and sex on the glial cell marker TSPO—a multicentre [11 C] PBR28 HRRT PET study. **46**, 2329–2338 (2019).
9. Billot, B., Bocchetta, M., Todd, E., Dalca, A.V., Rohrer, J.D. & Iglesias, J.E. Automated segmentation of the hypothalamus and associated subunits in brain MRI. *Neuroimage* **223**, 117287 (2020).
10. Paul, S., Gallagher, E., Liow, J.-S., Mabins, S., Henry, K., Zoghbi, S.S., Gunn, R.N., Kreisl, W.C., Richards, E.M. & Zanotti-Fregonara, P. Building a database for brain 18 kDa translocator protein imaged using [11C] PBR28 in healthy subjects. *Journal of Cerebral Blood Flow & Metabolism*, 0271678X18771250 (2018).
11. Kumar, A., Muzik, O., Shandal, V., Chugani, D., Chakraborty, P. & Chugani, H.T. Evaluation of age-related changes in translocator protein (TSPO) in human brain using 11 C-[R]-PK11195 PET. *Journal of neuroinflammation* **9**, 232 (2012).

12. Cagnin, A., Brooks, D.J., Kennedy, A.M., Gunn, R.N., Myers, R., Turkheimer, F.E., Jones, T. & Banati, R.B. In-vivo measurement of activated microglia in dementia. *The Lancet* **358**, 461–467 (2001).
13. Pauli, W.M., Nili, A.N. & Tyszka, J.M. A high-resolution probabilistic in vivo atlas of human subcortical brain nuclei. *Scientific data* **5**, 1–13 (2018).
14. Giannaccini, G., Betti, L., Palego, L., Pirone, A., Schmid, L., Lanza, M., Fabbrini, L., Pelosini, C., Maffei, M. & Santini, F.J.B.n. Serotonin transporter (SERT) and translocator protein (TSPO) expression in the obese ob/ob mouse. **12**, 1–8 (2011).
15. Holtze, S., Gorshkova, E., Braude, S., Cellerino, A., Dammann, P., Hildebrandt, T.B., Hoeflich, A., Hoffmann, S., Koch, P. & Terzibasi Tozzini, E. Alternative animal models of aging research. *Frontiers in Molecular Biosciences* **8**, 311 (2021).
16. Brinton, R.D., Yao, J., Yin, F., Mack, W.J. & Cadenas, E. Perimenopause as a neurological transition state. *Nature reviews endocrinology* **11**, 393–405 (2015).
17. El Khoudary, S.R., Aggarwal, B., Beckie, T.M., Hodis, H.N., Johnson, A.E., Langer, R.D., Limacher, M.C., Manson, J.E., Stefanick, M.L. & Allison, M.A. Menopause transition and cardiovascular disease risk: implications for timing of early prevention: a scientific statement from the American Heart Association. *Circulation* **142**, e506-e532 (2020).
18. Miller, V.M., Naftolin, F., Asthana, S., Black, D.M., Brinton, E.A., Budoff, M.J., Cedars, M.I., Dowling, N.M., Gleason, C.E. & Hodis, H.N. The Kronos early estrogen prevention study (KEEPS): what have we learned? *Menopause (New York, NY)* **26**, 1071 (2019).
19. McEwen, B. Estrogen actions throughout the brain. (2002).
20. Weiss, G., Skurnick, J.H., Goldsmith, L.T., Santoro, N.F. & Park, S.J. Menopause and hypothalamic-pituitary sensitivity to estrogen. *Jama* **292**, 2991–2996 (2004).
21. Downs, J.L. & Wise, P.M. The role of the brain in female reproductive aging. *Molecular and cellular endocrinology* **299**, 32–38 (2009).
22. Bacon, E.R., Mishra, A., Wang, Y., Desai, M.K., Yin, F. & Brinton, R.D. Neuroendocrine aging precedes perimenopause and is regulated by DNA methylation. *Neurobiology of aging* **74**, 213–224 (2019).
23. Gui, Y., Marks, J.D., Das, S., Hyman, B.T. & Serrano-Pozo, A. Characterization of the 18 kDa translocator protein (TSPO) expression in post-mortem normal and Alzheimer's disease brains. *Brain pathology* **30**, 151–164 (2020).
24. Chen, C., Kuo, J., Wong, A. & Micevych, P. Estradiol modulates translocator protein (TSPO) and steroid acute regulatory protein (StAR) via protein kinase A (PKA) signaling in hypothalamic astrocytes. *Endocrinology* **155**, 2976–2985 (2014).
25. Wimberley, C., Lavis, S., Hillmer, A., Hinz, R., Turkheimer, F., Zanotti-Fregonara, P.J.E.J.o.N.M. & Imaging, M. Kinetic modeling and parameter estimation of TSPO PET imaging in the human brain. 1–11 (2021).
26. Jenkinson, M., Bannister, P., Brady, M. & Smith, S. Improved optimization for the robust and accurate linear registration and motion correction of brain images. *Neuroimage* **17**, 825–841 (2002).

27. Smith, S.M., Jenkinson, M., Woolrich, M.W., Beckmann, C.F., Behrens, T.E., Johansen-Berg, H., Bannister, P.R., De Luca, M., Drobnjak, I. & Flitney, D.E. Advances in functional and structural MR image analysis and implementation as FSL. *Neuroimage* **23**, S208-S219 (2004).
28. Butler, T., Ichise, M., Teich, A.F., Gerard, E., Osborne, J., French, J., Devinsky, O., Kuzniecky, R., Gilliam, F. & Pervez, F.J.J.o.n. Imaging inflammation in a patient with epilepsy due to focal cortical dysplasia. *Neuroimage* **23**, 129–131 (2013).
29. Butler, T., Li, Y., Tsui, W., Friedman, D., Maoz, A., Wang, X., Harvey, P., Tanzi, E., Morim, S. & Kang, Y.J.E. Transient and chronic seizure-induced inflammation in human focal epilepsy. *Neuroimage* **57**, e191-e194 (2016).
30. Schubert, J., Tonietto, M., Turkheimer, F., Zanotti-Fregonara, P., Veronese, M.J.E.J.o.N.M. & Imaging, M. Supervised clustering for TSPO PET imaging. 1–12 (2021).
31. Avants, B.B., Tustison, N.J., Song, G., Cook, P.A., Klein, A. & Gee, J.C. A reproducible evaluation of ANTs similarity metric performance in brain image registration. *Neuroimage* **54**, 2033–2044 (2011).
32. Penny, W.D., Friston, K.J., Ashburner, J.T., Kiebel, S.J. & Nichols, T.E. *Statistical parametric mapping: the analysis of functional brain images*. (Elsevier, 2011).
33. Benjamini, Y. & Hochberg, Y. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of the Royal statistical society: series B (Methodological)* **57**, 289–300 (1995).
34. Fischl, B., Salat, D.H., Busa, E., Albert, M., Dieterich, M., Haselgrove, C., Van Der Kouwe, A., Killiany, R., Kennedy, D. & Klaveness, S. Whole brain segmentation: automated labeling of neuroanatomical structures in the human brain. *Neuron* **33**, 341–355 (2002).

Figures




Figure 1

Age-correlated TSPO expression in bilateral thalamus in a mixed-sex group of 43 subjects. Whole-brain statistical parametric analysis with brain age⁷ as the regressor of interest, controlling for sex and BMI, shows robust results in bilateral thalamus (peak MNI coordinates: X=20, Y=-14, Z=4 [t=7.05]; X=17, Y=-18, Z=-4 [t=6.1]) as well as frontal subcortical regions. T-map is displayed at $p_{\text{uncorrected}} < .001$. Results were similar when using chronologic age, which was strongly correlated with brain age ($r = .819, p < .001$) as the regressor of interest. Additional results of SPM analysis are presented in Supplementary Figure 1 and Table 2.




Figure 2

Partial correlation plot showing sex differences in the partial correlation (controlling for BMI) between brain age and hypothalamic TSPO expression, with the hypothalamus segmented in subjects' native space using a deep learning technique.⁹ There was a significant positive partial correlation in women ($r = 0.49; p = 0.03$) but not men ($r = -0.25, p = 0.24$.) Fisher Z-comparison showed that these correlations between brain age and hypothalamic TSPO expression different significantly between women and men ($p = 0.009$).



Figure 3

Sagittal view (right of midline; $X=8$) of whole-brain statistical z-map showing regions where brain age-correlated TSPO expression was greater in women than men, with prominent results in a region spanning both hypothalamus and thalamus (hypothalamus peak: $X=11, Y=-7, Z=-13$ [$z=4.2$]; thalamus peak: $X=4, Y=-6, Z=0$ [$z=5.1$]; center of gravity in hypothalamus: $X=10, Y=-5, Z=-9$; additional results of SPM analysis are presented in Supplementary Figure 2 and Table 5.) This sex difference corresponds to significant age-correlated TSPO expression in hypothalamus in women (peak: $X=8, Y=0, Z=-12$ [$t=3.3$; $p_{\text{corrected}} = 0.002$]) but not men. There were no areas of greater age-correlated TSPO expression in men as compared to women.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [butlersupplementaryResults20220313.docx](#)