

Population health and burden of disease profile in Sierra Leone from 1990 to 2017

Jolleen Zembe

University of Pretoria

Flavia Senkubuge

University of Pretoria

Tanita Cronje

University of Pretoria

Tom Achoki ([✉ tachoki@sloan.mit.edu](mailto:tachoki@sloan.mit.edu))

Massachusetts Institute of Technology

Research Article

Keywords: Non-communicable diseases (NCDs), communicable, maternal, neonatal, and nutritional disease (CMNNs), burden of disease, Sierra Leone

Posted Date: April 18th, 2022

DOI: <https://doi.org/10.21203/rs.3.rs-1547700/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background

Sierra Leone, in West Africa, is one of the poorest developing countries in the world. Sierra Leone has experienced several recent challenges namely, a civil war from 1991 to 2002, a massive Ebola outbreak from 2014 to 2016, followed by floods and landslides in 2017. In this study, we quantified the burden of disease in Sierra Leone over a 27-year period, from 1990 to 2017.

Methodology

In this cross-sectional analysis, we analysed secondary data from the Institute of Health Metrics and Evaluation, Global Burden of Disease (GBD) study. We quantified patterns of burden of disease, injuries, and risk factors in Sierra Leone. We report GBD metrics including mortality rates, years of life lost and risk factors for all ages and both sexes from 1990 to 2017.

Results

From 1990 to 2017, mortality rates for all ages and sexes have declined in Sierra Leone although mortality rates remain some of the highest when compared to other developing countries. The burden of communicable, maternal, neonatal, and nutritional (CMNN) diseases are greater than the burden of non-communicable diseases (NCDs) due to the prevalence of endemic diseases in Sierra Leone. The most important CMNNs associated with premature mortality included respiratory infections, neglected tropical diseases, malaria and HIV-Aids. Life expectancy has increased from 37 years to 52 years

Conclusion

Sierra Leone's health status is gradually improving following the civil war and Ebola outbreak. Sierra Leone has a double burden of disease with CMNNs leading and NCDs progressively increasing. Despite these challenges, Sierra Leone has promising initiatives and programs pursuing the Universal Health Coverage 2030 Sustainable Developmental Goals Agenda. There is need for accountability of available resources, clear rules and expected roles for non-governmental organisations to ensure a level playing field for all actors to rebuild the health system.

Background

The world is experiencing rapid health, demographic and epidemiologic transitions. Many developing countries are monitoring their health trends due to poor health outcomes from infectious diseases and an increase in chronic diseases. Sierra Leone is a developing country with a history grounded in the slave trade in the 18th century to a civil war, which lasted for ten years. Despite long periods of conflict, Sierra

Leone has made great strides towards achieving political stability from a history of long periods of conflict. Sierra Leone has an ailing economy which is slowly growing following the civil war which ended in 2002. Sierra Leone's gross domestic product is growing between 4% and 7% annually.¹ As a consequence of political instability, Sierra Leone has a dysfunctional health system which remains a challenge.² Sierra Leone has a population of about 7.4 million with a reported of growth rate of 2.18% in 2017.

The Institute for Health Metrics and Evaluation (IHME) has been measuring ongoing Global Burden of Disease (GBD) for 27 years. The GBD study measures the most important health problems in each country and how health systems are responding to their health problems.³ The GBD quantifies mortality caused by major health problems, injuries, risk factors by age and sex in 195 countries. The GBD study provides evidence and motivation for governments to allocate resources and set relevant health agendas.³⁻⁴

Sierra Leone made remarkable strides and laudable progress towards the implementation of the Millennium Developmental Goals from around 2002 despite the civil war and an Ebola virus outbreak in May 2014.⁵ Sierra Leone's government attempted to address the health needs of their population by increasing the healthcare finance budget by 34%, sourcing 86.5% of necessary funds from external partners for the Free Health Initiative.⁶ The Free Health Initiative 2010 for women and children increased and improved health access and coverage to address high morbidity and mortality in women and children.

In this study, we report on the GBD study focusing on Sierra Leone from 1990 to 2017 (27 years). In Sierra Leone, the burden of disease is characterised by a combination of persistent, emerging and re-emerging infectious diseases and increasing chronic conditions and injuries.⁷ Sierra Leone is experiencing a double burden of non-communicable diseases (NCDs) and communicable diseases, maternal, neonatal and nutritional disease (CMNNs). As in most developing countries, the burden of CMNNs seems to be decreasing but with fluctuations caused by persistent malaria. The burden of NCDs also seems to be steadily decreasing but hypertension, alcohol and substance abuse are prevalent in the society and NCDs are predicted to increase. We analyse the GBD data to describe changing burdens of CMNNs and NCDs over a 27-year period.

Methods

We conducted a cross sectional study using secondary data from IHME GBD database.³ Data from 1990 to 2017 were extracted on the causes of mortality and morbidity for all age groups and both genders. The GBD estimates burden of disease using quality-controlled, bias-corrected data sources, including country wide surveys, birth and death registration systems, census and disease surveillance which are released annually. The data are analysed using standardised statistical estimation and cross-validated to assess model performance.³ Sampling and non-sampling error in the data and model assumptions are

accounted for by reporting 95% uncertainty intervals (UIs) for all GBD estimates. The UIs are derived from the 2.5th and 97.5 percentiles of 1 000 draws.³ Complete information on the GBD data sources are available from the Global Health Data Exchange. Data can be explored and visualised on the IHME website. The GBD framework also classifies causes of health loss into mutually exclusive and collectively exhaustive categories organised in a four-level hierarchy.³ The causes of health loss are first organised into three primary categories namely CMNNs, NCDs and injuries. These broad categories are divided further into increasingly more detailed categories in a consistent and comprehensive manner.³ Standard estimates for different causes of health loss are produced for different sexes and age groups by country, enabling useful comparisons.

Mortality estimates

The IHME GBD estimates mortality rates of adults of all ages and both sexes including children under five.⁸ Data for children under five and adults are separated using Gaussian and spatiotemporal regressions. Cause-specific mortality is estimated using standard data sources which show cause of death including death registrations, reports from autopsies and surveillance.⁸ Data with no cause of death are allocated garbage codes and redistributed using standard algorithms. The Cause of Death Ensemble mode on the IHME website uses country-level covariates and builds models which are combined and tested to provide the most robust estimates for cause-specific mortality. Models for cause-specific mortality are combined and corrected to be internally consistent with estimates of all-cause mortality using the cause of death correction process, Cod Correct. Global burden of disease data are expressed as various metrics, including mortality rates, and years of life lost due to premature death (YLLs).³ In this study we will focus on YLLs and contributing risk factors.

Years of life lost

The Years of life lost summarizes years lost to premature death, at which age death occurred and the frequency of deaths.⁸⁻⁹ YLL is expressed per 100000 population.³ YLL is calculated using the formula: $YLL = N_{(causeofdeath+age+year)} \times L_{(sex+age)}$, where N=mortality and L=standard life expectancy at the age at which the death occurred.^{3,9} The formula was developed during consultations, collaboration and research with experts and is supported by the World Global Health Estimates. In 2010, the GBD study simplified the calculation. The values were acknowledged and adopted by the World Health Organisation (WHO).⁹

Results

Between 1990 and 2017, the burden of both CMNNs and NCDs declined for men and women. By 2017, Sierra Leone still had a larger burden of CMNNs than NCDs, although the burden of CMNNs had declined markedly since 1990 (Fig. 1). In 1990, men had a greater burden of NCDs, but by 2017 the gap between

men and women had narrowed. The burden of CMNNs dropped remarkably for both men and women (Fig. 1). Despite the declining burden of CMNNs, men were still more affected than women. We noted increases in CMNNs in 1997 and 2014, for both men and women, hinting at events that destabilised the health system.

Cause-specific mortality

YLLs declined for most causes between 1990–2017.

Trends in CMNNs and NCDs

The burden of respiratory infections peaked in 1990 accounting for an estimated 25000 deaths per 100000 (Fig. 2). The burden of respiratory infections declined steadily to about 8000 deaths per 100000 in 2017. In 2017, respiratory infections remained the 3rd largest contributor to YLLs.

Maternal and neonatal deaths were the 2nd largest contributor to YLLs in the period under review. Although maternal and neonatal deaths have declined steadily from 1996 to 2017, these deaths have remained an important contributor to total YLLs. In 2000, maternal and neonatal deaths (17000/100000) overtook respiratory infections and tuberculosis (16000/100000) (Fig. 2).

The most important contributor to YLLs in Sierra Leone over the study period were neglected tropical diseases including malaria (Fig. 2). In 1990, these diseases caused about 18000 deaths/100000 population, and steadily increased to about 20000 deaths/100000 population in 2000 and peaked at 24000 deaths per 100000 in 2004. This peak lasted until 2008, when YLL due to neglected tropical diseases and malaria started to decline (Fig. 2).

The burden of YLL due to enteric diseases and other infectious diseases have declined dramatically over the 27-year review period. Yearly lives lost due to HIV/AIDS and sexually transmitted infections gradually increased from 1000 per 100000 in 1990 to 4000 per 100000 in 2017.

Risk factors of CMNNs

Table 1 summarizes the risk factors for CMNNs. Throughout the study period, child and maternal malnutrition-related problems were ranked first. Secondly, contaminated drinking water, inadequate sanitation and a lack of handwashing facilities continue to be an issue in Sierra Leone. Exposure to air pollution was rated as the third most important risk factor for YLL. Between 2000 and 2017, the importance of cigarette consumption declined from fourth to sixth place. Over the 27-year study period, the importance of risky sexual practices grew from seventh to fourth. Fasting glucose levels were first placed sixth, but then dropped to eighth position. Intimate partner violence increased in prominence, increasing from eighth to seventh place (1990–2000). From 1990 to 2017, the importance of drug use remained constant, ranking ninth.

Table 1: Risk factors contributing to CMNNs in Sierra Leone from 1990 to 2017.

Risk	1990	1995	2000	2005	2010	2015	2016	2017	Average Rank
Child and maternal malnutrition	1	1	1	1	1	1	1	1	1
Unsafe water, sanitation, and handwashing	2	2	2	2	2	2	2	2	2
Air pollution	3	3	3	3	3	3	3	3	3
Alcohol use	5	5	5	5	5	5	5	5	4
Tobacco	4	4	4	6	6	6	6	6	5
Unsafe sex	7	7	6	4	4	4	4	4	5
High fasting plasma glucose	6	6	7	8	8	8	8	7	7
Intimate partner violence	8	8	8	7	7	7	7	8	7
Drug use	9	9	9	9	9	9	9	9	9

Risk Factors of NCDs

Table 2 displays the risk variables for NCDs in Sierra Leone between 1990 and 2017. The GBD study identifies 16 risk factors for noncommunicable diseases in Sierra Leone. During 1990 and 2017, the most important risk factor for NCDs were high systolic blood pressure and dietary hazards (Table 2). In 2000, fasting glucose became the third most important risk factor, a position it held for seventeen years.

Tobacco and alcohol usage have diminished in relevance as risk factors. Tobacco use was ranked third in 1990, fifth in 2000, and finally sixth in 2010. In 2010, alcohol consumption slipped from seventh to ninth place, where it remained until 2017. From 1990 to 2017, all drug consumption was rated tenth. From ninth place in 1990 to fourth place in 2017, a high body mass index increased in prominence as a risk factor. From 1990 to 2017, environmental risks and child and maternal malnutrition maintained consistent rankings (Table 2).

Table 2: Risk factors contributing to non-Communicable disease for 1990 -2017

Risk	Rank								Average Rank
	1990	1995	2000	2005	2010	2015	2016	2017	
High systolic blood pressure	1	1	1	1	1	1	1	1	1,00
Dietary risks	2	2	2	2	2	2	2	2	2,00
High fasting plasma glucose	6	6	3	3	3	3	3	3	3,86
Air pollution	5	4	4	4	4	4	4	5	4,25
Tobacco	3	3	5	5	5	6	6	6	4,46
Impaired kidney function	4	5	6	6	6	7	7	7	5,75
Alcohol use	7	7	7	7	7	9	9	9	7,36
High body-mass index	9	9	9	9	8	5	5	4	8,07
High LDL cholesterol	8	8	8	8	9	8	8	8	8,25
Other environmental risks	10	10	10	10	10	10	10	10	10,00
Unsafe sex	11	11	11	11	11	11	11	11	11,00
Low physical activity	12	12	12	12	12	12	12	12	12,00
Occupational risks	13	13	13	13	13	13	13	13	13,00
Drug use	14	14	14	14	14	14	14	14	14,00
Child and maternal malnutrition	15	15	15	15	15	15	15	15	15,00
Childhood maltreatment	16	16	16	16	16	16	16	16	16,00

Discussion

In this GBD study, we describe the burden of disease in Sierra Leone from 1990 to 2017. Overall, the burden of disease declined resulting in fewer YLLs. In Sierra Leone, CMNNs continue to be a problem due to the prevalence of endemic disease. CMNNs have a greater effect on male mortality than on female mortality. The most important CMNNs are respiratory infections, neglected tropical diseases and malaria, and maternal and neonatal disease. The burden of NCDs was low compared to CMNNs, a trend which is likely to change as the health system recovers and populations age. Sierra Leone faces a dual burden of disease, with NCDs contributing about 29% while CMNNs account for 65% of YLLs and 6% represent injuries.¹⁰⁻¹¹ The end of the Sierra Leone civil war brought the government, international partners, stakeholders and civil society together to start reconstructing the health system. The recovering health system and implementation of health policy interventions has resulted in the decline of YLLs due to CMNNs and NCDs. Our study shows that Sierra Leone has made progress in population health outcomes despite multiple drawbacks.

The Sierra Leone civil war took place from 1991 to 2002, lasted 11 years and left more than fifty thousand people dead. The civil war would have contributed to burden of disease in several ways, including an increase in injuries. Mortality rates due to NCDs peaked between 1990 and 1994, reflecting the potential impact of the civil war. The mortality rates of men due to NCDs at this time were also markedly higher than the mortality rates for women. Mortality rates have declined consistently after the end of the civil war, suggesting efforts to rebuild the health system. Men and women's mortality rates

equalised after 2010. In 2014, mortality increased slightly due to the Ebola outbreak which lasted for two years.

Sierra Leone had to address a greater burden of CMNNs than NCDs from 1990–2017. The burden of CMNNs is high when compared to other countries.³ Malaria and neglected tropical diseases, respiratory infections contribute to significantly to YLLs and disability.³ The burden of CMNNs peaked in 1990 and 1992 and can be attributed to persistent endemic malaria.¹² Sierra Leone health services treat approximately 2,240,000 outpatients annually for malaria and almost half of these are children under the age of five years.^{13–14} Malaria mortality was estimated at approximately 4.4% of pregnant women and 17% of children. Malaria contributes to 40% of hospitalised morbidity in all ages and 37% of children under five.¹³ In 2004, Sierra Leone launched their first National Malaria Strategic Plan (2004–2008), which was funded by the Global fund.¹³

In Sierra Leone, maternal and neonatal disorders remained high from 1990 to 2017. In Sierra Leone, children under five years suffer high mortality rates with 120 deaths per 1000 children. Maternal deaths are amongst the highest in the world at 1360 deaths per 100000 live births in 2015, which far exceeds the MDG targets of 450 deaths per 100,000 births.^{11,15} In this study, the burden of maternal and neonatal disorders dropped from 21833.31 per 100,000 in 1990 to 8 889.03 in 2017.³ In 2010, the Sierra Leone government launched the Free Health Initiative to reduce mortality and morbidity due maternal and neonatal disorders.¹¹

Aside from malaria and maternal and neonatal mortality, respiratory diseases and tuberculosis, enteric infections and HIV also contribute to YLLs in Sierra Leone. In 1990, the German Leprosy Relief Association assisted the Ministry of Health and Sanitation to establish the National Leprosy and Tuberculosis Control Programme to monitor the surveillance of tuberculosis control activities.¹⁶ In this study, respiratory disease YLLs decreased from 24255 fatalities per 100,000 population in 1990 to 7330 deaths per 100,000 population in 1997. Sierra Leone continues to have one of the highest tuberculosis burdens in the world despite the fact that treatment is free and readily available.¹⁷ Sierra Leone opened its first drug-resistant tuberculosis treatment centre at Lakka Government Hospital in 2017. Nonetheless, new recommendations, constant monitoring and surveillance of the National Tuberculosis Program remain necessary.¹⁷

The frequency of enteric infections has dropped from 17210 deaths per 100,000 persons in 1990 to 4558 deaths per 100,000 persons in 2017.³ Enteric diseases are most prevalent in children under the age of five and account for around 12% of all child deaths in Sierra Leone.¹⁸ Sierra Leone added the rotavirus vaccine to its immunization schedule to combat diarrhoeal infections on March 28, 2014, in an effort to address this issue.¹⁸

The burden of HIV/AIDS and sexually transmitted infections (STIs) increased from 1101 to 2124 cases per 100,000 population.³ In 2013 and 2014, commercial sex workers were responsible for 40% of newly

infected HIV patients.¹⁹ The Sierra Leone government is stepping up efforts to test, prevent, treat and increase awareness with the support of the WHO, Global Fund and many other partners. The Sierra Leone government has also implemented a national HIV AIDS strategic plan 2016–2020, including programmes such as Prevention of Mother to Child Transmission.¹⁷

As a developing country with a relatively young population, the risk factors associated with YLLs due to CMNNs are also linked to the health and wellbeing of younger age groups. The most important risk factors for CMMN YLLs were environmental risk factors including child and maternal nutrition, unsafe water and sanitation and exposure to air pollution. Less important risk factors included lifestyle risk factors such as alcohol and tobacco use, drug use and intimate partner violence. In Sierra Leone, environmental risk factors are being addressed on various fronts. In 2012, the presidency committed to the UNICEF Scaling Up Nutrition Movement and adopted a Food and Nutrition Security policy to secure the nutritional well-being of infants and women. Unsafe water and sanitation are other environmental risk factors being considered by the WHO and UNICEF who are trying to improve water and sanitation services by conducting risk assessments and improving health facilities that were destroyed during the civil war.

Additionally, lifestyle risk factors such as HIV/Aids and sexually transmitted diseases, as well as respiratory illnesses, contribute to CMMNs. In Sierra Leone, alcohol consumption is ranked in the fifth place, while cigarette consumption is placed sixth as risk factors for YLLs. These lifestyle risk variables were associated with a relatively small number of deaths; for example, cigarette smoking was associated with 5% of YLLs.³ Although the number of YLLs connected with these risk factors is still small, it is increasing and requires monitoring by local organizations. Sierra Leone signed the WHO Framework Convention on Tobacco Control in May 2009, with the objective of reducing tobacco consumption, and the Ministry of Health and Sanitation adopted a National Tobacco Control Strategic Plan in 2012.¹⁷

In this study, CMNN contributed significantly more to YLLs than NCDs from 1990–2017. NCDs are expected to grow by 25% globally by 2030.²⁰ Sierra Leone was also predicted to have an increase.^{20–22} In 2008, the WHO estimated that NCDs accounted for 18% of fatalities in Sierra Leone, followed by cardiovascular disease at 7%, cancer at 3%, diabetes at 1%, and chronic respiratory illness at 2%.²¹ In 2012, mortality from NCDs increased to 26%. Sierra Leone's government developed its first strategic plan and policy for NCDs in 2013, in response to the World Health Organization's global status report on NCDs. Sierra Leone's strategic plan, which ran from 2013 to 2017, attempted to mitigate the burden of NCDs such as cardiovascular disease, chronic lung disease, diabetes mellitus, obesity, cancer, sickle cell disease, mental disorders, and epilepsy.^{21,23} By 2014, the burden of NCDs had reduced across all age groups and sexes, with men accounting for 323 deaths per 100,000 and women accounting for 321 deaths per 100,000.²²

The burden of NCDs remained relatively constant between 2005 and 2017. In our study, most YLLs due to NCDs can be attributed to cardiovascular related diseases and neoplasms.³ Cardiovascular disease contributed to approximately 9% of NCD deaths (this study) and has been among the top ten causes of

YLLs since 1990.²¹ There is evidence that NCDs are increasing. In 1993, 68% of hospitalisations at Freetown hospital were admitted due stroke.²¹ In 1994, 25% of the population above 50 years of age were estimated to be hypertensive.²¹ A review of death certificates issued between 1983 and 1992, showed an increase in deaths related to hypertension in Sierra Leone.²¹ There is little information on the prevalence of cancer in Sierra Leone, even though our study reported that neoplasms were among the top ten causes of mortality.²¹ In Sierra Leone, recording and reporting of data on NCDs remains inconsistent even though there is a ministerial department responsible for NCDs.²¹

Sierra Leone suffered an Ebola outbreak in 2014 and 2015, which led to inadequate quality surveillance data on the incidences, cases and deaths of NCDs.²⁴ The Ebola outbreak occurred when the government was transitioning from hospital care for NCDs to management, treatment and care in primary health care facilities.²⁴ Following the Ebola outbreak, significant reporting systems focusing on morbidity and risk factors for NCDs were put in place. Although policies are being developed by the government, there seems to be little funding for treating and controlling NCDs.²⁴ The most dominant risk factors associated with NCDs include high blood pressure, dietary risks, high fasting plasma glucose, high body mass index and the use of alcohol and tobacco (this study). In Sierra Leone, DALYs due to high blood pressure have also increased (18%) from 2007 to 2017.

Dietary risks are also associated with YLLs due to NCDs in Sierra Leone. A nutritional survey done in 2014 revealed that more than 25% of children younger than five years old had stunted growth.^{21,24} Glucose has recently become an important risk factor NCD associated YLLs and is growing in importance. High fasting plasma glucose is an indicator of diabetes mellitus. The prevalence of diabetes in Sierra Leone has also increased from 2.4% in 1997 to 7% in 2014.²¹

Tobacco use is an important risk factor of NCDs, including cardiovascular disease, respiratory diseases and lung cancers.²¹ In Sierra Leone, 14.3% of men and 1.4% of women, comprising 34% of people, smoke more than 10 cigarettes a day.²¹ Consuming alcohol is an important risk factor for cardiovascular disease. In 2005, 6.5% of adults older than 15 years consumed alcohol. In 2014, 17% of adults between the ages of 25 and 64 years old were estimated to consume alcohol.²¹ Added to problems of hypertension, glucose and substance abuse is the fact that Sierra Leoneans engage in low to moderate physical activity. The importance of high body mass index as a risk factor jumped from 9th in 1990 to 5th place in 2017.³ The burden of NCDs remains low compared to CMNNs, which may contribute to few resources being allocated to preventing NCDs at this time.

Conclusion

We described the burden of disease profile in Sierra Leone. We described the trends and patterns of CMNNs and NCDs in Sierra Leone for the period 1990 to 2017. The burden of disease, expressed as YLLs, in Sierra Leone declined from 1990 to 2017. During this time, the most dramatic decreases were seen in YLLs attributed to CMNNs. YLLs due to CMNNS remain higher than NCDs due the presence of endemic

diseases including respiratory infections, neglected tropical diseases and malaria. Maternal and neonatal disease also contributed to YLLs. The high burden of these conditions is driven by environmental risk factors including inadequate nutrition, unsafe water and sanitation and air pollution. The burden of NCDs was represented by cardiovascular disease and to a lesser extent neoplasms. Although low compared to CMNNs, the Sierra Leonean government should monitor the impact of NCDs, to inform health promotion strategies. As the health system recovers from the civil war and the Ebola outbreak, the quality of health care will improve, and the population will age. As with other developing countries, aging populations are associated with a greater burden of NCDs. The end of the civil war brought together the government, international partners, stakeholders and civil society to start reconstructing the health system and implementing health policy interventions which have contributed to the decline of CMNNs and NCDs.

Abbreviations

AIDS Acquired Immune Deficiency Syndrome

CMNN Communicable Maternal Neonatal Nutritional Diseases

DALY Disability-Adjusted Life Years

DHIS District Health Information System

GDB Global Burden of Disease

HALE Health Adjusted Life expectancy

HC Health Centre

HIV Human Immunodeficiency Virus

HRH Human Resource for Health

IHME Institute for Health Metrics and Evaluation

MoH Ministry of Health

MDG Millennium Development Goal

NCD Non-Communicable Disease

NGO Non-Governmental Organization

SDG Sustainable Development Goal

TB Tuberculosis

UN United Nations

WHO World Health Organization

YLL Years of Life Lost

YLD Years of Life Lost to Disability

Declarations

Ethical Considerations

The protocol was tabled at the Academic Planning Committee (APC: SHSPH) meeting for approval. The protocol was sent to the University of Pretoria Ethics Committee after attending TNM 800 module for ethical approval and clearance to conduct the study. The ethics committee approved and allocated the Ethics (Protocol Number: 515/2019). An approval certificate is attached. The Data in the Institute for Health Metrics and Evaluation (IHME) site has unrestricted access for public use. All data were anonymous and there were no concerns about privacy

Competing interests

JZ: None declared

FS: None declared

TC: None declared

TA: None declared Authors' contributions

JZ: wrote the first and subsequent drafts of the manuscripts

FS: conceived and supervised the project and contributed to the subsequent writing and interpretation of the manuscript.

TC: participated in the analysis and interpretation of the results

TA: contributed to the subsequent writing of the manuscript

Acknowledgements

Dr. Cheryl Tosh for editing.

References

1. Edoka I, McPake B, Ensor T, Amara R, Edem-Hotah J. Changes in catastrophic health expenditure in post-conflict Sierra Leone: An oaxaca-blinder decomposition analysis. International Journal for Equity in Health 2017. p. 1–15.

2. Voors M VDWP Kostadis J,Papaioannou Bulte E. Resources and governance in Sierra Leone's civil war. *The Journal of Development Studies*. 2016; 53(2):278–94.
3. Institute for Health Metrics and Evaluation (IHME). Seattle WI, 2017. Global burden of disease Sierra Leone. IHME. 2017.
4. ACAPS. Sierra leone:Country profile. 2014:1–12.
5. Government of Sierra leone MoHaS. Interim millennium development goals report 2015 seirra Leone. February 2016.
6. Sanitation GoS MoHa. National health sector strategic plan 2010–2015 2009.
7. Group WB. Epidemiologic transition theory exception. World Bank. 2001:1–28.
8. WHO Dol, Evidence and Research WHO methods and data sources for global burdern of disease estimates 2000–2016. 2018; 4(4).
9. (WHO) WHO [Internet] About the global burden of disease project. 2017.
10. Organisation WH. WHO country cooperation strategy Sierra Leone, 2017–2021. 2017.
11. WHO U, UNPFA, WORLD BANK GROUP,UNITED NATIONS POPULATION DIVISION. Trends in maternal mortality :1990–2015. World Health Organization 2015.
12. Government of Sierra leone MoHaS. Annual health sector perfomance report 2016. 2016.
13. Government of Sierra leone MoHaS. Sierra Leone malaria control strategic plan 2016–2020. 2015.
14. Sierra Leone: A profile of malaria control and epidemiology. 2015.
15. Figueroa CA, Linhart CL, Beckley W, Pardosi JF. Maternal mortality in Sierra Leone: From civil war to ebola and the sustainable development goals. *International Journal of Public Health: International Journal of Public Health*. 2018; 63(4):431–2.
16. WHO. About us Sierra Leone. WHO Africa. 2017.
17. WHO. World health organisation Sierra Leone annual report 2017. WHO. 2017.
18. Government of Sierra leone MoHaS, Health education division 2017–2021. National health promotion strategy of Sierra Leone. 2016.
19. Government of Sierra leone MoHaS, Health Education Division National health strategy of Sierra Leone 2017–2021. 2016.
20. Bennett JE, Stevens GA, Mathers CD, Bonita R, Rehm J, Kruk ME, et al. Ncd countdown 2030: Worldwide trends in non-communicable disease mortality and progress towards sustainable development goal target 3.4. *The Lancet*. 2018; 392(10152):1072–88.
21. Idriss A WHR, Bertone M, et al. A scoping study on non- communicable diseases (ncd) in Sierra Leone 2018.
22. Islam SMS PT, Phuong NTA,et al. Non communicable diseases (ncds) in developing countries: A symposium report. *Globalization and health*. 2014.
23. Ministry of Health and Sanitation DoRacH. National ncd policy and strategic plan 2013–2017 2013.

Figures




Figure 1

Trends in burdens of CMNNs and NCDs by gender for Sierra Leone from 1990-2017.

Top 10 Diseases YLLs 1990 - 2017 (CMNN and NCD)




Figure 2

Trends (YLLs) for top CMNNs and NCDs in Sierra Leone from 1990 - 2017