

# Effect of a real-time automatic nosocomial infection surveillance system on hospital- acquired infection prevention and control

**Ruiling Wen** (✉ [ruiling622@126.com](mailto:ruiling622@126.com))

Department of Infection Management, Huizhou First People's Hospital

**Xinying Li**

Department of Infection Management, Huizhou First People's Hospital

**Tingting Liu**

Department of Infection Management, Huizhou First People's Hospital

**Guihong Lin**

Department of Infection Management, Huizhou First People's Hospital

---

## Research Article

**Keywords:** Hospital-acquired infections, multidrug resistant organisms, infections surveillance, infection control

**Posted Date:** May 17th, 2022

**DOI:** <https://doi.org/10.21203/rs.3.rs-1548386/v1>

**License:** © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

---

## Abstract

**Background:** The systematic collection of valid data related to hospital-acquired infections (HAIs) is considered effective for nosocomial infection prevention and control programs. New strategies to reduce HAIs have recently fueled the adoption of real-time automatic nosocomial infection surveillance systems (RT-NISSs). Although RT-NISSs have been implemented in some hospitals for several years, the effect of RT-NISS on HAI prevention and control needs to be further explored.

**Methods:** A retrospective, descriptive analysis of inpatients from January 2017 to December 2019 was performed. We collected hospital-acquired infection (HAI) cases and multidrug resistant organism (MDRO) infection cases by traditional surveillance in period 1 (from January 2017 to December 2017), and these cases were collected in period 2 (from January 2018 to December 2018) and period 3 (from January 2019 to December 2019) using a real-time nosocomial infection surveillance system (RT-NISS). The accuracy of MDRO infection surveillance results over the 3 periods was examined. The trends of antibiotic utilization rates and pathogen culture rates in periods 2 and 3 were also analysed.

**Results:** A total of 114647 inpatients, including 2242 HAI cases, were analysed. The incidence of HAIs in period 2 was significantly greater than that in period 1 (2.28% vs. 1.48%,  $\chi^2=61.963$ ,  $p<0.001$ ) and period 3 (2.28% vs. 2.05%,  $\chi^2=4.767$ ,  $p=0.029$ ). The incidence of five HAI sites, including respiratory infection, urinary tract infection (UTI), surgical site infection (SSI), bloodstream infection (BSI) and skin and soft tissue infection, was significantly greater in period 2 compared with period 1 (both  $p<0.05$ ) but was not significantly different from that in period 3. The incidence of hospital-acquired MDRO infections in period 3 was lower than that in period 2. The identification of MDRO infection cases using the RT-NISS achieved a high level of sensitivity (Se), specificity (Sp), positive predictive value (PPV) and negative predictive value (NPV), especially in period 3 (Se=100%, Sp=100%, PPV=100% and NPV=100%).

**Conclusion:** The adoption of a RT-NISS to adequately and accurately collect HAI cases is useful to prevent and control HAIs. Furthermore, RT-NISSs improve accuracy in MDRO infection case reporting, which can timely and accurately guide and supervise clinicians in implementing MDRO infection prevention and control measures.

## Background

Hospital-acquired infections (HAIs), also known as nosocomial infections (NIs), are currently one of the most important challenges for modern medicine [1, 2]. Patients with HAIs might have prolonged hospital stays and high mortality, thus not only threatening the safety of patients but also causing a significant waste of social and economic resources, representing an important public health problem threatening human health [3–6]. Nosocomial infection surveillance is an important basis for controlling the occurrence and development of HAIs [6–10]. Hospitals should detect HAI cases and outbreaks of HAIs in a timely manner, analyse causes and take effective prevention and control measures. Timely reporting of nosocomial infection cases is of great significance for preventing nosocomial infection outbreaks and improving the quality of hospital management. However, due to the wide coverage and complexity of HAIs, a large amount of data needs to be analysed statistically, and traditional manual surveillance is inaccurate and inefficient and has been unable to meet the actual needs of HAI management [9]. The real-time automatic nosocomial infection surveillance system (RT-NISS) is a real-time, automatic and effective monitoring system that has been developed methodologically and practically in a stepwise manner and is a reliable surveillance intelligent information technology tool for infection control physicians to systematically evaluate HAI information. A RT-NISS enhances case-finding efficiency by automatically and systematically selecting patients with the highest HAI probability. Once infection control physicians identify HAI cases, they can take strategies and measures to control HAI cases and accumulate experience to prevent subsequent HAI events [8, 11]. Therefore, the adoption of RT-NISS is one of the core components of infection prevention and control programs for modern medicine.

RT-NISS have been implemented in some hospitals for several years, but it remains an open question whether the use of a RT-NISS is effective in the reduction of nosocomial infections. Our study aimed to describe and analyse the effect of a RT-NISS on HAI prevention and control using surveillance data for a 3-year period from 2017 to 2019.

## Methods

### Patient population and study design

We conducted a retrospective study of all patients admitted to Huizhou First Peoples' Hospital (HZFH) from 2017 to 2019. Patients were categorized into three groups (Fig. 1): the first group (period 1) included patients admitted to HZFH from January 2017 to December 2017. We used traditional surveillance to collect hospital-acquired infection (HAI) cases and multidrug resistant organism (MDRO) infection cases. The second group (period 2) and the third group (period 3) included all patients admitted to HZFH from January 2018 to December 2018 and January 2019 to December 2019, respectively. In these two groups, we collected information regarding HAI cases, MDRO cases, the utilization rate of antibiotics and the rate of pathogen culture using the real-time nosocomial infection surveillance system (RT-NISS). Furthermore, all of

the MDRO infection cases were peer-reviewed by senior infection control physicians to examine the accuracy of the surveillance results of the 3 periods.

Traditional surveillance is a time-consuming manual inspection that is mainly based on daily assessment of laboratory results and drug prescriptions from the pharmacy service, regularly visiting patients' clinical histories and routinely receiving information on HAI cases reported by clinicians. The RT-NISS used in our study was systematically applied to monitor, diagnose, and control HAIs under the strict supervision of infection control physicians. The RT-NISS selects patients with the highest HAI probability by connecting other hospital information systems, including the hospital information system (HIS), laboratory information system (LIS), radiology information system (RIS) and anaesthesia operation system (AOS), and sets a filter to acquire the necessary infection information (Fig. 2).

## Definitions

Hospital-acquired infections (HAIs) are infections occurring 48 h after hospital admission and have no evidence of subclinical infection at the time of admission. HAIs are classified as respiratory infection, urinary tract infection (UTI), gastrointestinal tract infection, surgical site infection (SSI), bloodstream infection (BSI), skin and soft tissue infection and other infections according to the specific infection site. MDROs were defined as those bacteria that were resistant to at least 3 antimicrobial classes, including extended-spectrum beta-lactamase (ESBL)-producing Gram-negative bacilli (mainly *Escherichia coli* and *Klebsiella pneumoniae*), methicillin-resistant *Staphylococcus aureus* (MRSA), carbapenem-resistant *Acinetobacter baumannii* (CRAB), carbapenem-resistant *Pseudomonas aeruginosa* (CRPA), carbapenem-resistant *Enterobacteriaceae* (CRE) and vancomycin-resistant enterococci (VRE).

## Statistical Methods

The following parameters were assessed for MDRO cases: TP = true positive (the collected positive case is truly positive), FP = false positive (negative case is classified as positive), FN = false negative (positive case is classified as negative), TN = true negative (the collected negative case is truly negative), Se = sensitivity (the proportion of positive cases correctly classified), Sp = specificity (the proportion of negative cases correctly collected), PPV = positive predictive value (the proportion of positive predictions that are correct) and NPV = negative predictive value (the proportion of negative predictions that are correct). The  $\chi^2$ -test was used for the comparison of categorical variables. All statistical analyses were performed by using SPSS 22.0 software (IBM SPSS), and 2-tailed p values < 0.05 were considered statistically significant.

## Results

### A comparison between traditional surveillance and RT-NISS

We used traditional surveillance to collect 34197 patients, including 505 (1.48%) HAI cases in period 1. Of these HAI patients, 318 (0.93%) suffered respiratory infection, 84 (0.25%) suffered UTI, 25 (0.07%) suffered GTI, 30 (0.09%) suffered SSI, 48 (0.14%) suffered BSI and 40 (0.12%) suffered skin and soft tissue infection. In period 2, 38028 patients, including 866 (2.28%) HAI cases, were collected from the RT-NISS. Of these HAI patients, 526 (1.38%) suffered respiratory infection, 219 (0.58%) suffered UTI, 31 (0.08%) suffered GTI, 55 (0.14%) suffered SSI, 86 (0.23%) suffered BSI and 80 (0.21%) suffered skin and soft tissue infection. The incidence of HAIs in period 2 was significantly greater than that in period 1 ( $\chi^2=61.963$ ,  $p < 0.001$ ). With the exception of GTI, the incidences of different HAI sites, including respiratory infection, UTI, SSI, BSI and skin and soft tissue infection, in period 2 were also significantly greater than those in period 1 (Table 1). The incidence of hospital-acquired MDROs in period 1 was 0.56% (191 cases). However, in period 2, the incidence was 0.52% (197 cases), and no significant difference was noted compared with period 1. These results demonstrated that the adoption of RT-NISS improved the detection rate of HAI cases and eliminated the serious underreporting phenomenon of traditional surveillance.

Table 1  
Comparative results of HAIs reported by traditional surveillance during period 1 and RT-NISS in periods 2 and 3

Infection type	Period 1 (n = 34197)	Period 2 (n = 38028)	Period 3 (n = 42422)	$\chi^2_1$	P <sub>1</sub>	$\chi^2_2$	P <sub>2</sub>
HAIs cases (n/%)	505 (1.48)	866 (2.28)	871 (2.05)	61.963	< 0.001	4.767	0.029
Respiratory infection	318 (0.93)	526 (1.38)	538 (1.27)	32.033	< 0.001	2.031	0.154
UTI	84 (0.25)	219 (0.58)	202 (0.48)	47.008	< 0.001	3.831	0.050
GTI	25 (0.07)	31 (0.08)	41 (0.10)	0.164	0.685	0.513	0.474
SSI	30 (0.09)	55 (0.14)	87 (0.21)	5.232	0.022	3.847	0.050
BSI	48 (0.14)	86 (0.23)	89 (0.21)	7.155	0.007	0.247	0.619
Skin and soft tissue infection	40 (0.12)	80 (0.21)	78 (0.18)	9.470	0.002	0.719	0.397
Hospital-acquired MDROs (n/%)	191 (0.56)	197 (0.52)	173 (0.41)	0.552	0.457	5.322	0.021

UTI: urinary tract infection, GTI: gastrointestinal tract infection, SSI: surgical site infection, BSI: bloodstream infection,  $\chi^2_1$  and P<sub>1</sub> for comparison between period 1 and period 2,  $\chi^2_2$  and P<sub>2</sub> for comparison between period 2 and period 3

## The Effect Of Rt-niss On Hai Prevention And Control

In period 3, we used RT-NISS to collect 42422 patients, including 871 (2.05%) HAI cases. Of these HAI patients, 538 (1.27%) suffered respiratory infection, 202 (0.48%) suffered UTI, 41 (0.10%) suffered GTI, 87 (0.21%) suffered SSI, 89 (0.21%) suffered BSI and 78 (0.18%) suffered skin and soft tissue infection. The incidence of HAIs in period 3 was significantly lower than that in period 2 ( $\chi^2=4.767$ ,  $p = 0.029$ ). The incidences of four hospital-acquired infection sites, including respiratory infection, UTI, BSI and skin and soft tissue infection, in period 3 were slightly lower than those in period 2, but the difference was not statistically significant. The incidence of hospital-acquired MDROs was 0.41% (173 cases) in period 3, which was significantly lower than that in period 2 ( $\chi^2=5.322$ ,  $p = 0.021$ ). After RT-NISS implementation in period 2, we collected HAI cases and analysed the risk factors for HAIs. Then, we took these risk factors into consideration and implemented some interventions that could prevent and control HAI events in period 3. As shown, the adoption of RT-NISS coupled with the implemented interventions in period 3 could avoid HAI events to some extent and reduce the HAI rate.

## Rt-niss Improves Accuracy In Mdro Case Reporting

To further investigate the effect of the adoption of RT-NISS on certain HAI prevention and control programs, such as MDRO infection case surveillance and control, we examined the accuracy of MDRO infection case reporting by traditional surveillance and RT-NISS. As shown in Table 2, high numbers of FPs and FNs in both varieties of MDROs, including ESBL, MRSA, CRAB, CRPA and CRE, were noted in period 1. In periods 2 and 3, the number of FPs and FNs of both varieties of MDROs was significantly reduced. In particular, in period 3, these values were reduced to 0. As shown in Table 3, the Se, Sp, PPV and NPV of both 5 different varieties of MDROs, including ESBL, MRSA, CRAB, CRPA and CRE, in period 1 were significantly lower than those in period 2 and period 3. Furthermore, the Se, Sp, PPV and NPV of both varieties of MDROs in period 3 were all improved to 100%. In addition, no VRE cases were reported from 2017 to 2019 in HZFH. After the RT-NISS implementation in period 2, we collected causes of mistaken reports of MDRO infection cases, and we fixed the mistakes of the RT-NISS operation in period 3. In addition, we also monitored the utilization rate of antibiotics and the rate of pathogen culture by RT-NISS. As shown in Fig. 3, the trend of the utilization rate of antibiotics from January 2018 to December 2019 slightly decreased, whereas the rate of pathogen culture slightly improved.

Table 2

Respective comparisons of MDROs reported by traditional surveillance in period 1 and the RT-NISS in periods 2 and 3 by senior infection control physicians

Group	ESBL				MRSA				CRAB				CRPA				CRE			
	TP	FP	FN	TN	TP	FP	FN	TN	TP	FP	FN	TN	TP	FP	FN	TN	TP	FP	FN	TN
Period 1	325	82	192	562	85	91	21	114	44	23	15	106	25	52	19	236	0	0	29	1050
Period 2	388	14	13	557	101	0	19	139	50	0	7	119	38	24	5	228	0	0	14	0
Period 3	392	0	0	614	81	0	0	134	48	0	0	100	26	0	0	243	15	0	0	991

FP: true positive, FP: false positive, FN: false negative, TN: true negative

Table 3

NPV	97.31	0	100	
PPV	0	0	100	
Sp	100	100	100	
Se	0	0	100	
CRPA	NPV	92.55	97.85	100
	PPV	67.53	61.29	100
	Sp	81.94	90.48	100
CRAB	NPV	87.60	94.44	100
	PPV	65.67	100	100
	Sp	82.17	100	100
	Se	74.58	87.72	100
MRSA	NPV	84.44	87.97	100
	PPV	48.30	100	100
	Sp	55.61	100	100
	Se	80.19	84.17	100
ESBL	NPV	74.54	97.72	100
	PPV	79.85	96.52	100
	Sp	87.27	97.55	100
	Se	62.86	96.75	100
Group	Period 1	Period 2	Period 3	

## Discussion

Surveillance of HAIs is a fundamental and essential aspect of infection prevention and control programs [6–10]. Traditional surveillance is a manual time-consuming infection case-finding and reporting program that involves infection control physicians looking over lists of positive infection information of pathogen culture and patient histories daily and HAI case reports by clinicians. In this study, the results showed low infection incidences of HAIs (1.48%) and certain infection sites, including respiratory infection (0.93%), UTI (0.25%), SSI (0.09%), BSI (0.14%) and skin and soft tissue infection (0.12%), reported by traditional manual surveillance methods in period 1. These low reported infection incidences echo the inaccurate report demonstrated by other studies. Seifi’s study presented the results of under- and overreporting of HAIs by traditional surveillance [9]. Many hospitals have a large number of beds but insufficient numbers of infection control physicians, and infection control physicians need to collect large amounts of data daily, which will result in false negative infection reports. Many cases are excessive, or unnecessary antibiotic usage will lead to false-positive infection findings. In addition, many cases are not ordered for pathogen culture to cause

false negative infection reports. Moreover, the lack of knowledge regarding complicated infection cases of clinicians will lead to underreporting and mistaken reporting of infection events. Given their high workloads and need for multitasking, clinicians can make mistakes and lack sufficient time to report infection cases. All these factors make this type of manual surveillance unreliable and lack accuracy. There is no doubt that this frustrating situation would have serious consequences, including absent treatment for underreporting HAI patients, inappropriate treatment of false-positive infection patients, transmission of pathogens from HAI patients to other uninfected patients, and many HAIs and outbreaks that are not monitored and uncontrolled. Therefore, this predicament has catalysed the development and implementation of RT-NISSs.

Many studies in the past have demonstrated accurate and efficient electronic automatic surveillance systems [11–15]. Our analysis showed higher infection incidences of HAIs and certain infection sites, including respiratory infection, UTI, SSI, BSI and skin and soft tissue infection, reported by RT-NISS (both in period 2 and period 3) compared with those reported by traditional surveillance in period 1 (both  $p < 0.05$ ), which is consistent with the accurate surveillance results based on RT-NISSs reported in other studies. Mingmei's study presented the sensitivity and specificity of a RT-NISS with an approximately 200-fold time-saving capacity compared with manual surveys [12]. Studies in this field are more concerned with defects by traditional manual surveillance and the virtues by RT-NISSs on the surveillance of all HAIs. However, few analyses have evaluated the use of RT-NISSs for surveillance of different types of MDRO or have only focused on one type of MDRO at a time [16, 17]. In the present study, we demonstrated a more accurate surveillance result of MDRO reports by the RT-NISS in both periods 2 and 3 compared with traditional surveillance in period 1 (Tables 2 and 3). In period 1 of the present study, clinicians and infection control physicians obtained MDRO case information based on microbiology laboratory physicians' daily reporting. However, microbiology laboratory physicians underreported MDRO cases or reported cases in an untimely manner due to their high workload; thus, the MDRO data were inaccurate. We present the results of a lower number of FPs and FNs and higher Se, Sp, PPV and NPV of 5 different varieties of MDROs, including ESBL, MRSA, CRAB, CRPA and CRE, since RT-NISS implementation. Compared with period 1, the MDRO reporting process and surveillance changed, and the MROM information and data became reliable and accurate. MDROs are the main pathogenic bacteria of SSI, UTI, BSI and pneumonia, and these types of nosocomial infections are intractable and complex and the factor of high mortality for patients. Moreover, MDROs easily lead to nosocomial transmission and nosocomial cluster infection events [18–20]. Thus, timely and accurate monitoring of MDROs is an important targeted monitoring project for infection prevention and control programs [12]. The results of the present study about MDROs surveilled by RT-NISS could benefit and inform targeted prevention and control programs.

Another strength of our study was the investigation of the effect of the daily implementation of RT-NISS on HAI surveillance. After RT-NISS implementation in period 2, we analysed the long-term data collected by RT-NISS and acquired comprehensive knowledge about HAI risk factors. Then, we implemented some interventions, including monitoring the utilization rate of antibiotics and the rate of pathogen culture by RT-NISS, and punishment for doctors who did not implement prevention and control measures for MDROs, inappropriately administered antibiotics or without standard pathogen culture. The incidences of HAIs and hospital-acquired MDROs in period 3 were significantly lower than those in period 2 (0.41% vs. 0.52%,  $p = 0.021$ ). These results demonstrated that the adoption of RT-NISS coupled with the implemented interventions could reduce the rate of HAIs. In addition, regarding the results of the MDROs reported by RT-NISS in period 2, we further found some mistakes associated with the use of the RT-NISS, such as misclassifying CRE as ESBL and underreporting of MDROs when the LIS connection fails. After modifying the rule definition and correcting the data input, the number of FPs and FNs of 5 different varieties of MDROs, including ESBL, MRSA, CRAB, CRPA and CRE, in period 3 was reduced to 0, and the Se, Sp, PPV and NPV of 5 different varieties of MDROs in period 3 were both improved to 100%.

With an increasing number of studies in this field demonstrating the accuracy and efficiency of RT-NISS on HAI surveillance, health care authorities increasingly demand the installation and daily use of RT-NISSs as a part of quality management. Our data showed that RT-NISS could adequately and accurately collect HAI cases, which is in agreement with other studies [12, 15, 21]. Furthermore, RT-NISS improves the accuracy of MDRO infection case reports, and the adoption of a RT-NISS coupled with intervention implementations can reduce the infection incidences of HAIs and MDROs.

## Abbreviations

HAIs  
hospital-acquired infections  
RT-NISS  
real-time automatic nosocomial infection surveillance system  
MDROs  
multidrug resistant organisms  
UTI  
urinary tract infection  
SSI

surgical site infection  
BSI  
bloodstream infection  
HIS  
hospital information system  
LIS  
laboratory information system  
RIS  
radiology information system  
AOS  
anaesthesia operation system  
TP  
true positive  
FP  
false-positive  
FN  
false negative  
TN  
true negative  
Se  
sensitivity  
Sp  
specificity  
PPV  
positive predictive value  
NPV  
negative predictive value  
ESBL  
extended-spectrum beta-lactamase producing Gram-negative bacilli  
MRSA  
methicillin-resistant *Staphylococcus aureus*  
CRAB  
carbapenem-resistant *Acinetobacter baumannii*  
CRPA  
carbapenem-resistant *Pseudomonas aeruginosa*  
CRE  
carbapenem-resistant *Enterobacteriaceae*  
VRE  
vancomycin-resistant enterococci.

## Declarations

### Acknowledgements

All authors want to thank the departments of Huizhou First People's Hospital that strongly supported the infection prevention and control programs.

### Authors' contributions

RW provided insightful thoughts and design, analysed and interpreted the data and wrote the manuscript. XY supplied the support, resources, and supervision required to perform the study and revised the manuscript critically for important intellectual content. TL and GL obtained the data, curated the patient histories, and provided supervision. All authors read and approved the final manuscript.

### Funding

This work was supported by grants from the Science and Technology Project of Huizhou, Guangdong Province, China (2020Y060 to RW).

## Availability of data and materials

The datasets analysed during the current study are not publicly available due to privacy or ethical restrictions but are available from the corresponding author on a reasonable request.

## Ethics approval and consent to participate

This study was reviewed and approved by the Ethics Committee of Huizhou First People's Hospital. The ethics committee approved the waiver of informed consent given the retrospective nature of the review. All patient records were confirmed for eligibility for collection in accordance with the relevant guidelines and regulations.

## Consent for publication

The consent to publish images of the infection cases in an online, open-access journal was confirmed.

## Competing interests

The authors declare that they have no conflicts of interest.


## References

1. Boev C, Kiss E: **Hospital-Acquired Infections: Current Trends and Prevention**. Crit Care Nurs Clin North Am 2017, **29**(1):51–65.
2. Kevin W. Lobdell SS, Juan A. Sanchez: **Hospital-Acquired Infections**. Surgical Clinics of North America 2012, **92**(1):13.
3. Fukuda H, Lee J, Imanaka Y: **Variations in analytical methodology for estimating costs of hospital-acquired infections: a systematic review**. J Hosp Infect 2011, **77**(2):93–105.
4. Stone PW, Braccia D, Larson E: **Systematic review of economic analyses of health care-associated infections**. Am J Infect Control 2005, **33**(9):501–509.
5. Giraldi G, Montesano M, Sandorfi F, Iachini M, Orsi GB: **Excess length of hospital stay due to healthcare acquired infections: methodologies evaluation**. Ann Ig 2019, **31**(5):507–516.
6. Zingg W, Holmes A, Dettenkofer M, Goetting T, Secci F, Clack L, Allegranzi B, Magiorakos A-P, Pittet D: **Hospital organisation, management, and structure for prevention of health-care-associated infection: a systematic review and expert consensus**. The Lancet Infectious Diseases 2015, **15**(2):212–224.
7. Li Y, Gong Z, Lu Y, Hu G, Cai R, Chen Z: **Impact of nosocomial infections surveillance on nosocomial infection rates: A systematic review**. Int J Surg 2017, **42**:164–169.
8. Gastmeier P: **Nosocomial infection surveillance and control policies**. Curr Opin Infect Dis 2004, **17**(4):295–301.
9. Seifi A, Dehghan-Nayeri N, Rostamnia L, Varaei S, Akbari Sari A, Haghani H, Ghanbari V: **Health care-associated infection surveillance system in Iran: Reporting and accuracy**. Am J Infect Control 2019, **47**(8):951–955.
10. Estan-Capell J, Alarcon-Torres B, Bermudez JD, Martinez-Rodriguez L, Martinez-Costa C: **Effect of a surveillance system for decreasing neonatal nosocomial infections**. Early Hum Dev 2019, **131**:36–40.
11. Sips ME, Bonten MJM, van Mourik MSM: **Automated surveillance of healthcare-associated infections: state of the art**. Curr Opin Infect Dis 2017, **30**(4):425–431.
12. Mingmei Du YX, Jijiang Suo, Bowei Liu, Na Jia, Rui Huo, Chunping Chen, Yunxi Liu: **Real-time automatic hospital-wide surveillance of nosocomial infections and outbreaks in a large Chinese tertiary hospital**. BMC Med Inform Decis Mak 2014, **14**(9):8.
13. Tvardik N, Kergourlay I, Bittar A, Segond F, Darmoni S, Metzger MH: **Accuracy of using natural language processing methods for identifying healthcare-associated infections**. Int J Med Inform 2018, **117**:96–102.
14. Villamarin-Bello B, Uriel-Latorre B, Fdez-Riverola F, Sande-Meijide M, Glez-Pena D: **Gold Standard Evaluation of an Automatic HAIs Surveillance System**. Biomed Res Int 2019, **2019**:1049575.
15. Walter Koller AB, Claudia Bauer, Harald Mandl, Klaus-Peter Adlassnig: **Electronic surveillance of healthcare-associated infections with MONHCU—a clinical breakthrough compared to conventional surveillance systems**. Stud Health Technol Inform 2010, **160**(1):5.
16. Makoto Jones SLD, Joshua Spuhl, Matthew H Samore, Christopher Nielson, Michael Rubin: **Identification of methicillin-resistant Staphylococcus aureus within the nation's Veterans Affairs medical centers using natural language processing**. BMC Med Inform Decis Mak 2012, **12**(34):8.
17. Schonfeld V, Diercke M, Gilsdorf A, Eckmanns T, Walter J: **Evaluation of the statutory surveillance system for invasive MRSA infections in Germany, 2016–2017**. BMC Public Health 2018, **18**(1):1063.




18. Lin J, Gao X, Cui Y, Sun W, Shen Y, Shi Q, Chen X, Hu B: **Increased multidrug resistant isolates: new clinical burdens for 66 hospitals in Shanghai, 2015 to 2017.** *Ann Transl Med* 2020, **8**(4):112.
19. Blanco N, O'Hara LM, Harris AD: **Transmission pathways of multidrug-resistant organisms in the hospital setting: a scoping review.** *Infect Control Hosp Epidemiol* 2019, **40**(4):447–456.
20. Qureshi S, Maria N, Zeeshan M, Irfan S, Qamar FN: **Prevalence and risk factors associated with multi-drug resistant organisms (MDRO) carriage among pediatric patients at the time of admission in a tertiary care hospital of a developing country. A cross-sectional study.** *BMC Infect Dis* 2021, **21**(1):547.
21. Zhao X, Wang L, Wei N, Zhang J, Ma W, Zhao H, Han X: **Epidemiological and clinical characteristics of healthcare-associated infection in elderly patients in a large Chinese tertiary hospital: a 3-year surveillance study.** *BMC Infect Dis* 2020, **20**(1):121.

## Figures




**Figure 1**

Study design. From January 1, 2017 to December 2017 (period 1), HAIs and MDRO cases were collected by traditional surveillance. The adoption of a RT-NISS to monitor HAIs and MDROs was implemented in period 2 (January 2018-December 2018). The adoption of a RT-NISS coupled with some infection-control interventions was implemented in period 3 (January 2019-December 2019).



**Figure 2**

The infection surveillance operational process of the RT-NISS.



**Figure 3**

Monthly utilization rate of antibiotics and rate of pathogen culture in HZFH from January 2018 to December 2019. A, intervention implemented: encouraged and trained clinicians to appropriately use antibiotics and timely pathogen culture; B, intervention implemented: punishment for doctors who did not implement prevention and control measures for MDROs or inappropriate use of antibiotics or under standard rate of pathogen culture.