

In Vitro Effects of Annona Senegalensis Root Bark, Musa Sapientum L and Malus Pumila Peel Extracts on Xanthine Oxidase

Madalitso Mlozen (✉ mlozenim@mchs.adventist.org)

Malawi Adventist University

Elias Bonya

Malawi Adventist University

Adam M Nyanda

Malawi Adventist University

Charity Mkwanda

Malawi Liverpool Wellcome Trust

Alinafe Kululanga

Malawi Adventist University

Jonathan Majamanda

Malawi Adventist University

Wilfred Taika

Malawi Adventist University

Linly Linje

Malawi Adventist University

Martin Kalumbi

Malawi Adventist University

Patrick Chagwa

Malawi Adventist University

Robert Chinyama

Malawi Adventist University

Zefaniah Katuah

Malawi Adventist University

Chikondi Kamwendo

Malawi Adventist University

Blessings Katiniche

Malawi Adventist University

Exton Siyano

Malawi Adventist University

Research Article

Keywords: Flavonoids, Inhibition, Gout, Phytochemicals, Extraction, Uric acid

Posted Date: April 21st, 2022

DOI: <https://doi.org/10.21203/rs.3.rs-1550422/v1>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background

Xanthine Oxidase activity may increase plasma urates, superoxide radicals and hydrogen peroxide leading to gout, arthritis and cancer. Allopurinol, a known Xanthine Oxidase inhibitor, is noted to have various adverse effects. Many laboratories are in research projects to find alternative inhibitors of XO including plant sources. Plants are known to contain therapeutically effective agents. *A. senegalensis* and *M. sapientum L* are reported to contain phytochemicals with antioxidant, anti-inflammatory and enzyme inhibitory activities.

Methods

Aqueous extracts of Root bark of *A. senegalensis*, peels of *M. sapientum L* and *M. pumila* were assayed for their inhibitory effects on Xanthine oxidase in vitro

Results

All aqueous extracts exhibited the presence of flavonoids. *A. senegalensis* root bark and *M sapientum L* and *M pumila* peels were investigated for their effects on Xanthine Oxidase activity. *A. senegalensis* root bark, *M. sapientum L* and *M. Pumila* peel extracts inhibited Xanthine Oxidase activity by 83%, 90% and 61% respectively as which are significantly different ($p < 0.05$) from that of the positive control, allopurinol (65%)

Conclusions

The results obtained in this study suggest that the flavonoids found in *A. senegalensis* root bark and *M. sapientum L* and *M. pumila* peel extracts could be potential Xanthine Oxidase activity inhibitors.

1.0 Introduction

Plants have been known to be of medicinal use in many societies and cultures around the globe. They have served and still serve as alternatives for conventional medicine in homes as natural remedies for infections, inflammations and noncommunicable diseases such as diabetes mellitus, gout, and hypertension. In other circumstances induction of labour has been achieved by plants. Elsewhere, *Marantodes pumilum (Blume) Kuntz* is commonly used to treat parturition, flatulence, dysentery, dysmenorrhoea, gonorrhoea, and bone diseases (7).

Recently, there has been an increased interest in use of plant-based remedies either to find new drugs, employ cheaper sources of medicine, or even to take advantage of the claimed safety in plants (4, 14).

The use of plants as medicine has been done either through food, or special preparations such as infusions, smoothies, decoctions, or poultices. Therefore, many edible plants are part of the search for alternative medicines. However, there are still many plants whose mechanism of action is known (30).

Plants are also used as raw materials for pharmaceutical products. A major interest has been in the plant phytochemistry and their natural oils. Xanthine oxidase (XO) is a key enzyme in formation of uric acid from degradation of purine nucleotides in the last step of in humans. XO is associated with inflammation through production of free radicals. During re-oxidation of XO, molecular oxygen acts as an electron acceptor, producing superoxide radical (O_2^-) and hydrogen peroxide (H_2O_2). XO is part of an important biological source of superoxide radicals (33). Under favourable conditions especially when XO is overproduced, uric acid can crystallise in arthrosis (joints) and kidneys and cause inflammation known as arthritis or gout and renal calculi respectively. Uric acid is a marker for gout and several haemodynamic abnormalities (7).

Xanthine oxidase is a therapeutic target for Allopurinol and Febuxostat, the commonly available xanthine oxidase inhibitors (XOI). Xanthine oxidase inhibitors are associated with side effects including Steven Johnson Syndrome, fever, skin rash, eosinophilia, hepatitis, and renal toxicity (19). Both of these drugs are expensive, inaccessible to some developing countries. Such unmet medical needs and health hazards posed by these drugs require more effort in finding novel Xanthine oxidase inhibitors that are much effective and have a good safety profile. These findings indicate the necessity for the development and discovery of more precise Xanthine oxidase inhibitors aimed at improving the treatment of gout and a reduction of complications that arise due to hyperuricemia while realising fewer adverse effects profile (19). The use of plant-based products may be very efficient as they are easily available and generally safe for biological systems (30). *Musa sapientum L* is one of the species in the banana family, and is one of the common fruits in the world. Nearly all parts of a banana tree are commonly used as traditional medicine for treating diarrhoea, menorrhagia, diabetes, dysentery, and antiulcerogenic, hypoglycaemic, antilithic, hypolipidemic conditions, plus antioxidant actions, inflammation, pains and even snakebites (1).

Malus pumila is largely cultivated around the world in temperate regions. It is usually eaten as a fruit and flowers can be used as tea. Studies have demonstrated that the plant contains some medicinal properties which can be targeted against ageing, oxidative stress, cancers, and bacterial infections. The chemical constituents of *M. pumila* include flavonoids, terpenoids and organic acids. Its main chemical components are dihydrochalcone such as phlorizin, phloretin, and other flavonoids such as quercetin, kaempferol and rutin (17).

Annona senegalensis is commonly called wild custard apple, used as food or a food additive as all parts of the plant contain varying amounts of essential oils. According to some study, it contains major bioactive constituents including tannins, flavonoid, saponins, alkaloids, glycosides, steroids, volatile acids and anthocyanin (20). It has also been reported in literature that the plant contains various minerals such as calcium, potassium, magnesium, zinc, copper, manganese as well as ascorbic acid and amino acids

which makes it an important source of nutrients. The roots, root bark and leaves have been reported to have been used to treat malaria, tuberculosis (6, 20).

Annona senegalensis is used for both food and medicinal purposes. It has also been reported in literature that different parts of the plant are employed in traditional medicine and home remedies to cure some diseases such as tuberculosis, hernia, diabetes, gastritis, male sexual impotence, difficulty in swallowing, and snake bites (18). *Annona senegalensis* has also been reported to have anti-cancer properties (5). Again, some researchers reported on the potential of *A. senegalensis* in the treatment of a minimum of three COVID 19 symptoms such as cough, fever, myalgia, and the treatment of liver, breast, and colon cancers (18). In this study, aqueous extracts of *Musa sapientum* and *Malus pumila* peels, and *A. senegalensis* root bark were investigated for the inhibitory potential on xanthine oxidase activity.

2.0. Materials And Methods

2.1. Specimen collection, Identification, Authentication and preparation

The samples of *A. senegalensis* were collected from areas around Malamulo, Thyolo while *M. sapientum* L and *M. pumila* samples were bought at a local market in Makwasa, Thyolo and Limbe respectively. Samples were taken to Mulanje Mountain Conservation Trust for identification and National Herbarium and Botanical Gardens of Malawi where they were authenticated. *Annona senegalensis* roots were washed with clean tap water and shade dried for 2 weeks (16). *Musa sapientum* L and *M. pumila* samples were washed under running tap water, the surfaces were sterilised with 70% ethanol, rinsed with distilled water, *A. Senegalensis* samples were refrigerated until needed. Peels of *M. pumila* were removed and shade dried for 2 weeks.

2.2 Plant material extractions

2.2.1 *A. senegalensis* root bark extractions

After drying, *A. senegalensis* root barks were pounded to a fine powder using a mortar and pestle, active ingredients were obtained by using the extraction method as described in literature with slight modifications (21). Where 40 g of the pounded sample was soaked in 350 ml of distilled water in a sterile conical flask and left to stand for 24 hours with periodic mixing and then it was filtered with a filter paper (Whatman No.1) after which the filtrate was stored in a refrigerator for further investigations.

M sapientum L peels were taken and added to distilled water after it had just boiled, left to cool. After sometime the contents were mixed and then filtered to remove large, non-homogenised particles in order to get clear aqueous extract. The extract was then kept at 4 °C until the time it was ready for use (15). *M. pumila* peels extracts were obtained using the method as described elsewhere, where 40 g of the dried peels were soaked in distilled water for 24 hours at room temperature with periodic vortexing, after which

the mixture was filtered using a filter paper (Whatman, No. 1) the filtrate was stored in a refrigerator for further investigations (21).

2.3 Phytochemical screening

2.3.1 Test for Flavonoids

A. Senegalensis, *M. Sapiantum L* and *M. pumila* phytochemical analyses were done according to literature with slight modifications (21). Extracts (1 ml) was added into 2 ml of sodium hydroxide (NaOH) solution. The resulting appearance of a yellow solution disappeared upon adding hydrochloric acid, which indicated the presence of Flavonoids.

2.4 Xanthine Oxidase activity assay

XO activity determination was performed according to the method described in literature, where the substrate and the enzyme solutions were prepared immediately before use (11). The reaction mixture contained sodium phosphate buffer (50mM pH 7.5, 300 μ l), XO (100 μ l, 0.1U/l), the reaction mixture was pre- incubated at 37 $^{\circ}$ C for 15 minutes. Then 100 μ l of substrate solution (0.15mM of xanthine) was added into the mixture and incubated at 37 $^{\circ}$ C for 30 minutes. The reaction was stopped by adding HCl (0.5M, 20 μ l).

The absorption was read at 295 nm against an assay blank, checking for uric acid formation at 37 $^{\circ}$ C using a UV spectrophotometer. Enzyme activity was determined using the formulae;

$$\text{Enzyme activity} = (\Delta\text{abs} \cdot V_t) / (\epsilon \cdot t \cdot V_e) \quad (1)$$

Where Δabs is the change in absorbance; V_t is the total reaction volume (800 μ l); ϵ = the extinction coefficient of uric acid (12.56); t is the time in minutes; V_e is the volume of the extract which was added in the reaction mixture (100 μ l). The calculated results were expressed in $\text{U} \cdot \text{L}^{-1}$. One unit of enzyme activity was defined as the amount of enzyme that converts 1 μ mol of xanthine to uric acid per min under defined conditions (13).

2.5 Xanthine Oxidase Inhibitory assay

The inhibitory effects of the extracts on XO activity was measured spectrophotometrically at 295 nm using a UV spectrophotometer, measuring the uric acid formation under aerobic conditions, with some modifications according to the method described elsewhere (11). Prior to the assay, the enzyme and *A. senegalensis*, *M. sapiantum L* and *M. pumila* extracts were mixed in a ratio of 1:1 $^{v/v}$ to obtain a final enzyme concentration of 0.1 U/L. The reaction mixture contained sodium phosphate buffer (50mM pH 7.5, 200 μ l) and 200 μ l of XO-extract pre-mixture, the reaction mixture was pre- incubated at 37 $^{\circ}$ C for 15 minutes. Then 100 μ l of substrate solution (0.15mM of xanthine) was added into the mixture and incubated at 37 $^{\circ}$ C for 30 minutes. The reaction was stopped by adding HCl (0.5M, 200 μ l). The UV spectrophotometer was blanked with an inhibition assay blank prepared in the same way but the enzyme

solution was replaced with a phosphate buffer. XO inhibitory activity was calculated and expressed as a percentage inhibition of XO in the above assay.

$$\text{Inhibition \% (I\%)} = 100 \times \left(\frac{\text{ABS}_{\text{control}} - \text{ABS}_{\text{test}}}{\text{ABS}_{\text{control}}} \right) \quad (2)$$

2.6 Quality control

All assays were carried out in triplicates, an average absorbance was calculated and used for all enzyme activities and inhibition studies. Control assays were included, an assay blank and inhibition assay blank were used. A well-known XO inhibitor (100 ug/ml) was used as a standard for the XO inhibitory studies. Negative control (blank: 0% XO activity) was prepared containing only the assay mixture without extract.

3.0. Results

3.1 Plant extractions and phytochemical screening

Flavonoids were identified in all aqueous extracts as summarised in Table 1.

Table 1
Phytochemical screening

Plant name	Flavonoid test results
A. senegalensis	++
M. sapientum L	++
M. pumila	+
Key: (+) = low in abundance (++) = moderate in abundance	

3.2 XO Inhibition assay

The results of XO activity determination and XO inhibition studies for *A. senegalensis* and *M. sapientum L* are summarised in Table 2. XO had an activity of 20.9 U/L. The experimental data indicate that the extracts under study showed good to outstanding inhibitory effects towards XO. *A. senegalensis* reduced XO activity from 20.9 to 3.50U/L representing 83% activity inhibition. *M. sapientum L* exhibited a 91% inhibition by reducing XO activity to 3.50 U/L and *M. pumila* reduced XO activity to 5.8U/L representing 80% inhibition. Allopurinol, the positive control, reduced XO activity from 20.9U/L to 7.26U/L, representing 65% inhibitory effects, a summary is presented in Table 2 with graphical representation in Figs. 1 and Fig. 2 respectively. Statistical analysis is as summarized in Table 3.

Table 2
A summary of XO enzyme activity and in vitro inhibitory studies

Plant name	Avg Abs	Activity (U/L)	% inhibition
XO	0.324	0.206	0
<i>A. senegalensis</i>	0.055	0.035	83.0
<i>M. sapentium</i>	0.031	0.019	90.6
<i>M. Pumila</i>	0.092	0.058	84
Allopurinol	0.114	0.072	65.23

Table 3

The differences in mean absorbance between the positive control and the test sample; enzyme activity between the positive control and the test sample; and the inhibitory activity between the positive control and the test samples and their t-values and p-values at 95% confidence interval

Test Sample	In relation to mean ABS		In relation to mean enzyme activity		In relation to mean I%		Mean Inhibition difference (%)
	t-value	p-value	t-value	p-value	t-value	p-value	
<i>A. senegalensis</i>	29.4675	0.0000	28.5349	0.0000	-7.2259	0.0010	-18.229 (-17.77)
<i>M. sapentium</i>	26.2473	0.0000	25.2918	0.0000	-7.4072	0.0009	-25.569 (-25.37)
<i>M. pumila</i>	25.7729	0.0000	25.1366	0.0000	26.7655	0.0000	64.70 (-18.77)

4.0. Discussion

In the quest to search for alternative drugs for the cure of disease, and as a step towards identifying a novel medicinal agent, this study assessed three plants for their effect against the activity of XO. This study found slightly lower concentrations of flavonoids, which may be attributed to the type of extraction medium employed. Some literature reported that there are observed variations of phytochemical presence in medicinal plants owing to solvents used for extraction and extraction procedure (8). Water as a solvent for extraction is advantageous as it effectively extracts most polar compounds, cheap, nontoxic and nonflammable (2).

However it may affect the extraction efficiency and content and hydrolysis of compounds due to high heat requirements to concentrate extracts (31, 32). According to literature, low to no evidence of alkaloids was reported upon using water as a solvent (8).

Therefore the use of aqueous solvents might have contributed to the observed flavonoids test results in the current study.

Flavonoids, a member of a group of naturally occurring active compounds in plants, have been reported to possess tremendous health benefits (26). Medically important flavonoids are reported to be very potent antioxidants and thus have attracted a significant amount of interest among researchers as possible potent therapeutic agents for illnesses whose aetiologies and pathogenesis are associated with free radicals (26). Free radicals including hydroxyl radicals, superoxide anions, hydrogen peroxide, oxygen singlets, hypochlorite and nitric oxide are reported to play a key role in various inflammatory diseases; viz rheumatoid arthritis and gout (24, 27). XO catalyzes the formation of uric acid and hydrogen peroxide from purine degradation which are responsible for oxidative damage that causes gout, hyperuricemia, arthritis, vascular endothelium damage and ageing (10, 28).

Various parts of *M. sapientium*, *A. senegalensis* and *M. pumila* have been reported to contain active secondary metabolites active on various enzymes that effectively inhibit various enzymes including Glutathione-s-transferase, Acetylcholinesterase, Carboxylesterase and Xanthine oxidase (XO) α -glucosidase and α -amylase, angiotensin 1 converting enzyme (ACE) (3, 12, 29). The flavonoids observed XO inhibition as also reported by elsewhere, might be helpful in the prevention of slowing down the pathogenesis of gout (25).

Interestingly results obtained in the current research indicate that aqueous extracts of *M. Pumila* peels exhibited higher inhibitory effects as compared to those observed by some research fellows, whereby they reported that aqueous extracts of *M. pumila* exhibited no inhibition and methanolic extracts inhibited XO activity by 28% (23).

Annona senegalensis crude extracts are reported to inhibit several enzyme activities including XO, lower than observed in this study (12). This study also found that *A. senegalensis* together with other species of *Annona* inhibited xanthine oxidase activity by 25% which is also lower than that obtained in this study. This variation was suggested to arise from some interaction of compounds between the species that led to retardation of the inhibition (12).

There is limited information pertaining to the interaction of *Musa sapientium L* and XO to support its inhibitory activity, however, some researchers found that other antioxidative *Musa* species decrease uric acid levels by inhibiting the xanthine oxidase enzyme (9, 22).

The antioxidative properties of *M. sapientium L* peel, *M. pumila* peel and *A. senegalensis* root bark extracts have potential to qualify that they are effective anti-gout agents due to their ability to inhibit XO enzyme activity.

5. 0 Conclusion And Recommendations

The results of this study indicate that *A. senegalensis*, *M. sapientium L* peel and *M. pumila* aqueous extract possess significant inhibitory effects on xanthine oxidase activity. Further *in vitro* studies may be conducted on the effects of *A. senegalensis*, *M. sapientium* and *M. pumila* extracts obtained using various extraction solvents and methods. Further, purifications and identification of purified extract are

considered to identify exact active phytochemical(s) that exhibit the inhibitory effects observed in the current study.

List Of Abbreviations

XO	Xanthine oxidase
H ₂ O ₂	Hydrogen peroxide
O ₂ ⁻	Superoxide radical
XOI	Xanthine oxidase inhibitors
NaOH	Sodium Hydroxide
HCL	Hydrochloric acid
Δabs	Change in absorbance
V _t	Total reaction volume
Ve	Extract volume
U.L ⁻¹	Enzyme activity unit
UV	Ultraviolet
AS	<i>Anonna senegalensis</i>
MP	<i>Malus pumila</i>
MS	<i>Mussa sapientum</i>
AL	Allopurinol
ACE	Angiotensin converting enzyme

Declarations

Ethical approval

This research was approved by the National Health Sciences Research Committee (NHSRC) and Malawi Adventist University Research Committee. *A. senegalensis*, *M. pumila* and *M. sapientum* L were identified and authenticated by a Botanist at the National Herbarium and Botanical Gardens of Malawi, under authentication deposition numbers of 15053 and 1729 respectively. All methods were carried out in

relevant guidelines and regulations. National Health Sciences Research Committee (NHSRC) and Malawi Adventist University Research Committee gave permission to collect samples of *A. senegalensis*

Consent to participate

Not applicable

Consent for publication

Not applicable

Availability of data

The datasets used and/or analysed during the current study are available from the Corresponding author on reasonable request.

Competing interests

The authors declare that they have no competing interests

Funding

The authors received no funding from any other body, instead the project was self-funded.

Authors' contributions

MM, EB, ES, AMN, CM and AK: Data analysis and write up

MM, EB, JM, WT and MK: Literature review and write up

EB, MM, AMN, LL, PC, RC, ZK, CK and BK: Proof reading and discussion of results

MM, AMN, EB and ES: Data curation and editing.

All authors reviewed the manuscript.

Acknowledgements

The authors would like to acknowledge the Mulanje Mountain Conservation Trust and The National Herbarium and Botanical gardens of Malawi for the identification and authentication of the plants.

References

1. Abu Zarin, M., Tan, J.S., Murugan, P. and Ahmad, R. (2020), "Investigation of potential anti-urolithiatic activity from different types of Musa pseudo-stem extracts in inhibition of calcium oxalate

- crystallization”, BMC Complementary Medicine and Therapies, BioMed Central Ltd. 2020; 20(1):1–12.
2. Abubakar A.R and Haque M., 2020-JPharmBioallSci1211-1574732_042227.pdf.
 3. Adamson SS, Ganiyu O. Aqueous Extracts from Unripe Plantain (*Musa paradisiaca*) Products Inhibit Key Enzymes Linked with Type 2 Diabetes and Hypertension in vitro. *Jordan J Biol Sci.* 2012;5(4):239–46.
 4. Adjakpa, J. B., Ahoton, L. E., Obossou, F. K. & Ogoube, C., 2016. Ethnobotanical study of Senegal custard apple (*Annona senegalensis* Pers.) in Dassa-Zoumétownship, Republic of Benin. *International Journal of Biological and Chemical Sciences*, 10(5):2123–2137.
 5. Africanus Beauv Biseko, A.P. and Zacharia, E. Evaluation of anti-cancer potential of crude extracts of *Annona senegalensis* Pers. and *Allophylus africanus* P Beauv.”, NM-AIST, 2019. Available at: <https://dspace.nm-aist.ac.tz/handle/20.500.12479/1037> (Accessed on 30 March 2022).
 6. Ajaiyeoba, E. et al. *invivo* antimalarial and cytotoxic properties of *annona senegalensis* extract. *african journal traditional complementary and alternative medicine.* 2006;3(1):131–171.
 7. Aladdin, N.A., Husain, K., Jalil, J., Sabandar, C.W. and Jamal, J.A. Xanthine oxidase inhibitory activity of a new isocoumarin obtained from *Marantodes pumilum* var. *pumila* leaves: BMC Complementary Medicine and Therapies, BioMed Central Ltd, 2020;20(1):1–12.
 8. Assanga SBI, Luján LML, Espinoza CLL, Salido AAG, Angulo DF, Pino JLR, et al. Solvent effects on phytochemical constituent profiles and antioxidant activities, using four different extraction formulations for analysis of *Bucida buceras* L. and *Phoradendron californicum*. *BMC Res Notes.* 2015;1–14.
 9. Ayoola, I.O., Abberton, B.; Gueye, M.A., Sonibare, M.T. Antioxidant activity and acetylcholinesterase inhibition of field and in vitro grown *Musa* L. species: *Journal of Food Measurement and Characterization.* 2016.
 10. Aziz, N., & Jamil, R. T. *Biochemistry, Xanthine Oxidase.* StatPearls. 2021 <http://www.ncbi.nlm.nih.gov/pubmed/31424829>
 11. Azmi, S. M., Jamal, P. & Amid, A. Xanthine oxidase inhibitory activity from potential Malaysian medicinal plant as remedies for gout. *International food research journal.* 2012;19(1):151–165.
 12. Bangou MJ, et. al. Evaluation of Enzymes Inhibition Activities of Medicinal Plant Plant from Burkina Faso. *Pakistan Journal of Biological Sciences.* 2011;14(2):99–105
 13. Beyaztas, S. & Arsian, O. Purification of Xanthine Oxidase from Bovine Milk by Affinity Chromatography with a Novel Gel. *Journal of Enzyme Inhibition and Medicinal Chemistry.* 2014;30(3):442–447.
 14. Bhutkar MA, Bhingde SD, Randive DS, Wadkar GH. Hypoglycemic effects of *Berberis aristata* and *Tamarindus indica* extracts in vitro. *Bull Fac Pharm Cairo Univ.* 2017 Jun;55(1):91–94.
 15. Chabuck, Z. A. G., Al-Charrakh, A. H., Hindi, N. K. K. & Hindi, S. K. K. Antimicrobial Effect of Aqueous Banana Peel Extract, Iraq. *Research Gate: Pharmaceutical sciences.* 2013;1:73–75.

16. Chinyere, N. H., Milala, M. A. & Zannah, H. Effects of aqueous root extract of *annona senegalensis* on bitisarietans venom, protease and phospholipase A2 activities. *J pharma Biomed Sci.* 2016;06(08):469–473.
17. Cui, L., Hou, X., Li, W., Leng, Y., Zhang, Y., Li, X., Hou, Y., et al. Dynamic changes of secondary metabolites and tyrosinase activity of *Malus pumila* flowers: *BMC Chemistry*, BioMed Central Ltd. 2019;13(3):1–8.
18. Donhouédé, J.C.F., Salako, K.V., Gandji, K., Idohou, R., Tohoun, R., Hounkpèvi, A., Ribeiro, N., et al. Food and medicinal uses of *Annona senegalensis* Pers.: a country-wide assessment of traditional theoretical knowledge and actual uses in Benin, West Africa: *Journal of Ethnobiology and Ethnomedicine*. BioMed Central. 2022;18(1):1–15.
19. Duong, N. T. et al. Xanthine oxidase inhibitors from *Archidendron clypearia* (Jack). I.C. Nielsen: Results from systematic screening of Vietnamese medicinal plants. *Asian Pacific Journal of tropical Medicine*. 2017;10(6):1–9.
20. Edipofon, A. et al. *Annona Senegalensis* persoon: a review of its ethnomedicinal uses, biological activities and phytochemical compounds. *Journal of pharmacognosy and phytochemistry*. 2016;5(2):211–219.
21. Idayant, S. A., Samira, A. & Musliu, A. Extraction and Phytochemical Screening of root and leaves of *Annona Senegalesis*(Wild Custard Apple). *Academic Journal of Interdisciplinary Studies*. 2014;3(7).
22. Irawan, C. et al., 2021. Potential of Ethanolic Extract from Ripe *Musa balbisiana* Colla Fruit Using Ultrasound-Assisted Extraction as An Antioxidant and Anti-Gout. *Pharmacognosy Journal*, 13(6):1332–1340.
23. Lee E-H, Kim Y-J, Kwon S-I, Kim J-H, Kang I-K, Jung H-Y, et al. Anti-Oxidative, Health Functional, and Beauty Food Activities of Extract from Newly Bred Ruby S Apple (*Malus pumila* Mill.) Peel. *J Korean Soc Food Sci Nutr*. 2018 Nov 30;47(11):1093–102.
24. Lobo, V., Patil, A., Phatak, A., & Chandra, N. Free radicals, antioxidants and functional foods: Impact on human health. *Pharmacognosy Reviews*. 2010;4(8):118–126. doi.org/10.4103/0973-7847.70902
25. Ngbolua K-N, Mudogo V, Mpiana PT, Tshibangu DST, Tshilanda DD, Ashande MC. In vitro and in vivo anti-malarial and cytotoxic activities of ethanolic extracts of *Annona senegalensis* Pers (Annonaceae) from Democratic Republic of the Congo. 2(2):6.
26. Panche, A. N., Diwan, A. D., & Chandra, S. R. Flavonoids: An overview. In *Journal of Nutritional Science*. 2016;5 doi.org/10.1017/jns.2016.41
27. Pham-Huy, L. A., He, H., & Pham-Huy, C. (2008). Free Radicals, Antioxidants in Disease and Health. *International Journal of Biomedical Science*. 2008;4(2):89. /pmc/articles/PMC3614697/
28. Puddu, P., Puddu, G. M., Cravero, E., Vizioli, L., & Muscari, A. The relationships among hyperuricemia, endothelial dysfunction, and cardiovascular diseases: Molecular mechanisms and clinical implications. *Journal of Cardiology*. 2012;59(3):235–242. https://doi.org/10.1016/j.jjcc.2012.01.013
29. Ramu, R. et. al. Ramu R, Shirahatti PS, Zameer F, Ranganatha LV, Nagendra Prasad MN. Inhibitory effect of banana (*Musa* sp. var. Nanjangud rasa bale) flower extract and its constituents

Umbelliferone and Lupeol on α -glucosidase, aldose reductase and glycation at multiple stages. South Afr J Bot. 2014;95:54–63.

30. Shukor, N. A. A., Ablat, A., Muhamad, N. A. & Mohamad, J. In vitro antioxidant and in vivo xanthine oxidase inhibitory activities of *Pandanus amaryllifolius* in potassium oxonate induced hyperuricemic rats. International Journal of Food Science and Technology. 2018;53(1):1476–1485
31. Truong, D.H. et al. Evaluation of the use of different solvents for phytochemical constituents, antioxidants, and in vitro anti-inflammatory activities of *Severinia buxifolia*. Journal of Food Quality, 2019. doi:10.1155/2019/8178294.
32. Złotek, U, S. Mikulska, M. Nagajek, and M. Świeca. Effect of different solvents and number of extraction steps on the polyphenol content and antioxidant capacity of basil leaves (*Ocimum basilicum* L.) extracts,” Saudi Journal of Biological Sciences. 2016;23(5):628–633,
33. Zorov, D.B., Juhaszova, M. and Sollott, S.J. Mitochondrial Reactive Oxygen Species (ROS) and ROS-Induced ROS Release: Physiological Reviews, American Physiological Society. 2014;94(3):909.

Figures




Figure 1

A graph of mean absorbance and enzyme activity against different extracts (XO = Xanthine oxidase, AS = *A. senegalensis*, MP = *M. pumila*, MS = *M. sapientum*, AL = allopurinol)




Figure 2

A graph of inhibitory activity against different extracts (XO = Xanthine oxidase, AS = *A. senegalensis*, MP = *M. pumila*, MS = *M. sapientum*, AL = allopurinol)