Postnatal Mental Health, Breastfeeding Beliefs, and Breastfeeding Practices in Rural China

Qi Jiang

Stanford University Freeman Spogli Institute for International Studies

Evelyn Zhang

Stanford University Freeman Spogli Institute for International Studies

Nourya Cohen

Stanford University Freeman Spogli Institute for International Studies

Mika Ohtori

FSI: Stanford University Freeman Spogli Institute for International Studies

Sabrina Zhu

FSI: Stanford University Freeman Spogli Institute for International Studies

Yian Guo

Stanford University Freeman Spogli Institute for International Studies

Hannah Faith Johnstone

Stanford University Freeman Spogli Institute for International Studies

Sarah-Eve Dill (✉ sedill@stanford.edu)

Freeman Spogli Institute for International Studies, Stanford University <https://orcid.org/0000-0002-9321-6606>

Huan Zhou

West China School of Medicine: Sichuan University West China Hospital

Scott D Rozelle

Stanford University Freeman Spogli Institute for International Studies

Research article

Keywords: breastfeeding, breastfeeding self-efficacy, breastfeeding attitude, mental health, depression, anxiety, stress, rural China

Posted Date: January 28th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-155309/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Postnatal Mental Health, Breastfeeding Beliefs, and Breastfeeding Practices in Rural China

Qi Jiang¹; Evelyn Zhang¹; Nourya Cohen¹; Mika Ohtori¹; Sabrina Zhu¹; Yian Guo¹; Hannah Faith Johnstone¹; Sarah-Eve Dill^{1*}; Huan Zhou²; Scott D Rozelle¹

¹ Rural Education Action, Program (REAP), Freeman Spogli Institute for International Studies, Stanford University, Stanford, California, USA

² West China School of Public Health and University, Chengdu, China

* Correspondence:

Sarah-Eve Dill

Freeman Spogli Institute for International Studies, Stanford University

sedill@stanford.edu

Abstract

Background: The importance of breastfeeding in low- and middle- income countries is well recognized, yet the importance of postnatal mental health on breastfeeding practices and beliefs in these settings has been understudied. This study investigates the associations between maternal mental health problems and breastfeeding beliefs as well as practices in rural China.

Methods: Cross-sectional data were collected from 742 mothers of infants under six months old in rural Sichuan Province, China. Surveys collected data on maternal mental health problems (depression, anxiety, and stress symptoms), breastfeeding beliefs (attitudes and self-efficacy), and breastfeeding practices. Ordinary least squares regression, multiple logistic regression and heterogeneous effect analyses were used to determine the associations between maternal mental health and breastfeeding outcomes.

Results: Among all respondents, 13% showed symptoms of depression, 16% anxiety, and 9% stress. The prevalence of full breastfeeding was 59.3%. Breastfeeding attitude was significantly associated with symptoms of depression ($p=0.023$) and breastfeeding self-efficacy with symptoms of depression ($p=0.001$) and symptoms of stress ($p=0.020$). However, there were no significant associations between symptoms of mental health problems and full breastfeeding. The heterogeneous effects analyses revealed that full breastfeeding was negatively associated with stress symptoms when the infant was from a high-income family ($p=0.011$). In addition, full breastfeeding was negatively associated with the father having a higher education level ($p=0.026$, $p=0.048$, and $p=0.020$) and the infant being older than 2 months old ($p=0.000$, $p=0.000$, $p=0.00$), regardless of maternal mental health problem symptoms.

Conclusion: Symptoms of maternal mental health problems are significantly associated with breastfeeding attitude and self-efficacy, yet has less of an association with breastfeeding practices. To improve breastfeeding practices, interventions need a multi-dimensional approach that should not only focus on improving maternal mental well-being but also consider demographic background characteristics.

Keywords: breastfeeding, breastfeeding self-efficacy, breastfeeding attitude, mental health, depression, anxiety, stress, rural China

Introduction

It is well-established in the international literature that breastfeeding is crucial for the health and development of infants. Extensive research has shown that breastfeeding is associated with a number of positive infant health outcomes such as lower rates of respiratory infections and diarrhea (1–5), reduced risk of sudden infant death syndrome (6–8), and better immune system functioning (9). Breastfed infants have also been shown to have better health outcomes later in life, including improved cognitive performance and lower rates of chronic diseases such as diabetes and cardiovascular disease (8–10). Based on this evidence, the World Health Organization (WHO) recommends exclusive breastfeeding for the first 6 months and continued breastfeeding until the child is 2 years old (11).

Unfortunately, international literature has also shown that postnatal mental health problems among mothers can present a barrier to breastfeeding. Breastfeeding is a practice that involves heavy involvement from the mother, requiring her to be engaged in and willing to participate in the feeding process. For mothers facing mental health problems, these demands can feel burdensome and ultimately discourage her from breastfeeding. Studies have found that depressed mothers are less likely to exclusively breastfeed and more likely to terminate breastfeeding earlier (12–17). Similarly, studies have found that postpartum maternal anxiety is negatively associated with breastfeeding initiation, duration (18), and exclusivity (18,19), although these studies have only been conducted in high-income countries.

Postnatal mental health problems can also worsen breastfeeding beliefs, which have been found to be critical for breastfeeding practices. Existing global literature suggests that maternal depression affects breastfeeding beliefs, such as self-efficacy, which is defined as a mother's confidence in her ability to breastfeed her newborn, and attitude, which is defined as a mother's

viewpoint and stance on breastfeeding. Maternal depression and anxiety have been found to be associated with lower breastfeeding self-efficacy (20–22). Maternal depression and stress have also been shown to be associated with less favorable attitudes towards breastfeeding (23,24), such as being more skeptical or unsupportive of the practice (25). Having a positive attitude about breastfeeding is important since breastfeeding beliefs have been shown to not only be associated with higher rates of exclusive breastfeeding (25–28), but also other measures of breastfeeding outcomes, such as the duration the infant is breastfed (27–29), and the mother’s satisfaction with breastfeeding (30).

Increasing evidence shows that the prevalence of postnatal mental health problems in low- and middle-income countries (LMICs), including China, is particularly high (27). Given the association between mental health and breastfeeding, there is a concern that breastfeeding outcomes in these countries are being severely impacted. Like other LMICs, rural China has been shown to have a high prevalence of maternal mental health issues, as well as poor infant feeding practices. Postnatal depression rates have empirically been shown to range from 20% to 25% (28,29), which are high compared to the global rate of 13% (30). This high prevalence of maternal mental health is accompanied by low rates of exclusive breastfeeding, ranging from 4.2% to 28.7% (31–33). These rates are low compared to the global rate of exclusive breastfeeding among infants younger than 6 months, which is 37% (34). Yet, there have been no studies on the relationship between postnatal mental health and breastfeeding beliefs and practices in rural China. Considering the high prevalence of maternal mental health issues and poor infant feeding practices, the relationship between maternal mental health and breastfeeding outcomes in rural China needs to be better understood.

The goal of this study is to investigate the association between symptoms of postnatal mental health problems, breastfeeding beliefs (self-efficacy and attitudes), and breastfeeding practices in rural China. Specifically, we aim to pursue three objectives. First, we describe the prevalence of symptoms of mental health problems among new mothers in rural China and identify their breastfeeding practices and beliefs. Second, we examine the correlation between symptoms of postnatal mental health problems and breastfeeding beliefs. Finally, we determine the correlation between symptoms of postnatal mental health problems and breastfeeding practices.

Methods

A cross-sectional study was conducted in the rural areas of one prefecture in Sichuan Province. According to the National Bureau of Statistics of China, 48% of Sichuan's population are rural residents. The average per capita disposable income in the rural areas of Sichuan is 13,331 RMB (1906 US dollars), far lower than the national average of 28,228 RMB (4033 US dollars) (35). For these reasons, the study area can be considered relatively representative of rural southwestern China.

Sampling

The research team implemented a 3-step sampling protocol to select households for the study. First, from the nine counties within the sample prefecture, the four nationally-designated "poverty counties" (*pinkun xian*) were included, which are counties that have more than 2% of their population living under the national poverty line of 3,000 RMB (458 US dollars) (36). These counties were selected in order to sample the mental health and breastfeeding outcomes of low-income, rural mothers. Second, sample townships were chosen within each sample county.

The sampling frame included two exclusion criteria: non-rural townships and rural townships with populations less than 10,000. Of the remaining townships, 20 townships per county were randomly selected, resulting in a total of 80 rural townships.

The third step of the protocol selected mothers and households from the sample townships. After obtaining a list of all households with pregnant women beyond their second trimester or with infants under 6 months old from the local county-level Maternal and Child Hospital in each sample county, the research team randomly selected 25 families in each township. If there were not enough eligible households in a township, the team then included eligible households living in villages surrounding the township. In total, 1,296 households were identified as potential candidates for the study. For the purposes of this study, two groups of women were excluded: pregnant women and non-breastfeeding new mothers, since the focus of the study is specifically on the impact of mental health on breastfeeding mothers. This resulted in a final sample of 742 breastfeeding mothers.

Data collection

Data were collected from November to December 2019 by trained survey enumerators. During the training, enumerators were taught how to implement the survey instruments for each of the study's main components. Once the training was completed, enumerators conducted one-on-one survey interviews at each sample household. The survey data collected included: postnatal mental health (symptoms of depression, anxiety, and stress), breastfeeding beliefs (attitudes and self-efficacy), breastfeeding practices, and demographic characteristics.

Postnatal mental health

To measure the mental health of sample mothers, the research team used the Depression, Anxiety, and Stress Scale-21 (DASS-21). This is a 21-item short-form version of the DASS-42 that was originally created by Lovibond and Lovibond (37). The DASS-21 scale has been

validated in China (38,39). To complete the DASS-21 scale, enumerators asked mothers to rank each statement (7 for each subsection) from 0 to 3, depending on how much the statement applied to them in the past week. Each individual's DASS-21 score was then calculated by adding up the ranking for every sub-question in a specific section and multiplying the sum by 2. Hence, for each section, the total sum score could range from 0 to 42. Mothers who scored greater than or equal to 9 for depression, 7 for anxiety, and 14 for stress were considered to have symptoms of each respective mental health problem. It is important to note that the resulting score of the DASS-21 scale is not a clinical diagnosis but only a reasonable measurement of the severity of depression, anxiety, and stress symptoms.

Breastfeeding beliefs

Two measures for breastfeeding beliefs were collected: breastfeeding attitudes and breastfeeding self-efficacy. First, to measure maternal attitudes about breastfeeding, enumerators administered the Iowa Infant Feeding Attitude Scale (40), which has been used in Turkey (23), Ethiopia (41), the United Kingdom (42), Russia (42), and the United States (43). It has also been validated in China (44). Breastfeeding mothers were read 18 statements, such as “breastfeeding is more convenient than formula feeding” and asked to use a 5-point Likert scale to express if they agreed or disagreed with the statement (1 being strongly disagree and 5 being strongly agree). Total scores range from 18 to 90. Statements that were representative of worse attitudes were reverse coded so that a higher score correlates to a more positive attitude about breastfeeding.

Second, to determine the mother’s self-efficacy surrounding breastfeeding, enumerators administered the short form of the Breastfeeding Self-Efficacy Scale (45). It has previously been used in Kenya (46), Turkey (47), and Malaysia (48), and was validated in Chinese by Ip et al (49). Breastfeeding mothers were asked to rate how confident they were about 16 breastfeeding-

related items based on a 5-point Likert scale (1 being not at all confident to 5 being always confident). An example of an item is “I manage to keep up with my infant’s breastfeeding demands”. Total scores range from 16 to 80, where a higher score corresponds to higher levels of self-efficacy.

Breastfeeding practices

The mother’s breastfeeding practices were determined through a 24-hour dietary recall, in which enumerators asked mothers to list all the foods and liquids they fed their infant in the last 24 hours. Based on this data, mothers were then divided into two feeding categories: full breastfeeding and mixed breastfeeding. Full breastfeeding included both exclusive and predominant breastfeeding. As defined by the WHO, exclusive breastfeeding is breastfeeding while giving no other foods or liquids, while predominantly breastfeeding is feeding breastmilk along with other water or water-based liquids (broth or juice) (50). Mixed breastfeeding included infants who were fed breastmilk along with other milk substitutes (formula or animal milk), other liquids, or solids.

Demographic characteristics

Data on demographic characteristics of each sample family and infant were also collected by enumerators. Family characteristics included mother’s age, mother’s and father’s education levels, and family yearly income. Demographic characteristics of the infant were collected from the birth certificate, and included information such as the infant’s gender, age in months, whether the infant’s birth weight was low, and whether the infant was born prematurely.

Statistical Analysis

The statistical analysis for the study is composed of four parts. First, the research team calculated the overall prevalence of maternal depression, anxiety, and stress symptoms, as well as the overall prevalence of breastfeeding outcomes, which include breastfeeding practice and

beliefs. Second, an adjusted ordinary least squares (OLS) regression was used to identify the correlations between mental health and breastfeeding beliefs, while controlling for other potential confounders. The specification of the adjusted model is:

$$Belief_i = \beta_0 + \beta_1 Mental_health_i + Control_i + \epsilon_i \quad (1)$$

where $Belief_i$ refers to breastfeeding beliefs (breastfeeding attitude or breastfeeding self-efficacy). $Mental_health_i$ is the dummy variable for mental health (depression, anxiety, or stress symptoms), which takes the value of 1 when mother_i showed symptoms of mental health problems and takes the value of 0 when mother_i did not. $Control_i$ is a set of control variables, including infant gender and age, whether the infant was born prematurely, whether the infant has low birth weight, maternal age and education level, paternal education level, and family income.

Third, a multivariate logistic regression was run to examine the associations between mental health problems and breastfeeding practices. The specification of the regression model is:

$$Practice_i = \beta_0 + \beta_1 Mental_health_i + Control_i + \epsilon_i \quad (2)$$

where $Practice_i$ takes the value of 1 when mother_i was fully breastfeeding and takes the value of 0 when mother_i was mixed breastfeeding. In equation (2), the definitions of the variables, $Mental_health_i$ and $Control_i$ are the same as those in equation (1).

Finally, a heterogeneous analysis was conducted to measure the effects of symptoms of maternal mental health problems on breastfeeding practice by five demographic characteristics, including three socio-economic status (SES) characteristics (education levels of each parent and family income), maternal age, and infant age. The specification of the adjusted model is:

$$Practice_i = \beta_0 + \beta_1 Mp_low_i + \beta_2 Mp_high_i + \beta_3 Mh_low_i + Control_i + \epsilon_i \quad (3)$$

where Mp_low_i , Mp_high_i , and Mh_low_i are three binary variables. Mp_low_i takes the value of 1 when mother_i shows symptoms of mental health problems and is of low SES/low maternal

age/low infant age, and 0 otherwise. Mp_high_i takes the value of 1 when $mother_i$ shows symptoms of mental health problems and is of high SES/high maternal age/high infant age, and 0 otherwise. Mh_low_i takes the value of 1 when $mother_i$ does not have symptoms of mental health problems and is of low SES/low maternal age/low infant age, and 0 otherwise. $Practice_i$ and $Control_i$ are the same as those in equation (1).

Results

Table 1 shows the demographic characteristics of the study sample. Of the infants sampled, 416 (56.1%) were male and the average age was a little over two and a half months (2.7 months). Only 22 sample infants (3.0%) were born prematurely, and 22 (3.0%) had low birth weight. The average age of the 742 mothers sampled was approximately 27.9 years old. Additionally, mothers were slightly less educated than fathers, with 297 (40.0% of mothers) having graduated from high school compared to 341 (46.0% of the fathers) having graduated from high school.

Table 1. Descriptive statistics of demographic characteristics (n=742).

	N (%) or Mean (SD)
Infant gender (% male), N (%)	416 (56.1%)
Infant age (months), mean (SD)	2.7 (2.0)
Premature, N (%)	22 (3.0%)
Low birth weight, N (%)	22 (3.0%)
Maternal age (years), mean (SD)	27.9 (4.9)
Mother graduated high school, N (%)	297 (40.0%)
Father graduated high school, N (%)	341 (46.0%)
Family yearly income (ten thousand yuan), mean (SD)	7.4 (6.2)

The prevalence of the symptoms of mental health issues and descriptive data on breastfeeding practices and beliefs among sample mothers is shown in Table 2. According to the data, 96 mothers (13.0%) exhibited symptoms of depression, 119 (16.0%) exhibited symptoms of anxiety, and 67 (9.0%) exhibited symptoms of stress. Just over half of the sample mothers (440) fully breastfed their infants, which accounts for 59.3% of the sample. For breastfeeding attitude, mothers scored an average of 61.3, within a range of 18-90. For breastfeeding self-efficacy, mothers scored an average of 56.1, within a range of 16-80.

Table 2. Descriptive statistics of breastfeeding practices, knowledge, and beliefs (n=742).

	N (%) or Mean (SD)
<i>Panel A. Postnatal mental health</i>	
Symptoms of depression, N (%)	96 (13.0%)
Symptoms of anxiety, N (%)	119 (16.0%)
Symptoms of stress, N (%)	67 (9.0%)
<i>Panel B. Breastfeeding practice</i>	
Exclusive or predominantly breastfeeding, N (%)	440 (59.3%)
Any breastfeeding, N (%)	302 (40.7%)
<i>Panel C. Breastfeeding beliefs</i>	
Breastfeeding attitude score (range: 18-90), mean (SD)	61.3 (4.5)
Breastfeeding self-efficacy score (range: 16-80), mean (SD)	56.1 (8.8)

The OLS regression analysis for the correlations between mental health problems and breastfeeding beliefs are shown in Table 3. Mothers with symptoms of depression had significantly lower breastfeeding attitude scores ($p=0.023$). Additionally, mothers with symptoms of depression and symptoms of stress also had significantly lower breastfeeding self-efficacy scores ($p=0.001$ and 0.020 , respectively). Symptoms of anxiety were not significantly correlated with breastfeeding beliefs.

Table 3. Associations between postnatal mental health problems and breastfeeding beliefs (n=742).

	Breastfeeding attitude		Breastfeeding self-efficacy	
	β (SE)	p-value	β (SE)	p-value
Symptoms of depression	-1.11 (0.49)	0.023	-2.93 (0.89)	0.001
Symptoms of anxiety	-0.47 (0.46)	0.305	-1.58 (0.83)	0.057
Symptoms of stress	-0.71 (0.59)	0.228	-2.49 (1.07)	0.020

Notes: Controlled for demographic characteristics and breastfeeding practices (infant gender, infant age, premature birth status, whether the infant was born with a low birth weight, maternal age, maternal education level, paternal education level, family income, and breastfeeding practice).

239
240 Table 4 reports the logit regression analysis of the associations between mental health
241 problems and breastfeeding practices. None of the mental health problems measured were
242 significantly associated with the mother's breastfeeding practices. In order to determine if the
243 associations between mental health and breastfeeding practices differed among various
244 subgroups of the sample, a heterogeneous effects analysis was performed. The results are
245 presented in Table 5, where each panel shows adjusted odds ratio of a mother who practiced full
246 breastfeeding given a specific demographic variable and whether or not she had symptoms of
247 depression, anxiety, or stress.

Table 4. Associations between postnatal mental health problems and breastfeeding practices (n=742).

	Adjusted OR (95% CI)	p-value
Symptoms of depression	0.75 (0.47-1.74)	0.206
Symptoms of anxiety	1.23 (0.80-1.90)	0.346
Symptoms of stress	0.60 (0.35-1.03)	0.062

Notes: Mixed breastfeeding used as reference group. Controlled for demographic characteristics (infant gender, infant age, premature birth status, whether the infant was born with a low birth weight, maternal age, maternal education level, paternal education level, and family income).

Panel A shows the heterogeneous effects of family income on the association of postnatal mental health and breastfeeding practices. Compared to those without symptoms of stress and with high family income, mothers with stress symptoms and high family income (OR: 0.20, 95% CI: 0.07-0.55) and mothers without stress symptoms and low family income (OR: 0.70, 95% CI: 0.50-0.98) were significantly less likely to practice full breastfeeding. In terms of the differences between subgroups, among mothers experiencing symptoms of stress, those with high family income were significantly different than those with low family income ($p=0.011$).

Panel B shows the heterogeneous effects of maternal education level, which yielded no significant results in any of the subgroups. Panel C shows the heterogeneous effects of maternal age. Younger mothers with symptoms of anxiety were significantly more likely to fully breastfeed (OR: 1.94, 95% CI: 1.14-3.31) compared to older mothers without symptoms of anxiety. Younger mothers without symptoms of depression (OR: 1.45, 95% CI: 1.03-2.03) or stress (OR: 1.41, 95% CI: 1.01-1.96) were also more likely to fully breastfeed their infants compared to their older counterparts. Panel D shows the heterogeneous effects of paternal education level. Compared to infants who had fathers with higher education levels, infants whose fathers had lower education levels and mothers without symptoms of depression (OR: 1.48, 95% CI: 1.05-2.08), anxiety (OR: 1.42, 95% CI: 1.00-2.00), or stress (OR: 1.49, 95% CI: 1.06-2.09) were significantly more likely to be fully breastfed.

Panel E shows the heterogeneous effects of infant age. Mothers with younger infants and symptoms of anxiety were significantly more likely to practice full breastfeeding (OR: 2.68, 95% CI: 1.40-5.13) compared to mothers without anxiety symptoms who had older infants. Mothers with younger infants without symptoms of depression (OR: 1.87, 95% CI: 1.35-2.59), anxiety (OR: 1.60, 95% CI: 1.15-2.22), and stress (OR: 1.81, 95% CI: 1.32-2.49) were also more likely

to practice full breastfeeding compared to mothers with older infants. Among mothers with symptoms of anxiety, there is a significant difference between those with younger infants and those with older infants ($p=0.034$).

Discussion

The goal of this study was to examine the association between maternal mental health and breastfeeding beliefs and practices. From the sample of 742 mothers, this study found significant correlations between maternal mental health and breastfeeding beliefs. Symptoms of depression were negatively associated with breastfeeding attitude and self-efficacy, and symptoms of stress were negatively associated with self-efficacy. These results align with the international literature that has found that postnatal mental health can have a negative impact on breastfeeding beliefs (20–24).

However, the overall correlation between maternal mental health and breastfeeding practices was surprisingly not significant, contradicting previous findings in the international literature (12–17). The results did show nuanced correlations between mental health and SES, with stress associated with a lower likelihood of fully breastfeeding among high income mothers. In addition, various demographic characteristics (mother age, infant age, and paternal education) were significantly correlated with breastfeeding practices. While there are mixed results on the relationship between various demographic characteristics and breastfeeding practices in the global literature, these findings are supported by other studies in LMICs showing that mothers are more likely to breastfeed if they are younger (51), have younger infants (52,53), or have higher income status (52–54).

Interestingly, while maternal education was not significantly correlated with fully breastfeeding, there was a significant association with lower paternal education level. These results seem to contradict previous findings that higher paternal education levels are positively associated with breastfeeding (55,56). One possible explanation specific to the context of rural China is that more educated fathers are more likely to out-migrate for work than less educated fathers, leaving the mother and infant behind. Previous studies have shown that paternal support promotes exclusive breastfeeding (57,58), so it is possible that less educated fathers are more likely to be at home and provide the support the mother needs to fully breastfeed.

This study has limitations. First, the study only looks at cross sectional data and is therefore unable to draw conclusions on the directionality or causality of the relationship between mental health and breastfeeding outcomes. In addition, our assessment of mental health problems was based on a self-report assessment of symptoms rather than a professional diagnosis. Due to stigma against reporting symptoms of mental illness in LMICs, it is possible that the prevalence of depression, anxiety and stress symptoms are underestimated among women in the sample.

This study also has many strengths. This research fills a gap in the literature surrounding the associations between maternal mental health and breastfeeding beliefs and practices, specifically in rural China. Additionally, this study includes a large sample of women with newborns aged 0-6 months, a population which very few studies have focused on. Future studies could include multiple survey waves to determine directionality or perform mediation or interaction analysis between maternal mental health and breastfeeding. Further research should also investigate why the relationship between maternal mental health and breastfeeding beliefs is not associated with breastfeeding practices in this population.

316

317

Conclusions

318

319

320

321

322

323

324

325

326

Abbreviations

327

WHO: World Health Organization

328

LMICs: Low- and Middle-Income Countries

329

DASS-21: Depression, Anxiety, and Stress Scale (short form)

330

OLS: Ordinary Least Squares

331

SES: Socio-Economic Status

332

333

Declarations

334

Ethics approval and consent to participate

335

This study received ethical approval from the Stanford University Institutional Review Board

336

(Protocol # 44312). All participants provided oral consent for participation in this study.

337

Individuals found to have severe mental health problems were referred to the local hospital for

338

diagnosis and treatment.

339 ***Consent for Publication***

340 Not applicable.

341 ***Availability of data and materials***

342 Datasets used in the current study are available from the corresponding author upon reasonable
343 request.

344 ***Competing interests***

345 The authors declare that they have no competing interests.

346 ***Funding***

347 This research was supported by unrestricted gifts by anonymous donors. Funding sources played
348 no role in the design of the study, data collection, analysis, or interpretation, or in writing the
349 manuscript.

350 ***Authors' contributions***

351 QJ, YG, EZ, NC, MO, SZ, SD and SR contributed to the conceptualization of the study. QJ, YG,
352 EZ, NC, MO, SZ, SD, and SR contributed to the acquisition, analysis or interpretation of data.
353 YG, EZ, and SZ drafted the manuscript, and NC, MO, SD, HZ, HJ, and SR made substantial
354 revisions to the manuscript.

355 ***Acknowledgements***

356 We would like to thank Dr. Huan Zhou and her team at the Sichuan University West China
357 School of Public Health for their assistance in conducting this study. We would also like to thank
358 the Sichuan Provincial Center for Women and Children Health and the county-level Maternal
359 and Child Hospitals in our study area for their assistance in identifying and recruiting eligible
360 participants.

361

362

References

- 363 1. Khadivzadeh T, Parsai S. Effect of exclusive breastfeeding and complementary feeding on
364 infant growth and morbidity. *East Mediterr Health J Rev Sante Mediterr Orient Al-Majallah*
365 *Al-Sihhiyah Li-Sharq Al-Mutawassit*. 2004 May;10(3):289–94.
- 366 2. Duijts L, Ramadhani MK, Moll HA. Breastfeeding protects against infectious diseases
367 during infancy in industrialized countries. A systematic review. *Matern Child Nutr*. 2009
368 Jul;5(3):199–210.
- 369 3. Duijts L, Jaddoe VWV, Hofman A, Moll HA. Prolonged and exclusive breastfeeding
370 reduces the risk of infectious diseases in infancy. *Pediatrics*. 2010 Jul;126(1):e18-25.
- 371 4. Raisler J, Alexander C, O'Campo P. Breast-feeding and infant illness: a dose-response
372 relationship? *Am J Public Health*. 1999 Jan 1;89(1):25–30.
- 373 5. Scariati PD, Grummer-Strawn LM, Fein SB. A longitudinal analysis of infant morbidity
374 and the extent of breastfeeding in the United States. *Pediatrics*. 1997 Jun;99(6):E5.
- 375 6. Hauck FR, Thompson JMD, Tanabe KO, Moon RY, Vennemann MM. Breastfeeding and
376 Reduced Risk of Sudden Infant Death Syndrome: A Meta-analysis. *Pediatrics*. 2011 Jul
377 1;128(1):103–10.
- 378 7. Vennemann MM, Bajanowski T, Brinkmann B, Jorch G, Yücesan K, Sauerland C, et al.
379 Does breastfeeding reduce the risk of sudden infant death syndrome? *Pediatrics*. 2009
380 Mar;123(3):e406-410.
- 381 8. Allen J, Hector D. Benefits of breastfeeding. *New South Wales Public Health Bull*. 2005
382 Apr;16(3–4):42–6.
- 383 9. Stuebe AM, Schwarz EB. The risks and benefits of infant feeding practices for women and
384 their children. *J Perinatol Off J Calif Perinat Assoc*. 2010 Mar;30(3):155–62.
- 385 10. Binns C, Lee M, Low WY. The Long-Term Public Health Benefits of Breastfeeding. *Asia*
386 *Pac J Public Health*. 2016 Jan;28(1):7–14.
- 387 11. World Health Organization (WHO). *Global Strategy for Infant and Young Child Feeding*.
388 World Health Organization; 2003.
- 389 12. Henderson JJ, Evans SF, Straton JAY, Priest SR, Hagan R. Impact of postnatal depression
390 on breastfeeding duration. *Birth Berkeley Calif*. 2003 Sep;30(3):175–80.
- 391 13. Rahman A, Hafeez A, Bilal R, Sikander S, Malik A, Minhas F, et al. The impact of
392 perinatal depression on exclusive breastfeeding: a cohort study. *Matern Child Nutr*.
393 2016;12(3):452–62.

- 394 14. Sha T, Gao X, Chen C, Li L, Cheng G, Wu X, et al. A prospective study of maternal
395 postnatal depressive symptoms with infant-feeding practices in a Chinese birth cohort.
396 BMC Pregnancy Childbirth. 2019 Oct 28;19(1):388.
- 397 15. Mohamad Yusuff AS, Tang L, Binns CW, Lee AH. Breastfeeding and Postnatal
398 Depression: A Prospective Cohort Study in Sabah, Malaysia. J Hum Lact. 2016 May
399 1;32(2):277–81.
- 400 16. Silva CS, Lima MC, Sequeira-de-Andrade LAS, Oliveira JS, Monteiro JS, Lima NMS, et
401 al. Association between postpartum depression and the practice of exclusive breastfeeding
402 in the first three months of life. J Pediatr Versão Em Port. 2017 Jul 1;93(4):356–64.
- 403 17. Feldens CA, Vitolo MR, Rauber F, Cruz LN, Hilgert JB. Risk factors for discontinuing
404 breastfeeding in southern Brazil: a survival analysis. Matern Child Health J. 2012
405 Aug;16(6):1257–65.
- 406 18. Hoff CE, Movva N, Rosen Vollmar AK, Pérez-Escamilla R. Impact of Maternal Anxiety on
407 Breastfeeding Outcomes: A Systematic Review. Adv Nutr. 2019 Sep 1;10(5):816–26.
- 408 19. Adedinsewo DA, Fleming AS, Steiner M, Meaney MJ, Girard AW, MAVAN team.
409 Maternal anxiety and breastfeeding: findings from the MAVAN (Maternal Adversity,
410 Vulnerability and Neurodevelopment) Study. J Hum Lact Off J Int Lact Consult Assoc.
411 2014 Feb;30(1):102–9.
- 412 20. Dennis C-LE. Identifying predictors of breastfeeding self-efficacy in the immediate
413 postpartum period. Res Nurs Health. 2006 Aug;29(4):256–68.
- 414 21. Zubaran C, Foresti K. The correlation between breastfeeding self-efficacy and maternal
415 postpartum depression in southern Brazil. Sex Reprod Healthc Off J Swed Assoc
416 Midwives. 2013 Mar;4(1):9–15.
- 417 22. Ngo LTH, Chou H-F, Gau M-L, Liu C-Y. Breastfeeding self-efficacy and related factors in
418 postpartum Vietnamese women. Midwifery. 2019 Mar 1;70:84–91.
- 419 23. Duran S, Kaynak S, Karadaş A. The relationship between breastfeeding attitudes and
420 perceived stress levels of Turkish mothers. Scand J Caring Sci. 2020;34(2):456–63.
- 421 24. Khodabandeh M, Kashani L, Pirastehfar Z, Dashti Dargahloo S, Tabarestani M, Dordeh M,
422 et al. Relationship between Attitude towards Breastfeeding and Postpartum Depression in
423 Kerman, Iran. Int J Pediatr. 2020 Aug 1;8(8):11751–9.
- 424 25. Galler JR, Harrison RH, Ramsey F, Chawla S, Taylor J. Postpartum feeding attitudes,
425 maternal depression, and breastfeeding in Barbados. Infant Behav Dev. 2006
426 Apr;29(2):189–203.
- 427 26. Ku C-M, Chow SKY. Factors influencing the practice of exclusive breastfeeding among
428 Hong Kong Chinese women: a questionnaire survey. J Clin Nurs. 2010 Sep;19(17–
429 18):2434–45.

- 430 27. Wu DS, Hu J, McCoy TP, Efird JT. The effects of a breastfeeding self-efficacy intervention
431 on short-term breastfeeding outcomes among primiparous mothers in Wuhan, China. *J Adv*
432 *Nurs*. 2014;70(8):1867–79.
- 433 28. Cox KN, Giglia RC, Binns CW. The influence of infant feeding attitudes on breastfeeding
434 duration: evidence from a cohort study in rural Western Australia. *Int Breastfeed J*. 2015
435 Aug 21;10(1):25.
- 436 29. Sun K, Chen M, Yin Y, Wu L, Gao L. Why Chinese mothers stop breastfeeding: Mothers'
437 self-reported reasons for stopping during the first six months. *J Child Health Care Prof*
438 *Work Child Hosp Community*. 2017 Sep;21(3):353–63.
- 439 30. Kim S-H. Factors Explaining Mothers' Breastfeeding Satisfaction. *Korean J Women Health*
440 *Nurs*. 2009;15(4):270–9.
- 441 31. Fisher J, Mello MC de, Patel V, Rahman A, Tran T, Holton S, et al. Prevalence and
442 determinants of common perinatal mental disorders in women in low- and lower-middle-
443 income countries: a systematic review. *Bull World Health Organ*. 2012 Feb;90:139–49.
- 444 32. Zhang S, Dang R, Yang N, Bai Y, Wang L, Abbey C, et al. Effect of Caregiver's Mental
445 Health on Early Childhood Development across Different Rural Communities in China. *Int*
446 *J Environ Res Public Health*. 2018 Nov;15(11):2341.
- 447 33. Yue A, Gao J, Yang M, Swinnen L, Medina A, Rozelle S. Caregiver Depression and Early
448 Child Development: A Mixed-Methods Study From Rural China. *Front Psychol* [Internet].
449 2018 [cited 2020 Oct 6];9. Available from:
450 <https://www.frontiersin.org/articles/10.3389/fpsyg.2018.02500/full>
- 451 34. World Health Organization (WHO). Maternal Mental Health [Internet]. Mental Health and
452 Substance Use. [cited 2020 Dec 7]. Available from: [https://www.who.int/teams/maternal-](https://www.who.int/teams/maternal-newborn-child-adolescent-health-and-ageing/maternal-health/about/mental-health-and-substances-use)
453 [newborn-child-adolescent-health-and-ageing/maternal-health/about/mental-health-and-](https://www.who.int/teams/maternal-newborn-child-adolescent-health-and-ageing/maternal-health/about/mental-health-and-substances-use)
454 [substances-use](https://www.who.int/teams/maternal-newborn-child-adolescent-health-and-ageing/maternal-health/about/mental-health-and-substances-use)
- 455 35. China Statistical Yearbook 2018 [Internet]. [cited 2020 Oct 6]. Available from:
456 <http://www.stats.gov.cn/tjsj/ndsj/2018/indexeh.htm>
- 457 36. Wang H, Zhao Q, Bai Y, Zhang L, Yu X. Poverty and Subjective Poverty in Rural China.
458 *Soc Indic Res*. 2020 Jul 1;150(1):219–42.
- 459 37. Lovibond PF, Lovibond SH. The structure of negative emotional states: Comparison of the
460 Depression Anxiety Stress Scales (DASS) with the Beck Depression and Anxiety
461 Inventories. *Behav Res Ther*. 1995 Mar;33(3):335–43.
- 462 38. Chan RCK, Xu T, Huang J, Wang Y, Zhao Q, Shum DHK, et al. Extending the utility of
463 the Depression Anxiety Stress scale by examining its psychometric properties in Chinese
464 settings. *Psychiatry Res*. 2012 Dec 30;200(2–3):879–83.

- 465 39. Wang K, Shi H-S, Geng F-L, Zou L-Q, Tan S-P, Wang Y, et al. Cross-cultural validation of
466 the Depression Anxiety Stress Scale–21 in China. *Psychol Assess.* 2016;28(5):e88–100.
- 467 40. Mora A de la, Russell DW, Dungy CI, Losch M, Dusdieker L. The Iowa Infant Feeding
468 Attitude Scale: Analysis of Reliability and Validity1. *J Appl Soc Psychol.*
469 1999;29(11):2362–80.
- 470 41. Abdulahi M, Fretheim A, Argaw A, Magnus JH. Adaptation and validation of the Iowa
471 infant feeding attitude scale and the breastfeeding knowledge questionnaire for use in an
472 Ethiopian setting. *Int Breastfeed J.* 2020 Apr 9;15(1):24.
- 473 42. Yu J, Wei Z, Lukoyanova O, Borovik T, Fewtrell MS. Maternal Infant-Feeding Attitudes,
474 Infant Eating Behaviors, and Maternal Feeding Choice at 3 and 6 Months Postpartum: A
475 Comparative Multicenter International Study. *Breastfeed Med.* 2020 Jun 26;15(8):528–34.
- 476 43. Quinn P, Tanis SL. Attitudes, Perceptions, and Knowledge of Breastfeeding Among
477 Professional Caregivers in a Community Hospital. *Nurs Womens Health.* 2020 Apr
478 1;24(2):77–83.
- 479 44. Dai H-X, Guan X-D, Li X-M, You L-M, Lau Y. Psychometric properties of a mainland
480 Chinese version of the Iowa Infant Feeding Attitude Scale among postpartum women in
481 China. *Contemp Nurse.* 2013 Apr;44(1):11–20.
- 482 45. Dennis C-L. The breastfeeding self-efficacy scale: psychometric assessment of the short
483 form. *J Obstet Gynecol Neonatal Nurs JOGNN.* 2003 Dec;32(6):734–44.
- 484 46. Tuthill EL, Miller JD, Collins SM, Widen EM, Onono M, Young SL. HIV infection,
485 hunger, breastfeeding self-efficacy, and depressive symptoms are associated with exclusive
486 breastfeeding to six months among women in western Kenya: a longitudinal observational
487 study. *Int Breastfeed J.* 2020 Jan 16;15(1):4.
- 488 47. Aluř Tokat M, Okumuř H, Dennis C-L. Translation and psychometric assessment of the
489 Breast-feeding Self-Efficacy Scale—Short Form among pregnant and postnatal women in
490 Turkey. *Midwifery.* 2010 Feb 1;26(1):101–8.
- 491 48. Rashid AA, Shamsuddin NH, Devaraj NK. Breastfeeding Practice, Support, and Self-
492 Efficacy Among working Mothers in a Rural Health Clinic in Selangor. 2018;11.
- 493 49. Ip W-Y, Gao L-L, Choi K-C, Chau JP-C, Xiao Y. The Short Form of the Breastfeeding
494 Self-Efficacy Scale as a Prognostic Factor of Exclusive Breastfeeding among Mandarin-
495 Speaking Chinese Mothers. *J Hum Lact.* 2016 Nov 1;32(4):711–20.
- 496 50. World Health Organization (WHO). Indicators for assessing infant and young child feeding
497 practices, part 1: Definitions. World Health Organization (WHO); 2008.
- 498 51. Qiu L, Zhao Y, Binns CW, Lee AH, Xie X. Initiation of breastfeeding and prevalence of
499 exclusive breastfeeding at hospital discharge in urban, suburban and rural areas of Zhejiang
500 China. *Int Breastfeed J.* 2009 Jan 28;4(1):1.

- 501 52. Agho KE, Dibley MJ, Odiase JI, Ogbonmwan SM. Determinants of exclusive breastfeeding
502 in Nigeria. BMC Pregnancy Childbirth. 2011 Jan 11;11(1):2.
- 503 53. Alemayehu T, Haidar J, Habte D. Determinants of exclusive breastfeeding practices in
504 Ethiopia. Ethiop J Health Dev [Internet]. 2009 [cited 2020 Dec 8];23(1). Available from:
505 <https://www.ajol.info/index.php/ejhd/article/view/44832>
- 506 54. Ruan Y, Zhang Q, Li J, Wan R, Bai J, Wang W, et al. Factors associated with exclusive
507 breast-feeding: A cross-sectional survey in Kaiyuan, Yunnan, Southwest China. PLOS
508 ONE. 2019 Oct 9;14(10):e0223251.
- 509 55. Heck KE, Braveman P, Cubbin C, Chávez GF, Kiely JL. Socioeconomic Status and
510 Breastfeeding Initiation among California Mothers. Public Health Rep. 2006 Jan
511 1;121(1):51–9.
- 512 56. Ratnasari D, Paramashanti BA, Hadi H, Yugistiyowati A, Astiti D, Nurhayati E. Family
513 support and exclusive breastfeeding among Yogyakarta mothers in employment. Asia Pac J
514 Clin Nutr. 2017;26(Supplement):S31.
- 515 57. Tohotoa J, Maycock B, Hauck YL, Howat P, Burns S, Binns CW. Dads make a difference:
516 an exploratory study of paternal support for breastfeeding in Perth, Western Australia. Int
517 Breastfeed J. 2009 Nov 29;4(1):15.
- 518 58. Su M, Ouyang Y-Q. Father's Role in Breastfeeding Promotion: Lessons from a Quasi-
519 Experimental Trial in China. Breastfeed Med. 2016 Feb 2;11(3):144–9.
- 520
- 521

Table 5. Heterogeneous effects of postnatal mental health on breastfeeding practices (n = 742).

	Symptoms of depression	Symptoms of anxiety	Symptoms of stress
<i>Panel A: Family income (reference group: without MH problem & high family income)</i>			
(1) with MH problem & low family income	0.75 (0.43 - 1.32)	1.10 (0.62 - 1.96)	0.90 (0.45 - 1.80)
(2) with MH problem & high family income	0.65 (0.30 - 1.40)	1.32 (0.68 - 2.59)	0.20*** (0.07 - 0.55)
(3) without MH problem & low family income	0.78 (0.56 - 1.10)	0.81 (0.57 - 1.13)	0.70** (0.50 - 0.98)
(4) p-value of test (1) = (2)	0.753	0.656	0.011
(5) p-value of test (1) = (3)	0.878	0.269	0.467
<i>Panel B: Maternal education level (reference group: without MH problem & high education level)</i>			
(6) with MH problem & low education level	0.75 (0.41 - 1.40)	1.09 (0.61 - 1.97)	0.51 (0.26 - 1.02)

(7) with MH problem & high education level	0.60 (0.29 - 1.22)	1.20 (0.61 - 2.37)	0.81 (0.32 - 2.06)
(8) without MH problem & low education level	0.72 (0.50 - 1.05)	0.76 (0.52 - 1.20)	0.80 (0.56 - 1.14)
(9) p-value of test (6) = (7)	0.596	0.825	0.409
(10) p-value of test (6) = (8)	0.891	0.189	0.178
<i>Panel C: Maternal age (reference group: without MH problem & high maternal age)</i>			
(11) with MH problem & low maternal age	1.15 (0.66 - 2.01)	1.94** (1.14 - 3.31)	1.03 (0.55 - 1.93)
(12) with MH problem & high maternal age	0.96 (0.42 - 2.21)	1.02 (0.43 - 2.40)	0.57 (0.19 - 1.74)
(13) without MH problem & low maternal age	1.45** (1.03 - 2.03)	1.32 (0.94 - 1.85)	1.41** (1.01 - 1.96)
(14) p-value of test (11) = (12)	0.699	0.175	0.341
(15) p-value of test (11) = (13)	0.395	0.127	0.315
<i>Panel D: Paternal education level (reference group: without MH problem & high education level)</i>			
(16) with MH problem & low education level	0.98 (0.51 - 1.88)	1.66 (0.91 - 3.05)	0.83 (0.41 - 1.67)
(17) with MH problem & high education level	0.93 (0.47 - 1.84)	1.41 (0.72 - 2.77)	0.86 (0.36 - 2.07)
(18) without MH problem & low education level	1.48** (1.05 - 2.08)	1.42** (1.00 - 2.00)	1.49** (1.06 - 2.09)
(19) p-value of test (16) = (17)	0.899	0.705	0.952
(20) p-value of test (16) = (18)	0.218	0.611	0.100
<i>Panel E: Infant age (reference group: without mental health & high infant age)</i>			
(21) with MH problem & low infant age	1.11 (0.61 - 2.03)	2.68*** (1.40 - 5.13)	0.88 (0.43 - 1.83)
(22) with MH problem & high infant age	1.22 (0.64 - 2.31)	1.12 (0.62 - 2.00)	1.00 (0.47 - 2.13)
(23) without MH problem & low infant age	1.87*** (1.35 - 2.59)	1.60*** (1.15 - 2.22)	1.81*** (1.32 - 2.49)
(24) p-value of test (21) = (22)	0.829	0.034	0.799
(25) p-value of test (21) = (23)	0.189	0.239	0.124
Notes: *p<0.05, **p<0.01, ***p<0.001. Values shown as adjusted OR (95% CI). MH = mental health.			

522

523

524

525