

Patients' and clinicians' perspectives towards primary care consultations for shoulder pain: qualitative findings from the Prognostic And Diagnostic Assessment of the Shoulder programme

Benjamin Saunders (✉ b.saunders@keele.ac.uk)

Keele University

Claire Burton

Keele University

Danielle Van Der Windt

Keele University

Helen Myers

Keele University

Rachel Chester

University of East Anglia

Tamar Pincus

Royal Holloway University of London

Gwenllian Wynne-Jones

Keele University

Research Article

Keywords: Shoulder pain, primary care, general practice, diagnosis, prognosis, qualitative

Posted Date: April 19th, 2022

DOI: <https://doi.org/10.21203/rs.3.rs-1555137/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.
[Read Full License](#)

Additional Declarations: No competing interests reported.

Version of Record: A version of this preprint was published at BMC Musculoskeletal Disorders on January 2nd, 2023. See the published version at <https://doi.org/10.1186/s12891-022-06059-1>.

Abstract

Background

Clinical management of musculoskeletal shoulder pain can be challenging due to diagnostic uncertainty, variable prognosis and limited evidence for long-term treatment benefits. The UK-based PANDA-S programme (Prognostic And Diagnostic Assessment of the Shoulder) is investigating short and long-term shoulder pain outcomes. This paper reports linked qualitative research exploring patients' and clinicians' views towards primary care consultations for shoulder pain.

Methods

Semi-structured interviews were conducted with 24 patients and 15 primary care clinicians. Twenty-two interviews (11 patients, 11 clinicians) were conducted as matched patient-clinician 'dyads'. Data were analysed thematically.

Results

Clinicians reported attempts to involve patients in management decisions; however, there was variation in whether patients preferred treatment choice, or for decisions to be clinician-led. Some patients felt uncertain about the decisions made, due to a lack of discussion about available management options. Many GPs expressed a lack of confidence in diagnosing shoulder pain. Patients reported either not being given a diagnosis, or receiving different diagnoses from different professionals, resulting in confusion. Whilst clinicians reported routinely discussing prognosis of shoulder pain, patients reported that prognosis was not raised. Patients also expressed concern that their shoulder pain could be caused by serious pathology; however, clinicians felt that this was not a common concern for patients.

Conclusions

Findings showed disparities between patients' and clinicians' views towards shoulder pain consultations, indicating a need for improved patient-clinician communication. Findings will inform the design of an intervention to support treatment and referral decisions for shoulder pain that will be tested in a randomised controlled trial.

Background

Musculoskeletal (MSK) shoulder pain is common and associated with impacts for the individual, healthcare and society [1, 2]. Many shoulder pain problems are managed predominantly in primary care, and in the UK, approximately 3% of adults consult their general practitioner (GP) for shoulder pain annually [3]. Whilst many shoulder pain problems are self-limiting and of short duration, 40–50% of

people with shoulder pain continue to experience problems 6 to 12 months after first consulting their GP or physiotherapist [4, 5]. Individuals with persistent shoulder pain have reported negative impacts on daily activities including work, sleep, and self-care, e.g. dressing and bathing [6, 7].

There are many uncertainties regarding the clinical management of shoulder pain. Whilst research has highlighted modest short-term effects of commonly used treatments such as corticosteroid injection, therapeutic exercise and manual therapy, there is limited evidence for long-term benefits [8, 9, 10]. There is a degree of uncertainty in relation to shoulder pain prognosis and diagnosis. Shoulder pain prognosis is variable, and systematic reviews of shoulder physical examination tests have highlighted variation in performance and interpretation of these tests, resulting in low diagnostic accuracy. There is also limited evidence available regarding the usefulness of diagnostic imaging, such as USS (Ultrasound Scanning) or MRI (Magnetic Resonance Imaging) [11].

These uncertainties can result in challenges for clinicians when deciding on the most appropriate treatments for patients. These challenges have been highlighted by general practitioners (GPs) in the Netherlands, particularly in relation to diagnostic uncertainty [12], and UK GPs have reported a lack of confidence in managing MSK problems in general, including shoulder pain [13]. However, the impact that these uncertainties and challenges have on decision-making between primary care clinicians and patients in consultations has not been fully explored in the literature. Whilst clinicians' and patients' views have been explored separately in the literature, there has been no research directly comparing their views towards shoulder pain management in primary care settings. Gaining an in-depth understanding of clinicians' and patients' perspectives is an important step towards identifying ways to better support clinicians and patients to address the current challenges and uncertainties in order to optimise treatment and referral decisions, which in turn may improve patient outcomes. The aim of this paper is therefore to explore patients' and clinicians' perspectives towards primary care consultations for shoulder pain. This includes the influences on, and uncertainty regarding, decision-making and perspectives on the importance of diagnostic and prognostic information when negotiating treatment and referral options.

This study is linked to an 8-year research programme funded by the UK National Institute for Health Research (NIHR) and Versus Arthritis: Prognostic And Diagnostic Assessment of the Shoulder (PANDA-S). PANDA-S is investigating the short and long-term outcomes of shoulder pain, with the aim of informing the design of an intervention to support clinicians and patients to make optimal decisions regarding self-management, treatment and referral. A longitudinal cohort study is currently being undertaken of 492 patients with shoulder pain presenting to UK NHS general practices and physiotherapy services [14]. Participants invited to take part in the qualitative study were recruited from the cohort (see Recruitment and Sampling for details).

Methods

One-to-one semi-structured interviews were carried out at a single time-point with patients ($n = 24$) and clinicians ($n = 15$). Ten of the clinicians were GPs, 1 was an advanced practice physiotherapist working in

a general practice as a First Contact Practitioner (FCP) (FCPs are utilised in UK primary care as the first point-of-contact for MSK presentations), and 4 were physiotherapists. Interviews were carried out between May 2019 and June 2021, which included a 10-month break in recruitment to the qualitative study from March 2020 to January 2021 as a result of the COVID-19 pandemic. Twenty-two interviews (11 patients, 11 clinicians) were conducted as 'dyads', where the patient and the clinician they had consulted with for their shoulder pain were interviewed separately about the same consultation. This was to allow for direct comparison between patients' and clinicians' accounts of how decisions about treatments and referral were reached. The other 17 interviews (13 patients, 4 clinicians) were standalone interviews, i.e. they were not part of a 'dyad'.

The study settings were general practices and physiotherapy services in the West Midlands, North West and Thames Valley/South Midlands regions of England. The study received ethical approval from the NRES Committee Yorkshire & The Humber - Sheffield, 16/10/2018, ref: 18/YH/0346.

Recruitment and Sampling

Patients were recruited by invitation letter and then via phone, having consented to contact as part of their participation in the PANDA-S cohort study. Patients were purposively sampled from the baseline questionnaires completed in the cohort, to capture a range of characteristics; including, age, gender, pain duration, self-reported pain intensity and geographical location. Patients who agreed to take part in an interview were also asked for their permission to contact the clinician with whom they consulted for their shoulder pain, in order to invite them to a separate interview. Patients gave permission for this in all cases.

Clinicians were contacted via emailing their practice administrator or practice manager, followed by a telephone call or further email. In cases where patients could not remember the clinician's name, the practice manager first identified which clinician the patient had consulted, and then put the research team in touch with the clinician. The patients interviewed had all consulted with a GP except for one patient who had consulted with an FCP. These 'matched' clinicians were invited for a separate interview. 13 of the 'matched' clinicians either did not reply to the invitation, or declined citing lack of time, and 11 were interviewed (resulting in 11 'dyads'). In the case of the 13 patient interviews where the 'matched' clinician had not been interviewed, the patient's interview was included in the dataset as a standalone interview. Patient and GP interviews were conducted as near as possible to the time of the consultation in which the patient was identified for PANDA-S. In some cases, this was several weeks, but others were interviewed a few months after the consultation.

Given that all of the patients interviewed had consulted with a GP – as well as one FCP who was based in a general practice— at a later stage in the study recruitment it was decided to invite MSK physiotherapists to standalone interviews (i.e. not as part of a dyad), to also explore their views and experiences. Whilst the physiotherapists were invited from services participating in the cohort study, they had not consulted with patients recruited to the qualitative study. The intention was to explore their views about

management of shoulder pain more generally, both in treating patients who had self-referred to physiotherapy services and those patients referred to physiotherapy from general practice. Physiotherapists were contacted directly via email and invited to participate. All of the physiotherapists who were invited agreed to be interviewed.

Data collection

Of the 24 patient interviews, 11 took place at participants' homes; and 13 via telephone. Interviews lasted between 21 minutes and 53 minutes. Five clinician interviews took place at their practice/clinic, and 10 were via telephone. Clinician interviews lasted between 19 minutes and 38 minutes. All of the interviews conducted after the start of the COVID-19 pandemic (9 patients, 8 clinicians) were via telephone. All interviews were conducted by BS (male, PhD), an experienced qualitative researcher from a social science background.

All participants were given an information leaflet explaining the study prior to providing written informed consent (for in-person interviews) or audio-recorded consent (for telephone interviews) at the start of interview. Interviews were audio-recorded. Field notes were not made during interviews or focus groups or as it was felt this could negatively impact upon the rapport between researcher and participants. Consent was reaffirmed verbally at the end of each interview.

Separate topic guides were used for patient and clinician interviews, covering a range of areas relevant to the study aims (see supplementary files). The PANDA-S Clinical Advisory Group (CAG) and Patient and Public Involvement and Engagement (PPIE) group, helped to develop these topic guides. The PPIE group consisted of 4 patients with experience of receiving care for shoulder pain. The CAG group included 11 clinicians: 7 physiotherapists, 3 GPs and 1 rheumatologist. Topic guides were used flexibly within interviews, allowing the interviewer to follow up on any unexpected findings. To aid recall during interviews, several clinicians referred to the medical record from the consultation being discussed. However, only the parts of the medical record directly related to the patient's shoulder pain management were communicated to the interviewer. Early interview findings informed subsequent data collection, with the topic guides iteratively revised throughout the data collection process.

Data analysis

Audio-recordings of interviews were transcribed and anonymised. Transcripts were checked for accuracy against the audio, but were not returned to participants for comment or correction so as to avoid overburdening. Data were analysed thematically, using the constant comparison method [15], looking for connections within and across interviews, and across codes, highlighting data consistencies and variations. Analysis was an iterative process and data collection continued until saturation was judged to have been reached, defined as 'informational redundancy' – the point at which additional data no longer offers new insights [16].

Anonymised transcripts were first systematically coded on a line-by-line basis by BS, with the aid of the software program Nvivo 12, in order to identify recurrent concepts inductively. Coding was at first largely

descriptive, and later became more conceptual as interpretations of the data moved towards a higher level of theoretical abstraction. Coding was reflexive and recursive, with codes being revisited in light of the findings of subsequent data-collection. Analysis began with the patient data and then mapped the views and experiences of clinicians against that, to allow for comparison between the two datasets.

Impressions of the data based on early analysis were discussed among the research team, and with the PANDA-S programme's independent steering committee. Transcripts from 2 patient-clinician dyads, i.e. 2 patient transcripts and 2 clinician transcripts, were then independently coded by two other members of the team. Coders brought different disciplinary perspectives to the data (BS: social science; GWJ: nursing, epidemiology; CB: general practice). The aim was to understand cross-disciplinary perspectives on the data and, through discussion, to come to an agreement on shared meanings and interpretations. Based on these analyses, four main themes were developed. These themes, along with accompanying data extracts, were then presented to the study's CAG and PPIE groups in separate meetings, in order to gain their perspectives. Members of both groups expressed strong agreement with the themes developed and the research team's interpretations of these themes.

A further stage of analysis was then carried out in which identified themes were explored specifically in relation to the 11 patient-clinician dyads. This involved close analysis of each of the matched patient-GP transcripts, and the matched patient-FCP transcripts, in order to explore the key themes within the context of views towards individual consultations. Analysis was therefore both across and within cases, allowing for a broader understanding of patients' and clinicians' views, as well as fine-grained, idiographic investigation of patients' and clinicians' perspectives on individual consultations.

In what follows we outline the characteristics of the participant sample, before reporting the key themes.

Results

Patient characteristics

Eleven patients were female and 13 male, aged from 38 to 79 years (average age: 62). Eleven patients were currently in employment, representing a range of occupation types, 12 were retired, and 1 was currently not in work. Patients were located across each of the UK geographical regions recruited from for the cohort: West Midlands, North West, Thames Valley and South Midlands. Pain severity at its worst over the past week, measured at baseline in the cohort study varied between 3/10 and 10/10, with an average of 6.7/10. Duration of the current episode of shoulder pain measured at baseline ranged from 2–6 weeks to over >1 year. Table 1, below, summarises the characteristics of the 24 patients interviewed.

Table 1
Characteristics of the patients interviewed

Gender	Age	Geographical location	Duration of current pain episode	Pain severity at its worst over the past week (on a 0–10 scale, where 0 = no pain; 10 = worst pain)	Employment status
F	58	West Midlands	> 1 year	6/10	Not working due to ill-health
F	79	West Midlands	2–6 weeks	3	Retired
M	68	West Midlands	2–6 weeks	8	Retired
M	74	South Midlands	> 1 year	8	Retired
M	74	North West	6–12 weeks	4	Retired
M	58	South Midlands	2–6 weeks	7	Employed
M	46	North West	3–6 months	6	Employed
M	60	North West	6–12 weeks	5	Retired
F	67	West Midlands	6–12 weeks	8	Retired
F	68				Retired
F	55	South Midlands	6–12 weeks	10	Employed
M	70	South Midlands	6–12 weeks	7	Retired
M	69	West Midlands	2–6 weeks	10	Retired
F	38	West Midlands	6 m-1 year	7	Employed
M	57	North West	3–6 months	8	Employed
M	53	West Midlands	6m-1 year	8	Employed
F	39	Thames Valley	> 1 year	7	Employed

Gender	Age	Geographical location	Duration of current pain episode	Pain severity at its worst over the past week (on a 0–10 scale, where 0 = no pain; 10 = worst pain)	Employment status
M	65	Thames Valley	> 1 year	10	Employed
M	70	Thames Valley	2–6 weeks	5	Retired
F	64	Thames Valley	3–6 months	4	Employed
M	55	Thames Valley	6–12 weeks	7	Employed
F	79	Thames Valley	> 1 year	7	Retired
F	73	Thames Valley	2–6 weeks	8	Retired
F	41	Thames Valley	6 m–1 year	5	Employed

[Table 1 inserted here]

Clinician characteristics

Of the 15 clinicians interviewed, 8 were female and 7 male. Five GPs were female and 5 male; 2 MSK physiotherapists were female and 2 male, and the 1 FCP was female. The length of time clinicians had been practising ranged from 3 to 32 years. Clinicians were located across each of the geographical regions recruited from for the cohort.

Main themes

The four main themes identified were:

1. Decision-making and discussion of management options
2. Diagnosis of shoulder conditions
3. Discussion of prognosis
4. Giving and receiving reassurance

Theme 1: Decision-making and discussion of management options

Clinicians highlighted a number of considerations in the management of shoulder pain, such as the patient's age, occupation, pain history and the impact of pain on their daily lives. However, they indicated

that whilst these factors may influence their decisions, they routinely attempt to involve patients in decision-making through offering them a choice of treatments:

You need to take into account the patient demographics: age, sex, occupation and their presenting history, history of how the pain has started and presents...but I think it's got to be a shared decision process between you and the patient as to which route they want to take...ultimately it's patient choice.
(Physiotherapist, male, 8 years practising)

This correlates with the views of some patients who reported making the final decision between treatments offered to them by the clinician:

P: He said I could leave it a while and have an injection, or I could refer you to physiotherapy. So, I thought I'd try physio.

Int: So it sounds like the doctor gave you some choice of which you preferred?

P: Yeah, it was my choice...I thought the physio might be a better long-term solution to the underlying problem rather than injection (Female patient, aged 64)

Clinicians reported that, in the absence of 'red flag' symptoms, in most cases they favoured initial conservative management, such as self-management advice, analgesia or referral to physiotherapy. However, some clinicians reported offering more invasive options, such as corticosteroid injection, at any early stage if the pain was significantly impacting the patient's everyday life, or the patient was seeking short-term pain relief. This was the case in the following extract from a clinician-patient dyad:

So there was bicipital tendinosis, and [the patient] came back (for a follow-up consultation) and said the pain was still the same, it was still disturbing his sleep, what would be a quick fix. So we spoke about the pros and cons of injection, and he did have an injection...people do come with a pain, it's affecting their sleep, it's affecting their lifestyle. For people like this patient, who it does give them relief, then I think it's a service that's appreciated. (GP, female, 32 years practising)

The matched interview in this dyad shows consistency with the clinician's account, in that the patient also highlighted the impact of the pain on his sleep as being his main concern. The patient reported having been hesitant about an injection, but expressed a strong preference for the decision about his treatment to be led by the GP:

It was a few days after when it started to occur and I couldn't sleep. Oh, it was a nightmare. The whole of my arm ached...I couldn't care less about medical stuff. I'll leave that to the doctors. I'm not interested. All I want them to do is to treat me. So all I wanted, get rid of this pain in my arm, whatever way, let them tell me. I didn't fancy an injection in my shoulder. I got home and thought, 'What have I done?'. But do you know, from then on after that night it went easier. (Male patient, aged 64)

Clinicians similarly highlighted this preference of some patients not to be involved in their treatment decisions and instead wanting the clinician to take a directive approach, deferring to their clinical

knowledge:

I do try and do shared management decision-making as much as possible with people and some patients are very amenable to that, and other people say 'Well you're the doctor, what should I do? What would you recommend?'. And then you think 'Right okay, I think we probably should do this first and this next'. (Female GP, 3 years practising)

However, some patients' views challenged the dichotomy between patient choice and clinician-led decision-making. They indicated that, even if they preferred the eventual decision to be taken by the clinician, they still wanted to engage in a form of shared decision-making involving discussion about available options and for the clinician to outline a potential treatment pathway. However, for many patients they reported that these discussions had not taken place:

He basically told me what it was [i.e. bursitis], felt along my shoulder and explained where things were and that there was the fluid sac or something, so basically that was it...if other treatment is available like physio or if I need another scan or anything like that, I would have liked to have discussed that because I don't just want to be told, 'oh well it's this, learn to live with it'. (Female patient, aged 38)

Some patients also reported a lack of information from the clinician about why the recommended treatment was the most suitable and how it would treat their symptoms. They seemed to indicate a reluctance to raise these concerns during the consultation, but then felt dissatisfied following the consultation:

Basically the doctor asked me, would I like to have the steroid injections and I asked him 'would it do me any good?'; which he said he thought it would, made my appointment...and that was basically the end of the consultation...when you come away you think, 'what did I learn then?'. It was nothing, I learnt nothing, I didn't know what the injection would do or why. (Male patient, aged 70)

Theme 2: Diagnosis of shoulder conditions

Receiving a diagnosis for their shoulder pain was important to patients, partly to reassure them that the pain was not caused by serious pathology (this is explored further in the 'Giving and receiving reassurance' theme, later), but also this was seen to be key to receiving the right management:

It's nice to be told what it is [i.e. bursitis]...they could actually tell me what I can take to relieve the pain and not just keep saying 'oh we don't know, try this, it's just something you are going to have to live with'... I know now that if it gets worse I can go to the doctors and they are aware of what it is and maybe how they can treat it. (Female patient, aged 38)

However, several patients reported that their GP either did not offer them a diagnosis or explanation of possible causes of their shoulder pain, or that they conveyed uncertainty. In many cases this resulted in the patient being referred for further investigation:

The GP thought that it was potentially something to do with the ligaments, just something's worn out there and she thinks the bones are actually scratching on each other and that's why it's causing this sharp pain. It's like the nerve's been damaged there, that's what she thinks has been going on...she will see on the ultrasound. She can't say 100% but that's what she's guessing. (Female patient, aged 39)

This diagnostic uncertainty was reflected in clinicians' views. Many GPs reported lacking confidence in diagnosing shoulder conditions due to a perceived lack of skill in shoulder examination, leading them to commonly send patients for investigations such as an ultrasound scan:

I'm conscious that I do probably far too many ultrasounds...and I probably need to brush up on specific shoulder examination skills in order to be able to more specifically diagnose a specific tendonitis or bursitis or impingement or whatever, rather than just thinking 'ooh it could be this that or the other', without having to image; unless there were any red flags obviously. (GP, female. 32 years practising)

There was variation in the data, however, as all of the physiotherapists and a few GPs reported having greater confidence in diagnosing shoulder conditions. For many of these clinicians, providing a diagnosis was perceived to be a key part of their professional role:

If my role isn't to diagnose people, then what am I doing? Of course it is...I'm perfectly happy in that part of my role. But I also am aware sadly of things like NICE [National Institute for Health and Care Excellence] guidance which seems a lot more investigation-focused, and I do think maybe some of my younger colleagues are more inclined to sort of hedge before making a definitive diagnosis, whereas I'm quite happy to put the code in, subacromial and move on. So yes, I think that's an absolutely crucial part of my role. (GP, male, 28 years practising)

The majority of clinicians reported that during the initial stages of the COVID-19 pandemic, the need to conduct consultations remotely via telephone or video call had made diagnosing shoulder conditions more difficult. This was because in many cases physical contact with the patient was felt necessary for assessment and diagnosis. Additionally, examinations were limited by the patient's ability to follow the clinician's instructions in carrying out arm movements, as well as ensuring that these movements were captured on the screen:

To assess strength appropriately in different movements and different directions, you're reliant on the patient having something suitable in their home to be able to do it for starters. Secondly, then to be able to position the phone in such a way that you can see them doing it, which sometimes isn't the case. You also then need to be able to guide them with the technique...one of the things then you're looking at with your rehab folk is trying to normalise that movement pattern. Sometimes that does require physical feedback, not just verbal feedback. So I think it's a challenge to do over video. It's not impossible, but it's immensely difficult.

(Physiotherapist, male, 11 years practising)

This view was mirrored by patients, who felt that remote examinations were less thorough than in-person, hands-on examination, and also reported difficulty performing movements in line with the clinician's instructions over video. As a result, patients reported feeling less confident when a diagnosis was provided:

You don't get a full examination unless you meet the person I'd imagine. I think you can only do so much on the telephone or on video. I'm disappointed really that she was taking my [previous] diagnosis without examining me...I've had a video call from her and it was comical, absolutely comical. 'Right put your phone in front of you, lift your arm up so I can see it'. [I replied] 'Look, if I put my phone there I can't see it, so I can't see what you can see, so I can't do the exercise'. (Male patient, aged 53)

Several patients reported receiving different diagnoses from different professionals, leading them to experience confusion. This is exemplified in the following clinician-patient dyad. The patient reported feelings of uncertainty, having been told by his GP that he had arthritis, but by another professional that he had subacromial shoulder pain (i.e. impingement):

They've been telling me different things. The first doctor who rang said it's just arthritis...they referred me to physiotherapy...but the man who did my scan at the hospital told me it was quite a bad impingement. So I didn't know which one to believe. (Male patient, aged 65)

The First Contact Practitioner (FCP) that the patient consulted with explained that, based on the patient's X-ray and her examination, both diagnoses were indicated:

He had an X-ray of his shoulder which showed calcific rotator cuff tendonitis and subacromial bursitis and then the ultrasound showed some moderate OA (osteoarthritis) changes in his acromioclavicular joint. Normal long head of biceps, no rotator cuff tear but some thickening of the bursa which was felt likely to be representative of impingement. (FCP, female, 24 years practising)

Theme 3: Discussion of prognosis

Patients reported concerns about how long their shoulder pain might last and whether it could worsen or recur. In some cases, patients appeared to catastrophize about the future:

The pain is getting worse and there will be a point where I won't be able to move my arm beyond that point [signals waist height]. And so that's obviously going to limit my movement, which will impact my job severely. And who knows?" (Male patient, aged 46)

However, the majority of patients reported that their shoulder pain prognosis had not been discussed with the clinician. Many perceived that this was because the clinician was unlikely to know how their pain may progress, particularly if a diagnosis had not been established. Some patients appeared accepting of this uncertainty so long as a clear management plan was in place:

Int: Did the GP give you any idea about the future, whether it might be something that could be resolved, or...?

P: No, no, nothing, the GP didn't really say, but nobody knows...she doesn't really know exactly what's wrong with me, she can't give me promises, because then I could be, 'oh yeah, she said it'll be fine, you know after this injection', and then if it isn't, I'd be like well 'why did she say that?' So I think it's more not giving me false hope, she's like 'okay, let's see what we can do, we can offer you this and that, and if none of this works, we will try something else'. (Female patient, aged 49)

In contrast to this, other patients reported that they would have liked their prognosis to have been discussed in the consultation:

Int: Did they talk to you at all about the future outlook, the prognosis?

P: No, nothing of the sort. No, we didn't talk about that.

Int: Is that something you'd liked to have discussed with the clinician?

P: Absolutely yeah. Managing expectations yeah, it's a big thing isn't it? Just to give me an idea if you're going to get better or have I just got to live with it. (Male patient, aged 53)

Clinicians' accounts showed discrepancy with these views, as all clinicians reported that they routinely discuss shoulder pain prognosis in consultations. The perceived this to be important for encouraging patients to engage with a management plan:

Int: Would you discuss with patients how their pain might progress?

P: Yeah all the time. I usually say, 'right, for this sort of problem, from previous experiences and from the research out there it looks like it will take X amount of time to get better and if you engage with your physio programme the aim is to reduce that time scale of course'...just give them that advice on time scales and what we do next should they not improve within that timescale.

(Physiotherapist, female, 6 years practising)

Some clinicians reported a more cautious approach whereby they give patients a general idea of a timeframe for recovery rather than a specific prognosis. They also highlighted the importance of ensuring patients have realistic expectations about their recovery:

I want people to be realistic, so I don't want them to expect it to get hugely better immediately and then be upset that it isn't. So I will often say something like, 'Shoulder problems can go on for months' because I want them to have some realistic expectations. But I would then try and counter it by saying, 'But if we do this and that and the other we'll aim to make it feel better along the way and improve more quickly.'...because if you tell someone with a frozen shoulder the natural history is 18 months without any treatment, I mean that's just depressing, but also realistic. So I kind of hedge my bets in the middle a

little...what I don't want them to do is ring me in a week and say, 'It's not better yet.' So it's just getting that balance really. (GP, female, 15 years practising)

The disparity in patients' and clinicians' accounts of discussing prognosis in consultations is exemplified in patient-clinician dyads. The following patient reported that neither diagnosis nor prognosis were discussed with his GP, and that his expectation was that his shoulder pain may never resolve:

The doctor didn't actually say what he thought it was, he just thought the way to go forward was physiotherapy... my feeling is that I've probably got to live with it. I don't really know what to expect. I'm not optimistic because I'm not 100% sure it'll ever go. (Male patient, aged 74)

This contrasts with the GP's account in which he reports having communicated to the patient a diagnosis of frozen shoulder and outlined the likely timeframe for recovery:

Int: Did you discuss [the pain condition] with him in the consultation?

P: Yes, yes. I always do, I always give explanation to patients that what I think it is, in lay terms, a frozen shoulder, and I gave him an idea of what it was about and then I referred him to physio...prognosis is vitally important. You need to give them an idea of timeframe because otherwise they're not getting better. You've got to tell them the likelihood, how long it's going to last before getting better, otherwise they come back and they don't have faith in your management. (Male GP, 34 years practising)

Theme 4: Giving and receiving reassurance

Clinicians reported that patients commonly needed reassurance about the impact of shoulder pain on their function and daily activities. However, they felt that, unlike spinal pain, patients do not tend to have concerns about serious pathology related to their shoulder:

I don't see necessarily that patients have that mentality and worrying about it being something nasty. However, I do see a lot more that they worry about whether they're allowed to do certain things. So are they allowed to lift their arm if it's painful? So I see a lot more worry about the pain or how long did it last rather than it being due to a non-MSK or a more sinister pathology.

(Physiotherapist, male, 11 years practising)

Patients' accounts showed some discrepancy with these views, however, as patients reported concerns that something serious may be causing their shoulder pain, such as cancer, or a serious injury to the bones or ligaments. They therefore reported feeling reassured if they received a diagnosis:

Looking for exercises to help it, you go onto Google and it comes back you've got cancer straight away, isn't it? Everything, 'Oh yeah, points to cancer'... Once I knew it was tendonitis I was happy. I wasn't happy, but I was happy if that makes sense.

(Male patient, aged 46)

The disparity between clinicians' and patients' views is reflected in the following dyad. The patient expressed concern that something serious may be wrong with his shoulder, but this appears to have either not been communicated, or not identified by the clinician:

You always think these things, is it some sort of bone cancer? Or, I'm not sure what can be done on ligaments or muscles that would be more serious, I'm not that familiar with things that go wrong in your shoulder. But yes you always wonder could it be something very serious, have I done some serious injury and not realised it? Is it some disease that's eating away at my muscle or bone? (Male patient, aged 60)

I don't think it was anxiety as such. It was more like he said, 'I'd like not to have this pain please, if you can?' rather than anything else. I certainly wouldn't describe anxiety as a presenting feature with this chap. (GP male, 24 years practising)

In some cases, when asked in interviews patients reported that they had not in fact expressed these concerns to their clinician:

P: When you get things like this you think the worst. I'm a bit of a worrier and I just think oh, cancer or something like that.

Int: Was she able to reassure you that it wasn't?

P: No, no, no.

Int: Did you express those concerns to her?

P: Well I didn't, no. I didn't actually, no. (Female patient, aged 56)

Some patients who had not received a diagnosis for their shoulder pain suggested a tension whereby on the one hand they felt a sense of reassurance through having confidence in the expertise of their clinician, particularly where there was a strong therapeutic rapport, but at the same time they felt a lack of reassurance with regard to the cause of their pain:

She allows you time to go through everything and talks to you about everything, she's a fabulous doctor. But in this situation she wouldn't be able to tell me exactly what it was...so it was sort of reassuring but it wasn't, at the same time.

(Female patient, aged 68)

A few GPs displayed a recognition of this tension, and highlighted the challenge they faced in attempting to reassure patients within the frame of diagnostic uncertainty. They reported addressing this through being honest about these uncertainties, and involving the patient in discussing management:

I think [the patient] was happy that the X-ray didn't show any arthritis particularly, but I think he was then worried to say 'Well what is it? What's causing it?'. And I think that's the difficult point where you say 'Well

it could be any number of things'... I believe very much so that we need to be honest with our patients if we don't know the answer. What I often will say is 'Look, these are the possibilities and these are the things I'm going to do to rule in or rule out those possibilities'. And I try and have a conversation about it and do shared management decision-making as much as possible. (Female GP, 6 years practising)

Clinicians also highlighted the need, in some instances, to provide reassurance about the relative value and risks of treatment options; for instance, reassuring patients that physiotherapy will not cause them harm or worsen their pain, even if the cause of pain is unclear:

Sometimes patients are nervous to go to physio if they don't know what's going on and we do find that's sometimes a bit of barrier. Some of these people find it very reassuring to have something in terms of a scan or an X-ray that says 'This is what's happening'. Some people seem to worry that the physio will make things worse and I try to explain to a lot of patients that physio is very unlikely to make anything worse; chances are it'll make probably make things better. (Male GP, 20 years practising)

Discussion

The findings show some consistencies between the views of patients and primary care clinicians towards shoulder pain management, but also a number of disparities, which could reflect miscommunication or misunderstanding within consultations, or a lack of recall following the consultation. Both clinician and patient data indicated that there was variation in patients' preferences towards shared decision-making. However, some patients reported leaving the consultation with a lack of clarity about their management, as they did not feel that available management options had been discussed, nor a clear treatment pathway outlined. Diagnostic uncertainty was apparent in the accounts of some clinicians and patients, and as a result patients expressed concerns that there may be something seriously wrong with their shoulder. Clinicians perceived these types of concerns to be uncommon for patients with shoulder pain, and therefore did not always provide the 'cognitive reassurance', i.e. information about aetiology [17] that patients sought. However, it appears that this may be due to patients' reluctance to raise these concerns with their clinician. Some patients highlighted a tension between feeling reassured on the one hand due to the confidence and trust they have in their clinician, i.e. 'affective reassurance' [17], yet at the same time feeling a lack of cognitive reassurance if they had not received a diagnosis to alleviate their concerns about serious pathology or injury. Discrepancy was also observed in that patients reported that their prognosis had not been discussed in their consultation, whereas clinicians reported that they routinely outlined a timeframe for recovery, which they perceived to be important for patient outcomes. It may be that this was either not clearly communicated, or not comprehended within consultations, or that patients and/or clinicians had difficulty reliably recalling these discussions.

Comparison with other literature

Some of our findings show similarity with previous qualitative research on shoulder pain management. The importance that patients placed on receiving a diagnosis for informing the management of their

shoulder pain corroborates Cuff and Littlewood's [18] findings on patients with subacromial impingement syndrome. However, Cuff and Littlewood reported that patients in their study were able to give a clear and accurate explanation of their diagnosis, which differs from our finding that some patients reported confusion about the cause of their pain even when they had been given a diagnosis. This difference may be explained by Cuff and Littlewood having recruited patients referred for physiotherapy at an orthopaedic shoulder clinic, who may have received a more detailed explanation of their condition in this setting. Our findings perhaps align more with Ottenheijm et al.'s [12] study of Dutch GPs, who reported patients' confusion about their shoulder pain diagnosis due to diagnostic disagreements between different professionals. We also observed patient confusion due to having received different diagnoses. However, this was not caused by cross-professional disagreement, but was due either to a lack of explanation to the patient that different diagnoses can co-occur, or patients not recalling that this was explained to them. Another similarity between ours and Ottenheijm et al.'s [12] findings relates to GPs' uncertainty about shoulder diagnosis; some GPs in their study also felt that their diagnostic knowledge was insufficient in relation to the shoulder. This is corroborated by a UK-wide survey of 714 GPs which found that diagnostic uncertainty commonly resulted in GPs referring patients for further investigations such as blood tests, radiographs and USS [19].

Our finding that patients experienced concerns and anxiety about their shoulder pain shows similarity with other studies. However, the patient concerns about serious underlying pathology such as cancer, or serious injury, that we identified were not observed in these studies. Instead, patient anxiety was found to be a result of delays in receiving a diagnosis [20], functional limitations of shoulder pain [7], and fears about symptom exacerbation [6]. Similar to our study, a qualitative review by Cheung and Soundy [21] highlighted the importance to patients with shoulder pain that their worries and concerns are addressed through clinicians providing both affective and cognitive reassurance [17]. However, this review did not draw out the tension we observed that patients experience when one form of reassurance is provided but not the other, nor the challenges clinicians face in reassuring patients in the absence of a diagnostic explanation.

The issues and concerns identified among patients and clinicians about shoulder pain consultations being carried out remotely via telephone or video, show both similarities and differences with other literature. Donaghy et al [22] found that GPs and patients responded positively to video consultations, generally finding them helpful and convenient. However, their study looked at primary care consultations more broadly, and not specifically shoulder pain. It was also carried out prior to the COVID-19 pandemic; therefore, at this time clinicians and patients would have had more choice as to which types of problems could be most effectively managed via remote consultations. A later, post-COVID-19 study by Murphy et al [23], showed mixed findings, with GPs highlighting that for certain conditions there is a need to examine the patient in-person in order to visualise signs and symptoms close-up. This shows similarity with our findings, as both patients and clinicians indicated a strong preference for hands-on, in-person examination of the shoulder.

Strengths and limitations

A strength of this study is the parallel investigation of the views of both patients with shoulder pain and primary care clinicians. This included 11 patient-clinician dyads (i.e. 22 matched interviews), which represents a novel aspect of the study, allowing for direct comparison of views towards the same consultation. The multi-disciplinary team involved in data analysis is also a strength, as well as the input from our PPIE group and Clinical Advisory Group, which increases the trustworthiness of the findings presented. A further strength is that whilst this study was conducted in a UK NHS setting, because of the focus on the primary care consultation itself, many of these findings could also be applicable to other non-UK settings.

A limitation of the study is that participants were interviewed several weeks after the consultation took place, therefore recall will likely have been a factor in some cases. GPs were able to use their clinical notes to aid their memories, but these notes would likely have included only a brief record of the decisions made. Therefore, where differences were identified between patients' and clinicians' accounts of what was discussed in the consultation, this disparity could have been due, in part, to participants misremembering, or being unable to recall certain details. However, this does not negate the usefulness of the findings, because if patients were unable to recall crucial information about diagnosis, prognosis or treatment decisions, or if clinicians recorded or recalled information differently from the way in which it was discussed, this can represent a key barrier to positive patient experiences of care and possibly outcomes.

When interpreting the findings, it is important to acknowledge the influence of the researcher on participants' responses in interviews; however, a reflexive approach was adopted throughout, in which the researcher attended to, and acknowledged any biases and preconceptions. Patients and clinicians were made aware that the researcher conducting interviews was a social scientist and was not from a clinical background, and that the team were interested in understanding both positive and negative aspects of their experiences.

Conclusions And Implications For Clinical Practice

In this paper we have presented findings on patients' and clinicians' views towards primary care consultations for shoulder pain. This included, influences on, and uncertainty regarding, decision-making and the importance of diagnostic and prognostic information when negotiating treatment and referral options.

There are a number of implications that can be drawn out for future clinical practice and research. This is particularly in relation to the disparities identified between patients' and clinicians' accounts of the consultation and its outcomes. These disparities point to apparent miscommunication and/or miscomprehension and lack of recall of key information following the consultation. This may have led to many patients reporting uncertainty about their shoulder pain diagnosis, prognosis or the available treatment options. It is therefore important that, in line with UK National Institute for Health and Care Excellence (NICE) guidelines for shoulder pain management [24], clinicians should consistently provide

clear information about diagnosis and prognosis, including any uncertainties, as well as explicitly checking patients' comprehension of this information. There is also a need to make patients aware of the possibility that their pain may have more than one cause; for instance, arthritis may be present alongside other diagnoses such as subacromial shoulder pain. This can avoid any patient confusion in receiving more than one diagnostic label. Linked to this is the importance of fostering better cross-disciplinary communication regarding the information patients are given about their diagnosis and prognosis, to minimise the chance of patients receiving what they perceive to be conflicting information.

With regard to decision-making about treatment and referral options, training for clinicians should address the nuances around shared decision-making for shoulder pain. Whilst clinicians were aware of the importance of shared decision-making, this appeared to be dichotomised in terms of patients either wanting to be offered a choice of treatments such as physiotherapy or corticosteroid injection, or preferring the decision to be clinician-led. Clinicians should therefore be aware that even when the patient displays an apparent lack of willingness to engage with shared decision-making focused on patient choice, they may still wish to engage in a different form of shared decision-making. Clinicians should therefore be encouraged to discuss all of the available options with patients and outline a clear treatment pathway. Additionally, patients' worries about their shoulder pain being related to serious underlying pathologies such as cancer were not identified by clinicians, perhaps due to patients' reluctance to voice these concerns to their clinician. It is therefore important that clinicians are made aware of the need to always ask patients if they have any specific concerns about their shoulder pain, in line with the ICE consultation model (ideas, concerns, expectations) [25]. They should also look to provide cognitive reassurance to patients, in the form of reassurance about the absence of any indicators of serious pathology, even if these concerns are not explicitly raised by the patient.

The findings presented here will directly inform the design of a clinical intervention as part of the PANDA-S programme, that will be tested in a future RCT. The intervention will address the issues raised in these findings, with the intention of helping clinicians to identify and address the needs, concerns and priorities of patients with shoulder pain, and to improve communication about treatment and referral decisions. Through this we will aim to optimise the management of shoulder pain within primary care consultations, with the goal of improving patient outcomes.

List Of Abbreviations

GP

General Practitioner/ General Practice

RCT

Randomised Controlled Trial

NHS

National Health Service

PANDA-S

Prognostic and Diagnostic Assessment of the Shoulder

Declarations

Ethics approval and consent to participate

The study received ethical approval from the NRES Committee Yorkshire & The Humber - Sheffield, 16/10/2018, ref: 18/YH/0346. All participants provided written informed consent prior to participating in the study. All methods reported were performed in accordance with the relevant guidelines and regulations as outlined in the Declaration of Helsinki (2013).

Consent for publication

Written consent for publication was gained from all participants as part of consenting to participate in the study.

Availability of data and materials

In line with the Standard Operating Procedures in place at Keele School of Medicine, where this study was conducted, data are archived at a dedicated location within the Keele University's network. A request to access archived data can be made by completion of a Data Transfer Request form, which can be accessed by contacting: Primary Care Centre Versus Arthritis, School of Medicine, Keele University, Staffordshire, ST5 5BG, UK; Tel: +44 (0) 1782 733905.

Competing interests

The other authors declare that they have no competing interests.

Funding

This study is funded by the National Institute for Health Research (NIHR) Programme Grants for Applied Research programme in collaboration with Versus Arthritis (RP-PG-0615-20002). The views expressed in this publication are those of the authors and not necessarily those of the NHS, the NIHR or the Department of Health and Social Care.

Authors' contributions

BS, CB, GWJ, DvW, HM and TP conceptualised and developed the idea for the study. BS collected all of the data. BS, CB and GWJ contributed to analysis of the data. BS wrote the first draft of the manuscript. BS, CB, GWJ, DvW, TP, HM and RC critically edited the manuscript, and read and approved the final manuscript.

Acknowledgements

The authors would like to thank the study participants. We would also like to thank the wider PANDA-S study team, and members of the study's PPIE group and Clinical Advisory Group for their valuable contributions to the study.

References

1. Paloneva J, Koskela S, Kautiainen H, et al. Consumption of medical resources and outcome of shoulder disorders in primary health care consulters. *BMC Musculoskelet Disord.* 2013;14:1–7.
2. Virta L, Joranger P, Brox JI, et al. Costs of shoulder pain and resource use in primary health care: a cost-of-illness study in Sweden. *BMC Musculoskelet Disord.* 2021;13:17.
3. Jordan KP, Kadam UT, Hayward R, et al. Annual consultation prevalence of regional musculoskeletal problems in primary care: an observational study. *BMC Musculoskelet Disord.* 2010;11:144.
4. Chester R, Jerosch-Herold C, Lewis J, et al. Psychological factors are associated with the outcome of physiotherapy for people with shoulder pain: a multicentre longitudinal cohort study. *Br J Sports Med.* 2018;52:269–75.
5. Laslett M, Steele M, Hing W, et al. Shoulder pain patients in primary care—part 1: Clinical outcomes over 12 months following standardized diagnostic workup, corticosteroid injections, and community-based care. *J Rehabil Med.* 2014;46:898–907.
6. Maxwell C, Robinson K, McCreesh K. Understanding Shoulder Pain: A Qualitative Evidence Synthesis Exploring the Patient Experience. *Physical Therapy.* 2021;101:1–15.
7. Page MJ, O'Connor DA, Malek, M. Patients' experience of shoulder disorders: a systematic review of qualitative studies for the OMERACT Shoulder Core Domain Set. *Rheumatology.* 2019;58:1410–1421.
8. Page MJ, Green S, McBain B. Manual therapy and exercise for rotator cuff disease. *Cochrane Database Syst Rev.* 2016;6:CD012224.
9. Steuri R, Sattelmayer M, Elsig S, et al. Effectiveness of conservative interventions including exercise, manual therapy and medical management in adults with shoulder impingement: a systematic review and meta-analysis of RCTs. *Br J Sports Med.* 2017;51:1340–7.
10. Wang W, Shi M, Zhou C, et al. Effectiveness of corticosteroid injections in adhesive capsulitis of shoulder: a meta-analysis. *Medicine.* 2017;96:e7529.
11. Dunn WR, Kuhn JE, Sanders R, et al. Symptoms of pain do not correlate with rotator cuff tear severity: a cross-sectional study of 393 patients with a symptomatic atraumatic full-thickness rotator cuff tear. *J Bone Joint Surg Am.* 2014;96:793–800.
12. Ottenheijm RPG, Hesselmans NJJM, Kemper A, et al. GPs' perspectives on the diagnostic work-up in patients with shoulder pain: a qualitative study. *J Eval Clin Pract.* 2014;20:239–45.
13. Saunders B, Bartlam B, Foster NE, et al. General Practitioners' and patients' perceptions towards stratified care: a theory informed investigation. *BMC Fam Pract.* 2016;17:125.
14. Wynne-Jones G, Myers H, Hall A, et al. Predicting pain and function outcomes in people consulting with shoulder pain: the PANDA-S clinical cohort and qualitative study protocol. *BMJ Open.* 2021;11(9).
15. Glaser B, Strauss A. *Discovery of Grounded Theory: Strategies for Qualitative Inquiry.* Aldine, Chicago; 1967.

16. Sandelowski, M. Theoretical saturation. In: In: Given, L.M. (Ed.), The SAGE Encyclopaedia of Qualitative Research Methods vol. 2. Sage. Thousand Oaks. 2008, pp. 875–876.
17. Pincus T, Holt N, Vogel S et al. Cognitive and affective reassurance and patient outcomes in primary care: a systematic review. *Pain*. 2013;154(11):2407–16.
18. Cuff A, Littlewood C. Subacromial impingement syndrome – What does this mean to and for the patient? A qualitative study. *Musclo Sci and Pract*. 2018;33:24–28.
19. Artus M, van der Windt DA, Afolabi EK, et al. Management of shoulder pain by UK general practitioners (GPs): a national survey. *BMJ Open*. 2017;7:e015711.
20. Jones S, Hanchard N, Hamilton S, et al. A qualitative study of patients' perceptions and priorities when living with primary frozen shoulder. *BMJ Open*. 2013;3(9).
21. Cheung L, Soundy A. The Impact of Reassurance on Musculoskeletal (MSK) Pain: A Qualitative Review. *Behav. Sci.* 2021;11:150.
22. Donaghy E, Atherton H, Hammersley V et al. Acceptability, benefits, and challenges of video consulting: a qualitative study in primary care. *Br J Gen Pract*. 2019;69: 686.
23. Murphy M, Scott LJ, Salisbury C et al. Implementation of remote consulting in UK primary care following the COVID-19 pandemic: a mixed-methods longitudinal study. *Br J Gen Pract*. 2021;71: 704.
24. NICE (National Institute for Health and Care Excellence) Guidance.
<https://cks.nice.org.uk/topics/shoulder-pain/>. Accessed April 2022.
25. Silverman JD, Kurtz SM, Draper J. Skills for communicating with patients; Radcliffe Publishing: Oxford/San Francisco; 2016.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SupplementaryfileApatienttopicguide.docx](#)
- [SupplementaryfileBcliniciantopicguideGPs.docx](#)
- [SupplementaryfileCcliniciantopicguidephysio.docx](#)