Facility and Resident Characteristics Associated With Variation in Nursing Home Transfers: Evidence From the OPTIMISTIC Demonstration Project

Justin Blackburn (✉ jblackb@iu.edu)

Indiana University Richard M. Fairbanks School of Public Health

Casey P. Balio

Indiana University Richard M. Fairbanks School of Public Health

Jennifer L. Carnahan

Indiana University School of Medicine

Nicole R. Fowler

Indiana University School of Medicine

Susan E. Hickman

Indiana University School of Nursing

Greg A. Sachs

Indiana University School of Medicine

Wanzhu Tu

Indiana University School of Medicine

Kathleen T. Unroe

Indiana University School of Medicine

Research Article

Keywords: nursing facility, resident characteristics, Medicare, long-term care, avoidable hospitalizations

Posted Date: February 10th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-156211/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background

Centers for Medicare and Medicaid Services (CMS) funded demonstration project to evaluate financial incentives for nursing facilities providing care for 6 clinical conditions to reduce potentially avoidable hospitalizations (PAHs). The Optimizing Patient Transfers, Impacting Medical Quality, and Improving Symptoms: Transforming Institutional Care (OPTIMISTIC) site tested payment incentives alone and in combination with the successful nurse-led OPTIMISTIC clinical model. Our objective was to identify facility and resident characteristics associated with transfers, including financial incentives with or without the clinical model.

Methods

This was a longitudinal analysis from April 2017 to June 2018 of transfers among nursing home residents in 40 nursing facilities, 17 had the full clinical + payment model (1,726 residents) and 23 had payment only model (2,142 residents). Multilevel logit models estimated the likelihood of all-cause transfers and PAHs (based on CMS claims data and ICD-codes) associated with facility and resident characteristics using CMS claims data, the Minimum Data Set, and Nursing Home Compare.

Results

The clinical + payment model was associated with 4.1 percentage point (pp) lower risk of all-cause transfers (95% confidence interval [CI] -6.2 to 2.1). Characteristics associated with lower PAH risk included residents aged 95+ years (-2.4 pps; 95% CI -3.8 to -1.1), dual eligible (-2.5 pps; 95% CI -3.3 to -1.7), advanced and moderate cognitive impairment (-3.3 pps; 95% CI -4.4 to -2.1; -1.2 pps; 95% CI -2.2 to -0.2). CHESS score above most stable (CHESS score 4) increased the risk of PAH by 7.3 pps (95% CI 1.5 to 13.1).

Conclusion

Multiple resident and facility characteristics are associated with transfers. Facilities with the clinical + payment model demonstrated lower risk of all-cause transfers compared to those with payment only, but not for PAHs.

Introduction

Long-term nursing home residents have an increased risk of transfers to the hospital including emergency department (ED) visits, observation stays, and hospital admissions.¹⁻³ Adverse consequences of hospitalizations have been well-documented among older adults, including cognitive and functional decline, risk of iatrogenic disease, and added cost—over \$14 billion dollars in 2011.^{1,4-6} Furthermore, up to 60% of hospitalizations are considered potentially avoidable hospitalizations (PAHs), meaning transfer could have been prevented through early detection or improved management of the condition within the

nursing home.⁷⁻⁹ Although different methodologies are used to classify the avoidability of transfer, hospital discharge-based algorithms have determined congestive heart failure, pneumonia, urinary tract infections, sepsis, skin infections, and dehydration have a high probability of being PAHs.^{7,8} Several factors are associated with PAHs, including resident comorbidities (e.g. renal disease, diabetes), resident race (i.e. Black), lower facility staffing levels, and lower reimbursement rates.^{8,10,11} A greater understanding of resident and facility characteristics is needed to guide efforts aimed at reducing resident transfers.

The Centers for Medicare and Medicaid Services (CMS) Innovations Center launched a clinical demonstration project in 2012 to test models to reduce PAHs.¹² Seven project sites implemented unique clinical models and the Indiana site was called Optimizing Patient Transfers, Impacting Medical Quality, and Improving Symptoms: Transforming Institutional Care (OPTIMISTIC). The OPTIMISTIC clinical model deployed registered nurses (RNs) and nurse practitioners (NPs) in participating facilities to provide targeted monitoring and assessment of residents and to provide staff training. A robust, multi-site evaluation of the overall CMS initiative found that all-cause transfers and PAHs were reduced without increasing mortality.¹³⁻¹⁵ Among participating facilities in Indiana from 2014 to 2016, OPTIMISTIC reduced the annual probability of all-cause hospitalizations and PAHs by 5.1 and 3.9 percentage points, respectively.¹⁶

A second phase began in October 2016 among six of the original CMS-funded sites, including OPTIMISTIC. Special Medicare billing codes (Healthcare Common Procedure Coding System [HCPCS] codes G9679–G9684) were introduced enabling additional reimbursement for acute care provided onsite to facilities and medical providers (approximately \$218 per day).¹⁷ Facilities could bill for six procedure codes corresponding to pneumonia, COPD/asthma, dehydration, congestive heart failure, skin infections, and urinary tract infections. Codes were used if residents experienced a change in condition, met condition-specific clinical criteria, and were certified by a medical provider for in-facility care up to 7 days. As with the 19 facilities originally recruited and completed Phase 1, specific criteria were required of the 23 facilities recruited to join Phase 2.¹⁷ All facilities demonstrated tools, processes, and services were in place to provide onsite care. Facilities that participated in Phase 1 of OPTIMISTIC received billing codes in addition to the established clinical model; facilities joining in Phase 2 had access to the billing codes but did not implement the clinical model.^{17,18}

Results from OPTIMISTIC Phase 1 indicated facility-level variation in the observed reductions of hospitalizations.¹⁹ However, additional analyses are needed to elucidate the source of this variation, including facility and resident characteristics. Furthermore, characteristics associated with the risk of transfers within the context of the financial incentives introduced in Phase 2 of the OPTIMISTIC Project may offer insights into successful implementation of new care and payment models.

Methods

Sample and Setting

Although Phase 2 officially began in October 2016, to account for variation in the implementation, this study focuses on eligible residents in participating nursing facilities between April 2017 and June 2018. Eligible residents were defined by CMS as long-stay residents (> 100 days in facility) enrolled in Medicare Parts A and B fee-for-service (i.e. without Medicare Advantage). Residents were censored if they died or became ineligible because of discharge from the facility, enrolling in Medicare Advantage plan, or being admitted to hospice. Residents were analyzed based on eligibility periods designated by three-month quarters according to calendar year. Resident's observations were included only for the quarters in which they were eligible for at least one day during the quarter. Among 40 total facilities, 23 were in the payment only group and 17 in the clinical + payment group.

Data

Demographic and chronic condition information were derived from the Master Beneficiary Summary File (MBSF). Medicare claims data were used to identify all transfers, including the carrier line file, inpatient, and outpatient files. Data were supplemented by the Minimum Data Set (MDS) assessments for measures of health, functional, and cognitive status. Nursing homes are required to complete an MDS assessment on all residents at the time of admission and quarterly thereafter, or after a significant change in status. Quarter in which death occurred was identified from the MBSF or MDS. Nursing Home Compare data, including Provider Files for facility staffing and quality measures, and the Point of Service (POS) Files for facility characteristics including rural status, for-profit status, and whether the facility was part of a chain.

Outcomes

The two primary outcomes of focus for our analyses were all-cause transfers, including hospitalization, ED, or observation, and PAHs. Both are primary targets for reduction by the OPTIMISTIC clinical model and the financial incentive payments. However, to put our outcomes into context, we considered a variety of transfers consistent with the national, multi-site evaluation of the demonstration project, including all-cause ED visits, all-cause hospitalizations, PAHs, potentially avoidable EDs, hospitalization for any of the six conditions corresponding to financial incentive billing codes..¹⁹ Only those treated and released from the ED were considered a visit; admissions from the ED were considered hospitalizations. Outcomes were not mutually-exclusive, such that PAHs were nested within all-cause hospitalizations. Outpatient and carrier files were used to identify ED visits with revenue center codes 450–459, or 981 that did not result in admission. Hospitalizations were any consecutive stays as in inpatient, regardless of whether transferred to another hospital during the stay. Observation stays were identified as claims with a revenue center code 0762 that lacked a corresponding overnight stay.²⁰ The identification of potentially avoidable for PAHs and ED visits was done using International Classification of Disease version 9 or 10 (ICD-9 / ICD-10) codes consistent with the larger evaluation of the demonstration project.¹⁹ Due to the sparse data representing repeated transfers among residents within a quarter, counts of transfers were

aggregated to binary indicators for whether at least one of each type of transfer occurred within a given quarter.

Measures- resident & facility characteristics

Potential resident and facility characteristics were included based on a priori hypotheses and assessment of the bivariate associations. Resident demographic and health status information included chronic conditions identified from the annually-updated MBSF Chronic Conditions Warehouse (CCW) segment including Alzheimer's/Dementia, chronic kidney disease, COPD, congestive heart failure, diabetes, ischemic heart disease, depression, osteoporosis, stroke/transient ischemic attack, and hypertension. Other health status measures were derived quarterly from the MDS including cognitive functioning scale (CFS) (range cognitively intact [1]-severe impairment [4]), Changes in Health, End-stage disease and Symptoms and Signs (CHESS) score (range most stable [0]-least [5]), and activities of daily living (ADL) (range complete independence [0]-complete dependence on staff [28]).²¹⁻²³

Facility characteristics included an indicator for which OPTIMISTIC intervention the facility received: payment only model versus the full clinical + payment model. Additionally, from the Nursing Home Compare Provider Files, quarterly categorical indicators for staffing (total licensed hours per resident per day), bed size, and overall star rating (1–5) for the last month of the quarter. Binary indicators for rural setting, for-profit/non-profit status, and chain-owned were obtained from the CMS POS files from baseline (December 2017).

Analysis

We used mixed-effects logistic regression to account for repeated resident observations over time and nesting within facilities. This multi-level approach has been used previously when resident nesting within facilities occurs.²⁴⁻²⁶ Our modeling strategy estimated residents' outcome risk over time for each characteristic, while accounting for all others. Furthermore, we estimate the amount of variance in the outcome attributed to the resident and facility, and estimate the association of resident and facility characteristics using a similar modeling strategy as Herrin et al.²⁷ The intraclass correlation coefficient for null multi-level model was compared to the full model (with covariates) to quantify the variance explained at each level and by the addition of resident and facility characteristics. Marginal effects were estimated and reported with 95% confidence intervals, which can be interpreted as the absolute risk different in the outcome holding all other characteristics at the mean, expressed as a percentage point change. Interactions were considered for facilities in the payment only group or the clinical + payment group to determine whether the OPTIMISTIC model or financial incentives only modified the risk of transfer.

Results

During the study period, 3,868 unique residents in OPTIMISTIC facilities contributed 12,787 resident-quarter observations. At least one transfer was observed during 2,290 (17.9%) resident-quarters. Among

them, we observed 1,439 hospitalizations (11.2% of resident-quarters), 1,204 all-cause ED visits (9.4% of resident-quarters), 487 PAHs 487 (3.8%of resident-quarters), 308 hospitalizations for any of the six conditions (2.4% of resident-quarters) and 256 potentially avoidable ED visits (2.0% of resident-quarters).

Figure 1 displays the percentage of residents with at least one transfer by each type over time, stratified by intervention group. All-cause transfers increased over time among residents in payment only facilities, ranging from 16.8% in the first quarter to 19.6% one year later, while clinical + payment facilities remained stable, and consistently lower, at approximately 17% of residents. Similar differences were observed for all-cause ED visits and all-cause hospitalizations. The percent of residents with PAHs ranged between 3% and 4% for the duration of the study period, with no apparent difference by intervention group.

Residents were mostly female (69.2%), white (84.8%), dual eligible (75.1%), with some form of Alzheimer's Disease/Dementia (81.3%) and depression (74.5%, Table 1). Fifty-seven percent of the resident-quarter observations were in payment only facilities and 43% in payment + clinical facilities. Statistically significant bivariate associations were observed between risk of all-cause transfers and race, ADL score, hypertension, facility intervention group, and for-profit status; however, none of these characteristics were associated with PAHs at conventional significance levels.

Table 1

Overall cohort description (N = 12,787 resident-quarters) as well as bivariate statistics for all-cause transfer (n = 2,290).

Individual-level Characteristics	Overall (N = 12,787)	All-cause Transfer (n = 2,290)		Potentially Avoidable Hospitalization (n = 487)	
	%	%	P	%	P
Aged $\leq$ 64 years	9.9	15.9	< 0.001	15.4	< 0.001
Aged 65–74 years	16.6	19.1		20.3	
Aged 75–84 years	28.7	29.7		33.3	
Aged 85–94 years	35.8	29.5		26.9	
Aged $\geq$ 95 years	9.0	5.7		4.1	
Male	30.8	37.0	< 0.001	37.4	0.001
Female	69.2	63.0		62.6	
White	84.8	82.6	0.001	81.1	0.067
Black	13.7	16.1		17.3	
Other	1.5	1.3		1.6	
Medicare Eligible Only	24.9	29.6	< 0.001	33.5	< 0.001
Dual Eligible	75.1	70.4		66.5	
Cognitive Functioning Scale 1	30.9	39.0	< 0.001	40.7	< 0.001
Cognitive Functioning Scale 2	22.9	24.5		25.5	
Cognitive Functioning Scale 3	38.9	30.7		30.8	
Cognitive Functioning Scale 4	7.3	5.9		3.1	
CHESS Score 0	35.2	28.8	< 0.001	23.6	< 0.001

CHESS Score 1	34.7	34.3		34.5	
CHESS Score 2	22.7	26.1		29.9	
CHESS Score 3	6.4	9.1		10.7	
CHESS Score 4	1.0	1.8		2.3	
ADLs 1st Quartile (≤ 17)	29.6	27.6	0.010	28.1	0.479
ADLs 2nd Quartile (18–19)	25.1	26.4		27.5	
ADLs 3rd Quartile (20–21)	32.7	31.9		30.8	
ADLs 4th Quartile (22–28)	12.6	14.1		13.6	
Acute Myocardial Infarction	9.2	12.6	< 0.001	16.2	< 0.001
Alzheimer's Disease / Dementia	81.3	75.2	< 0.001	77.6	0.035
Chronic Kidney Disease	57.9	68.0	< 0.001	71.5	< 0.001
Chronic Obstructive Pulmonary Disease	42.1	52.0	< 0.001	57.3	< 0.001
Congestive Heart Failure	56.7	63.3	< 0.001	68.6	< 0.001
Diabetes	50.1	56.2	< 0.001	61.4	< 0.001
Ischemic Heart Disease	62.9	66.6	< 0.001	73.1	< 0.001
Depression	74.5	76.1	0.050	78.4	0.042
Osteoporosis	38.0	32.7	< 0.001	30.6	0.001
Stroke / Transient Ischemic Attack	36.3	38.6	0.012	41.1	0.026
Hypertension	77.1	88.2	0.046	87.5	0.176
Never Married	13.1	14.6	< 0.001	14.8	0.014
Married	20.8	21.5		23.8	

	Widowed	44.5	40.5		37.4	
	Not Married	21.6	23.4		24.0	
Facility-level Characteristics	Payment Only Facility	56.6	59.3	0.004	58.7	0.325
	Payment + Clinical Model Facility	43.4	40.7		41.3	
	Staffing 1st Quartile ($\leq$ 3.31 NHPPD)	23.8	25.8	< 0.001	26.7	0.004
	Staffing 2nd Quartile (3.32–3.55 NHPPD)	24.3	24.1		23.4	
	Staffing 3rd Quartile (3.56–4.15 NHPPD)	24.9	27.2		29.2	
	Staffing 4th Quartile ($\geq$ 4.15 NHPPD)	27.1	22.9		20.7	
	Beds 1st Quartile ($\leq$ 115 beds)	23.9	26.4	< 0.001	26.1	0.003
	Beds 2nd Quartile (116–147 beds)	27.5	28.8		27.5	
	Beds 3rd Quartile (148–169 beds)	23.8	23.9		28.1	
	Beds 4th Quartile ($\geq$ 170 beds)	24.8	20.9		18.3	
	Overall Star Rating 1	4.5	8.2	< 0.001	7.0	0.022
	Overall Star Rating 2	10.4	12.2		12.1	
	Overall Star Rating 3	20.7	20.0		17.3	
	Overall Star Rating 4	28.5	26.9		28.1	
	Overall Star Rating 5	35.9	32.7		35.5	
	Rural	27.2	26.5	0.367	30.4	0.109
	For-profit Facility	15.7	17.6	0.004	16.0	0.820
	Multi-facility Ownership	76.3	78.7	0.002	80.5	0.026
	2017 Quarter 2	21.0	20.0	0.269	18.3	0.672
	2017 Quarter 3	20.4	19.4		21.4	
	2017 Quarter 4	20.3	21.3		20.5	

2018 Quarter 1	19.4	20.3	19.7
2018 Quarter 2	19.0	19.0	20.1
Abbreviations: CHESS = Changes in Health, End-stage disease and Symptoms and Signs; ADLs = Activities of Daily Living; NHPPD = nursing hours per patient day Note: Results are provided for the resident-quarter level. Residents were included in the analyses if they were eligible for at least one day of the quarter. Results provide descriptions of the population by the percent of resident-quarter observations for given characteristics, and by residents experiencing at least one all-cause transfer or PAH during a given quarter.			

The absolute risk differences of characteristics associated with all-cause transfers and PAHs, estimated using multilevel logit regression, are shown in Table 2. Notable characteristics associated with PAH risk include residents aged 95 years or older (-2.4 percentage points [pps]; 95% CI -3.8 to -1.1), history of hypertension (-3.7 pps; 95% CI -5.2 to -2.1), dual eligible (-2.5 pps; 95% CI -3.3 to -1.7), advanced cognitive impairment (CFS category 4) (-3.3 pps; 95% CI -4.4 to -2.1) and moderate impairment (CFS category 3) (-1.2 pps; 95% CI -2.2 to -0.2). Higher CHESS scores relative to the most stable were associated with increased risk of PAH including CHESS score of 4 (7.3 pps; 95% CI 1.5 to 13.1), CHESS score 3 (4.3 pps; 95% CI 2.0 to 6.6), and CHESS score 2 (2.4 pps; 95% CI 1.2 to 3.5). Each characteristic was also associated with all-cause transfer risk.

Table 2

Estimated risk difference for characteristics associated with resident transfers (N = 12,787 resident-quarters).

		All-cause Transfer (n = 2,290)		Potentially Avoidable Hospitalization (n = 487)	
Individual-level Characteristics	Aged ≤ 64 years	8.6	(5.0, 12.2)	1.5	(-0.2, 3.3)
	Aged 65–74 years	Reference			
	Aged 75–84 years	-1.0	(-3.5, 1.5)	0.1	(-1.1, 1.3)
	Aged 85–94 years	-4.5	(-7.1, -2.0)	-1.4	(-2.5, -0.2)
	Aged ≥ 95 years	-7.7	(-10.9, -4.4)	-2.4	(-3.8, -1.1)
	Female	-2.6	(-4.4, -0.8)	-0.1	(-0.9, -0.7)
	White	Reference			
	Black	3.4	(0.9, 6.0)	1.3	(0.1, 2.6)
	Other	-4.6	(-9.9, 0.8)	0.3	(-2.6, 3.3)
	Dual Eligible	-6.7	(-8.4, -5.1)	-2.5	(-3.3, -1.7)
	Cognitive Functioning Scale 1	Reference			
	Cognitive Functioning Scale 2	-0.8	(-2.9, 1.3)	-0.3	(-1.4, 0.8)
	Cognitive Functioning Scale 3	-4.5	(-6.6, -2.5)	-1.2	(-2.2, -0.2)
	Cognitive Functioning Scale 4	-6.9	(-10.1, -3.8)	-3.3	(-4.4, -2.1)
	CHESS Score 0	Reference			
	CHESS Score 1	1.9	(0.1, 3.6)	1.1	(0.3, 1.9)
	CHESS Score 2	4.7	(2.4, 7.0)	2.4	(1.2, 3.5)
	CHESS Score 3	9.3	(5.4, 13.1)	4.3	(2.0, 6.6)

		All-cause Transfer (n = 2,290)		Potentially Avoidable Hospitalization (n = 487)	
	CHESS Score 4	13.8	(5.6, 22.0)	7.3	(1.5, 13.1)
	ADLs 1st Quartile (≤ 17)	Reference			
	ADLs 2nd Quartile (18–19)	3.8	(1.9, 5.8)	0.8	(-0.2, 1.8)
	ADLs 3rd Quartile (20–21)	2.6	(0.6, 4.6)	-0.4	(-1.3, 0.6)
	ADLs 4th Quartile (22–28)	4.1	(1.1, 7.0)	-0.1	(-1.4, 1.3)
	Acute Myocardial Infarction	3.2	(0.6, 5.8)	1.3	(0.2, 2.4)
	Alzheimer's Disease / Dementia	-3.0	(-5.2, -0.7)	0.4	(-0.1, 1.5)
	Chronic Kidney Disease	6.2	(4.3, 8.0)	1.6	(0.6, 2.5)
	Chronic Obstructive Pulmonary Disease	4.6	(2.8, 6.3)	1.0	(0.2, 1.9)
	Ischemic Heart Disease	2.2	(0.2, 4.1)	1.3	(0.3, 2.3)
	Hypertension	-6.9	(-10.1, -3.7)	-3.7	(-5.2, -2.1)
Facility-level Characteristics	Payment Only Facility	Reference			
	Payment + Clinical Model Facility	-4.1	(-6.2, -2.1)	-0.5	(-1.6, 0.4)
	Beds 1st Quartile (≤ 115 beds)	Reference			
	Beds 2nd Quartile (116–147 beds)	-3.4	(-5.6, -1.1)	-0.9	(-1.9, 0.2)
	Beds 3rd Quartile (148–169 beds)	-0.2	(-2.7, 2.2)	0.7	(-0.5, 0.2)
	Beds 4th Quartile (≥ 170 beds)	-3.4	(-6.0, -0.1)	-0.9	(-2.1, 0.3)
	Overall Star Rating 1	13.0	(8.3, 17.6)	2.4	(-0.2, 4.9)
	Overall Star Rating 2	4.6	(1.7, 7.4)	0.6	(-0.9, 2.1)

	All-cause Transfer (n = 2,290)		Potentially Avoidable Hospitalization (n = 487)	
Overall Star Rating 3	1.7	(-0.5, 3.9)	-0.7	(-1.7, 0.3)
Overall Star Rating 4	0.4	(-1.5, 2.3)	-0.2	(-1.2, 0.7)
Overall Star Rating 5	Reference			
2017 Quarter 2	Reference			
2017 Quarter 3	1.3	(-0.7, 3.2)	1.2	(0.2, 2.1)
2017 Quarter 4	2.5	(0.5, 4.5)	0.9	(-0.1, 1.9)
2018 Quarter 1	3.2	(1.1, 5.2)	1.1	(0.1, 2.1)
2018 Quarter 2	1.9	(-0.3, 4.0)	1.2	(0.1, 2.3)
Abbreviations: ADLs = Activities of Daily Living; NHPPD = nursing hours per patient day Model also controls for residents with congestive heart failure, diabetes, ischemic heart disease, depression, osteoporosis, stroke / transient ischemic attack and marital status, as well as facility characteristics staffing level, rural status, for-profit status, and multi-facility ownership.				

Notable resident characteristics associated with all-cause transfers but not PAHs included ADL scores above 17 (most independent) including ADL 18–19 (3.8 pps; 95% CI 1.9 to 5.8), ADL 20–21 (2.6 pps; 95% CI 0.6 to 4.6), ADL 22–28 (4.1 pps; 95% CI 1.1 to 7.0) and residents with Alzheimer’s Disease/dementia (-3.0 pps; 95% CI -5.2 to -0.7). Notable facility characteristics associated with all-cause transfers, but not PAHs, included the presence of the clinical + payment model (-4.1 pps; 95% CI -6.2 to -2.1), 116 to 147 beds in the facility (-3.4 pps; 95% CI -5.6 to -1.1), and overall star rating of 1 (13.0 pps; 95% CI 8.3 to 17.6) and overall star rating of 2 (4.6 pps; 95% CI 1.7 to 7.4).

From the intraclass correlation coefficient estimates, we observed 20% of the variance in all-cause transfers was explained by residents within the same facilities (i.e. at the facility level) and 32% by residents over time (i.e. at the resident-level). Similar levels of nesting were observed for PAHs, with 20% of variance explained at the facility-level and 35% at the resident-level. Introducing covariates reduced the unexplained variance at the facility-level by approximately 10 pps and at the resident-level by approximately 6 pps (7 pps and 13 pps reductions among PAHs, respectively). Resident and facility characteristics demonstrated heterogeneity among resident-quarters by clinical + payment or payment only facilities including race, CHESS Scale, and bed size. Although statistical evidence of interaction for the presence of the clinical + payment model was not definitive, stratified results are presented in the Appendix.

Discussion

In facilities participating in a demonstration project to reduce PAHs, we identified age, dual eligibility, cognitive function, CHESS, and history of certain comorbidities were associated with the risk of all-cause transfer, including PAHs. These characteristics generally reflect limitations to residents' functional status and a history of debilitating comorbidities. However, some characteristics were inconsistently associated with the outcomes. Resident ADL score, history of Alzheimer's Disease/dementia, CMS Star Rating, facility bed count, and participation in the clinical + payment model was associated with the risk of all-cause transfer, but not with the risk of PAHs. Although variation in the risk of transfer was explained at both the resident- and facility-level, resident characteristics explained a greater percentage. These results add to the existing knowledge regarding how to reduce transfers, specifically PAHs, among nursing home residents. Furthermore, risk stratification of nursing home residents by specific characteristics may be necessary for consideration in planning and evaluation of future efforts to reduce PAHs.

The oldest residents (≥ 95 years), the most cognitively impaired (CFS 4), and dual eligible residents had the lowest adjusted risk of both all-cause transfers and PAHs. The lower risk of hospitalization for older residents is equivocal in other studies and may be dependent on how well functional status and comorbidities are measured and controlled for.²⁸ Lower risk in the oldest age group may be attributable to informative censoring, wherein residents in this age group remaining eligible for OPTIMISTIC are overall healthier than those who lose eligibility due to hospice enrollment. To avoid this, **some studies** exclude observations within 12 months of residents' death.¹¹ However, due to concerns about sample size reductions and the importance of including events that will enhance our understanding of PAHs, this approach was not feasible. Alternatively, resident characteristics may be confounded by completion of advance care planning/advance directives wherein transfer is inconsistent with their goals of care. Residents in clinical + payment facilities had consistent access to structured Advance Care Planning through the OPTIMISTIC nurses and all facilities had access to tailored educational materials related to palliative care and incorporating goals of care into treatment plans. This may also explain the lower risk of transfer among cognitively impaired residents, and is consistent with other studies of residents with advanced illness and limited life expectancy.²⁹ Although residents with advanced age, cognitive impairment, and/or comorbidities may have clinical features consistent with hospital-level care, decisions to transfer are complex and mitigated by multiple factors that are not captured by the MDS.^{30–33}

In contrast to lower risk of transfer for the oldest residents, residents in the youngest age category—those under 64 years old and thus qualifying for Medicare based on disability—had an increased risk of all-cause transfers, but not PAHs. This finding is consistent with another study where younger age also had functional impairment.³⁴ Furthermore, the lack of an association with PAHs may indicate that hospitalizations among younger residents tend not to be considered potentially avoidable. Moreover, risk of PAHs may not be uniform across all age groups and warrants further investigation.

We observed an increased risk of transfer with higher CHESS scores, which may seem counter-intuitive. One potential explanation is that the CHESS Scale reflects recent health instability, some of which may

appear treatable.³⁵ Although CHESS Scale is a predictor of mortality, it may also be sensitive to acute resident changes resulting in transfers, including acute mental status change, dehydration, and pressure ulcers.²¹ Furthermore, there may be multicollinearity with CHESS Scale and other resident characteristics included in our model, including cognitive impairment. However, variance inflation factors from our models were less than 3 for all variables included.

Facility characteristics including bed size, CMS quality ranking, and presence of the clinical + payment model was associated with reduced all-cause transfers, the associations were not statistically significant with PAHs. Nationally, the overall risk of hospital transfers among nursing home residents has decreased since 2011, coinciding with multiple initiatives to reduce hospitalizations.²⁹ Previous analyses from the OPTIMISTIC project, which has similar components as the INTERACT Program, has demonstrated benefits to improving the management of acute changes in residents' condition and observed reductions in transfers.^{13,14,17,18,36,37} Although we do not observe a decrease in transfer risk during our study period, residents in OPTIMISTIC clinical + payment model facilities maintained a consistently lower risk of all-cause transfer than payment only facilities, which increased slightly over time. The OPTIMISTIC clinical model was first introduced in 2012. Thus, nearly half of the facilities represented in our analyses were focused on reducing transfers, specifically PAHs, using specially trained nurses for four years prior to the release of the payment reimbursement codes. The payment component was introduced in the fourth quarter of 2016 and thus the payment only facilities may have had a more heterogeneous experience reducing PAHs.³⁶ Furthermore, the payment only facilities lacked on-site nurses trained to detect and treat conditions to reduce PAHs, instead relevant staff were trained on the use of billing codes for the six conditions.

The absence of an association between PAHs and the clinical + payment model is notable. This may be attributable to limitations of the identification of PAHs from claims data. First, we cannot exclude the possibility that coding practices designed to maximize hospital reimbursement may affect whether a hospitalization was subsequently determined to be a PAH; it is known that administrative claims data provides only a partial picture of the actual clinical experience.³⁸ Furthermore, claims data do not provide direct insights into the primary reason for a transfer—this may be different than what was determined as the final principal diagnosis code for the hospital stay. Avoidability of transfers is difficult to assess, even by nursing staff within the facility, adding to the challenge of making a post-discharge determination using claims-based algorithms.^{17,39,40} Finally, claims diagnoses reflect the full clinical stay which may include infections or other events that were not present on admission. Nonetheless, our findings that facilities with the clinical + payment model had lower rates of transfers highlights the importance of increasing capacity of facilities to treat patients in place with a proven clinical model such as OPTIMISTIC, and to support practice changes in addition to payment reform.

Although facility characteristics explained a substantial portion of the variation in transfer risk, we observed few with statistical associations. Among them, lower CMS Star Ratings were associated with increased transfers. Notably, initial recruitment of new facilities in Phase 2 of the OPTIMISTIC

demonstration project required a minimum 3-star rating. However, facility rating fluctuates quarterly and maintenance was not a requirement, therefore we observed variation within participating facilities over time. It is possible that a decreasing Star Rating over time is indicative of decreasing overall quality of care.

This study has some limitations. First, we recognize this is a non-randomized intervention in 40 facilities within a single state. We do not have the benefit of randomization which could introduce selection bias and limit the generalizability of our results. Specifically, the selection of facilities to the clinical + payment group was based on several important characteristics, including a minimum CMS Star Rating, and were located in largely urban and suburban Central Indiana, may have resulted in bias and as compared to the payment only group. We lack a true control group and instead used individuals' risk over time to account for within-person confounding as well as differences in the intervention facility to contrast risk. The lack of control for some potentially important variables, such as the presence of newly diagnosed terminal diseases, family/resident preferences, or other relevant unmeasurable variables, could result in omitted variable bias. In such a case, the attribution of unobserved factors such as a do not resuscitate orders could bias the associations of other characteristics. We note this as an alternative explanation for the lower risk of transfer among residents with advanced cognitive impairment. However, prior work within this same demonstration project found that associations between advance care planning and potentially avoidable hospitalizations were attenuated after accounting for facility clustering and resident characteristics.³⁵

Conclusions

Overall, our study contributes to a growing body of evidence around facility and resident characteristics associated with risk of transfers. In particular, although we observe considerable variation at the facility-level, although few of the facility variables included were associated with resident transfers. However, facilities with the OPTIMISTIC clinical + payment model in place were able to demonstrate a lower risk of all-cause transfers as compared to those with payment only. We also raise questions about the ability to assess PAHs from claims data. Nonetheless, our findings provide evidence for risk stratification for facilities in reducing transfers and also highlights the potential benefit of additional staff in a proven clinical model supporting efforts to limit transfers for some potentially avoidable transfers.

Declarations

Ethics approval and consent to participate:

OPTIMISTIC was conducted as a demonstration project under contract to the Centers for Medicare and Medicaid Services (CMS) in the United States utilizing a suite of interventions, staff education, and quality improvement strategies already proven effective. It was explicitly deemed by CMS to not be a research study and did not require individual informed consent of participants. Per CMS regulations and guidance, residents of nursing facilities, or their legally authorized representatives (proxies) were

informed of the project by nursing facility leaders via letter and were provided an opportunity to opt-out of the demonstration project. The OPTIMISTIC demonstration project also underwent an independent review and was approved by the Indiana University Institutional Review Board (IRB). The IRB similarly determined that as an evaluation of a program for federal Medicare beneficiaries, informed consent was not required from individual participants and granted a waiver of informed consent. All project methods were carried out in accordance with relevant guidelines and regulations.

Consent for publication: N/A

Availability of data materials: Limited versions of the datasets used and analyzed during the current study are available from the corresponding author on reasonable request and subject to approval from the sponsor.

Competing interests: Kathleen Unroe is CEO and Founder of Probari, Inc., a program to train nurses to reduce nursing home hospital transfers. All other authors have no conflicts of interest to disclose.

Funding: This work was supported by the U.S. Department of Health and Human Services, Centers for Medicare and Medicaid Services [Funding Opportunity 1E1CMS331488] and an internal grant from Regenstrief Institute. The opinions expressed in this article are the authors' own and do not reflect the view of the U.S. Department of Health and Human Services, Centers for Medicare and Medicaid Services.

Author contribution: KTU, NRF, SEH, GS, JB developed the research question. JB, KTU, NRF, CPB conceptualized the research design. GS, JLC, KTU provided clinical input. WT provided input on statistical methods and oversight of analyses. JB, CPB performed analyses. JB, CPB wrote first draft of the manuscript. All authors reviewed and made edits to the manuscript.

Acknowledgement: N/A

References

1. Grabowski DC, Stewart KA, Broderick SM, Coots LA. Predictors of nursing home hospitalization: a review of the literature. *Med Care Res Rev.* 2008;65(1):3-39.
2. Graverholt B, Forsetlund L, Jamtvedt G. Reducing hospital admissions from nursing homes: a systematic review. *BMC Health Serv Res.* 2014;14:36.
3. Graverholt B, Riise T, Jamtvedt G, Ranhoff AH, Kruger K, Nortvedt MW. Acute hospital admissions among nursing home residents: a population-based observational study. *BMC Health Serv Res.* 2011;11:126.
4. Creditor MC. Hazards of hospitalization of the elderly. *Ann Intern Med.* 1993;118(3):219-223.
5. Ehlenbach WJ, Hough CL, Crane PK, et al. Association between acute care and critical illness hospitalization and cognitive function in older adults. *JAMA.* 2010;303(8):763-770.

6. Services USDoHaH. Medicare Nursing Home Resident Hospitalization Rates Merit Additional Monitoring. In: General Ootl, ed. <https://oig.hhs.gov/oei/reports/oei-06-11-00040.asp>2013.
7. Ouslander JG, Lamb G, Perloe M, et al. Potentially avoidable hospitalizations of nursing home residents: frequency, causes, and costs: [see editorial comments by Drs. Jean F. Wyman and William R. Hazzard, pp 760-761]. *J Am Geriatr Soc*. 2010;58(4):627-635.
8. Walsh EG, Wiener JM, Haber S, Bragg A, Freiman M, Ouslander JG. Potentially avoidable hospitalizations of dually eligible Medicare and Medicaid beneficiaries from nursing facility and Home- and Community-Based Services waiver programs. *J Am Geriatr Soc*. 2012;60(5):821-829.
9. Ouslander JG, Naharci I, Engstrom G, et al. Root Cause Analyses of Transfers of Skilled Nursing Facility Patients to Acute Hospitals: Lessons Learned for Reducing Unnecessary Hospitalizations. *J Am Med Dir Assoc*. 2016;17(3):256-262.
10. Spector WD, Limcangco R, Williams C, Rhodes W, Hurd D. Potentially avoidable hospitalizations for elderly long-stay residents in nursing homes. *Med Care*. 2013;51(8):673-681.
11. O'Malley AJ, Caudry DJ, Grabowski DC. Predictors of nursing home residents' time to hospitalization. *Health Serv Res*. 2011;46(1 Pt 1):82-104.
12. Unroe KT, Nazir A, Holtz LR, et al. The Optimizing Patient Transfers, Impacting Medical Quality, and Improving Symptoms: Transforming Institutional Care approach: preliminary data from the implementation of a Centers for Medicare and Medicaid Services nursing facility demonstration project. *J Am Geriatr Soc*. 2015;63(1):165-169.
13. Feng Z, Ingber MJ, Segelman M, et al. Nursing Facilities Can Reduce Avoidable Hospitalizations Without Increasing Mortality Risk For Residents. *Health Aff (Millwood)*. 2018;37(10):1640-1646.
14. Ingber MJ, Feng Z, Khatutsky G, et al. Initiative To Reduce Avoidable Hospitalizations Among Nursing Facility Residents Shows Promising Results. *Health Aff (Millwood)*. 2017;36(3):441-450.
15. Vadnais AJ, Vreeland E, Coomer NM, Feng Z, Ingber MJ. Reducing Transfers among Long-Stay Nursing Facility Residents to Acute Care Settings: Effect of the 2013-2016 Centers for Medicare and Medicaid Services Initiative. *J Am Med Dir Assoc*. 2020.
16. Ingber MJ, Feng Z, Khatutsky G, et al. *Evaluation of the Initiative to Reduce Avoidable Hospitalizations among Nursing Facility Residents: Final Report*. 2017.
17. Unroe KT, Fowler NR, Carnahan JL, et al. Improving Nursing Facility Care Through an Innovative Payment Demonstration Project: Optimizing Patient Transfers, Impacting Medical Quality, and Improving Symptoms: Transforming Institutional Care Phase 2. *J Am Geriatr Soc*. 2018;66(8):1625-1631.
18. Blackburn J, Stump TE, Carnahan JL, et al. Reducing the Risk of Hospitalization for Nursing Home Residents: Effects and Facility Variation From OPTIMISTIC. *J Am Med Dir Assoc*. 2020;21(4):545-549 e541.
19. CMS.gov. Evaluation of the Initiative to Reduce Avoidable Hospitalizations among Nursing Facility Residents—Payment Reform. Centers for Medicare and Medicaid Services.

<https://downloads.cms.gov/files/cmimi/rahnfr-phasetwo-secondannrpt.pdf>. Published 2019. Accessed April 23, 2020.

20. Sheehy AM, Shi F, Kind AJH. Identifying Observation Stays In Medicare Data: Policy Implications of a Definition. *J Hosp Med*. 2019;14(2):96-100.
21. Ogarek JA, McCreedy EM, Thomas KS, Teno JM, Gozalo PL. Minimum Data Set Changes in Health, End-Stage Disease and Symptoms and Signs Scale: A Revised Measure to Predict Mortality in Nursing Home Residents. *J Am Geriatr Soc*. 2018;66(5):976-981.
22. Thomas KS, Dosa D, Wysocki A, Mor V. The Minimum Data Set 3.0 Cognitive Function Scale. *Med Care*. 2017;55(9):e68-e72.
23. Kruse RL, Petroski GF, Mehr DR, Banaszak-Holl J, Intrator O. Activity of daily living trajectories surrounding acute hospitalization of long-stay nursing home residents. *J Am Geriatr Soc*. 2013;61(11):1909-1918.
24. Intrator O, Zinn J, Mor V. Nursing home characteristics and potentially preventable hospitalizations of long-stay residents. *J Am Geriatr Soc*. 2004;52(10):1730-1736.
25. Xing J, Mukamel DB, Temkin-Greener H. Hospitalizations of nursing home residents in the last year of life: nursing home characteristics and variation in potentially avoidable hospitalizations. *J Am Geriatr Soc*. 2013;61(11):1900-1908.
26. Hass Z, Woodhouse M, Kane R, Arling G. Modeling Community Discharge of Medicaid Nursing Home Residents: Implications for Money Follows the Person. *Health Serv Res*. 2018;53 Suppl 1:2787-2802.
27. Herrin J, St Andre J, Kenward K, et al. Community factors and hospital readmission rates. *Health Serv Res*. 2015;50(1):20-39.
28. Hoffmann F, Allers K. Age and sex differences in hospitalisation of nursing home residents: a systematic review. *BMJ Open*. 2016;6(10):e011912.
29. McCarthy EP, Ogarek JA, Loomer L, et al. Hospital Transfer Rates Among US Nursing Home Residents With Advanced Illness Before and After Initiatives to Reduce Hospitalizations. *JAMA Intern Med*. 2019.
30. Jennings LA, Zingmond D, Louie R, et al. Use of the Physician Orders for Life-Sustaining Treatment among California Nursing Home Residents. *J Gen Intern Med*. 2016;31(10):1119-1126.
31. Mukamel DB, Ladd H, Caprio T, Temkin-Greener H. Prototype End-of-Life Quality Measures Based on MDS 3 Data. *Med Care*. 2016;54(11):1024-1032.
32. Nakashima T, Young Y, Hsu WH. Are Hospital/ED Transfers Less Likely Among Nursing Home Residents With Do-Not-Hospitalize Orders? *J Am Med Dir Assoc*. 2017;18(5):438-441.
33. Trahan LM, Spiers JA, Cummings GG. Decisions to Transfer Nursing Home Residents to Emergency Departments: A Scoping Review of Contributing Factors and Staff Perspectives. *J Am Med Dir Assoc*. 2016;17(11):994-1005.
34. Brown RT, Diaz-Ramirez LG, Boscardin WJ, Lee SJ, Williams BA, Steinman MA. Association of Functional Impairment in Middle Age With Hospitalization, Nursing Home Admission, and Death.

JAMA Intern Med. 2019;179(5):668-675.

35. Heckman GA, Hirdes JP, Hebert PC, et al. Predicting Future Health Transitions Among Newly Admitted Nursing Home Residents With Heart Failure. *J Am Med Dir Assoc.* 2019;20(4):438-443.
36. Hickman SE, Unroe KT, Ersek M, et al. Systematic Advance Care Planning and Potentially Avoidable Hospitalizations of Nursing Facility Residents. *J Am Geriatr Soc.* 2019;67(8):1649-1655.
37. Kane RL, Huckfeldt P, Tappen R, et al. Effects of an intervention to reduce hospitalizations from nursing homes: a randomized implementation trial of the INTERACT Program. *JAMA Intern Med.* 2017;177(9):1257-1264.
38. Adler-Milstein J, Jha AK. No evidence found that hospitals are using new electronic health records to increase Medicare reimbursements. *Health Aff (Millwood).* 2014;33(7):1271-1277.
39. Unroe KT, Carnahan JL, Hickman SE, Sachs GA, Hass Z, Arling G. The Complexity of Determining Whether a Nursing Home Transfer Is Avoidable at Time of Transfer. *J Am Geriatr Soc.* 2018;66(5):895-901.
40. McAndrew RM, Grabowski DC, Dangi A, Young GJ. Prevalence and patterns of potentially avoidable hospitalizations in the US long-term care setting. *Int J Qual Health Care.* 2016;28(1):104-109.

Figures

Figure 1

Observed percentage of OPTIMISTIC nursing home residents experiencing at least one of each transfer type by quarter, stratified by payment only facilities and clinical + payment. Note: All-cause transfers include hospitalizations, emergency department (ED) visits, and observation stays. ED visits that ended in a hospitalization are included in all-cause transfers and hospitalizations but not in ED visits. The 6 conditions include pneumonia, COPD/asthma, dehydration, congestive heart failure, skin infections, and urinary tract infections.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Appendix.docx](#)