Human hepatocyte PNPLA3 148M exacerbates rapid non-alcoholic fatty liver disease development in chimeric mice

Mohammad Kabbani

Rockefeller University <https://orcid.org/0000-0001-9556-384X>

Eleftherios Michailidis

The Rockefeller University, Laboratory of Virology and Infectious Disease <https://orcid.org/0000-0002-9907-4346>

Sandra Steensels

Weill Cornell Medicine

Clifton Fulmer

The Cleveland Clinic

Joseph Luna

The Rockefeller University

Jérémie Le Pen

The Rockefeller University

Matteo Tardelli

Weill Cornell Medicine

Brandon Razooky

The Rockefeller University

Inna Ricardo-Lax

The Rockefeller University

Chenhui Zou

The Rockefeller University

Briana Zeck

NYU Langone

Ansgar Stenzel

The Rockefeller University

Corrine Quirk

The Rockefeller University

Lander Foquet

Yecuris Corporation

Alison Ashbrook

The Rockefeller University

William Schneider

The Rockefeller University

Serkan Belkaya

The Rockefeller University

Gadi Lalazar

The Rockefeller University

Yupu Liang

The Rockefeller University

Meredith Pittman

Weill Cornell Medicine

Lindsey Devisscher

Ghent University

Hiroshi Suemizu

Central Institute for Experimental Animals

Neil Theise

NYU Langone

Luis Chiriboga

NYU Langone

David Cohen

Division of Gastroenterology & Hepatology, Weill Cornell Medical College, New York, NY 10021

Robert Copenhaver

Yecuris Corporation

Markus Grompe

Yecuris (United States)

Philip Meuleman

Ghent University

Baran Ersoy

Weill Cornell Medicine

Charles Rice

The Rockefeller University

Ype de Jong (✉ ydj2001@med.cornell.edu)

Weill Cornell Medicine <https://orcid.org/0000-0001-6357-2370>

Article

Keywords: advanced non-alcoholic fatty liver disease (NAFLD), patatin-like phospholipase domain-containing protein 3 (PNPLA3-I148M),

Posted Date: February 10th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-156250/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.
[Read Full License](#)

1 **Human hepatocyte PNPLA3 148M exacerbates rapid non-alcoholic fatty liver disease**
2 **development in chimeric mice**

3

4 Mohammad Kabbani^{1,2,*}, Eleftherios Michailidis^{1,*}, Sandra Steensels³, Clifton G. Fulmer^{4,5},
5 Joseph M. Luna¹, Jérémie Le Pen¹, Matteo Tardelli³, Brandon Razooky¹, Inna Ricardo-Lax¹,
6 Chenhui Zou^{1,3}, Briana Zeck⁶, Ansgar F. Stenzel^{1,7}, Corrine Quirk¹, Lander Foquet⁸, Alison W.
7 Ashbrook¹, William M. Schneider¹, Serkan Belkaya⁹, Gadi Lalazar^{3,10}, Yupu Liang¹¹, Meredith
8 Pittman⁴, Lindsey Devisscher¹², Hiroshi Suemizu¹³, Neil D. Theise⁶, Luis Chiriboga⁶, David E.
9 Cohen³, Robert Copenhaver⁸, Markus Grompe^{8,14}, Philip Meuleman¹⁵, Baran A. Ersoy³, Charles
10 M. Rice¹ and Ype P. de Jong^{1,3, #}.

11

- 12 1. Laboratory of Virology and Infectious Disease, The Rockefeller University, New York,
13 NY10065.
- 14 2. Department of Gastroenterology, Hepatology and Endocrinology, Hannover Medical School,
15 Hannover, Germany.
- 16 3. Division of Gastroenterology and Hepatology, Weill Cornell Medicine, New York, NY10065.
- 17 4. Department of Pathology, Weill Cornell Medicine, New York, NY10065.
- 18 5. Robert J. Tomsich Pathology and Laboratory Medicine Institute, The Cleveland Clinic,
19 Cleveland, Ohio, OH 44195.
- 20 6. Department of Pathology, NYU Langone, New York, NY10028.
- 21 7. Department of Infectious Diseases, Molecular Virology, Heidelberg University, Heidelberg,
22 Germany.
- 23 8. Yecuris Corporation, Tualatin, OR97062.
- 24 9. St. Giles laboratory of Human Genetics of Infectious Diseases, The Rockefeller University,
25 New York, NY10065.

- 26 10. Laboratory of Cellular Biophysics, The Rockefeller University, New York, NY10065.
- 27 11. Center for Clinical and Translational Science, The Rockefeller University, New York,
- 28 NY10065.
- 29 12. Department of Basic and Applied Medical Sciences, Gut-Liver Immunopharmacology Unit,
- 30 Ghent University, Ghent, Belgium
- 31 13. Central Institute for Experimental Animals, Kanagawa, Japan.
- 32 14. Department of Pediatrics, Oregon Stem Cell Center, Oregon Health and Science University,
- 33 Portland, OR97239.
- 34 15. Laboratory of Liver Infectious Diseases, Ghent University, Ghent, Belgium.

35 *These authors contributed equally.

36 # Lead contact.

37

38 **Correspondence:** Ype de Jong, Division of Gastroenterology and Hepatology, Weill Cornell
39 Medicine, 1305 York Ave, New York, NY 10065, Tel: 646-962-4000, Fax: 646-962-0355, Email:
40 ydj2001@med.cornell.edu.

41

42 **Short title:** Hepatocyte PNPLA3-148M worsens steatohepatitis

43 **SUMMARY**

44 Advanced non-alcoholic fatty liver disease (NAFLD) is a rapidly emerging global health problem
45 associated with pre-disposing genetic polymorphisms, most strikingly an isoleucine to methionine
46 substitution in patatin-like phospholipase domain-containing protein 3 (PNPLA3-I148M). Here, we
47 study how human hepatocytes with PNPLA3 148I and 148M variants engrafted in the livers of
48 chimeric mice respond to hypercaloric diets. As early as 4 weeks, mice developed dyslipidemia,
49 impaired glucose tolerance, and steatohepatitis selectively in the human graft, followed by
50 pericellular fibrosis after 8 weeks of hypercaloric feeding. The PNPLA3 148M variant, either from
51 a homozygous 148M human donor or overexpressed in a homozygous 148I donor background,
52 caused microvesicular and more severe steatosis. In these livers hepatocytes displayed frequent
53 ballooning degeneration, resulting in more active steatohepatitis than in 148I livers. We conclude
54 that PNPLA3 148M in human hepatocytes exacerbates NAFLD. These models will facilitate
55 mechanistic studies into human genetics associated with advanced fatty liver diseases.

56 INTRODUCTION

57 Non-alcoholic fatty liver disease (NAFLD) has become the most prevalent liver disease in
58 many countries, affecting approximately 25% of the world population.¹ NAFLD is associated with
59 obesity, insulin resistance, dyslipidemia and hypertension.² A subset of patients with NAFLD
60 develop non-alcoholic steatohepatitis (NASH), which is widely believed to drive fibrosis
61 progression towards advanced liver disease including cirrhosis and hepatocellular carcinoma
62 (HCC).³

63 The complex nutritional and environmental factors leading to NAFLD remain to be better
64 defined.⁴ In addition, a number of genetic variants have been associated with advanced NAFLD.⁵
65 The first genetic polymorphism identified in a population with increased hepatic steatosis was
66 rs738409 in patatin-like phospholipase domain containing protein-3 (*PNPLA3*).⁶ This
67 polymorphism encodes for an isoleucine to methionine substitution at codon position 148 in
68 *PNPLA3*. Since its identification by Hobbs and colleagues, the *PNPLA3*-I148M variant has been
69 globally associated with advanced NAFLD and other liver diseases including HCC.⁷

70 The mechanisms by which *PNPLA3*-148M drives advanced NAFLD remains an active area
71 of investigation.^{8, 9} Although *PNPLA3* is expressed in hepatocytes, hepatic stellate cells and
72 adipocytes, accumulating evidence from mouse models indicates that hepatocyte-specific
73 *PNPLA3*-148M promotes the pathogenesis of NAFLD. Transgenic overexpression of human
74 *PNPLA3*-148M in mouse hepatocytes but not adipocytes exacerbated hepatic steatosis.¹⁰ And
75 while increased hepatic fat content was observed in *Pnpla3*-148M knock-in mice,¹¹ this was
76 reversed by silencing *Pnpla3* in hepatocytes.¹² Functionally, *PNPLA3* exhibits hydrolase activity
77 towards triglycerides and acyltransferase activity for polyunsaturated fatty acids in phospholipids.⁸
78 *PNPLA3*-148M differs from 148I in several ways: *PNPLA3*-148M was shown to have decreased
79 hydrolase activity against triglycerides,¹³ increased acyltransferase activity for lysophosphatidic
80 acid,¹⁴ resistance to ubiquitylation and proteasomal degradation,¹⁵ and inhibition of adipose

81 triglyceride lipase on lipid droplets via interaction with α/β -hydrolase domain containing 5.^{16, 17}
82 How these and other mechanistic differences drive NAFLD progression remains to be further
83 elucidated.

84 Despite this vast disease burden there are no approved pharmacological therapies for
85 NASH and sustained weight loss remains the only proven intervention. Given its complex
86 pathogenesis, many different pathways are under pre-clinical and clinical investigation, some of
87 which have resulted in compounds that progressed to Phase 3 trials.¹⁸ Several of these pathways
88 were identified in murine NASH models that share limited features with human disease.¹⁹
89 Furthermore, investigational strategies that target subsets of NASH patients, e.g. those with
90 genetic variants, are further complicated by species differences. In the case of PNPLA3, there is
91 low sequence homology between the human and mouse orthologues. The human protein is 97
92 amino acids longer than the mouse PNPLA3. The relative mRNA expression starkly differs
93 between adipose tissue and liver in each species.^{20, 21} and it remains unclear if the same
94 transcription factors regulate its expression in murine and human hepatocytes.^{22, 23} Elucidation of
95 the precise mechanisms by which human genetic variants in hepatocytes affect steatohepatitis
96 would be aided by minimizing species-specific variances in a physiologically relevant
97 experimental setting.

98 Here, we established a human hepatocyte NAFLD model. We applied this model to study
99 human hepatocyte PNPLA3-148M, either from a 148M homozygous human hepatocyte donor
100 and by overexpressing PNPLA3 variants in a 148I homozygous donor background. Our results
101 show that human hepatocytes swiftly develop steatohepatitis in response to hypercaloric feeding
102 and that NAFLD activity is higher when hepatocytes express PNPLA3-148M. These models will
103 allow for mechanistic studies into human genetic variants implicated in advanced NAFLD.

104

105

106 **RESULTS**

107 **Human hepatocytes in chimeric mice on Western diet rapidly develop steatosis**

108 To investigate how the human graft in liver chimeric mouse models responds to a
109 hypercaloric diet, we first created humanized mice by transplanting primary human hepatocytes
110 (PHH) into preconditioned immunocompromised *Fah^{-/-}* NOD *Rag1^{-/-}* *Il2rg^{null}* (FNRG) mice as
111 described previously ^{24,25}. Cycling the protective drug nitisinone ²⁶ resulted in at least 50%
112 humanization after which mice were subjected to an *ad libitum* Western-style diet (WD), which
113 consists of 60% fat in diet and 10% sucrose in drinking water (**Fig 1a**). WD feeding did not result
114 in graft loss as determined by persistently high ²⁷⁻²⁹ human serum albumin (hAlb) levels (**Suppl**
115 **Fig S1a**). Periodic examination of huFNRG mouse livers showed steatosis³⁰ as early as 4 weeks
116 on WD as evidenced by hematoxylin and eosin (H&E) staining (**Fig 1b**) and Oil Red O staining
117 (**Suppl Fig S1b**). Steatosis almost exclusively affected the human graft whereas the remaining
118 mouse hepatocytes, which appear larger with darker cytoplasm after H&E staining, rarely
119 developed steatosis. The contrast in steatosis between mouse and human hepatocytes was
120 further evident by using a human-specific marker (**Fig 1c**). Livers from highly humanized mice
121 with severe steatosis showed sparing around periportal regions (**Fig 1b**), reminiscent of zonal
122 steatosis in adult NAFLD patients. The zonal predominance was further highlighted by glutamine
123 synthetase staining, which is mostly expressed in zone 3 hepatocytes (**Fig 1d**). In order to quantify
124 steatosis in huFNRG mice, the human areas of chimeric livers were scored blindly according to
125 clinical criteria.³⁰ All animals developed mild to severe steatosis as early as 4 weeks on WD.
126 Steatosis persisted for 20 weeks on WD, which contrasted with huFNRG mice on control chow
127 (chow) that never developed more than 5% steatosis (**Fig 1e**). Clinical scoring was confirmed by
128 total triacylglycerol (TAG) quantification in huFNRG livers, which was 7 to 11-fold higher in WD
129 than chow mice (**Fig 1f**), and liver total cholesterol (**Suppl Fig S1c**). Overall, the accumulation of
130 TAG in huFNRG livers correlated to the steatosis score of the human graft (**Suppl Fig S1d**).

131 These findings show that human hepatocytes in huFNRG mice rapidly developed moderate
132 steatosis in response to WD feeding.

133 We next tested if the marked steatosis difference between human and mouse hepatocytes
134 was due to the NOD mouse background, the intermittent nitisinone used to protect *Fah*^{-/-} mice
135 from liver injury³¹, or the hepatocyte transplantation biology of the FNRG model. First, we
136 subjected immunodeficient NOD *Rag1*^{-/-} *Il2rg*^{null} (NRG) mice to WD feeding and nitisinone cycling.
137 Even after 12 weeks on WD NRG livers developed minimal steatosis (**Suppl Fig S1e**), consistent
138 with previous reports that T cell deficient mice are protected from experimental NAFLD.³²⁻³⁴ To
139 test if FNRG liver injury and/or hepatocyte repopulation physiology made hepatocytes susceptible
140 to hypercaloric feeding we transplanted NRG mouse hepatocytes into FNRG mice. These animals
141 were designated 'murinized' FNRG (muFNRG) mice. After 4 weeks on WD muFNRG livers did
142 not display steatosis, but small areas of hepatocyte steatosis were present after 12 weeks of WD
143 feeding (**Fig 1g**) which was in line with liver TAG levels (**Suppl Fig S1f**). These results show that
144 human hepatocytes responded differently to WD feeding than mouse hepatocytes in these
145 models.

146 We then tested if rapid steatosis was specific to this human hepatocyte donor (PHH1) and
147 to the FNRG liver injury model, extending our protocols to other donors and liver chimeric mouse
148 models. We first transplanted PHH2 into FNRG mice and thymidine kinase transgenic (TK-NOG)
149 liver injury mice.³⁵ Despite reaching lower levels of humanization in both models, human
150 hepatocyte islands developed steatosis in response to WD feeding but not on chow (**Suppl Fig**
151 **S1g**). We also transplanted a third PHH donor (PHH3) into urokinase plasminogen activator
152 (uPA)/SCID mice³⁶ and FNRG mice. Interestingly, humanized (hu-)uPA/SCID mice showed mild
153 steatosis in the human graft on chow, which contrasted to the absence of steatosis in huFNRG
154 livers on chow (**Suppl Fig S1h**). In the huFNRG model, PHH3 hepatocytes developed steatosis
155 after 4 weeks of WD feeding. These data illustrate that humanized livers rapidly became steatotic

156 following WD feeding regardless of the PHH donor, with different requirements for hypercaloric
157 feeding between liver injury models.

158 Combined, these results demonstrate that human hepatocytes but not mouse hepatocytes
159 in liver chimeric mice developed moderate steatosis as early as 4 weeks on WD.

160

161 **Western diet affects systemic metabolic parameters of chimeric FNRG mice**

162 NAFLD clinically overlaps with obesity, dyslipidemia, insulin resistance and possibly
163 hypertension.² To determine whether WD affects metabolic parameters in liver chimeras, we first
164 measured body weight accumulation in adult huFNRG mice on an *ad libitum* WD or control chow
165 (chow) diet. Compared to chow, WD caused nearly 20% increase in body weight within 4 weeks
166 on the diet and maintained this difference through 12 weeks on WD (**Fig 2a**). Furthermore, the
167 gonadal fat fraction (**Fig 2b**) increased 2- to 2.4-fold in huFNRG mice on WD compared to chow.
168 These data demonstrate that WD feeding resulted in weight gain and gonadal fat accumulation
169 in huFNRG mice.

170 We then tested plasma lipids in chimeric mice. Hepatocytes are central to lipid
171 homeostasis, and humanized mice have plasma cholesterol profiles shifted toward lower density
172 lipoproteins compared to wild type mice.³⁷ Similar to previous reports,³⁸⁻⁴⁰ huFNRG on chow
173 contained more LDL than HDL cholesterol particles compared to muFNRG mice (**Fig 2c**). Four
174 weeks of WD feeding resulted in stark increases in VLDL and LDL cholesterol in huFNRG plasma.
175 Notably, the HDL fraction in huFNRG mice decreased with WD feeding whereas it rose in WD-
176 fed muFNRG mice compared to chow (**Suppl Fig S2a**). Relative to chow, WD increased fasting
177 serum triglycerides fractions, mostly in VLDL, and to a lesser extent in LDL fractions in huFNRG
178 mice (**Fig 2d, Suppl Fig S2b**). These data show that WD feeding of huFNRG mice increased
179 lower density lipoproteins and decreased HDL.

Given the strong clinical overlap between NAFLD and type 2 diabetes, we tested the effect of WD on glucose tolerance in huFNRG mice relative to muFNRG animals. While humanized animals started at a higher baseline, 4 weeks on WD resulted in a 2.2-fold increase in fasting blood glucose both in huFNRG and muFNRG mice (**Fig 2e**). Fasting mouse insulin levels on chow appeared higher in humanized than murinized FNRG animals, and increased only slightly in huFNRG animals after 4 weeks on WD (**Fig 2f**). To further characterize glucose metabolism, fasting animals were subjected to glucose, insulin or pyruvate tolerance tests. After 4 weeks on WD, huFNRG mice displayed severely impaired glucose clearance with 3.1-fold higher cumulative blood glucose than chow, whereas this difference was 1.67-fold higher in muFNRG mice (**Fig 2g**). By contrast huFNRG mice on WD remained responsive to insulin challenge (**Fig 2g**). To assess gluconeogenesis in these animals, a pyruvate injection after fasting for 16 hours resulted in a sharper rise in blood glucose in huFNRG on a WD than chow (**Fig 2g**). Eight weeks of WD feeding further prolonged time for glucose clearance in huFNRG mice (**Suppl Fig S2c**) without affecting endogenous insulin secretion (**Suppl Fig S2d**). These data show that WD feeding of chimeric FNRG mice affected glucose homeostasis without altering insulin-mediated glucose uptake.

Together these data show that WD in huFNRG mice causes increased body weight, dyslipidemia and impaired glucose metabolism compared to chow diet.

Western diet causes mild steatohepatitis in huFNRG mice

Non-alcoholic steatohepatitis (NASH) frequently causes elevated serum transaminases⁴¹ and is diagnosed histologically by the presence of hepatocyte ballooning and mononuclear infiltrates in the setting of >5% hepatocyte steatosis.³⁰ Because the human graft rapidly developed steatosis, we tested whether huFNRG mice on WD also developed features associated with human NASH.

We quantified the activities of serum transaminases ALT and AST, which were elevated in huFNRG mice 4 weeks on WD compared to chow. With prolonged WD feeding, ALT activity remained high while AST increased further (**Fig 3a**). To determine whether the ALT was produced by human or mouse hepatocytes, human-specific ALT protein was quantified and this correlated with serum ALT activity (**Suppl Fig S3a**). The human origin of transaminase activity was further supported by the lack of ALT activity and only minimal increases in AST activity in muFNRG mice on WD (**Suppl Fig S3b**). These data show that WD increased human transaminases in huFNRG serum.

We next assessed whether human hepatocytes in huFNRG mice on WD displayed ballooning degeneration, a key characteristic of NASH.³⁰ As early as 4 weeks on WD, steatotic hepatocytes in huFNRG mice showed ballooning degeneration that persisted for 20 weeks (**Fig 3b**, **Suppl Fig S3c**). Hepatocyte ballooning was specific to the human graft (**Fig 3c**) and not donor hepatocyte-specific (**Suppl Fig S3d**). The fraction of huFNRG mice with ballooning degeneration did not increase with prolonged WD feeding (**Fig 3d**) but rather associated with the severity of steatosis grade (**Fig 3e**). Ballooning degeneration, which remains less well-defined in mouse hepatocytes,¹⁹ was not observed in either NRG or muFNRG mice on WD. Another key feature of NASH is lobular inflammation, which is dominated by neutrophils and macrophages.⁴² Despite their broad immunodeficient background inflammatory infiltrates were detected in huFNRG mouse livers after 4 weeks on WD (**Fig 3b**, **Suppl Fig S3c**). Lobular inflammation was observed in approximately one third of huFNRG mice up to 12 weeks on WD but was not observed at 20 weeks (**Suppl Fig S3e**). The livers of huFNRG mice on WD were scored according to the NAFLD activity score (NAS)⁴³ within the human graft. After 4 weeks on WD, 7 out of 8 huFNRG mice developed a NAS of 3 or more. Further durations of WD did not increase the fraction of mice with $\text{NAS} \geq 3$ (**Fig 3f**). These data show that huFNRG mice developed features of steatohepatitis following WD feeding.

230 Pericellular fibrosis can accompany NASH and is believed to result from ongoing
231 steatohepatitis.⁴⁴ To test if WD feeding of huFNRG mice caused fibrosis, livers were stained for
232 collagen with Picrosirius Red (PSR). Pericellular collagen depositions were first observed in a few
233 mice after 8 weeks on WD (**Fig 3g**), then in a larger fraction at 12 weeks and finally more prevalent
234 after 20 weeks on WD (**Fig 3h**). Trichrome staining confirmed collagen depositions around human
235 hepatocytes, which are smaller with paler cytoplasm compared to mouse hepatocytes (**Suppl Fig**
236 **S3f**). Fibrosis was not PHH1 specific since mice humanized with donor PHH2 showed a similar
237 phenotype after 10 weeks on WD (**Suppl Fig S3g**). These data indicate that prolonged WD
238 feeding results in mild fibrosis around human hepatocytes.

239 Overall, these findings show that huFNRG mice developed features of steatohepatitis with
240 mild activity as early as 4 weeks on WD, which was accompanied by pericellular fibrosis in the
241 human graft after 8 weeks of WD feeding.

242

243 **WD differentially affects human and mouse liver transcriptomes over time**

244 In huFNRG chimeras that were created with PHH the non-parenchymal cells and
245 leukocytes remained murine (²⁷ and unpublished observations). To determine how WD affected
246 human and mouse gene expression we performed RNA-sequencing (RNA-seq) on livers from
247 fasting huFNRG mice. STAR Alignment mapped >85% of transcripts uniquely to a human
248 reference genome (**Suppl Fig S4a**).

249 Transcripts aligned to human genome (human reads) in mice that were fed WD for 4 weeks
250 were compared to animals on chow, which identified 662 differentially expressed transcripts (**Fig**
251 **4a**). Among mice on WD human transcripts changed progressively less over time, totaling 154 at
252 8 vs. 4 weeks, 127 at 12 vs. 8 weeks, and 95 at 20 vs. 12 weeks (**Fig 4a**).

253 We then aligned transcripts of these highly humanized livers to mouse reference genome
254 (mouse reads). In contrast, only 2 mouse transcripts were statistically different between huFNRG

255 mice that were fed WD or chow for 4 weeks. Transcriptional changes in the mouse reads
256 remained low for the first 12 weeks of WD feeding, but a large number of differentially expressed
257 transcripts were observed by 20 weeks (**Fig 4b**).

258 To further investigate changes in the gene expression profile we performed gene ontology
259 (GO) analysis using the GO, Kyoto Encyclopedia of Genes and Genomes (KEGG), REACTOME
260 (REAC), and Transcription Factor (TF) databases. GO (**Fig 4c**), KEGG and REAC (**Suppl Fig**
261 **S4b**) alignments of human reads revealed mostly downregulation of metabolic and catabolic
262 pathways in mice 4 weeks on WD compared to chow, while transcription factors were upregulated
263 (**Suppl Fig S4b**). A notable exception to many downregulated GO pathways was the Extracellular
264 Matrix pathway, which was upregulated at 4 weeks of WD feeding. Comparing 8 vs. 4 weeks on
265 WD identified upregulation of human GO pathways involved in cell death and NF- κ B signaling.
266 Later timepoints revealed fewer and statistically less significant pathways. Few mouse GO
267 pathways were changed by WD feeding at early time points. At 20 weeks many pathways involved
268 in cellular development and metabolism changed (**Fig 4d**, **Suppl Fig S4c**). Notably several
269 metabolic pathways that were downregulated in the human transcriptome at 4 weeks were
270 upregulated in the mouse transcriptome at 20 weeks.

271 Taken together, these data indicate that many metabolic pathways were transcriptionally
272 downregulated in the human graft as early as 4 weeks on WD, while the mouse transcriptome of
273 huFNRG mice remained largely silent until 20 weeks of WD feeding.

274

275 **Mice humanized with a homozygous PNPLA3-148M donor develop steatohepatitis**

276 Having established a new NAFLD model in chimeric mice, we set out to characterize how
277 the PNPLA3-148M polymorphism in hepatocytes affected the phenotype. Thus far huFNRG mice
278 were humanized with PHH donors that were homozygous for PNPLA3-148I (148I-huFNRG). We
279 next tested how chimeric mice humanized with the PNPLA3-148M variant responded to WD. To

280 this end *Fah^{-/-} Rag2^{-/-} Il2rg^{null}* (FRG) mice²⁶ were humanized with a homozygous PNPLA3-148M
281 donor (148M-huFRG) and given WD for 4 weeks. As expected WD caused metabolic changes,
282 including raised fasting blood glucose, a minimal increase in fasting insulin and generally higher
283 plasma triglycerides (**Suppl Fig S5a-c**). Livers of 148M-huFRG mice on WD had moderate
284 (>33%) and severe (>66%) steatosis in contrast to <5% steatosis that was observed in chow-fed
285 mice (**Fig 5a-b**). Histological findings were confirmed by 5.3-fold increase in TAG concentrations
286 in the livers of 148M-huFRG mice on WD compared to chow-fed controls (**Suppl Fig S5d**). In
287 contrast to previous 148I-huFNRG studies, the human graft showed not only macrovesicular but
288 also microvesicular steatosis in all 148M-huFRG mice on WD (**Fig 5c**). In addition, 148M-huFRG
289 livers contained Mallory-Denk bodies, high grade of ballooning degeneration and lobular
290 inflammation, resulting in 4 out of 5 mice with a NAS of 5 or above (**Fig 5d, Suppl Fig S5e-f**). These
291 data demonstrate that 148M-huFRG mice subjected to WD for 4 weeks showed features of
292 steatohepatitis with moderate activity.

293 To investigate if 148M-huFRG mice transcriptionally responded differently to WD feeding
294 than 148I-huFNRG animals, RNA-seq analyses was performed. The human graft in 148M-huFRG
295 mice responded to 4 weeks on WD with numerous transcriptomic changes compared to chow,
296 with 1458 downregulated and 1294 upregulated genes (**Fig 5e**). Pathway analyses revealed that
297 metabolic and catabolic pathways were mostly downregulated by WD in 148M-huFRG mice,
298 similar to previous results in 148I-huFNRG mice. Despite the similarities between both models,
299 there were pathways, including oxidative phosphorylation and mitochondrial function, which were
300 downregulated following WD only in the livers of 148M-huFRG mice but not in 148I-huFNRG. In
301 addition, some pathways were differentially upregulated by WD between these two models (**Fig**
302 **5f**). Comparing individual genes from livers of 148M-huFRG mice and 148I-huFNRG, there were
303 2329 differently expressed transcripts on chow and 3268 on WD (**Suppl Fig S5g**). However,

304 when comparing transcripts that changed by WD feeding over chow in each model (Δ WD), there
305 were only 108 genes differentially regulated between these models (**Fig 5g**).

306 Taken together, these findings show that 148M-huFRG mice developed microvesicular
307 steatosis and steatohepatitis with moderate activity after 4 weeks on WD. While few pathways
308 were transcriptionally different between 148M-huFRG and 148I-huFNRG mice on the same diet,
309 several metabolic pathways were uniquely downregulated in 148M hepatocytes.

310

311 **PNPLA3-148M overexpression in hepatocytes exacerbates steatosis in mice on WD**

312 In addition to the PNPLA3 genotype, 148M-huFRG and 148I-huFNRG mice were distinct
313 in human hepatocyte donor background and mouse strain differences. To better characterize if
314 phenotypic differences were due to the PNPLA3 variant, we transduced mouse-passaged PHH
315 cultures from a 148I homozygous donor (PHH1) with PNPLA3-148M and red fluorescent protein
316 (RFP), 148I and RFP or RFP only before re-transplantation into mice ²⁵. Humanization kinetics
317 did not differ between the PNPLA3 variants or RFP controls as determined by serum hAlb values
318 (**Suppl Fig S6a**). Once hAlb serum levels plateaued, huFNRG mice with transduced PNPLA3-
319 148M hepatocytes (td148M), td148I or tdRFP mice were subjected to 4 weeks on WD or chow
320 (**Fig 6a**). In the WD groups lentiviral PNPLA3 transduction resulted in 61-fold higher mRNA levels
321 compared to endogenous levels in tdRFP mice, while in the chow groups td148I had 2-fold higher
322 levels than td148M mice. In tdRFP mice WD caused 3.7-fold higher endogenous PNPLA3 mRNA
323 levels compared to chow (**Suppl Fig S6b**).

324 WD induced human hepatocyte steatosis in all three groups of mice, with only td148M
325 livers containing widespread microvesicular steatosis (**Fig 6b-c**). Strikingly, even on chow td148M
326 livers contained rare areas with human microvesicular steatosis, which was not observed in
327 td148I or tdRFP livers (**Suppl Fig S6c**). Lipid quantification in livers from mice on WD showed
328 36% more triglycerides in td148M livers compared to td148I or tdRFP, with only minimal

329 differences in mice on chow (**Fig 6d**). Despite these phenotypic changes, the NAS was not
330 statistically different in td148M mice from td148I or tdRFP animals (**Fig 6e, Suppl Fig 6d-e**).
331 These data demonstrate that PNPLA3 overexpressing huFNRG mice on WD developed
332 steatohepatitis in the human graft, with td148M livers containing widespread microvesicular
333 steatosis and increased hepatic triglycerides.

334 Since RNA-seq pathway analyses did not identify many differences between the 148M-
335 huFRG and 148I-huFNRG models (**Fig 5g**), we performed qRT-PCR for 80 different human
336 genes associated with fatty liver diseases (**Table 1**). Liver tissue from td148I and td148M mice
337 that were fed chow revealed 20 genes that were expressed differently, with lipoprotein lipase
338 showing the greatest difference (**Suppl Fig S6f**). Comparing both groups following WD we
339 identified only 8 genes that were affected by 2-fold or less, of which *CYP2E1* was most
340 transcriptionally upregulated in td148M livers (**Fig 6f**). These results illustrate the modest
341 expression differences between mice overexpressing the two PNPLA3 variants on WD.
342 Combined, these data show that PNPLA3-148M overexpression in hepatocytes on a homozygous
343 148I background caused widespread microvesicular steatosis and increased liver triglycerides
344 after 4 weeks of WD feeding.

345

346 **A high fat diet causes more active steatohepatitis in PNPLA3-148M overexpressing mice**

347 We hypothesized that the minimal differences between tdRFP, td148I and td148M
348 mice was caused by the severe steatosis phenotype of WD feeding in lentivirally transduced livers
349 of huFNRG mice. Therefore, we challenged PNPLA3 overexpressing huFNRG mice with a high
350 fat diet (HFD, 60% fat) only, excluding the sucrose drinking water portion of WD. After 4 weeks
351 of HFD, PNPLA3-148M overexpression resulted in more steatosis (**Fig 7a-b**), which was
352 confirmed by hepatic triglyceride quantification (**Suppl Fig S7a**). In td148M mice the HFD resulted
353 in widespread microvesicular steatosis and Mallory Denk bodies (**Fig 7c**), which were not

observed in either td148I or tdRFP mice. Ballooning degeneration in mice fed a HFD trended toward being more severe in the td148M group than in tdRFP and td148I ($p=0.063$, **Suppl Fig S7b**) while lobular inflammation was not affected by PNPLA3 overexpression (**Suppl Fig S7c**). Driven by steatosis and ballooning degeneration grades, the NAS was higher in td148M than in td148I or tdRFP mice (**Fig 7d**). These data show that HFD caused steatohepatitis in PNPLA3 transduced human grafts, with 148M transduction resulting in microvesicular steatosis, Mallory Denk bodies and higher activity compared to 148I.

PNPLA3 mRNA levels in liver pieces of td148I and td148M mice on HFD were similar (**Suppl Fig S7d**). Since lentiviral PHH transduction results in heterogeneous expression²⁵ we next analyzed whether steatosis correlated to the transgene expression, using the RFP reporter as a surrogate marker.⁴⁵ Low magnification fluorescence imaging of liver cross-sections stained for neutral lipids suggested that td148M livers on HFD contained more lipids than td148I mice (**Fig 7e**). At higher magnification steatosis appeared more severe in RFP positive areas of td148M livers whereas lipid distribution in td148I mice was not associated with RFP. We then analyzed RFP and neutral lipid intensity from individual small hepatocytes, calibrated to a highly humanized huFNRG mouse whose graft was severely steatotic (**Suppl Fig S7e**). Using all mice from an experiment with high hAlb serum values, a two-dimensional density plot of individual hepatocytes from three td148M animals showed that hepatocytes with higher RFP intensity contained more lipids, whereas lipids were equally distributed along RFP intensity in hepatocytes from four td148I mice (**Fig 7f**). These findings suggest that 148M but not 148I overexpression, as determined by a RFP reporter, exacerbated steatosis.

Combined, these findings show that PNPLA3-148M overexpression in hepatocytes from a homozygous 148I donor increased steatosis and steatohepatitis activity in chimeric mice on HFD.

DISCUSSION

In this study, we have established systems to model NAFLD genetics in human hepatocytes. The most striking phenotype in the 'standard' 148I-huFNRG model was the rapid onset of human steatosis on WD. This contrasted to the minimal steatosis in the residual mouse hepatocytes of huFNRG mice or in muFNRG controls. Both resembled non-humanized mouse models that require prolonged WD feeding to develop moderate steatosis.⁴⁶ Impaired cross-talk between human hepatocytes and murine cells likely contributed, as we previously showed for FGF15 and FGF19 signaling,⁴⁷ yet mouse and human hepatocytes may also inherently respond differently to overnutrition. This is supported by the clinical observations of how rapidly hepatic fat content changes depending on the type and amount of food intake. For example, a three-week overfeeding study increased intrahepatic triglycerides up to 55%⁴⁸ and a carbohydrate-restricted diet reduced hepatic fat by 44% in two weeks.⁴⁹ These clinical data illustrate that hepatic steatosis dynamically responds to nutritional variations. huFNRG mice can provide a platform to explore how hepatocytes rapidly respond to hypercaloric diets.

Hepatocytes are central to systemic lipid and glucose regulation. Similar to previous reports³⁸⁻⁴⁰ huFNRG on chow had starkly different lipid profiles than muFNRG, in part due to mice lacking cholesterol ester transfer protein.⁴⁰ In addition, we show that fasting glucose and possibly fasting insulin levels were higher in huFNRG mice than muFNRG mice on chow. Species incompatibilities may contribute to the elevated fasting glucose of huFNRG mice on chow, as was illustrated by studies in the uPA liver chimeric model.⁵⁰ While some human donors develop spontaneous steatosis in these chimeras, similar to what we observed with PHH3, administration of human growth hormone reversed this phenotype, possibly through IGF-1 signaling in the liver.^{50, 51} We speculate that hepatocytes in huFNRG are likewise predisposed to steatosis, and therefore may only require a short duration of hypercaloric feeding to develop moderate steatosis. Interestingly, muFNRG controls without hepatic steatosis developed glucose intolerance on WD

that was proportionally similar to huFNRG mice with moderate steatosis. Uncoupling hepatocyte steatosis from glucose homeostasis in these models may advance insights into clinically distinct forms of NAFLD with and without insulin resistance.⁵²

Despite their broad immunodeficiency huFNRG mice displayed the features that define human NASH, including ballooning degeneration and lobular infiltrates. This illustrates that steatohepatitis and pericellular fibrosis can be initiated without lymphocytes. Previous NAFLD studies in T cell deficient mouse strains showed these to be resistant to steatosis development,^{32, 33} which are in line with our muFNRG negative controls and further highlight species differences between human and mouse hepatocytes. The cellular immune contributions to human NASH remain to be better defined. In human NAFLD, portal immune infiltrates contain macrophages and T cells whereas lobular infiltrates are dominated by macrophages and neutrophils.⁴² How these cell types and their subsets are involved in the immunopathogenesis of human NASH remains under investigation. Proposed mechanisms are largely based on animal models with unclear translational value.⁵³⁻⁵⁵ Some compounds that effectively interfered with the development of murine steatohepatitis failed in NASH patients,⁵⁶ and no immunological interventions have yet shown clear benefits in later phase clinical trials. Furthermore, species differences preclude investigations into non-conserved pathways, e.g. IL-32 that lacks a mouse orthologue. This cytokine was recently associated with advanced NAFLD⁵⁷ and *IL32* was transcriptionally upregulated in 148I-huFNRG livers after 12 weeks and 148M-huFRG livers after 4 weeks of WD feeding. Going forward, studies dissecting the roles of T cells and other lymphocytes will require immune competent mouse models while human immune system mice can be combined with liver chimeras^{27, 58} for investigations into pathways that are not conserved between mice and humans.

Transcriptomic characterization of huFNRG livers showed a large number of human metabolic pathways that were transcriptionally downregulated after 4 weeks of WD feeding and only a few pathways that were upregulated. Eight weeks of WD feeding caused upregulation of a

427 small number of cellular stress and cell death pathways, and remarkably few human pathways
428 transcriptionally changed after that. Equally surprising were the small number of mouse
429 transcripts that changed during the first 12 weeks on WD. This may be partly explained by under-
430 sampling of mouse hepatocytes at earlier timepoints, as illustrated by changes in mouse
431 metabolic pathways after 20 weeks WD when humanization histologically appeared to decline.
432 The murine transcriptome, however, was also largely unperturbed by WD feeding. Whether
433 changes from WD were masked by cyclical nitisinone withdrawal or if non-parenchymal cells did
434 not transcriptionally respond to signals from the human graft will require further studies, including
435 in other chimera models without cyclical liver injury.

436 These systems will now allow for mechanistic studies into genetic variants that drive
437 NAFLD. Many polymorphisms associated with elevated transaminases⁵⁹ and most variants
438 implicated in advanced NAFLD⁶⁰ are in genes expressed in hepatocytes. As a proof-of-principle,
439 we modeled the PNPLA3-148M polymorphism.⁶ Its prevalent allele frequency allowed for the
440 identification of a high-quality homozygous 148M donor to create human chimeras. PNPLA3
441 overexpression in homozygous 148I hepatocytes then controlled for genetic differences between
442 these two PHH donors. Only PNPLA3-148M overexpression resulted in microvesicular steatosis,
443 suggesting that this phenotype in 148M-huFRG mice is associated with its PNPLA3 genotype.
444 Differences between 148I and 148M had previously been identified in mouse hepatocytes and
445 human cell lines,^{10, 11, 13, 15, 16, 61} none of which displayed microvesicular steatosis. By contrast,
446 hepatocytes in arguably the most physiological mouse model, namely in PNPLA3 148M knockin
447 mice, contained increased lipid droplet sizes on hypercaloric diet.¹¹ This suggests that the
448 microvesicular phenotype, which clinically correlates with advanced NASH,⁶² may be a human-
449 specific 148M hepatocyte response to hypernutrition.

450 Taken together these results show the utility of liver chimeras for studies of pathways that
451 are genetically implicated in human fatty liver diseases. Notwithstanding their broad implications

452 to advance human hepatocyte research, several technical challenges remain. Identifying high-
453 quality PHH donors with rare allele variants, e.g. in *TM6SF2* or *MBOAT7*,^{63, 64} will remain difficult.
454 And creating isogenic controls for more frequent variants, e.g. in *PNPLA3* and *HSD17B13*,⁶⁵ will
455 require improvements in gene editing efficiency of primary human materials, either to disrupt
456 genes or prime edit polymorphisms.⁶⁶ An alternative strategy is to prime edit renewable cells such
457 as pluripotent stem cells. However, hepatocyte-like cells derived from these sources have thus
458 far failed to reliably and reproducibly humanize mouse chimera models.^{67, 68} We here used an
459 overexpression system to control for human hepatocyte donor differences. The lentiviral LTR
460 promotor⁶⁹ in our models failed to recapitulate the complexities of dietary regulation of *PNPLA3*
461 transcription and translation.²⁰ This likely resulted in supraphysiological protein amounts
462 particularly of the 148M variant with its known resistance to ubiquitylation,^{11, 12, 15} and may explain
463 the steatotic areas observed in td148M mice on chow. Human and mouse *PNPLA3* expression
464 are regulated by sterol regulatory binding protein 1c (*SREBP1c*),^{20, 22, 23} while mouse *Pnpla3*
465 transcription may additionally be under control of carbohydrate response element binding protein
466 (*ChREBP*).²² Creating human liver chimeric mice with *PNPLA3* expression under *SREBP1c*
467 regulation would become a more physiological system to study *PNPLA3*-148M in hepatocytes
468 with appropriate controls. Going forward, overcoming technical challenges with primary
469 hepatocyte gene editing and hepatocyte-like cell engraftment would further advance the models
470 presented here and generate ever more physiologically relevant systems to dissect the
471 pathophysiology of human fatty liver diseases.

472

REFERENCES

1. Younossi, Z.M. et al. Global epidemiology of nonalcoholic fatty liver disease-Meta-analytic assessment of prevalence, incidence, and outcomes. *Hepatology* **64**, 73-84 (2016).
2. Yki-Järvinen, H. Non-alcoholic fatty liver disease as a cause and a consequence of metabolic syndrome. *Lancet Diabetes Endocrinol* **2**, 901-910 (2014).
3. Diehl, A.M. & Day, C. Cause, Pathogenesis, and Treatment of Nonalcoholic Steatohepatitis. *N Engl J Med* **377**, 2063-2072 (2017).
4. Buzzetti, E., Pinzani, M. & Tsochatzis, E.A. The multiple-hit pathogenesis of non-alcoholic fatty liver disease (NAFLD). *Metabolism* **65**, 1038-1048 (2016).
5. Romeo, S., Sanyal, A. & Valenti, L. Leveraging Human Genetics to Identify Potential New Treatments for Fatty Liver Disease. *Cell Metab* **31**, 35-45 (2020).
6. Romeo, S. et al. Genetic variation in PNPLA3 confers susceptibility to nonalcoholic fatty liver disease. *Nat Genet* **40**, 1461-1465 (2008).
7. Trépo, E., Romeo, S., Zucman-Rossi, J. & Nahon, P. PNPLA3 gene in liver diseases. *J Hepatol* **65**, 399-412 (2016).
8. Pingitore, P. & Romeo, S. The role of PNPLA3 in health and disease. *Biochim Biophys Acta Mol Cell Biol Lipids* **1864**, 900-906 (2019).
9. Tardelli, M., Bruschi, F.V. & Trauner, M. The role of metabolic lipases in the pathogenesis and management of liver disease. *Hepatology* (2020).
10. Li, J.Z. et al. Chronic overexpression of PNPLA3I148M in mouse liver causes hepatic steatosis. *J Clin Invest* **122**, 4130-4144 (2012).
11. Smagris, E. et al. Pnpla3I148M knockin mice accumulate PNPLA3 on lipid droplets and develop hepatic steatosis. *Hepatology* **61**, 108-118 (2015).
12. Lindén, D. et al. Pnpla3 silencing with antisense oligonucleotides ameliorates nonalcoholic steatohepatitis and fibrosis in Pnpla3 I148M knock-in mice. *Mol Metab* **22**, 49-61 (2019).

498 13. Huang, Y., Cohen, J.C. & Hobbs, H.H. Expression and characterization of a PNPLA3
499 protein isoform (I148M) associated with nonalcoholic fatty liver disease. *J Biol Chem* **286**,
500 37085-37093 (2011).

501 14. Kumari, M. et al. Adiponutrin functions as a nutritionally regulated lysophosphatidic acid
502 acyltransferase. *Cell Metab* **15**, 691-702 (2012).

503 15. BasuRay, S., Smagris, E., Cohen, J.C. & Hobbs, H.H. The PNPLA3 variant associated
504 with fatty liver disease (I148M) accumulates on lipid droplets by evading ubiquitylation.
505 *Hepatology* **66**, 1111-1124 (2017).

506 16. Wang, Y., Kory, N., BasuRay, S., Cohen, J.C. & Hobbs, H.H. PNPLA3, CGI-58, and
507 Inhibition of Hepatic Triglyceride Hydrolysis in Mice. *Hepatology* **69**, 2427-2441 (2019).

508 17. Yang, A., Mottillo, E.P., Mladenovic-Lucas, L., Zhou, L. & Granneman, J.G. Dynamic
509 interactions of ABHD5 with PNPLA3 regulate triacylglycerol metabolism in brown
510 adipocytes. *Nat Metab* **1**, 560-569 (2019).

511 18. Wattacheril, J., Issa, D. & Sanyal, A. Nonalcoholic Steatohepatitis (NASH) and Hepatic
512 Fibrosis: Emerging Therapies. *Annu Rev Pharmacol Toxicol* **58**, 649-662 (2018).

513 19. Farrell, G. et al. Mouse Models of Nonalcoholic Steatohepatitis: Toward Optimization of
514 Their Relevance to Human Nonalcoholic Steatohepatitis. *Hepatology* **69**, 2241-2257
515 (2019).

516 20. Huang, Y. et al. A feed-forward loop amplifies nutritional regulation of PNPLA3. *Proc Natl*
517 *Acad Sci U S A* **107**, 7892-7897 (2010).

518 21. Lake, A.C. et al. Expression, regulation, and triglyceride hydrolase activity of Adiponutrin
519 family members. *J Lipid Res* **46**, 2477-2487 (2005).

520 22. Dubuquoy, C. et al. Distinct regulation of adiponutrin/PNPLA3 gene expression by the
521 transcription factors ChREBP and SREBP1c in mouse and human hepatocytes. *J Hepatol*
522 **55**, 145-153 (2011).

523 23. Perttilä, J. et al. PNPLA3 is regulated by glucose in human hepatocytes, and its I148M
 524 mutant slows down triglyceride hydrolysis. *Am J Physiol Endocrinol Metab* **302**, E1063-
 525 1069 (2012).
 526 24. de Jong, Y.P. et al. Broadly neutralizing antibodies abrogate established hepatitis C virus
 527 infection. *Sci Transl Med* **6**, 254ra129 (2014).
 528 25. Michailidis, E. et al. Expansion, in vivo-ex vivo cycling, and genetic manipulation of primary
 529 human hepatocytes. *Proc Natl Acad Sci U S A* **117**, 1678-1688 (2020).
 530 26. Azuma, H. et al. Robust expansion of human hepatocytes in Fah^{-/-}/Rag2^{-/-}/Il2rg^{-/-} mice.
 531 *Nat Biotechnol* **25**, 903-910 (2007).
 532 27. Billerbeck, E. et al. Humanized mice efficiently engrafted with fetal hepatoblasts and
 533 syngeneic immune cells develop human monocytes and NK cells. *J Hepatol* **65**, 334-343
 534 (2016).
 535 28. Kawahara, T. et al. Factors affecting hepatocyte isolation, engraftment, and replication in
 536 an in vivo model. *Liver Transpl* **16**, 974-982 (2010).
 537 29. Vanwolleghem, T. et al. Factors determining successful engraftment of hepatocytes and
 538 susceptibility to hepatitis B and C virus infection in uPA-SCID mice. *J Hepatol* **53**, 468-476
 539 (2010).
 540 30. Yeh, M.M. & Brunt, E.M. Pathological features of fatty liver disease. *Gastroenterology* **147**,
 541 754-764 (2014).
 542 31. Grompe, M. et al. Pharmacological correction of neonatal lethal hepatic dysfunction in a
 543 murine model of hereditary tyrosinaemia type I. *Nat Genet* **10**, 453-460 (1995).
 544 32. Bhattacharjee, J. et al. Role of immunodeficient animal models in the development of
 545 fructose induced NAFLD. *J Nutr Biochem* **25**, 219-226 (2014).
 546 33. Sun, G. et al. OX40 Regulates Both Innate and Adaptive Immunity and Promotes
 547 Nonalcoholic Steatohepatitis. *Cell Rep* **25**, 3786-3799.e3784 (2018).

548 34. Liu, X. et al. Lack of mature lymphocytes results in obese but metabolically healthy mice
549 when fed a high-fat diet. *Int J Obes (Lond)* **39**, 1548-1557 (2015).

550 35. Hasegawa, M. et al. The reconstituted 'humanized liver' in TK-NOG mice is mature and
551 functional. *Biochem Biophys Res Commun* **405**, 405-410 (2011).

552 36. Meuleman, P. et al. Morphological and biochemical characterization of a human liver in a
553 uPA-SCID mouse chimera. *Hepatology* **41**, 847-856 (2005).

554 37. Grompe, M. & Strom, S. Mice with human livers. *Gastroenterology* **145**, 1209-1214 (2013).

555 38. Andreo, U. et al. Analysis of Hepatitis C Virus Particle Heterogeneity in Immunodeficient
556 Human Liver Chimeric. *Cell Mol Gastroenterol Hepatol* **4**, 405-417 (2017).

557 39. Bissig-Choisat, B. et al. Development and rescue of human familial hypercholesterolaemia
558 in a xenograft mouse model. *Nat Commun* **6**, 7339 (2015).

559 40. Minniti, M.E. et al. Insights From Liver-Humanized Mice on Cholesterol Lipoprotein
560 Metabolism and LXR-Agonist Pharmacodynamics in Humans. *Hepatology* **72**, 656-670
561 (2020).

562 41. Ma, X. et al. Proportion of NAFLD patients with normal ALT value in overall NAFLD
563 patients: a systematic review and meta-analysis. *BMC Gastroenterol* **20**, 10 (2020).

564 42. Gadd, V.L. et al. The portal inflammatory infiltrate and ductular reaction in human
565 nonalcoholic fatty liver disease. *Hepatology* **59**, 1393-1405 (2014).

566 43. Kleiner, D.E. et al. Design and validation of a histological scoring system for nonalcoholic
567 fatty liver disease. *Hepatology* **41**, 1313-1321 (2005).

568 44. Schuppan, D., Surabattula, R. & Wang, X.Y. Determinants of fibrosis progression and
569 regression in NASH. *J Hepatol* **68**, 238-250 (2018).

570 45. Kane, M. et al. Identification of Interferon-Stimulated Genes with Antiretroviral Activity. *Cell*
571 *Host Microbe* **20**, 392-405 (2016).

572 46. Ito, M. et al. Longitudinal analysis of murine steatohepatitis model induced by chronic
573 exposure to high-fat diet. *Hepatol Res* **37**, 50-57 (2007).

574 47. Naugler, W.E. et al. Fibroblast Growth Factor Signaling Controls Liver Size in Mice With
575 Humanized Livers. *Gastroenterology* **149**, 728-740.e715 (2015).

576 48. Luukkonen, P.K. et al. Saturated Fat Is More Metabolically Harmful for the Human Liver
577 Than Unsaturated Fat or Simple Sugars. *Diabetes Care* **41**, 1732-1739 (2018).

578 49. Mardinoglu, A. et al. An Integrated Understanding of the Rapid Metabolic Benefits of a
579 Carbohydrate-Restricted Diet on Hepatic Steatosis in Humans. *Cell Metab* **27**, 559-
580 571.e555 (2018).

581 50. Tateno, C. et al. Growth hormone-dependent pathogenesis of human hepatic steatosis in
582 a novel mouse model bearing a human hepatocyte-repopulated liver. *Endocrinology* **152**,
583 1479-1491 (2011).

584 51. Yakar, S. et al. Inhibition of growth hormone action improves insulin sensitivity in liver IGF-
585 1-deficient mice. *J Clin Invest* **113**, 96-105 (2004).

586 52. Luukkonen, P.K. et al. Hepatic ceramides dissociate steatosis and insulin resistance in
587 patients with non-alcoholic fatty liver disease. *J Hepatol* **64**, 1167-1175 (2016).

588 53. Cai, J., Zhang, X.J. & Li, H. The Role of Innate Immune Cells in Nonalcoholic
589 Steatohepatitis. *Hepatology* **70**, 1026-1037 (2019).

590 54. Kazankov, K. et al. The role of macrophages in nonalcoholic fatty liver disease and
591 nonalcoholic steatohepatitis. *Nat Rev Gastroenterol Hepatol* **16**, 145-159 (2019).

592 55. Sutti, S. & Albano, E. Adaptive immunity: an emerging player in the progression of NAFLD.
593 *Nat Rev Gastroenterol Hepatol* **17**, 81-92 (2020).

594 56. Ratzl, V. et al. Cenicriviroc Treatment for Adults With Nonalcoholic Steatohepatitis and
595 Fibrosis: Final Analysis of the Phase 2b CENTAUR Study. *Hepatology* (2020).

596 57. Baselli, G.A. et al. Liver transcriptomics highlights interleukin-32 as novel NAFLD-related
597 cytokine and candidate biomarker. *Gut* **69**, 1855-1866 (2020).

598 58. Wilson, E.M. et al. Extensive double humanization of both liver and hematopoiesis in
599 FRGN mice. *Stem Cell Res* **13**, 404-412 (2014).

600 59. Chambers, J.C. et al. Genome-wide association study identifies loci influencing
601 concentrations of liver enzymes in plasma. *Nat Genet* **43**, 1131-1138 (2011).

602 60. Eslam, M., Valenti, L. & Romeo, S. Genetics and epigenetics of NAFLD and NASH: Clinical
603 impact. *J Hepatol* **68**, 268-279 (2018).

604 61. Luukkonen, P.K. et al. Human PNPLA3-I148M variant increases hepatic retention of
605 polyunsaturated fatty acids. *JCI Insight* **4** (2019).

606 62. Tandra, S. et al. Presence and significance of microvesicular steatosis in nonalcoholic fatty
607 liver disease. *J Hepatol* **55**, 654-659 (2011).

608 63. Kozlitina, J. et al. Exome-wide association study identifies a TM6SF2 variant that confers
609 susceptibility to nonalcoholic fatty liver disease. *Nat Genet* **46**, 352-356 (2014).

610 64. Mancina, R.M. et al. The MBOAT7-TMC4 Variant rs641738 Increases Risk of Nonalcoholic
611 Fatty Liver Disease in Individuals of European Descent. *Gastroenterology* **150**, 1219-
612 1230.e1216 (2016).

613 65. Abul-Husn, N.S. et al. A Protein-Truncating HSD17B13 Variant and Protection from
614 Chronic Liver Disease. *N Engl J Med* **378**, 1096-1106 (2018).

615 66. Anzalone, A.V. et al. Search-and-replace genome editing without double-strand breaks or
616 donor DNA. *Nature* **576**, 149-157 (2019).

617 67. Chen, C., Soto-Gutierrez, A., Baptista, P.M. & Spee, B. Biotechnology Challenges to
618 In Vitro Maturation of Hepatic Stem Cells. *Gastroenterology* **154**, 1258-1272 (2018).

619 68. Luce, E., Messina, A., Duclos-Vallée, J.C. & Dubart-Kupperschmitt, A. Advanced
620 techniques and awaited clinical applications for human pluripotent stem cell differentiation
621 into hepatocytes. *Hepatology* (2021).

622 69. Schoggins, J.W. et al. A diverse range of gene products are effectors of the type I interferon
623 antiviral response. *Nature* **472**, 481-485 (2011).

624 70. Santoro, N. et al. A common variant in the patatin-like phospholipase 3 gene (PNPLA3) is
625 associated with fatty liver disease in obese children and adolescents. *Hepatology* **52**,
626 1281-1290 (2010).

627 71. Donati, B. et al. The rs2294918 E434K variant modulates patatin-like phospholipase
628 domain-containing 3 expression and liver damage. *Hepatology* **63**, 787-798 (2016).

629 72. Steensels, S. et al. Acot9 traffics mitochondrial short-chain fatty acids towards de novo
630 lipogenesis and glucose production in the liver. *Hepatology* (2020).

631 73. Hyogo, H., Roy, S., Paigen, B. & Cohen, D.E. Leptin promotes biliary cholesterol
632 elimination during weight loss in ob/ob mice by regulating the enterohepatic circulation of
633 bile salts. *J Biol Chem* **277**, 34117-34124 (2002).

634 74. Alves-Bezerra, M. et al. Thioesterase Superfamily Member 2 Promotes Hepatic VLDL
635 Secretion by Channeling Fatty Acids Into Triglyceride Biosynthesis. *Hepatology* **70**, 496-
636 510 (2019).

637 75. Guilbert, M.C., Therrien, A., Soucy, G., Trudel, D. & Nguyen, B.N. Nodular Regenerative
638 Hyperplasia: Expression Pattern of Glutamine Synthetase and a Potential Role for Hepatic
639 Progenitor Cells. *Appl Immunohistochem Mol Morphol* **28**, 243-248 (2020).

640 76. Schindelin, J. et al. Fiji: an open-source platform for biological-image analysis. *Nat*
641 *Methods* **9**, 676-682 (2012).

642 77. Deroulers, C. et al. Analyzing huge pathology images with open source software. *Diagn*
643 *Pathol* **8**, 92 (2013).

- 644 78. Ritchie, M.E. et al. limma powers differential expression analyses for RNA-sequencing and
645 microarray studies. *Nucleic Acids Res* **43**, e47 (2015).
- 646 79. Raudvere, U. et al. g:Profiler: a web server for functional enrichment analysis and
647 conversions of gene lists (2019 update). *Nucleic Acids Res* **47**, W191-W198 (2019).
- 648 80. Dobin, A. et al. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-21 (2013).
- 649 81. Anders, S., Pyl, P.T. & Huber, W. HTSeq--a Python framework to work with high-
650 throughput sequencing data. *Bioinformatics* **31**, 166-169 (2015).
- 651 82. Liberzon, A. et al. The Molecular Signatures Database (MSigDB) hallmark gene set
652 collection. *Cell Syst* **1**, 417-425 (2015).

661 **FIGURE LEGENDS**

662 **Figure 1: Human hepatocytes in chimeric mice on WD rapidly develop steatosis**

663 **a)** Experimental timeline of FNRG mice that were transplanted with primary human hepatocytes
664 (PHH), cycled off nitisinone to create huFNRG mice, and then subjected to chow or WD for up to
665 12 weeks.

666 **b)** H&E staining on livers from huFNRG mice on chow and after 4 or 12 weeks on WD.

667 **c)** Staining against human nuclear mitotic apparatus-1 in livers of huFNRG mice on chow and 4
668 or 12 weeks on WD. Areas with predominantly mouse hepatocytes are outlined.

669 **d)** Staining against human glutamine synthetase on livers from huFNRG mice on chow and 4
670 weeks on WD. Central vein (CV), portal tract (PT).

671 **e)** Liver steatosis score of the human graft in huFNRG mice on chow and after 4 to 20 weeks on
672 WD. Symbols individual mice, bars are median, unpaired *t*-test with ***p*<0.005, ****p*<0.0005 and
673 *****p*<0.0001.

674 **f)** Hepatic triglyceride levels in livers of huFNRG mice on chow and after 4 to 12 weeks on WD.
675 Symbols individual mice, bars are median, unpaired *t*-test with ****p*<0.0005 and *****p*<0.0001.

676 **g)** H&E staining on livers from muFNRG mice on chow and after 4 or 12 weeks on WD.

677

678 **Suppl Figure S1: Human hepatocyte steatosis in chimeric models**

679 **a)** Serum human albumin (hAlb) levels in huFNRG mice during the course of chow and WD
680 feeding.

681 **b)** Oil Red O staining on livers from huFNRG mice on chow and after 4 weeks on WD.

682 **c)** Hepatic cholesterol quantification from huFNRG mice on chow and after 4 to 12 weeks on WD.
683 Symbols individual mice, bars are median, unpaired *t*-test with **p*<0.05, ***p*<0.005.

684 **d)** Hepatic triglyceride levels versus clinical steatosis scoring of the human graft in huFNRG mice
 685 on chow and after 4 to 12 weeks on WD. Symbols individual mice, line is simple linear regression,
 686 r^2 Pearson correlation coefficient.

687 **e)** H&E staining on livers from NRG mice on chow and after 12 weeks on WD.

688 **f)** Hepatic triglyceride quantification from muFNRG mice on chow, and after 4 and 12 weeks on
 689 WD. Symbols individual mice, bars are median.

690 **g)** Liver sections from PHH2 huFNRG mice on chow and after 10 weeks on WD stained for human
 691 nuclear mitotic apparatus-1 (left). Liver sections from PHH2 huTK-NOG mice on chow and after
 692 10 weeks on WD stained for H&E (right).

693 **h)** H&E staining on livers from PHH3 huFNRG and PHH3 hu-uPA/SCID mice livers. High
 694 magnification images of a huFNRG mouse liver on chow (upper left); after 4 weeks on WD (lower
 695 left); low and high magnification images of a hu-uPA/SCID mouse liver on chow (right).

697 **Figure 2: Systemic metabolic effects of WD in chimeric FNRG mice**

698 **a)** Body weight in huFNRG mice during time course of chow or WD feeding. Symbols individual
 699 mice, line is mean, two-way ANOVA Sidak pairwise comparison * $p < 0.05$, ** $p < 0.005$, *** $p < 0.0005$.

700 **b)** Gonadal fat fraction of total body weight in huFNRG mice on chow and after 4 to 20 weeks on
 701 WD. Symbols individual mice, bars are median, unpaired t -test with * $p < 0.05$, ** $p < 0.005$,
 702 *** $p < 0.0005$.

703 **c)** Cholesterol in plasma lipoprotein fractions from huFNRG and muFNRG mice on chow and
 704 after 4 weeks on WD. Pooled plasma (3 mice/group), symbols are technical means.

705 **d)** Triglycerides in plasma lipoprotein fractions from huFNRG and muFNRG mice on chow and
 706 after 4 weeks on WD. Pooled plasma (3 mice/group), symbols are technical means.

707 **e)** Fasting blood glucose from huFNRG and muFNRG mice on chow and after 4 weeks on WD.
 708 Symbols individual mice, lines are median, unpaired t -test **** $p < 0.0001$.

709 **f)** Fasting serum mouse insulin from huFNRG and muFNRG mice on chow and after 4 weeks on
 710 WD. Symbols individual mice, lines are median, unpaired *t*-test with **p*<0.05, ***p*<0.005.
 711 **g)** Intraperitoneal glucose, insulin and pyruvate tolerance testing in huFNRG and muFNRG mice
 712 on chow and after 4 weeks on WD. Blood glucose values measured at baseline (0) and at
 713 indicated timepoints after i.p. injections with each agent. Symbols are mean ± SEM of 5-7 mice
 714 per group, two-way ANOVA Sidak pairwise comparison of huFNRG on chow and WD (p-values
 715 on top) or muFNRG on chow and WD (p-values in bottom), **p*<0.05, ***p*<0.005, ****p*<0.0005 and
 716 *****p*<0.0001

717

718 **Suppl Figure S2: Systemic metabolic effects of WD feeding**

719 **a)** Cholesterol quantification in plasma lipoprotein fractions from huFNRG and muFNRG mice on
 720 chow and after 4 weeks on WD. Pooled plasma (3 mice/group), column bars are technical means
 721 ± SEM.
 722 **b)** Triglyceride quantification in plasma lipoprotein fractions from huFNRG and muFNRG mice on
 723 chow and after 4 weeks on WD. Pooled plasma (3 mice/group), column bars are technical means
 724 ± SEM.
 725 **c)** Intraperitoneal glucose tolerance testing in huFNRG on chow and after 8 weeks on WD.
 726 Symbols are mean ± SEM of 5-8 mice per group, two-way ANOVA Sidak pairwise comparison
 727 with **p*<0.05, ***p*<0.005, ****p*<0.0005 and *****p*<0.0001

728 **d)** Mouse insulin secretion during glucose tolerance testing in huFNRG mice on chow and after
 729 8 weeks on WD. Symbols are mean ± SEM of 5-8 mice per group.

730

731 **Figure 3: huFNRG mice on WD develop mild steatohepatitis**

732 **a)** ALT and AST activity in serum from huFNRG mice on chow and after 4 to 12 weeks on WD.
 733 Symbols individual mice, bars are median, unpaired *t*-test ***p*<0.005, *****p*<0.0001.

734 **b)** H&E staining from a huFNRG mouse liver after 8 weeks on WD. Black arrow indicates
 735 hepatocyte ballooning degeneration and white arrow indicates lobular inflammation.

736 **c)** Staining against human nuclear mitotic apparatus-1 in the liver of a huFNRG mouse after 4
 737 weeks on WD. Black arrow indicates hepatocyte ballooning.

738 **d)** Fractions of huFNRG mice with human hepatocyte ballooning degeneration on chow and WD
 739 over time. Number of mice with ballooning per group at bottom of bars.

740 **e)** Fractions of huFNRG mice with human hepatocyte ballooning degeneration by grade of
 741 steatosis. Number of mice with ballooning per group at bottom of bars.

742 **f)** NAFLD activity score (NAS) in the human graft of huFNRG mice on chow and after 4 to 20
 743 weeks on WD. Symbols individual mice, bars are median, unpaired *t*-test ***p*<0.005,
 744 *****p*<0.0001.

745 **g)** Picosirius Red staining for collagen in livers from huFNRG mice on chow and after 4 to 20
 746 weeks on WD.

747 **h)** Fibrosis stages in the human graft of huFNRG liver on chow and after 4 to 20 weeks on WD.
 748 Mouse numbers at bottom of bars.

749

750 **Suppl Figure S3: Mild steatohepatitis in huFNRG mice on WD**

751 **a)** Human ALT protein versus ALT activity in serum of huFNRG mice on chow and WD. Symbols
 752 individual mice, line is simple linear regression, r^2 Pearson correlation coefficient.

753 **b)** ALT and AST activity in serum of muFNRG mice on chow and after 4 to 12 weeks on WD.
 754 Symbols individual mice, bars are median, unpaired *t*-test with ***p*<0.005, *****p*<0.0001.

755 **c)** H&E staining on livers from huFNRG mice after 4, 12 and 20 weeks on WD. Black arrow
 756 indicates hepatocyte ballooning degeneration and white arrow indicated lobular inflammation.

757 **d)** H&E staining on liver from a huFNRG mouse engrafted with PHH3 after 4 weeks on WD. Black
 758 arrow indicates hepatocyte ballooning degeneration.

759 **e)** Fractions of huFNRG mice with lobular inflammation on chow and after 4 to 20 weeks on WD.
760 Number of mice with lobular inflammation per group at bottom of bars.
761 **f)** Masson trichrome staining for collagen on livers from huFNRG mice on chow and after 12 and
762 20 weeks on WD.
763 **g)** Picosirius Red staining for collagen on a liver of huFNRG mouse engrafted with PHH2 after 10
764 weeks on WD.

765

766 **Figure 4: Transcriptional changes in huFNRG mice on WD**

767 **a)** Volcano plots of liver transcripts mapped to human genome (human reads) comparing
768 huFNRG mice 4 weeks on WD to chow, and longer WD durations to the previous WD timepoint.
769 RNA-sequencing of 3-4 mice per group, red symbols FDR <0.05, grey denotes not significant.
770 **b)** Volcano plots of liver transcripts mapped to mouse genome (mouse reads) comparing
771 huFNRG mice 4 weeks on WD to chow, and longer WD durations to the previous WD timepoint.
772 RNA-sequencing of 3-4 mice per group, red symbols FDR <0.05, grey denotes not significant.
773 **c)** Gene-ontology (GO) pathway analysis of human reads from livers of huFNRG mice 4 weeks
774 on WD to chow, and longer WD durations to the previous WD timepoint. Blue symbols are down-
775 and red symbols are upregulated pathways, grey denotes no significant changes, symbol size
776 indicates statistical significance.
777 **d)** Gene-ontology (GO) pathway analysis of mouse reads from livers of huFNRG mice 4 weeks
778 on WD to chow, and longer WD durations to the previous WD timepoint. Blue symbols are down-
779 and red symbols are upregulated pathways, grey denotes no significant changes, symbol size
780 indicates statistical significance.

781

782 **Suppl. Figure S4: Transcriptional changes in huFNRG mice on WD**

783 **a)** Spliced Transcripts Alignment to a Reference (STAR) Scores of transcripts from fasting
784 huFNRG mice livers on chow and after 4 to 20 weeks on WD aligned to human reference genome.
785 Numbers on left indicate individual mouse tags.

786 **b)** Manhattan plots comparing human pathways in livers from huFNRG mice 4 weeks on WD to
787 chow, and longer WD durations to the previous WD timepoint. Gene-ontology (GO) with GO-MF
788 (Molecular Function), GO-CC (Cellular Component), GO-BP (Biological Process), Kyoto
789 Encyclopedia of Genes and Genomes (KEGG), REACTOME (REAC) and Transcription Factor
790 (TF). Symbol size indicates percentage of statistically significant differently expressed transcripts
791 aligned to a certain pathway, only pathways statistically significant enriched for a group of mice
792 are displayed.

793 **c)** Manhattan plots comparing mouse pathways in livers from huFNRG mice 4 weeks on WD to
794 chow, and longer WD durations to the previous WD timepoint. Gene-ontology (GO) with GO-MF
795 (Molecular Function), GO-CC (Cellular Component), GO-BP (Biological Process), Kyoto
796 Encyclopedia of Genes and Genomes (KEGG), REACTOME (REAC) and Transcription Factor
797 (TF). Symbol size indicates percentage of statistically significant differently expressed transcripts
798 aligned to a certain pathway, only pathways statistically significant enriched for a group of mice
799 are displayed.

800

801 **Figure 5) PNPLA3-148M huFRG mice on WD develop steatohepatitis**

802 **a)** H&E staining on livers from 148M-huFRG mice on chow and after 4 weeks on WD.

803 **b)** Clinical steatosis score of the human graft on livers from 148M-huFNRG mice on chow and
804 after 4 weeks WD. Symbols are individual mice, bars median, unpaired *t*-test ****p*<0.0005.

805 **c)** H&E staining on livers from 148M-huFRG mice 4 weeks on WD. Grey arrow indicates Mallory
806 Denk body, black arrow ballooning degeneration and white arrow lobular inflammation.

807 **d)** NAFLD Activity Score (NAS) in the human graft of 148M-huFRG mice on chow and after 4
808 weeks WD. Symbols individual mice, bars are median, unpaired *t*-test ****p*<0.0005.

809 **e)** Volcano plots of liver transcripts mapped to human genome (human reads) comparing 148M-
810 huFRG mice after 4 weeks WD vs. chow. RNA-sequencing of 3-4 mice per group, red symbols
811 FDR <0.05, grey denotes not significant.

812 **f)** Gene-ontology (GO) pathway analysis of human reads comparing 148I-huFNRG and 148M-
813 huFRG livers after 4 weeks WD vs. chow. Blue symbols are down- and red symbols are
814 upregulated pathways, grey denotes no significant changes, symbol size indicates statistical
815 significance.

816 **g)** Volcano plots of human reads differentially expressed after 4 weeks WD compared to chow (Δ
817 WD) in 148M-huFRG livers versus 148I-huFNRG livers. Red symbols FDR <0.05, grey denotes
818 not significant.

819

820 **Suppl Figure S5: 148M-huFRG mice develop steatohepatitis**

821 **a)** Fasting blood glucose from 148M-huFRG mice on chow and after 4 weeks on WD. Symbols
822 individual mice, bars are median, unpaired *t*-test, ***p*<0.005.

823 **b)** Fasting serum mouse insulin in 148M-huFRG mice on chow and after 4 weeks on WD.
824 Symbols individual mice, bars are median.

825 **c)** Fasting plasma triglycerides in 148M-huFRG mice on chow and after 4 weeks on WD.
826 Symbols individual mice, bars are median.

827 **d)** Hepatic triglyceride quantification in 148M-huFRG mice on chow and after 4 weeks on WD.
828 Symbols individual mice, lines are median, unpaired *t*-test, ***p*<0.005.

829 **e)** Left: Fractions of huFRG mice with human hepatocyte ballooning degeneration on chow and
830 4 weeks on WD. Number of mice with ballooning per group at bottom of bars. Right: Ballooning

831 score in the human graft of 148M-huFRG mice on chow and after 4 weeks of WD. Symbols
832 individual mice, bars are median, unpaired *t*-test ***p*<0.005.

833 **f)** Left: Fractions of 148M-huFRG mice with lobular inflammation on chow and 4 weeks on WD.
834 Number of mice with lobular inflammation per group at bottom of bars. Right: Lobular inflammation
835 score in the human graft of 148M-huFRG mice on chow and after 4 weeks of WD. Symbols
836 individual mice, bars are median, unpaired *t*-test.

837 **g)** Volcano plots showing differently expressed transcripts mapped to human reference genome
838 (human reads) from 148M-huFRG versus 148I-huFNRG livers on chow (left) and after 4 weeks
839 on WD (right). Red symbols FDR <0.05, grey denotes not significant.

840

841 **Figure 6: PNPLA3-148M overexpression in hepatocytes exacerbates steatosis in mice on**
842 **WD**

843 **a)** Experimental timeline of creating PNPLA3 overexpressing mice and diet challenge. Mouse
844 passaged primary human hepatocytes (PHH) cultures were transduced with lentiviral vectors
845 expressing PNPLA3 variants and re-transplanted. Following humanization, huFNRG with
846 PNPLA3 transduced hepatocytes (td) were exposed to 4 weeks of WD.

847 **b)** H&E staining on livers from tdRFP, td148I and td148M mice after 4 weeks on WD.

848 **c)** Liver steatosis score of the human graft in tdRFP, td148I and td148M mice on chow and after
849 4 weeks on WD. Symbols individual mice, bars are median, unpaired *t*-test with ***p*<0.005,
850 ****p*<0.0005 and *****p*<0.0001.

851 **d)** Hepatic triglyceride quantification in tdRFP, td148I and td148M mice on chow and after 4 weeks
852 on WD. Symbols individual mice, bars are median, unpaired *t*-test, **p*<0.05.

853 **e)** NAFLD Activity Score (NAS) in the human graft of tdRFP, td148I and td148M mice on chow
854 and after 4 weeks on WD. Symbols individual mice, bars are median, unpaired *t*-test, **p*<0.05,
855 ***p*<0.005, ****p*<0.0005.

856 f) Volcano plot of 80 human genes expressed in livers of td148I versus td148M mice after 4 weeks
857 on WD. Expression by qRT-PCR normalized to 11 housekeeping genes, n=3-4 mice per group,
858 burgundy symbols $p < 0.05$, grey symbols not statistically significant.

859

860 **Suppl Figure S6: PNPLA3-148M overexpression in hepatocytes exacerbates steatosis in**
861 **mice on WD**

862 a) Serum human albumin (hAlb) levels in FNRG mice after transplantation with transduced PHH.
863 Symbols are mean $\pm$ SEM of 3-16 mice.

864 b) human PNPLA3 mRNA levels in livers of tdRFP, td148I and td148M mice on chow and 4 weeks
865 on WD. Expression by qRT-PCR normalized to 11 housekeeping genes. Symbols individual mice,
866 bars are median, unpaired *t*-test, ** $p < 0.005$, *** $p < 0.0005$ and **** $p < 0.0001$.

867 c) H&E staining on livers from tdRFP, td148I and td148M mice on chow. White arrow indicates
868 macrovesicular and black arrow microvesicular steatosis.

869 d) Left: Fractions of tdRFP, tdr148I and td148M mice with human hepatocyte ballooning
870 degeneration on chow and 4 weeks on WD. Number of mice with ballooning per group at bottom
871 of bars. Right: Ballooning score in the human graft on chow and after 4 weeks of WD. Symbols
872 individual mice, bars are median, unpaired *t*-test ** $p < 0.005$ and **** $p < 0.0001$

873 e) Left: Fractions of tdRFP, tdr148I and td148M mice with lobular inflammation on chow and 4
874 weeks on WD. Number of mice with lobular inflammation per group at bottom of bars. Right:
875 Lobular inflammation score in the human graft on chow and after 4 weeks of WD. Symbols
876 individual mice, bars are median, unpaired *t*-test.

877 f) Volcano plot of 80 human genes expressed in livers of td148I versus td148M mice on chow.
878 Expression by qRT-PCR, normalized to 11 housekeeping genes, n=3-4 mice per group, green
879 symbols $p < 0.05$, grey symbols not statistically significant.

880

881 **Figure 7: A HFD causes more active steatohepatitis in PNPLA3-148M overexpressing mice**

- 882 **a)** H&E staining on livers from tdRFP, td148I and td148M mice after 4 weeks on HFD.
- 883 **b)** Liver steatosis score of the human graft in tdRFP, td148I and td148M mice on chow and after
884 4 weeks on HFD. Symbols individual mice, bars are median, unpaired *t*-test with ***p*<0.005.
- 885 **c)** H&E staining on liver from a td148M mouse after 4 weeks of HFD. White arrow indicates lobular
886 inflammation, black arrow microvesicular steatosis and grey arrow Mallory-Denk body.
- 887 **d)** NAFLD Activity Score (NAS) in the human graft of tdRFP, td148I and td148M mice on chow
888 and after 4 weeks on HFD. Symbols individual mice, bars are median, unpaired *t*-test, **p*<0.05.
- 889 **e)** Low and high magnification fluorescent images of livers from td148I and td148M mice after 4
890 weeks of HFD stained for neutral lipids (green, BODIPY) and nuclei (blue, DAPI).
- 891 **f)** Two-dimensional density plots of neutral lipids (BODIPY) and RFP of 355,161 small
892 hepatocytes in livers from four td148I mice and 183,289 small hepatocytes in livers from three
893 td148M mice after 4 weeks of HFD. Density plots are displaying RFP in X-, and BODIPY
894 intensities in Y-Axis, both in arbitrary units (AU). Dashed lines display cell population group
895 quadrants, with lower left quadrant harboring double negative and upper right quadrant harboring
896 double positive cells. Solid line is simple linear regression with equation displayed underneath,
897 r^2 is Pearson correlation coefficient.

898
899 **Supplement Figure S7: A HFD causes more active steatohepatitis in PNPLA3-148M**
900 **overexpressing mice**

- 901 **a)** Hepatic triglyceride quantification in td148I and td148M mice after 4 weeks on HFD. Symbols
902 individual mice, bars are median, unpaired *t*-test, **p*<0.05.
- 903 **b)** Left: Fractions of tdRFP, tdr148I and td148M mice with human hepatocyte ballooning
904 degeneration 4 weeks on HFD. Number of mice with ballooning per group at bottom of bars.

Right: Ballooning score in the human graft and after 4 weeks of HFD. Symbols individual mice, bars are median, unpaired *t*-test.

c) Left: Fractions of tdRFP, td148I and td148M mice with lobular inflammation 4 weeks on HFD. Number of mice with lobular inflammation per group at bottom of bars. Right: Lobular inflammation score in the human graft after 4 weeks of HFD. Symbols individual mice, bars are median, unpaired *t*-test.

d) human PNPLA3 mRNA levels in livers of td148I and td148M mice 4 weeks on HFD. Expression by qRT-PCR normalized to 11 housekeeping genes. Symbols individual mice, bars are median, unpaired *t*-test

e) H&E staining on liver of non-transduced huFNRG mouse with severe steatosis (left). Right panel shows two-dimensional density plot of neutral lipids (BODIPY) and RFP of 54,249 small hepatocytes in liver from one mouse displayed on left. Densitograms are displaying RFP in X-, and BODIPY intensities in Y-Axis, both in arbitrary units (AU). Lower left quadrant are double negative, upper right quadrant are double positive cells. Number in corners display fraction of total cell population.

TABLE LEGENDS

Table 1) qRT-PCR mRNA expression in transduced huFNRG animals - average fold change over housekeeping

Shown are fold changes of the expression of 80 human genes over 11 housekeeping genes (GAPDH, HPRT1, GUSB, ACTB, B2M, HMBS, IPO8, PGK1, RPLP0, TBP and TFRC) from livers of tdRFP, td148I and td148M mice on chow or on WD. Expression by qRT-PCR, n=3-4 mice per group.

929 **ACKNOWLEDGEMENTS**

930 This work was supported by the National Institutes of Health Grants R37DK048873,
931 R01DK056626 and R01DK103046 (to DEC), R01DK085713 (to CMR), K08DK090576 and
932 R01AA027327 (to YPJ) and the Starr Foundation (to CMR). PM was supported by grants from
933 Ghent University (Special Research Fund – ICOH expert center) and the Research Foundation –
934 Flanders (grants VirEOS 30981113 and G047417N). MK received funding from the Deutsche
935 Forschungsgemeinschaft under grant number KA4688/1-1. The project was co-sponsored by the
936 Center for Basic and Translational Research on Disorders of the Digestive System through the
937 generosity of the Leona M. and Harry B. Helmsley Charitable trust (to EM). The NYULH Center
938 for Biospecimen Research and Development, Histology and Immunohistochemistry Laboratory
939 (RRID:SCR_018304) is supported in part by the Laura and Isaac Perlmutter Cancer Center
940 Support Grant; NIH/NCI P30CA016087. The RU Center for Clinical and Translational Science
941 Bioinformatics program is supported by the Clinical and Translational Award (CTSA) and the
942 National Center for Advancing Translational Sciences (NCATS), part of the National Institutes of
943 Health. We thank Branka Brukner Dabovic (NYULH – Department of Cell Biology Research) for
944 help with imaging fluorescence samples. We thank the Rockefeller University High-Throughput,
945 Bioimaging and Genomics Resource Centers.

946
947

MATERIAL AND METHODS

Mice

Male and female *Fah*^{-/-} NOD *Rag1*^{-/-} *Il2rg*^{null} (FNRG) mice²⁴ were preconditioned with retrorsine²⁵ and male *Fah*^{-/-} *Rag2*^{-/-} *Il2rg*^{-/-} (FRG) mice²⁶ with adenoviral urokinase plasminogen activator (uPA) before transplantation with primary human hepatocytes (PHH). NOD *Rag1*^{-/-} *Il2rg*^{null} (NRG) mice were obtained from Jackson Labs (Bar Harbor, ME) and received retrorsine and nitisinone cycling similarly to FNRG mice. Seven week old male thymidine kinase transgenic mice on the NOD SCID *Il2rg*^{null} background (TK-NOG)³⁵ were obtained from Taconic Biosciences (Germantown, NY) and preconditioned with retrorsine²⁵ and 48 hours 0.1mg/ml valganciclovir (Sigma) in drinking water 14 days prior to transplantation. Male transgenic uPA^{+/+}/SCID mice were transplanted as previously described.³⁶ The following PHH donors were used: cryopreserved or mouse-passaged (mp)PHH1 (HUM4188, Lonza), cryopreserved or mpPHH2 (HUM4129, Lonza), cryopreserved PHH3 (HFCP940, BD Biosciences) and cryopreserved PHH4 (HHF13022, Yecuris). Hepatocytes were genotyped only for the PNPLA3 rs734809 variant. Genotyping was performed by Sanger sequencing using forward 5'-GCCCTGCTCACTTGGAGAAA-3' and reverse primer 5'-TGAAAGGCAGTGAGGCATGG-3' as reported.⁷⁰ To make PNPLA3 overexpressing mice, PNPLA3-148I or -148M (both containing the K434E variant⁷¹) were cloned into pSCRPSY⁶⁹ (GenBank: KT368137.1) to generate VSVg pseudotyped lentiviral particles.²⁵ Driven by HIV-LTR promoter, pSCRPSY allows a gene of interest (in our case either PNPLA3 variants or none in "tdRFP" controls) to be expressed from the completely spliced "early" HIV-1 mRNA, while TagRFP (a surrogate marker for transduction) is expressed from an unspliced late HIV-1 mRNA.⁴⁵ mpPHH1 were isolated from huFNRG, plated in 6 well collagen-coated plates and transduced with pSCRPSY vectors by 1000g spinoculation. Three to 5 days after transduction ~5x10⁵ cells/mouse were retransplanted into retrorsine preconditioned FNRG animals as described.²⁵ Mice were housed in a 12-hour light/dark cycle at 21°C and 50% humidity. All

973 experiments were conducted under animal use protocols approved by Rockefeller University,
974 Ghent University and Yecuris corporation. huFNRG and huTK-NOG mice were humanized at
975 Rockefeller University, huFRG mice at Yecuris corporation and hu-uPA/SCID mice at Ghent
976 University.

977

978 **Diet challenges**

979 One week prior and during dietary studies, huFNRG, huFRG, NRG and muFNRG mice were
980 changed from oral nitisinone (CuRx, Yecuris cat#20-0028) to intraperitoneal (i.p.) injections at
981 1mg/kg body weight for three days every 2 weeks. Injection solution was 0.5mg/ml nitisinone
982 dissolved in 500mM NaHCO₃ pH 7.4 and diluted 1:10 in PBS-/- (Gibco, Thermo Fisher) to final
983 concentration of 50µg/ml. During humanization or 'murinization' the FNRG, FRG and TK-NOG
984 lines were maintained on an amoxicillin containing chow diet (modified PicoLab Mouse Diet 20,
985 5058 - 5B1Q, 0.12% Amoxicillin, irradiated; TestDiet, Richmond, IN) and autoclaved water. One
986 week prior dietary challenge, amoxicillin medicated chow was replaced with regular chow diet
987 (PicoLab Rodent Diet 20, 5053, irradiated; LabDiet, Fort Worth, TX) lacking amoxicillin. For High
988 Fat Diet (HFD) studies, mice were given a diet with 60% kcal fat, 20% carbohydrates and 20%
989 protein (D12492i irradiated; Research Diets, New Brunswick, NJ) and autoclaved water. For
990 Western-style Diet (WD) studies, HFD was combined with 10% w/v sucrose (Fisher Scientific
991 cat#S5-3; Fair Lawn, NJ) in autoclaved water. Sucrose drinking water was freshly made once a
992 week. uPA^{+/+}/SCID mice were fed irradiated standard breeding chow (Mouse Breeding complete
993 feed for mice (GE 16.7MJ/kg; 14.0 MJ/kg), Ssniff Spezialdiäten GmbH, Soest, Germany). All diets
994 and drinking water were given *ad libitum*. Diet challenges in huFNRG, huFRG, NRG, muFNRG
995 and huTK-NOG mice were done at Rockefeller University.

996

997 **Blood and tissue harvest**

998 Serum was obtained from tail veins, retroorbital or submental venipuncture. Human albumin was
999 quantified as previously described²⁴ and serum mouse insulin quantified using Ultra-Sensitive
1000 Mouse Insulin ELISA Kit (Crystal Chem cat#90080, Elk Grove Village, IL) following
1001 manufacturer's instructions. In mice with hAlb serum levels >8mg/ml, ALT and AST activity
1002 (Elabscience cat#E-BC-K235 and K236, China), which is capable to detect mouse activity as per
1003 manufacturer, and human ALT protein (Abcam cat#ab234578) were determined following
1004 manufacturers instruction. Serum was obtained from submental venipuncture 2 to 3 days after
1005 last nitisinone injection cycle to minimize mouse transaminase activity. Tolerance tests to glucose
1006 (GTT), insulin (ITT) and pyruvate (PTT) were performed in FNRG mice in the weeks with no NTBC
1007 cycling injections similar as described.⁷² Briefly, age matched mice 4 to 12 weeks after dietary
1008 challenge with WD or chow were fasted with free access to water for 6 h prior to GTT and ITT
1009 and for 16 h prior to PTT. Approximately 5µl blood was collected from tail bleed before and at
1010 indicated intervals for up to 180 min following i.p. injection with 2.5 g/kg body weight glucose
1011 (Sigma Aldrich cat#G7021), 1 U/kg body weight human insulin analogue (Eli Lilly, Humulin R U-
1012 100), or 2 g/kg body weight pyruvate (Sigma Aldrich cat#P2256), each dissolved in isotonic PBS-
1013 /- (Gibco, Thermo Fisher Scientific). Blood glucose concentrations were determined using a
1014 GE100 Blood Glucose Monitor (GE, Ontario, CA). Prior to terminal blood and tissue collection (2
1015 to 3 days after last nitisinone injection in *Fah*^{-/-} strains), mice were fasted for 6 hours during light
1016 cycle with access to water. Blood was collected by submental and/or retroorbital venipuncture
1017 and tissue samples were weighed and stored in fixative or frozen on dry-ice and stored at -80°C.
1018 Blood was centrifuged with 16,000 x *g* for 15 min at 4°C and plasma was collected and stored at
1019 -80°C. Hepatic lipids were extracted from frozen liver tissue as described⁷² and liver and plasma
1020 TG, NEFA, total and free cholesterol, and phospholipids were quantified according to
1021 manufacturer's instructions (Wako Diagnostics, Mountain View, CA). Equal volumes of plasma
1022 were pooled from three mice and lipoproteins were fractionated by fast protein liquid

1023 chromatography (ÄKTA pure FPLC system, GE Healthcare, Pittsburgh, PA, USA) and triglyceride
1024 and cholesterol in fractions quantified with reagent kits (Wako Diagnostics, Mountain View, CA).⁷³
1025 ⁷⁴
1026

1027 **Liver histology and quantification**

1028 Neutral buffered formalin-fixed (Thermo Fisher Scientific cat#5725) paraffin-embedded liver
1029 samples were sectioned at 5µm onto charged slides (Fisher Scientific cat#22-042-924). Slides
1030 were dried for 1 hour at 60°C, deparaffinized in xylene, rehydrated through a graded series of
1031 ethanol, and rinsed in distilled water for 5 minutes. Sections were then stained with Hematoxylin
1032 (Richard-Allan Scientific cat #7211, USA) and Eosin (Leica cat#3801619) using our routine
1033 laboratory method described in [Carson, F.L., *Histotechnology: a self instructional text*. 2009,
1034 Chicago: ASCP Press., 114-123], and with Picro Sirius Red Stain (PSR) (Abcam cat#ab150681)
1035 and Masson's Trichrome (Newcomer Supply cat#9179A, USA) in accordance with manufacturer's
1036 supplied procedures. For chromogenic immunohistochemistry staining, unconjugated, polyclonal
1037 rabbit anti-Human Nuclear Mitotic Apparatus protein (Abcam Cat# 97585 Lot# GR268490 RRID
1038 AB_10680001),²⁷ and unconjugated, mouse anti-Human Glutamine Synthetase (Ventana
1039 Medical Systems Lot# V0001337 RRID AB_2861318) clone GS-9⁷⁵ were used.⁷⁵ Slides were
1040 scanned using a Leica Aperio AT2 System and digitally archived via eSlide Manager. To preserve
1041 lipids with and without fluorescent proteins for histopathological analysis, fresh liver samples
1042 haven been fixed in 4% paraformaldehyde (Electron Microscopy Sciences) for 18 hours,
1043 cryoprotected at 4°C with 10% and 18% sucrose w/v for 24 hours each before OCT (Sakura
1044 Finetek) embedding and storage at -80°C. OCT-embedded frozen specimens were cryosectioned
1045 at 5 µm onto charged slides (Fisher Scientific cat# 22-042-924). Slides were air-dried at room
1046 temperature overnight and fixed in 10% neutral buffered formalin for 1 hour at room temperature.
1047 Slides were then rinsed in distilled water, stained with BODIPY (Invitrogen cat# D3922) and Oil

1048 Red O (Sigma-Aldrich cat# O1391). For BODIPY staining, slides were incubated in Reaction
1049 Buffer (Ventana cat# 950-300) for 10 minutes and stained with BODIPY (1:1000 dilution) and
1050 DAPI (1:1000 dilution) in PBS for 60 minutes, and covered using ProLong™ Diamond Antifade
1051 Mountant (Invitrogen, cat# P36961). BODIPY, RFP, and DAPI were imaged on a Hamamatsu
1052 Nanozoomer 2.0 HT and digitally archived via SlidePath. For Oil Red O staining, sections were
1053 stained in Oil Red O solution for 10 minutes, rinsed with distilled water for 5 minutes,
1054 counterstained with hematoxylin (Richard-Allan Scientific cat# 7211,) for 1 minute and rinsed in
1055 distilled water for 1 minute. Samples were blued in 1.2% ammonium hydroxide solution for 1
1056 minute, rinsed in distilled water for 1 minute, and coverslipped using a 50% glycerol solution.
1057 Slides were scanned using a Leica Aperio AT2 System and digitally archived via eSlide Manager.
1058 For fluorescence image analyses SVS image files derived from Hamamatsu Nanozoomer 2.0 HT
1059 were extracted and saved as uncompressed multiple 40x magnification tile images in TIFF format
1060 using FIJI (<https://imagej.net/Fiji/Downloads>)⁷⁶ and NDPITools plugin.⁷⁷ TIFF files of tile images
1061 were split into fluorescence channels (DAPI, BODIPY or RFP) and saved as 8-bit grayscale
1062 images. Masks were generated around human hepatocytes nuclei approximated by diameter size
1063 in DAPI images, enlarged three folds to approximate the size of human hepatocytes in chimeric
1064 livers and used for measuring BODIPY and RFP signal intensities. RFP and BODIPY intensities
1065 per approximated human hepatocyte were plotted as two-dimensional density plots using Python.
1066 Histological scores were conducted from H&E stained samples by clinical pathologist (CF and
1067 MP), using the NAS system from NASH-CRN.⁴³ Steatosis grade (0 :< 5%, 1: 5-33%, 2: >33-66%
1068 and 3:> 66%), lobular inflammation (0: no foci, 1: <2 foci / 200x field, 2: 2-4 foci, 3: >4 foci) and
1069 ballooning degeneration (0: none, 1: few ballooning cells, 2: many cells / prominent ballooning)
1070 were used to generate an overall score (NAS). Fibrosis staging was evaluated separately in PSR
1071 stained samples using the same system,⁴³ with the following scores 0: none, 1: perisinusoidal or

1072 periportal, 1A: mild, zone 3, perisinusoidal, 1B: moderate, zone 3, perisinusoidal, 1C:
1073 portal/periportal.

1074

1075 **Liver tissue transcriptional analyses**

1076 Around 25 mg of freshly harvested liver tissue was placed into 1 ml 4°C TRIzol (Life Technologies,
1077 USA) with 500µl of 1 mm glass beads (BioSpec Products cat# 11079110) and homogenized using
1078 MagNA lyser (Roche, USA), weighed and stored at -80°C. For RNA extraction, 500µl of liver/Trizol
1079 mixture was used for chloroform extraction using MaXtract™ (Qiagen, cat# 129056) and
1080 RNeasy® Mini Kit (Qiagen, cat# 74104) coupled with on-column DNaseI treatment (Qiagen
1081 cat#79254) following manufacturer's instructions. Extracted RNA was stored at -80°C. For qRT-
1082 PCR, cDNA was generated from liver RNA using SuperScript IV VILO (Thermo Fisher Scientific
1083 cat#11766050) following manufacturer's instructions. qRT-PCR was performed in 96-well Human
1084 Fatty Liver Array plates (Thermo Fisher Scientific cat # 4391524) on a QuantStudio 3 System
1085 (Applied Biosystem) for the following genes: *ABCA1, ACACA, ACADL, ACLY, ACOX1, ACSL5,*
1086 *ACSM3, ADIPOR1, ADIPOR2, AKT1, APOA1, APOB, APOC3, APOE, ATP5C1, CASP3, CD36,*
1087 *CEBPB;CEBPB-AS1, CNBP, CPT1A, CPT2, CYP2E1, CYP7A1, DGAT2, FABP1, FABP3,*
1088 *FABP5, FAS, FASN, FOXA2, FOXO1, G6PC, G6PD, GCK, GK, GSK3B, HMGCR, HNF4A, IGF1,*
1089 *IGFBP1, IL1B, INSR, IRS1, LDLR, LEPR, LPL, MAPK1, MAPK8, MLXIPL, MTOR, NDUFB6,*
1090 *NFKB1, NR1H2, NR1H3, NR1H4, PCK2, PDK4, PIK3CA, PIK3R1, PKLR, PNPLA3, PPA1,*
1091 *PPARA, PPARG, PPARGC1A, PRKAA1, PTPN1, RBP4, RXRA, SCD, SERPINE1, SLC27A5,*
1092 *SLC2A1, SLC2A2, SLC2A4, SOCS3, SREBF1, SREBF2, STAT3, XBP1.* Transcriptional changes
1093 were calculated over mean of housekeeping genes: *GAPDH, HPRT1, GUSB, ACTB, B2M,*
1094 *HMBS, IPO8, PGK1, RPLP0, TBP, TFRC.* Data were analyzed and graphed in R (version 4.0.2).
1095 For bulk RNA sequencing (RNAseq) the quantity and integrity of extracted liver RNA were
1096 assessed using an Agilent 2100 Bioanalyzer (Agilent Technologies, Palo Alto, CA). Library

1097 construction and sequencing were performed by Novogene USA (Sacramento, CA), using
1098 Illumina S4 flowcell on Novaseq 6000 platform with a paired-end read length of 150 bp.
1099 Downstream RNAseq analysis was performed using a combination of programs. Alignments were
1100 parsed using STAR and differential expression was determined through limma/voom.⁷⁸
1101 Alignments were parsed using STAR and differential expression was determined through
1102 limma/voom.⁷⁸ GO, KEGG, REAC and TF enrichment were determined with the g:profiler R
1103 package.⁷⁹ Reference genome and gene model annotation files were downloaded from genome
1104 website browser (NCBI/UCSC/Ensembl) directly. Indices of the reference genome were built
1105 using STAR⁸⁰ and paired-end clean reads were aligned to human or mouse reference genome
1106 using STAR (v2.5), setting the outFilterMismatchNmax argument to 2, from which gene level
1107 count tables were built using HTseq-count.⁸¹ Gene-level read counts were then processed using
1108 the limma suite of tools⁷⁸ first with a voom with quality weights transformation, followed by linear
1109 model fitting to determine differentially expressed genes. The P values for individual genes were
1110 adjusted using the Benjamini & Hochberg method. Corrected P-value of 0.05 and absolute
1111 foldchange of 1 were set as the threshold for significantly differential expression. Geneset testing
1112 on the entire dataset was performed using the camera() function to determine geneset direction
1113 using genesets extracted from MsigDB.⁸² Select genesets were plotted as dotplots from the
1114 camera output. Gene Ontology (GO), Kyoto Encyclopedia of Genes and Genomes (KEGG), and
1115 REACTOME database enrichment analysis of differentially expressed genes was further
1116 implemented with the g:profiler R package,⁷⁹ in which gene length bias was corrected. Terms with
1117 corrected P-values less than 0.05 were considered significant. The high-throughput sequencing
1118 data from this study have been submitted to the NCBI Sequence Read Archive (SRA) under
1119 accession number #####.

1120

1121 **Data plotting and statistical analysis software**

1122 Unless stated otherwise, statistical analyses were performed in GraphPad Prism (San Diego, Ca).
1123 Statistics are indicated in graphs if $p < 0.05$.

1124

1125 **AUTHOR CONTRIBUTIONS:**

1126 MK, EM, SS, BE, CMR and YPJ designed research; MK, EM, SS, CGF, JML, JLP, MT, BR, IRL,
1127 C.Z., BZ, AFS, CQ, LF, AWA, WMS, SB, GL, LC and YPJ performed experiments; MK, EM, SS,
1128 CGF, JML, JLP, MT, BR, IRL, CZ, BZ, AFS, CQ, LF, AWA, YL, DEC, MP, LD, MG, PM, BAE,
1129 CMR and YPJ analyzed data; HS, NDT and RC provided new reagents/analytic tools; DEC, PM,
1130 CMR and YPJ secured funding; MK, EM, and YPJ wrote the paper; YPJ directed the study.

1131

Figure 1: Human hepatocytes in chimeric mice on WD rapidly develop steatosis

Figure 2: Systemic metabolic effects of WD in chimeric FNRG mice

Figure 3: huFNRG mice on WD develop mild steatohepatitis

Figure 4: Transcriptional changes in huFNRG mice on Western Diet

Figure 5: PNPLA3-148M huFRG mice on WD develop steatohepatitis

Figure 6: PNPLA3-148M overexpression in hepatocytes exacerbates steatosis in mice on WD

a

b

c

d

e

f

Figure 7: PNPLA3-148M overexpression worsens steatohepatitis on HFD

Supplemental Figure S1: Human hepatocyte steatosis in chimeric models

Supplemental Figure S2: Systemic metabolic effects of WD feeding

Supplemental Figure S3: Mild steatohepatitis in huFNRG mice on WD

Supplemental Figure S4: Transcriptional changes in huFNRG mice on Western Diet

a

b

human pathways

c

mouse pathways

Supplemental Figure S5: 148M-huFRG mice develop steatohepatitis

Supplemental Figure S6: PNPLA3 transgenic huFNRG mice

Figure S7: PNPLA3-148M overexpression worsens steatohepatitis on HFD

Table 1) qRT-PCR mRNA expression in transduced huFNRG animals - average fold change over housekeeping

Gene	Average fold change over housekeeping - chow			Average fold change over housekeeping - WD		
	tdRFP	td148I	td148M	tdRFP	td148I	td148M
ABCA1	0.368883198	0.267786579	0.5519846	0.6727912	0.707126819	0.449371439
ACACA	0.035656185	0.038552958	0.071810784	0.083214261	0.083744866	0.057474499
ACADL	0.110949301	0.101655054	0.14558227	0.199057575	0.170211114	0.188835791
ACLY	0.436496378	0.567439241	0.52725121	0.461086724	0.497637348	0.653387539
ACOX1	2.517925637	2.849879635	2.231809941	4.178923865	3.903006829	3.150915163
ACSL5	1.798792185	1.662574057	1.742976738	2.444098549	1.72579322	1.822920344
ACSM3	3.176615241	4.975173094	3.533113634	4.922244267	3.695856521	5.951371558
ADIPOR1	0.433077354	0.428142874	0.482731662	0.513066692	0.503029756	0.537061101
ADIPOR2	0.833701393	0.851964307	0.911126783	1.078684175	1.05454579	0.831764838
AKT1	0.238494562	0.272889999	0.25423422	0.325361344	0.323277443	0.272738552
APOA1	101.5835357	91.40737221	104.0803431	181.4814453	183.2887107	205.1508232
APOB	34.41468389	34.00765282	40.64313743	54.92296325	49.88464552	53.8572082
APOC3	90.65886756	75.41545032	105.3560986	128.6632013	128.3388654	132.7015488
APOE	14.36420394	12.37371322	15.63639277	22.92161029	21.05691049	19.42476157
ATP5C1	1.883365035	2.025512372	1.77809252	2.120542084	1.699034415	1.970308135
CASP3	0.14222822	0.138794303	0.153920158	0.178177748	0.169408443	0.173943887
CD36	1.109397909	0.964592474	1.11226837	2.520850775	2.150983894	2.280733948
CEBPB;CEBPA-AS1	0.131348551	0.150504893	0.230620196	0.345507675	0.234725837	0.180984119
CNBP	1.681373003	1.631270687	1.4649865	1.764718361	1.404109004	1.880379885
CPT1A	1.573508999	1.988468625	1.018277411	1.336678777	1.157082822	0.946755159
CPT2	0.860227974	0.828165	0.66481186	1.060712454	0.996925551	1.027784841
CYP2E1	48.13382884	52.98359038	50.65613294	98.46466494	71.22366345	100.4290926
CYP7A1	0.045502776	0.056265985	0.845162194	1.302704706	0.418776654	0.873183109
DGAT2	1.785244728	1.27057123	2.648681604	4.05056992	2.756586266	3.585261025
FABP1	53.76484075	60.30053405	32.83167347	73.11974507	69.42785371	65.05980676
FABP3	0.030896467	0.060774155	0.014648243	0.01640808	0.019624525	0.012184946
FABP5	0.051886321	0.044810817	0.03213479	0.032659698	0.026736462	0.030655666
FAS	0.111562643	0.086802296	0.13182671	0.154132372	0.163275093	0.152697716
FASN	0.559186083	0.570116112	1.225495927	1.890858451	2.355251867	1.136443648
FOXA2	0.103035877	0.10728501	0.142773019	0.177348528	0.158534305	0.160535032
FOXO1	0.120158742	0.148166229	0.083402316	0.089057933	0.067803239	0.068487858
G6PC	12.92404672	12.05760327	9.598385617	6.984889807	6.97351766	9.122014223
G6PD	0.009800051	0.014160332	0.010899059	0.01559439	0.013913829	0.016862318
GCK	8.3343E-05	9.51975E-05	0.003181612	0.001282137	0.003028274	0.001825874
GK	0.143295743	0.171232355	0.144058926	0.212581321	0.183416553	0.210181879
GSK3B	0.207578414	0.276604229	0.228618434	0.281228441	0.257897685	0.241680432
HMGCR	0.15452151	0.165345921	0.212269244	0.13726636	0.147313581	0.112117061
HNF4A	3.996306081	3.943692904	4.753606452	5.331060791	4.639432231	5.688434883
IGF1	0.010426567	0.004866301	0.020359608	0.00306642	0.003730813	0.002496448
IGFBP1	12.44027242	13.34406143	10.08440111	9.935696835	7.082208029	9.074199535
IL1B	8.3343E-05	9.51975E-05	0.000153504	0.000171541	0.000355026	0.00059247
INSR	0.183237252	0.195447769	0.175270148	0.222821746	0.180882517	0.177165927
IRS1	0.205456659	0.246516791	0.237258747	0.24300858	0.238575745	0.240730114
LDLR	0.074089788	0.071929665	0.150844332	0.076538055	0.08279066	0.054625664
LEPR	0.391030638	0.420141832	0.210327956	0.118092904	0.137903343	0.11766551
LPL	0.000305595	0.003088181	8.52238E-05	0.000824859	0.001619555	0.000269268
MAPK1	0.216294145	0.24540175	0.202622028	0.29092258	0.2612777	0.240152682
MAPK8	0.440281802	0.442547892	0.437249105	0.673947769	0.541084062	0.548329977
MLXIPL	0.981203864	1.023470324	1.673085147	1.834284935	1.644262293	2.133155067
MTOR	0.17906682	0.182718973	0.172394208	0.29297052	0.213826751	0.196764821
NDUFB6	2.083029561	1.521543606	1.555686354	1.911428134	1.58798678	1.968871463
NFKB1	0.106767086	0.102503414	0.111532414	0.152805067	0.151704325	0.147149618
NR1H2	0.114479501	0.110169063	0.104124579	0.137119984	0.130467613	0.134134529
NR1H3	0.247463587	0.287596294	0.268097208	0.400125993	0.338054687	0.433478265
NR1H4	0.201521892	0.207478544	0.292981797	0.440805966	0.34670219	0.418768654
PCK2	4.062199455	3.364462956	4.718561546	4.49595215	4.841278589	4.233615961
PDK4	1.112647549	1.749907845	0.555580537	1.678436521	1.128671641	0.977784582
PIK3CA	0.054329087	0.057465596	0.053603142	0.070732218	0.061532335	0.068284189
PIK3R1	0.448423342	0.564011593	0.681658373	0.734037504	0.701808132	0.564470116
PKLR	0.449193735	0.517960062	0.604507878	1.085207053	1.076839318	1.392419297
PNPLA3	0.113741573	42.36533239	19.63691814	0.41869652	25.73235028	25.07571438
PPA1	3.355371555	3.922263647	2.695813337	2.900275121	2.583081448	4.431660341
PPARA	0.721905106	0.909120989	0.938163465	1.131783972	0.943743907	0.946628258
PPARG	0.050849692	0.052560543	0.053859743	0.118730057	0.111007887	0.103434765
PPARGC1A	1.281363665	1.791011309	1.195756063	1.032135858	1.224376566	0.709771484
PRKAA1	0.123766016	0.136691229	0.134870277	0.121180919	0.132680989	0.136364862
PTPN1	0.058334826	0.045048939	0.056797798	0.072605707	0.063509352	0.070857878
RBPA	47.4800469	47.0333149	53.55747394	55.1223592	56.63062069	59.44587287
RXRA	0.47999656	0.456585111	0.594743968	1.003442925	0.886003198	0.783381395
SCD	1.508170175	1.266519099	3.007627677	8.603733964	9.775838241	6.078768955
SERPINE1	0.096467782	0.08181576	0.179991825	0.694545795	0.710336881	0.324698447
SLC27A5	3.237334403	2.818938018	4.087306836	4.617318743	4.981965166	3.976890967
SLC2A1	0.011493495	0.010800383	0.020086921	0.00544164	0.00680493	0.006026506
SLC2A2	2.953556671	2.595344705	4.167931432	7.107000571	6.062238245	5.953809267
SLC2A4	0.005322738	0.004416619	0.005518313	0.009949256	0.010087355	0.006353036
SOCS3	8.3343E-05	9.51975E-05	0.000832534	0.00011565	8.5228E-05	0.000137509
SREBF1	0.624108037	0.65410058	0.72348495	1.386376188	1.341956211	0.949654207
SREBF2	0.089736092	0.070610763	0.145507148	0.1244603	0.132038588	0.103745968
STAT3	0.700090113	0.810089398	0.848016211	0.849374371	0.849885479	0.649254289
XBP1	2.455458823	2.230970953	3.150858997	3.678411889	3.944201872	3.625131871

Figures

Figure 1

Human hepatocytes in chimeric mice on WD rapidly develop steatosis a) Experimental timeline of FNRG mice that were transplanted with primary human hepatocytes (PHH), cycled off nitisinone to create huFNRG mice, and then subjected to chow or WD for up to 12 weeks. b) H&E staining on livers from

huFNRG mice on chow and after 4 or 12 weeks on WD. c) Staining against human nuclear mitotic apparatus-1 in livers of huFNRG mice on chow and 4 or 12 weeks on WD. Areas with predominantly mouse hepatocytes are outlined. d) Staining against human glutamine synthetase on livers from huFNRG mice on chow and 4 weeks on WD. Central vein (CV), portal tract (PT). e) Liver steatosis score of the human graft in huFNRG mice on chow and after 4 to 20 weeks on WD. Symbols individual mice, bars are median, unpaired t-test with $**p<0.005$, $***p<0.0005$ and $****p<0.0001$. f) Hepatic triglyceride levels in livers of huFNRG mice on chow and after 4 to 12 weeks on WD. Symbols individual mice, bars are median, unpaired t-test with $***p<0.0005$ and $****p<0.0001$. g) H&E staining on livers from muFNRG mice on chow and after 4 or 12 weeks on WD.

Figure 2

Systemic metabolic effects of WD in chimeric FNRG mice a) Body weight in huFNRG mice during time course of chow or WD feeding. Symbols individual mice, line is mean, two-way ANOVA Sidak pairwise comparison * $p < 0.05$, ** $p < 0.005$, *** $p < 0.0005$. b) Gonadal fat fraction of total body weight in huFNRG mice on chow and after 4 to 20 weeks on WD. Symbols individual mice, bars are median, unpaired t-test with * $p < 0.05$, ** $p < 0.005$, *** $p < 0.0005$. c) Cholesterol in plasma lipoprotein fractions from huFNRG and

muFNRG mice on chow and after 4 weeks on WD. Pooled plasma (3 mice/group), symbols are technical means. d) Triglycerides in plasma lipoprotein fractions from huFNRG and muFNRG mice on chow and after 4 weeks on WD. Pooled plasma (3 mice/group), symbols are technical means. e) Fasting blood glucose from huFNRG and muFNRG mice on chow and after 4 weeks on WD. Symbols individual mice, lines are median, unpaired t-test **** $p < 0.0001$. f) Fasting serum mouse insulin from huFNRG and muFNRG mice on chow 709 and after 4 weeks on WD. Symbols individual mice, lines are median, unpaired t-test with * $p < 0.05$, ** $p < 0.005$. g) Intraperitoneal glucose, insulin and pyruvate tolerance testing in huFNRG and muFNRG mice on chow and after 4 weeks on WD. Blood glucose values measured at baseline (0) and at indicated timepoints after i.p. injections with each agent. Symbols are mean SEM of 5-7 mice per group, two-way ANOVA Sidak pairwise comparison of huFNRG on chow and WD (p-values on top) or muFNRG on chow and WD (p-values in bottom), * $p < 0.05$, ** $p < 0.005$, *** $p < 0.0005$ and **** $p < 0.0001$

Figure 3

huFNRG mice on WD develop mild steatohepatitis a) ALT and AST activity in serum from huFNRG mice on chow and after 4 to 12 weeks on WD. Symbols individual mice, bars are median, unpaired t-test **p<0.005, ****p<0.0001. b) H&E staining from a huFNRG mouse liver after 8 weeks on WD. 734 Black arrow indicates hepatocyte ballooning degeneration and white arrow indicates lobular inflammation. c) Staining against human nuclear mitotic apparatus-1 in the liver of a huFNRG mouse after 4 weeks on

WD. Black arrow indicates hepatocyte ballooning. d) Fractions of huFNRG mice with human hepatocyte ballooning degeneration on chow and WD over time. Number of mice with ballooning per group at bottom of bars. e) Fractions of huFNRG mice with human hepatocyte ballooning degeneration by grade of steatosis. Number of mice with ballooning per group at bottom of bars. f) NAFLD activity score (NAS) in the human graft of huFNRG mice on chow and after 4 to 20 weeks on WD. Symbols individual mice, bars are median, unpaired t-test ** $p < 0.005$, **** $p < 0.0001$. g) Picosirius Red staining for collagen in livers from huFNRG mice on chow and after 4 to 20 weeks on WD. h) Fibrosis stages in the human graft of huFNRG liver on chow and after 4 to 20 weeks on WD. Mouse numbers at bottom of bars.

Figure 4

Transcriptional changes in huFNRG mice on WD a) Volcano plots of liver transcripts mapped to human genome (human reads) comparing huFNRG mice 4 weeks on WD to chow, and longer WD durations to the previous WD timepoint. RNA-sequencing of 3-4 mice per group, red symbols FDR < 0.05, grey denotes not significant. b) Volcano plots of liver transcripts mapped to mouse genome (mouse reads) comparing huFNRG mice 4 weeks on WD to chow, and longer WD durations to the previous WD

timepoint. RNA-sequencing of 3-4 mice per group, red symbols FDR <0.05, grey denotes not significant. c) Gene-ontology (GO) pathway analysis of human reads from livers of huFNRG mice 4 weeks on WD to chow, and longer WD durations to the previous WD timepoint. Blue symbols are down and red symbols are upregulated pathways, grey denotes no significant changes, symbol size indicates statistical significance. d) Gene-ontology (GO) pathway analysis of mouse reads from livers of huFNRG mice 4 weeks on WD to chow, and longer WD durations to the previous WD timepoint. Blue symbols are down and red symbols are upregulated pathways, grey denotes no significant changes, symbol size indicates statistical significance.

Figure 5

PNPLA3-148M huFRG mice on WD develop steatohepatitis a) H&E staining on livers from 148M-huFRG mice on chow and after 4 weeks on WD. b) Clinical steatosis score of the human graft on livers from 148M-huFNRG mice on chow and after 4 weeks WD. Symbols are individual mice, bars median, unpaired t-test ***p<0.0005. c) H&E staining on livers from 148M-huFRG mice 4 weeks on WD. Grey arrow indicates Mallory Denk body, black arrow ballooning degeneration and white arrow lobular inflammation. d) NAFLD

Activity Score (NAS) in the human graft of 148M-huFRG mice 807 on chow and after 4 weeks WD. Symbols individual mice, bars are median, unpaired t-test *** $p < 0.0005$. e) Volcano plots of liver transcripts mapped to human genome (human reads) comparing 148M huFRG mice after 4 weeks WD vs. chow. RNA-sequencing of 3-4 mice per group, red symbols FDR < 0.05 , grey denotes not significant. f) Gene-ontology (GO) pathway analysis of human reads comparing 148I-huFNRG and 148M huFRG livers after 4 weeks WD vs. chow. Blue symbols are down- and red symbols are upregulated pathways, grey denotes no significant changes, symbol size indicates statistical significance. g) Volcano plots of human reads differentially expressed after 4 weeks WD compared to chow (Δ WD) in 148M-huFRG livers versus 148I-huFNRG livers. Red symbols FDR < 0.05 , grey denotes not significant.

Figure 6

PNPLA3-148M overexpression in hepatocytes exacerbates steatosis in mice on WD a) Experimental timeline of creating PNPLA3 overexpressing mice and diet challenge. Mouse passaged primary human hepatocytes (PHH) cultures were transduced with lentiviral vectors expressing PNPLA3 variants and re-transplanted. Following humanization, huFNRG with PNPLA3 transduced hepatocytes (td) were exposed to 4 weeks of WD. b) H&E staining on livers from tdRFP, td148I and td148M mice after 4 weeks on WD. c)

Liver steatosis score of the human graft in tdRFP, td148I and td148M mice on chow and after 4 weeks on WD. Symbols individual mice, bars are median, unpaired t-test with ** $p < 0.005$, *** $p < 0.0005$ and **** $p < 0.0001$. d) Hepatic triglyceride quantification in tdRFP, td148I and td148M mice on chow and after 4 weeks on WD. Symbols individual mice, bars are median, unpaired t-test, * $p < 0.05$. e) NAFLD Activity Score (NAS) in the human graft of tdRFP, td148I and td148M mice on chow and after 4 weeks on WD. Symbols individual mice, bars are median, unpaired t-test, * $p < 0.05$, ** $p < 0.005$, *** $p < 0.0005$. f) Volcano plot of 80 human genes expressed in livers of td148I versus td148M mice after 4 weeks on WD. Expression by qRT-PCR normalized to 11 housekeeping genes, $n = 3-4$ mice per group, burgundy symbols $p < 0.05$, grey symbols not statistically significant.

Figure 7

A HFD causes more active steatohepatitis in PNPLA3-148M overexpressing 881 micea) H&E staining on livers from tdRFP, td148I and td148M mice after 4 weeks on HFD. b) Liver steatosis score of the human graft in tdRFP, td148I and td148M mice on chow and after 4 weeks on HFD. Symbols individual mice, bars are median, unpaired t-test with ** $p < 0.005$. c) H&E staining on liver from a td148M mouse after 4 weeks of HFD. White arrow indicates lobular inflammation, black arrow microvesicular steatosis and grey

arrow Mallory-Denk body. d) NAFLD Activity Score (NAS) in the human graft of tdRFP, td148I and td148M mice on chow and after 4 weeks on HFD. Symbols individual mice, bars are median, unpaired t-test, $*p < 0.05$. e) Low and high magnification fluorescent images of livers from td148I and td148M mice after 4 weeks of HFD stained for neutral lipids (green, BODIPY) and nuclei (blue, DAPI). f) Two-dimensional density plots of neutral lipids (BODIPY) and RFP of 355,161 small hepatocytes in livers from four td148I mice and 183,289 small hepatocytes in livers from three td148M mice after 4 weeks of HFD. Density plots are displaying RFP in X-, and BODIPY intensities in Y-Axis, both in arbitrary units (AU). Dashed lines display cell population group quadrants, with lower left quadrant harboring double negative and upper right quadrant harboring double positive cells. Solid line is simple linear regression with equation displayed underneath, r^2 is Pearson correlation coefficient.