Detection of copy number variants in African goats using whole genome sequence data

CURRENT STATUS: UNDER REVIEW

BMC Genomics BMC Series

Wilson Nandolo

Lilongwe University of Agriculture and Natural Resources

Gábor Mészáros

University of Natural Resources and Life Sciences, Vienna, Austria

Maria Wurzinger

University of Natural Resources and Life Sciences, Vienna, Austria

Liveness J. Banda

Lilongwe University of Agriculture and Natural Resources

Timothy N. Gondwe

Lilongwe University of Agriculture and Natural Resources

Henry A. Mulindwa

National Livestock Resources Research Institute

Helen N. Nakimbugwe

Kyambogo University

Emily L. Clark

The University of Edinburgh The Roslin Institute

M. Jennifer Woodward-Greene

USDA-ARS National Agricultural Library

Mei Liu

USDA-ARS Animal Genomics and Improvement Laboratory

George E. Liu

USDA-ARS Animal Genomics and Improvement Laboratory

Curtis P. Van Tassell

USDA-ARS Animal Genomics and Improvement Laboratory

Benjamin D. Rosen

"USDA-ARS Beltsville Agricultural Research Center"

✉ ben.rosen@usda.gov *Corresponding Author*
ORCID: <https://orcid.org/0000-0001-9395-8346>

Johann Sölkner

University of Natural Resources and Life Sciences, Vienna, Austria

DOI:

10.21203/rs.3.rs-15747/v1

SUBJECT AREAS

Epigenetics & Genomics

KEYWORDS

African goats, copy number variations, whole genome sequence, structural variations

Abstract

Background

Copy number variations (CNV) are a significant source of variation in the genome, as such they are essential to the genetic characterization of animal breeds. The aim of this study was to develop a fine-scaled copy number variation map for African goats. We used sequence data from multiple breeds and from multiple African countries.

Results

A total of 253,553 CNV (244,876 deletions and 8,677 duplications) were identified, corresponding to an overall average of 1,393 CNV per animal. The mean CNV length was 3.3 kb, with a median of 1.3 kb. There was substantial differentiation between the populations for some CNV, suggestive of the effect of population-specific selective pressures. A total of 6,231 global CNV regions (CNVR) were found across all animals, representing 59.2 Mb (2.4%) of the goat genome. About 1.6% of the CNVR were present in all 34 breeds and 28.7% were present in all 5 geographical areas across Africa, where animals had been sampled. The CNVR had genes that were highly enriched in important biological functions, molecular functions, and cellular components including retrograde endocannabinoid signaling, glutamatergic synapse and circadian entrainment.

Conclusions

This study presents the first fine CNV map of African goat based on WGS data and adds to the growing body of knowledge on the genetic characterization of goats.

Background

Structural variations (SV) are an important source of genetic variation [1–4]. SV are broadly categorized into 4 groups: deletions, insertions, translocations and inversions. Deletions and insertions are referred to as unbalanced SV because they result in changes in the length of the genome, and can result in copy number variations (CNV). CNV are typically considered when these insertions or deletions in the genome are at least 50-1000 base-pairs (bp) long [5–10]. CNV are not as abundant as single nucleotide polymorphisms (SNP) in quantity, but because of their larger sizes, they may have a dramatic effect on gene expression in individuals [11]. The level to which the gene is

expressed may vary between individuals. Duplication or deletion in or near the gene or the regulatory region of a gene may lead to loss or gain of gene function. CNV cover about 4.5–9.8% of the human genome [12] and are associated with many Mendelian disorders [11]. Girirajan et al. [13] found that CNV significantly determine the severity and prognosis of many genetic disorders. Approximately 14% of diseases in children with intellectual disability are caused by CNV [14]. On the other hand, some CNV have been found to be associated with adaptive fitness of individuals, such as adaptation to starch diets associated in the gene encoding α -amylase [12].

Traditionally, microarray-based comparative genomic hybridization (array CGH) or SNP genotyping arrays are used to detect CNV. Several studies have been carried out using these methods to detect and map CNV in the goat genome, including studies by Fontanesi et al. [15] in four goat breeds; Nandolo et al. [16] in 13 East African goat breeds; and Liu et al. [17] in a global goat population. Detecting CNV using array CGH and SNP genotyping arrays suffer from shortcomings that include hybridization noise, limited coverage of the genome, low resolution, and difficulty in detecting novel and rare mutations [18–20]. The development of whole-genome sequencing (WGS) technologies has made it possible for more rigorous and accurate detection of CNV.

According to Mills et al. [21], WGS-based CNV detection methods fall into four major approaches, namely, methods based on paired-end mapping (PEM), split reads (SR), read depth (RD) and de novo assembly of a genome (AS). The PEM and SR methods are useful for detection of small-scale CNV [22], and several algorithms are loosely based on them, including BreakDancer [23], Pindel [24], and Delly [25]. RD approaches are very useful for detection of larger CNV. Algorithms using this approach include CNV-Seq [26], CNVnator [27] and the event-wise testing approach (EWT) developed by Yoon et al. [28]. The methods can also be combined. For example, LUMPY [29] is able to combine two or more of the previous approaches to refine SV detection. Assembly-based approaches are computationally intensive and are therefore not generally used with WGS data [22, 30]. Most of these SV-detection algorithms have been extensively reviewed [1, 30–33].

LUMPY implements a breakpoint prediction framework, where a breakpoint is defined as a pair of genomic regions that are adjacent in a sample, but not in the reference genome. The location of the

breakpoint is determined using a probability function that considers different sources of evidence supporting the existence of a breakpoint, including information from discordant read pairs and split reads. A discordant read pair occurs when sequence from two ends of an insert are inconsistent when compared to the reference genome. These inconsistencies result from differences between mapping distance or the orientation between the pairs of sequences. [34, 35]. Split reads are sequences that map to the reference genome on one end only, and, as explained by Ye and Hall [32], such reads can indicate the location of a breakpoint with a high degree of certainty. There are similar algorithms that rely heavily on the use of breakpoints to determine genome rearrangements at single-nucleotide resolution, including Delly [25] and Pindel [24].

Like LUMPY, Manta [36] incorporates use of PEM and SR methods. However, Manta also uses AS analysis. Manta overcomes the computational expense of AS methods by splitting the work into many smaller workflows which can be carried out in parallel. Manta scans the genome for SV and then scores, genotypes and filters the SV based on diploid germline and somatic biological models [36]. Manta can detect all structural variant types that are identifiable in the absence of copy number analysis and large-scale de-novo assembly, which is why this approach is also a good candidate for joint analysis of small sets of diploid individuals, tumor samples, and similar analyses. Both LUMPY and Manta are good at identifying SV break points with high resolution.

Many studies have been carried out to detect CNV using WGS data in various domesticated species: cattle [37], cats [38], chickens [39], dogs [40], etc. So far, there is no report of goat CNV discoveries using WGS data. The goal of this study was to identify CNV in the goat genome through the intersection of LUMPY and Manta outputs as a part of the characterization of African goats in conjunction with the ADAPTmap project [41]. Goats are a very important farm animal genetic resource for the livelihoods of African smallholders, and a deeper understanding of the goat genome is necessary to facilitate the improvement of goats in the region. This study aimed to generate a fine-scale CNV map for the goat genome.

Results

Number and distribution of CNV

LUMPY detected 8,563 duplications and 230,497 deletions while Manta detected 24,088 duplications and 320,374 deletions. A combined data set with 244,876 deletions and 8,677 duplications (totaling 253,553, translating into an average of 1,393 CNV per animal) was derived from the intersection of the LUMPY and Manta sets after removal of variants shorter than 50 bp or longer than 3 Mb. The combined data set had more observations than the LUMPY data set (which had fewer raw CNV) because for some individuals, a large number of short CNV from Manta intersected with few long CNV from LUMPY.

The CNV were distributed across the 29 autosomes as shown in Fig. 1. A vast majority of the CNV (96.6%) were losses. This is not unexpected, because all CNV detection methods suffer from an inherent deficiency in detecting insertions. In the case of CNV detection using WGS data, this limitation is even more pronounced with PEM methods, because they detect insertions when the mapped reads are at a distance shorter than the fragment length, so they are not able to detect insertions larger than the insert size of the reference library [42].

Overall, the mean CNV length was about 3.3 kb, with a median of 1.3 kb. The distribution of the lengths of the CNV for each population are shown in Fig. 2 by CNV length category. A summary of the descriptive statistics of the CNV for the populations are given in Table 1. Most of the CNV losses (99.92%) were less than 100 kb long while 6.3% of CNV gains were longer than 100 kb. Despite the overwhelming proportion of losses over gains, there were more CNV gains observed over 100 kb than losses. Similarly, only 1.04% of the loss CNV were longer than 10 kb, while almost one-quarter (22.99%) of all gain CNV were over 10 kb. As a result, CNV gains were longer than CNV losses and had larger range in length. Deletions and duplications averaged about 2.3 and 31.5 kb long, with median lengths of 1.3 and 1.4 kb, respectively. There were no significant differences in the distribution of CNV across the five populations as shown in the percentile and sample QQ plots in Fig. 3.

Table 1
Descriptive statistics of CNV and CNV length for each population

Population	Number of samples	CNV		CNV length (bp)			
		State	Number	Mean	Median	Minimum	Maximum
BOE	9	Loss	9079	2227.1	1326	67	254129
		Gain	331	20165.9	1500	161	631262
		Overall	9410	2858.1	1330	67	631262
EAF	80	Loss	108051	2244.7	1293	52	2161018
		Gain	3544	30979.2	1316.5	118	2777398
		Overall	111595	3157.2	1293	52	2777398
MAD	27	Loss	31426	2475.3	1295	84	2069909
		Gain	1078	28384.1	1446	84	1660243
		Overall	32504	3334.6	1296	84	2069909
SAF	44	Loss	67099	2368.9	1285	51	2539701
		Gain	2514	31000.7	1192	101	1959154
		Overall	69613	3402.9	1283	51	2539701
WAF	22	Loss	29221	2491.4	1280	52	2457795
		Gain	1210	40255.3	1234	65	2788546
		Overall	30431	3993	1280	52	2788546

Population CNV differentiation

V_{ST} analysis showed that several CNV were highly differentiated between and across the populations. Some of these CNV overlapped with genes of importance in goats. Results for the pairwise population V_{ST} tests and the V_{ST} test across all the populations with their respective 99th percentile CNV V_{ST} thresholds are given in Supplementary Table 1 (Additional file 1). V_{ST} values for the pairwise tests are given in Supplementary Figs. 1–10 (Additional file 2). The V_{ST} values for genes that were in CNV that were highly differentiated across all populations are shown in Fig. 4. The gene DST was in a CNV with a very high V_{ST} threshold across all the populations. DST has been associated with herpes virus and respiratory disease (BRD) in cattle [43]. Some CNV were highly differentiated both between and across populations. CNV with high differentiation between only some populations include the CNV corresponding to the genes BCO2, CCSER1 (FAM190A), COL24A1, CPNE4, CWC22, IMMP2L, KBTBD12, LAMA3, NAALADL2, RFX3, SEMA3D, SLC2A13, STPG2 (C4orf37), TAFA2 (FAM19A2), TMEM117, TMEM161B and VPS13B. The rest of the genes were in CNV that were highly differentiated across all populations.

Number and distribution of CNVR

The lists of CNVR by population are given in Supplementary Table 2 (Additional file 1) and their locations on the goat genome are shown in Fig. 5. Plots of the CNVR for each breed (with more than 2 animals) are given in Supplementary Figs. 11 to 39 (Additional file 2). Descriptive statistics of the CNVR for each population is given in Supplementary Table 3 (Additional file 1) while a distribution of

CNVR by size and populations is given in Fig. 6. Over 92% of the CNVR were copy losses. There was a wide variation in the number and sizes of the CNVR between and among the populations. The fraction of copy gains or gains and losses was highest in the group of CNVR of at least 10kbp, with 25% copy gains and 19% for losses/gains (Fig. 6). There were differences between breeds, with the Boer (BOE) having the longest median (1.3 kbp and 1.4 kbp) and the shortest mean (2.2 kbp and 10.0kbp) CNVR for the loss and the gain classes, respectively. Also, notable, was the very long loss/gain CNVR for all the breeds except Boer. This mixed African populations all had CNVR > 1 Mbp, while the longest Boer CNVR observed was around 0.5 Mbp.

Number and distribution of global CNVR

A total of 6,231 global CNVR were found across all animals. A list of the global CNVR is given in Supplementary Table 4 (Additional file 1) and a summary is given in Table 2. There were 5,742 losses, 280 gains and 209 regions had both losses and gains in different individuals. The locations of the global CNVR are given in Fig. 7. CNVR with both gains and losses were much longer (mean 185.8 kb) and constituted a significant proportion of the total CNVR coverage (65.6%). Sixteen of these were longer than 1 Mb (on chromosomes 1, 2, 6, 7, 12, 14 (two regions), 17, 19, 21, 23 (two regions), 27 and 29).

Overall, the CNVR covered about 59.2 Mb of the goat genome. Previous work on genome-wide CNV discovery in goats using SNP data done by Liu et al. [17] showed that CNVR cover approximately 262 Mb of the goat genome. Of the 978 CNVR reported in that study, 540 CNVR intersected with 819 CNVR identified in our study. The amount of the overlap between the CNVR in the two studies was 217.1 Mb, covering 38.6 Mb (65.1%) in this study, and 194.2 Mb (74.1%) in the other study.

Table 2

CNVR summary statistics for each CNV state based on CNV occurring in at least 2 individuals.

Copy state	Number of CNVR	Length (bp)				CNVR coverage (bp)
		Mean	Median	Minimum	Maximum	
Loss	5742	3041.3	1140.5	52	1177087	17463236
Gain	280	10377.9	1008.0	302	236347	2905806
Both	209	185755.2	1731.0	616	2956746	38822839
Overall	6231	9499.6	1157.0	52	2956746	59191881

Common and rare CNVR

Most of the CNVR (> 95.9%) were found in at least 2 breeds. Out of the 6,231 CNVR, 98 (1.6%) were

present in all the 34 breeds and 1,790 (28.7%) were present in all the populations (Fig. 8A and Fig. 8B). The most frequent CNVR observed was on chromosome 6 from 115,822,332 bp to 115,825,687 bp with a frequency of 96.2%. There were 259 CNVR private to 30 breeds, and 1,018 private to all 5 populations, distributed as shown in Fig. 8C and Fig. 8D. BOE (Tanzania and Zimbabwe), KEF (Ethiopia) and MLY (Tanzania) breeds had the highest numbers of private CNVR (20, 21 and 31, respectively).

Functional annotation and gene enrichment analysis

Functional annotation was carried out for genes in CNV that showed high levels of differentiation between and among populations and for genes in global and private CNVR.

Functional annotation for highly differentiated CNV

Gene ontology classifications are shown in Fig. 9 for biological processes, cellular components and molecular functions that were associated with genes contained in CNV that were highly differentiated based on both the pairwise V_{ST} tests between populations and the V_{ST} test across all the populations according to functional classification in the Panther Classification System using the *Bos taurus* gene set. The bovine classification was used because the goat gene set is not currently hosted. The top two biological processes represented were cellular processes and localization, represented in 27% and 14% of the processes. The other categories (biological regulation, response to stimulus, multicellular organism process and metabolic process) were represented at levels ranging from 5–11%. The top three cell components were cells, organelles, and membranes. Similarly, the top three molecular functions were catalytic activity, (cell) binding, and molecular transducer activity.

Functional annotation for global CNVR

Up to 2,980 genes overlapped with the 6,321 CNVR identified in this study. Up to 755 of these genes formed 24 clusters, with enrichment scores ranging from 0.0 to 1.89. Higher enrichment scores imply higher overrepresentation of the genes in the gene set for the gene enrichment term [44]. The top 3 clusters with the highest enrichment scores are given in Table 3 while the full list is given in Supplementary Table 5 (Additional file 1). The most significant GO terms identified in the analysis included retrograde endocannabinoid signaling; glutamatergic synapse; circadian entrainment; dopaminergic synapse; gastric acid secretion; long-term potentiation; salivary secretion; and calcium

signaling pathway.

Table 3

Functional annotation clusters of the genes found in the global CNVR based on analysis in DAVID

Cluster (Enrichment score, database)	Enrichment term	Gene count	p-value
1 (1.89, KEGG_PATHWAY)	chx04020: Calcium signaling pathway	31	0.001
	chx04970: Salivary secretion	16	0.011
	chx04022: cGMP-PKG signaling pathway	25	0.016
	chx04270: Vascular smooth muscle contraction	19	0.037
	chx04261: Adrenergic signaling in cardiomyocytes	21	0.046
2 (1.62, KEGG_PATHWAY)	chx04723: Retrograde endocannabinoid signaling	24	0.000
	chx04724: Glutamatergic synapse	24	0.000
	chx04713: Circadian entrainment	21	0.001
	chx04728: Dopaminergic synapse	25	0.002
	chx04971: Gastric acid secretion	16	0.004
	chx04720: Long-term potentiation	14	0.009
	chx04970: Salivary secretion	16	0.011
	chx04925: Aldosterone synthesis and secretion	15	0.014
	chx04730: Long-term depression	13	0.014
	chx04540: Gap junction	16	0.019
	chx04750: Inflammatory mediator regulation of TRP channels	18	0.020
	chx04921: Oxytocin signaling pathway	23	0.028
	chx04922: Glucagon signaling pathway	17	0.028
	chx04972: Pancreatic secretion	16	0.033
	chx04270: Vascular smooth muscle contraction	19	0.037
	chx04725: Cholinergic synapse	18	0.043
	chx04911: Insulin secretion	14	0.053
	chx04726: Serotonergic synapse	17	0.072
	chx04915: Estrogen signaling pathway	15	0.091
	chx04961: Endocrine and other factor-regulated calcium reabsorption	8	0.094
	chx04912: GnRH signaling pathway	13	0.131
	chx04918: Thyroid hormone synthesis	11	0.137
	chx04924: Renin secretion	10	0.161
	chx04611: Platelet activation	16	0.273
	chx04916: Melanogenesis	11	0.458

3 (1.14, KEGG_PATHWAY)	chx04310: Wnt signaling pathway	14	0.599
	chx05204: Chemical carcinogenesis	12	0.033
	chx00980: Metabolism of xenobiotics by cytochrome P450	10	0.075
	chx00982: Drug metabolism - cytochrome P450	10	0.075
	chx00830: Retinol metabolism	9	0.152

Supplementary Fig. 40 (Additional file 2) shows the biological processes, cellular components and molecular functions associated with the genes in the global CNVR according to functional classification in the Panther Classification System using the *Bos taurus* gene set. The top three biological processes were cellular processes, metabolic processes and biological regulation. The top three cell components were cells, organelles and membranes. Similarly, the top three molecular functions were catalytic activity, (cell) binding and transporter activity.

Functional annotation for CNVR private to breeds and populations

CNVR private to populations and breeds overlapped with 172 and 620 genes, respectively. The GO terms associated with these genes based on functional analysis are listed in Supplementary Table 6 (Additional file 1). The genes that overlapped with the CNVR private to breeds were not significantly enriched in biological processes, molecular functions and cellular components, while the ones that overlapped with the CNVR private to populations were significantly enriched ($P \leq 0.05$) with such terms as aldosterone synthesis and secretion; glucagon signaling pathway; insulin secretion; glutamatergic synapse; thyroid hormone synthesis; gastric acid secretion and phosphatidylinositol signaling system. The most common CNVR (chr6:115,822,332 – 115,825,687) includes the gene TMEM129 (transmembrane protein 129) that has been reported to be responsible for ubiquitination and proteasome-mediated degradation of misformed or unassembled proteins in the cytosol [45–47], and belongs to a network responsible for cellular assembly and organization, cellular function and maintenance, and cell cycle [48]. Supplementary Fig. 41 (Additional file 2) shows the biological processes, cellular components and molecular functions associated with the genes in the CNVR that were private to breeds and populations according to functional classification in the Panther Classification System using the *Bos taurus* gene set. The top three terms for biological processes, cell components and molecular functions were the same as for the genes in the global CNVR with slightly

different proportions. It was noted that the biological process “response to stimulus” (GO:0050896) had a higher representation for highly differentiated CNV than for global or private CNVR.

Discussion

This study identified CNV and CNVR in the goat genome using WGS data. Use of WGS for CNV detection is highly encouraged, because it overcomes many of the shortcomings of the other CNV detection methods such as the ones using array CGH and SNP data [18–20]. Genome-wide studies to discover CNV have already been done in other domesticated species, such as in *Sus scrofa* [49], *Bos taurus* [37, 50] and *Felis catus* [38]. Here we provide a first glimpse of the goat genome CNV map at a dense genome coverage, using animals from 34 diverse breeds from the African continent. This addition is an important contribution, as goats are an important source of income and high-quality animal protein for small holder farmers in Africa.

We used two software suites (LUMPY [29] and Manta [36]) for detecting SV to increase our confidence in the SV calls. Both software packages use split read and read-pair methods. They complement each other in that LUMPY makes use of read depth methods, while Manta draws heavily on genome assembly methods. Taking the intersection of SV calls from the two methods gives us confidence that the number of false positives in the SV calls was kept to a minimum, although this means that some true SV were possibly filtered out.

This study has shown that there are wide variations in the number and sizes of CNV in the goat genome between chromosomes, individuals and breeds. However, considering the small and variable numbers of samples within breeds, breed comparisons are not particularly meaningful. The results suggest that there are negligible differences in the sizes of CNV between populations. Some of the CNV displayed large differences between populations, suggestive of population-specific selective pressures.

A large proportion of the global CNVR identified in this study (65.1%) are within the CNVR reported by Liu et al. [17]. The remaining 34.9% may comprise false positive CNVR and CNVR that were missed by the PennCNV algorithm used in the other study, considering the limitation of CNV detection using SNP data, which include limited coverage for genome, low resolution, and difficulty in detecting novel and

rare mutations. The CNVR coverage of 2.4% (59.2 Mb of about 2,466 Mb of autosomal genome) found in this study is lower than the 4.8–9.5% SV coverage in the human genome [12], comparable to 55.6 Mb (2.0%) reported for cattle [37], later revised to 87.5 Mb (3.1%) [51].

V_{ST} analysis showed that several CNV were highly differentiated among and across the populations. The genes in the highly differentiated CNV included BCO2 (Madagascar vs West African population differentiation), CCSER1 (FAM190A) (Boer vs East African), FAM155A (across all populations), GNRHR (Boer vs Madagascar; Boer vs West African), IMMP2L (East vs Southern African), LAMA3 (East African vs Madagascar), NAALADL2 (East vs Southern African), TAFA2 (FAM19A2) (East vs Southern African) and TOMM70 (across all the populations). Våge and Boman [52] reported that BCO2 is associated with the accumulation of carotenoids in the adipose tissue of sheep, leading to the yellow fat syndrome. The quality of semen (including total sperm motility, average path velocity and beat cross frequency) in Holstein-Friesian bulls has been associated with CCSER1 (FAM190A) as well as FAM155A [53]. GNRHR has been associated with number of days to first service after calving in dairy cattle [54] while IMMP2L is associated with cow conception rate [55]. The partial deletion of LAMA3 is responsible for epidermolysis bullosa in horses [56]; NAALADL2 is believed to be responsible for immune homeostasis [57], and TAFA2 (FAM19A2) is believed to be responsible for the regulation of feed intake and metabolic activities in mice [58]. Yamano et al [59] reported that TOMM70 is responsible for integral mitochondrion proteins and for metabolism.

Functional annotation and clustering analysis revealed that the CNVR identified in the study have genes that are significantly enriched with many biological processes, molecular functions and cellular components, some of the most significant of which are retrograde endocannabinoid signaling, circadian entrainment and long-term potentiation. The retrograde endocannabinoid signaling system is a complex and diverse regulator of synaptic function [60], and is responsible for many diseases in the nervous system and peripheral organs. In the human genome, this system is widely considered as a potential target for treating conditions such as alcoholism [61]. A CNVR in the cannabinoid receptor 2 (CNR2) region has been reported in the human genome, but its effect has not been fully characterized [62]. Zajkowska et al. [63] suggested that there is need to explore genetic variation in

the system from the perspective of copy number of variations.

Circadian entrainment is an important aspect of animal behavior and adaptation, especially considering the wide range of environmental conditions the animals are exposed to. An example of goat adaptation to the environment is their ability to rapidly change the size of their foreguts in response to changes in the environment [64]. Goats tend to be active during some parts of the day only [65], and this varies with season [65], suggesting a considerable amount of circadian entrainment. The increased importance of the biological process “response to stimulus” (GO:0050896) in the highly differentiated CNV may also support the hypothesis of the importance of circadian entrainment in goats.

Conclusions

This study presents the first fine CNV map of the African goats based on WGS data. This information will prove invaluable for further improvement of goats, especially on African continent, as more phenotype data becomes available, through CNV or CNVR association analyses and other approaches.

Methods

Sample description

The data used in this study was generated from 182 goats representing 34 breeds from 9 Sub-Saharan African countries (Ethiopia, Kenya, Madagascar, Malawi, Mali, Mozambique, Tanzania, Uganda, and Zimbabwe), and these countries were grouped into four populations based on geographic locations and a fifth population of Boer goats obtained in Tanzania and Zimbabwe. The Boer goat is a special breed widely used in Africa and much of the world [66]. The samples were previously genotyped using the Illumina Goat SNP50 BeadChip [67] as described by Bertolini et al. [68], Cardoso et al. [69] and Colli et al. [70], and some of them were also used for detection of CNV using 50 K SNP chip data, as reported by Liu et al. [17]. A list of the breeds, populations and samples sizes used in the analysis is given in Table 4 while a full list of individual samples can be found in Supplementary Table 7 (Additional file 1).

Sample processing was done by Edinburgh Genomics using the Edinburgh Clinical Genomics method. This approach uses Illumina SeqLab products and services, including, Illumina TruSeq library preparation, Illumina cBot2 cluster generation, Illumina HiSeqX sequencing, Hamilton Microlab STAR

integrative automation, and Genologics Clarity LIMS X Edition as outlined in Supplementary Table 8

(Additional file 1).

Table 4
List of the breeds used in the analysis

Population	Population Code	Breed name	Breed Code	Country	Number of samples
Boer	BOE	Boer	BOE	Tanzania	2
Boer	BOE	Boer	BOE	Zimbabwe	7
East African	EAF	Abergelle	ABR	Ethiopia	6
East African	EAF	Galla	GAL	Kenya	7
East African	EAF	Gogo	GOG	Tanzania	7
East African	EAF	Gumez	GUM	Ethiopia	4
East African	EAF	Keffa	KEF	Ethiopia	7
East African	EAF	Landin	LND	Mozambique	5
East African	EAF	Maasai	MAA	Tanzania	7
East African	EAF	Manica	MAN	Mozambique	3
East African	EAF	Malya	MLY	Tanzania	7
East African	EAF	Norwegian	NRW	Tanzania	3
East African	EAF	Pare White	PRW	Tanzania	6
East African	EAF	Saanen	SAA	Tanzania	4
East African	EAF	Small East African	SEA	Kenya	7
East African	EAF	Small East African	SEA	Mozambique	6
East African	EAF	Sonjo	SNJ	Tanzania	2
East African	EAF	Woyito Guji	WYG	Ethiopia	7
Madagascar	MAD	Androy	AND	Madagascar	4
Madagascar	MAD	Diana	DIA	Madagascar	3
Madagascar	MAD	Menabe	MEN	Madagascar	7
Madagascar	MAD	Sofia	SOF	Madagascar	6
Madagascar	MAD	SudOuest	SOU	Madagascar	7
Southern African	SAF	Balaka-Ulongwe	BAW	Malawi	2
Southern African	SAF	Dedza	DZD	Malawi	4
Southern African	SAF	Lilongwe	LGW	Malawi	3
Southern African	SAF	Mashona	MSH	Zimbabwe	7
Southern African	SAF	Matebele	MTB	Zimbabwe	7
Southern African	SAF	Nsanje	NSJ	Malawi	6
Southern African	SAF	Thyolo	THY	Malawi	7
West African	WAF	Guerra	GUE	Mali	6
West African	WAF	Maure	MAU	Mali	1
West African	WAF	Naine	NAI	Mali	5
West African	WAF	Peulh	PEU	Mali	1
West African	WAF	Soudanaise	SDN	Mali	7
West African	WAF	Targui	TAR	Mali	2

Sequence alignment

Sequence alignment was done using the Burrow-Wheeler Alignment (BWA) tool version 0.7.13-r1126 with the maximal exact matches (MEM) “mem” option [71]. The reads were aligned to the ARS1 Capra hircus (goat) reference assembly (https://www.ncbi.nlm.nih.gov/assembly/GCF_001704415.1/) [72].

The aligned reads were processed into binary sequence alignment map (BAM) format using SAMTools version 1.8 [73].

Detection of SV

SV were detected using LUMPY version 0.2.13-85-gc1bcea1 and Manta version 1.5.1, which are two of

the most commonly used algorithms for detecting SV. In LUMPY, the “lumpyexpress” script was used. This script runs automated breakpoint detection for standard analyses. It uses SAMBLASTER [74] to extract split and discordant reads from BWA-MEM-aligned Binary Sequence Alignment Map (BAM) files. Default options were used, including minimum non-overlap and minimum sample weight set to 20 and 4, respectively. In Manta, the “configManta.py” script was used to process each sample, with default options including minimum variant candidate size (8); minimum candidate spanning count (3); minimum scored variant size (50); minimum diploid variant score (10); minimum diploid variant score pass point (20); minimum somatic score (10); and minimum somatic score pass point (30). The “runWorkflow.py” scripts were run in parallel to extract the SV for each sample.

Post-processing of SV

SV from LUMPY were genotyped with svtyper version 0.6.1 [75], which uses a Bayesian maximum likelihood algorithm to determine the most likely genotype of each base-pair. Variant call format (VCF) files from the two software packages were converted to browser extensible data (BED) format for downstream analysis using svtools version 0.5.0 [76]. In both LUMPY and Manta, SV with unknown or imprecise break points were discarded. A minimum of a sum of five paired-end reads (PR) and five split reads (SR) were required as evidence supporting an SV. Consensus SV were obtained by identifying the intersection of the SV from LUMPY and Manta using BEDTools version 2.26.0 [77] with default settings.

Derivation of copy number variations

CNV were defined as SV duplications and deletions longer than 50 bp [78]. SV longer than 3 Mb were also filtered out, because putative CNV in the goat genome are usually much shorter than this length. Visualization of the SV was done using R [79] package circlize version 0.4.7 [80].

Population CNV differentiation

A measure of population differentiation (V_{ST}) as described by Redon et al. [10] was computed based on normalized read count values for each CNV, similar to the method used in PECNV as described by Liu et al. [81], which was in turn based on clustering algorithms described by Cridland et al. [82] and transposable element detection algorithms described by Rogers et al. [83]. Read count values were corrected for size of the consensus CNV, batch effect, variable GC content and genomic mappability

as described by Liu et al. [81]. Regional and batch effect correction was done by computing reads per kb per million mapped reads (RPKM) as described by Mortazavi et al. [84], where $RPKM = \frac{10^9 * RC}{TRC * S}$, where RC is the read count of a region, S is the size of the region and TRC is the total number of mapped reads in the library. GC content and mappability correction was done on the RPKM using the formula used by Yoon et al. [28], where adjusted read count is given by $\frac{RPKM * m}{m_{GC}}$ where m_{GC} is the median GC content of all regions with the same read count and m is the median GC of all regions. This approach is similar to the read depth approaches used in CNVnator [27] and in CNVcaller [85]. The normalized read count values were treated as proxies of log R ratio (LRR) values normally obtained from array analysis. As defined by Redon et al [10], V_{ST} was computed as $\frac{V_T - V_S}{V_T}$, where V_T is the variance in LRR among all unrelated individuals and V_S is the average variance in LRR within each population. CNV V_{ST} testing was done pairwise (for each combination of two populations) and (separately) across all the 5 populations. CNV with V_{ST} values above the 99th percentile of all V_{ST} values for each comparison were treated as being highly differentiated. We searched for these highly differentiated CNV in the Golden Helix Genome Browse® software (version 3.0.0) (<https://www.goldenhelix.com/>) using the ARS1 caprine genome reference assembly to identify the genes in the CNV. We also carried out functional analysis of the genes in DAVID and in the Panther Classification System [86].

Determination of CNV regions

CNV regions (CNVR) were obtained by merging CNV that overlapped by at least 1 bp within populations (population CNVR) and across all the individuals (global CNVR) using the “merge” function in BEDTools version 2.26.0 [77].

CNVR functional annotation and gene enrichment analysis

A list of genes for the goat genome was downloaded from the NCBI website (<https://www.ncbi.nlm.nih.gov/gene>). To identify if genes in the CNVR have significant biological, cellular or molecular function, version 6.8 [87–89]. Functional annotation and clustering analyses were used to identify overrepresented Gene Ontology (GO) [90] annotation categories and Kyoto Encyclopedia of Genes and Genomes (KEGG) pathways [91] associated with the CNVR. The following gene databases

were used in the analysis: GO [biological process (GOTERM_BP_DIRECT), cellular component (GOTERM_CC_DIRECT), and molecular function (GOTERM_MF_DIRECT)], INTERPRO [92], KEGG_PATHWAY and UP_KEYWORDS [93]. Further information about various genes was obtained from the GeneCards (www.genecards.org) and NCBI (<https://www.ncbi.nlm.nih.gov/gene>) gene databases. Functional analysis was done using default parameters, with significance of enriched terms determined at $P \leq 0.05$. Functional classification was also carried out using the Bos taurus gene set in the Panther Classification System [86].

Abbreviations

Abbreviations

ABR	Abergelle
AND	Androy
BAM	Binary Sequence Alignment Map
BAW	Balaka-Ulongwe
BED	Browser Extensible Data
BOE	Boer
BWA	Burrow-Wheeler Alignment
CGH	comparative genomic hybridization
CNV	copy number variation
CNVR	copy number variation region
DAVID	Database for Annotation, Visualization, and Integrated Discovery
DIA	Diana
DZD	Dedza
EAF	East African
EWT	event-wise testing
GAL	Galla

GO	Gene Ontology
GOG	Gogo
GSEA	gene set enrichment analysis
GUE	Guerra
GUM	Gumez
KEF	Keffa
KEGG	Kyoto Encyclopedia of Genes and Genomes
LGW	Lilongwe
LND	Landin
MAA	Maasai
MAD	Madagascar
MAN	Manica
MAU	Maure
MEM	maximal exact matches
MEN	Menabe
MLY	Malya
MSH	Mashona
MTB	Matebele
NAI	Naine
NRW	Norwegian
NSJ	Nsanje
PEM	paired-end mapping
PEU	Peulh
PR	paired-end reads
PRW	Pare White
RD	read depth
RPKM	reads per kb per million mapped reads

SAA	Saanen
SAF	Southern African
SDN	Soudanaise
SEA	Small East African
SNJ	Sonjo
SNP	single nucleotide polymorphisms
SOF	Sofia
SOU	SudOuest
SR	split reads
SV	structural variation
TAR	Targui
THY	Thyolo
VCF	Variant call format
WAF	West African
WGS	whole genome sequencing
WYG	Woyito Guji

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable

Competing interests

The authors declare that there are no competing interests.

Funding

This research was done with funding from the Feed the Future project: Improving Livestock Productivity through Enhanced Breeding Programs (Agricultural Research Service/USDA Project No.: 8042-31000-001-02S). The data were collected by the African Goat Improvement Network (AGIN) (<https://www.ars.usda.gov/office-of->

international-research-programs/ftf-livestock-improvement/).

This research used resources provided by the SCINet project of the USDA Agricultural Research Service, ARS project number 0500-00093-001-00-D.

Authors' Contributions

CPVD, BDR, GM and JS conceived the idea of this work. ELK and the VarGoats Consortium carried out the sequencing of the samples and pre-processing of the sequence data. WN, BDR, CPVT, GM and JS carried out the analysis and drafted the manuscript. WN, GM, MW, LJB, TNG, HAM, HNN, ELK, MJWG, ML, GEL, CPVT, BDR and JS reviewed the manuscript.

Corresponding author

Benjamin D. Rosen (Ben.Rosen@usda.gov)

The VarGoats Consortium

<http://www.goatgenome.org/vargoats.html>

Affiliations

University of Natural Resources and Life Sciences, Vienna, Austria

Wilson Nandolo

Gábor Mészáros

Maria Wurzinger

Johann Sölkner

Lilongwe University of Agriculture and Natural Resources, Lilongwe, Malawi

Wilson Nandolo

Liveness Jessica J. Banda

Timothy N. Gondwe

National Livestock Resources Research Institute, Tororo, Uganda

Henry Aaron A. Mulindwa

Kyambogo University, Kyambogo, Uganda

Helen N. Nakimbugwe

The Roslin Institute, University of Edinburgh, Edinburgh, Scotland, United Kingdom

Emily L. Clark

Animal Genomics and Improvement Laboratory, USDA-ARS, Beltsville, MD, USA

1. Jennifer Woodward-Greene

Mei Liu

George E. Liu

Curtis P. Van Tassell

Benjamin D. Rosen

Acknowledgements

Whole genome sequencing libraries were prepared and sequenced by Edinburgh Genomics and funded via Biotechnology and Biological Sciences Research Council research grant (BBS/OS/GC/000012F) ‘Reference genome and population sequencing of African goats’ awarded to The Roslin Institute.

Author information

N/A

Availability of data and materials

The dataset supporting the conclusions of this article is available at the National Center for Biotechnology Information Sequence Read Archive as detailed in Additional file 1 (Supplementary Table 7)

References

1. Escaramís G, Docampo E, Rabionet R. A decade of structural variants: Description, history and methods to detect structural variation. *Brief Funct Genomics*. 2015;14:305–14.
doi:10.1093/bfgp/elv014.
2. Matsuzaki H, Wang PH, Hu J, Rava R, Fu GK. High resolution discovery and confirmation of copy number variants in 90 Yoruba Nigerians. *Genome Biol*. 2009;10:1–18.
3. Fan S, Meyer A. Evolution of genomic structural variation and genomic architecture in the adaptive radiations of African cichlid fishes. *Front Genet*. 2014;5 JUN:163.
doi:10.3389/fgene.2014.00163.
4. Conrad DF, Hurler ME. The population genetics of structural variation. *Nat Genet*. 2007;39:S30–

6. doi:10.1038/ng2042.
5. Sebat J, Levy DL, McCarthy SE. Rare structural variants in schizophrenia: one disorder, multiple mutations; one mutation, multiple disorders. *Trends Genet.* 2009;25:528–35.
doi:10.1016/j.tig.2009.10.004.
6. Sebat J, Lakshmi B, Troge J, Alexander J, Young J, Lundin P, et al. Large-scale copy number polymorphism in the human genome. *Science* (80-). 2004;305:525–8.
doi:10.1126/science.1098918.
7. Iafrate AJ, Feuk L, Rivera MN, Listewnik ML, Donahoe PK, Qi Y, et al. Detection of large-scale variation in the human genome. *Nat Genet.* 2004;36:949–51. doi:10.1038/ng1416.
8. Feuk L, Marshall CR, Wintle RF, Scherer SW. Structural variants: changing the landscape of chromosomes and design of disease studies. *Hum Mol Genet.* 2006;15 Spec No suppl_1:R57–66.
doi:10.1093/hmg/ddl057.
9. Huang CRL, Schneider AM, Lu Y, Niranjan T, Shen P, Robinson MA, et al. Mobile interspersed repeats are major structural variants in the human genome. *Cell.* 2010;141:1171–82.
doi:10.1016/j.cell.2010.05.026.
10. Redon R, Ishikawa S, Fitch KR, Feuk L, Perry GH, Andrews TD, et al. Global variation in copy number in the human genome. *Nature.* 2006;444:444–54. doi:10.1038/nature05329.
11. Beckmann JS, Estivill X, Antonarakis SE. Copy number variants and genetic traits: Closer to the resolution of phenotypic to genotypic variability. *Nat Rev Genet.* 2007;8:639–46.
doi:10.1038/nrg2149.
12. Zarrei M, MacDonald JR, Merico D, Scherer SW. A copy number variation map of the human genome. *Nat Rev Genet.* 2015;16:172–83. doi:10.1038/nrg3871.
13. Girirajan S, Rosenfeld JA, Coe BP, Parikh S, Friedman N, Goldstein A, et al. Phenotypic Heterogeneity of Genomic Disorders and Rare Copy-Number Variants. *N Engl J Med.* 2012;367:1321–31. doi:10.1056/NEJMoa1200395.

4. Cooper GM, Coe BP, Girirajan S, Rosenfeld JA, Vu TH, Baker C, et al. A copy number variation morbidity map of developmental delay. *Nat Genet.* 2011;43:838–46. doi:10.1038/ng.909.
5. Fontanesi L, Martelli PL, Beretti F, Riggio V, Dall’Olio S, Colombo M, et al. An initial comparative map of copy number variations in the goat (*Capra hircus*) genome. *BMC Genomics.* 2010;11:639. doi:10.1186/1471-2164-11-639.
6. Nandolo W, Lamuno D, Banda L, Gondwe T, Mulindwa H, Nakimbugwe H, et al. Distribution of copy number variants in the genomes of East African goat breeds. In: *Proceedings of the World Congress on Genetics Applied to Livestock Production. World Congress on Genetics Applied to Livestock Production; 2018.* p. 470. <http://www.wcgalp.org/proceedings/2018/distribution-copy-number-variants-genomes-east-african-goat-breeds>. Accessed 27 Aug 2018.
7. Liu M, Zhou Y, Rosen BD, Van Tassell CP, Stella A, Tosser-Klopp G, et al. Diversity of copy number variation in the worldwide goat population. *Heredity (Edinb).* 2018.
8. Snijders AM, Nowak N, Segraves R, Blackwood S, Brown N, Conroy J, et al. Assembly of microarrays for genome-wide measurement of DNA copy number. *Nat Genet.* 2001;29:263–4. doi:10.1038/ng754.
9. Shendure J, Ji H. Next-generation DNA sequencing. *Nat Biotechnol.* 2008;26:1135–45. doi:10.1038/nbt1486.
10. Zhao M, Wang Q, Wang Q, Jia P, Zhao Z. Computational tools for copy number variation (CNV) detection using next-generation sequencing data: features and perspectives. *BMC Bioinformatics.* 2013;14 Suppl 11:S1. doi:10.1186/1471-2105-14-S11-S1.
11. Mills RE, Walter K, Stewart C, Handsaker RE, Chen K, Alkan C, et al. Mapping copy number variation by population-scale genome sequencing. *Nature.* 2011;470:59–65. doi:10.1038/nature09708.
12. Yang H, Chen G, Lima L, Fang H, Jimenez L, Li M, et al. HadoopCNV- A dynamic programming imputation algorithm to detect copy number variants from sequencing data. *bioRxiv.*

- 2017;;124339. doi:10.1101/124339.
13. Fan X, Abbott TE, Larson D, Chen K. BreakDancer: Identification of genomic structural variation from paired-end read mapping. In: *Current Protocols in Bioinformatics*. Hoboken, NJ, USA: John Wiley & Sons, Inc.; 2014. p. 15.6.1-15.6.11. doi:10.1002/0471250953.bi1506s45.
 14. Ye K, Schulz MH, Long Q, Apweiler R, Ning Z. Pindel: A pattern growth approach to detect break points of large deletions and medium sized insertions from paired-end short reads. *Bioinformatics*. 2009;25:2865–71. doi:10.1093/bioinformatics/btp394.
 15. Rausch T, Zichner T, Schlattl A, Stütz AM, Benes V, Korbel JO. DELLY: Structural variant discovery by integrated paired-end and split-read analysis. *Bioinformatics*. 2012;28:i333–9. doi:10.1093/bioinformatics/bts378.
 16. Xie C, Tammi MT. CNV-seq, a new method to detect copy number variation using high-throughput sequencing. *BMC Bioinformatics*. 2009;10:80. doi:10.1186/1471-2105-10-80.
 17. Abyzov A, Urban AE, Snyder M, Gerstein M. CNVnator: An approach to discover, genotype, and characterize typical and atypical CNVs from family and population genome sequencing. *Genome Res*. 2011;21:974–84. doi:10.1101/gr.114876.110.
 18. Yoon S, Xuan Z, Makarov V, Ye K, Sebat J. Sensitive and accurate detection of copy number variants using read depth of coverage. *Genome Res*. 2009;19:1586–92. doi:10.1101/gr.092981.109.
 19. Layer RM, Chiang C, Quinlan AR, Hall IM, Quinlan AR. LUMPY: A probabilistic framework for structural variant discovery. *Genome Biol*. 2014;15:R84. doi:10.1186/gb-2014-15-6-r84.
 20. Pirooznia M, Goes F, Zandi PP. Whole-genome CNV analysis: Advances in computational approaches. *Front Genet*. 2015;6 MAR:138. doi:10.3389/fgene.2015.00138.
 21. Tattini L, D’Aurizio R, Magi A. Detection of Genomic Structural Variants from Next-Generation Sequencing Data. *Front Bioeng Biotechnol*. 2015;3:92. doi:10.3389/fbioe.2015.00092.
 22. Ye K, Hall G, K Y, G H. Structural Variation Detection from Next Generation Sequencing. *J Next*

- Gener Seq Appl. 2015;01. doi:10.4172/2469-9853.S1-007.
3. Guan P, Sung WK. Structural variation detection using next-generation sequencing data: A comparative technical review. *Methods*. 2016;102:36–49. doi:10.1016/j.ymeth.2016.01.020.
 4. Liu B, Conroy JM, Morrison CD, Odunsi AO, Qin M, Wei L, et al. Structural variation discovery in the cancer genome using next generation sequencing: Computational solutions and perspectives. *Oncotarget*. 2015;6:5477–89. doi:10.18632/oncotarget.3491.
 5. Arthur JG, Chen X, Zhou B, Urban AE, Wong WH. Detection of complex structural variation from paired-end sequencing data. *bioRxiv*. 2018;:200170. doi:10.1101/200170.
 6. Chen X, Schulz-Trieglaff O, Shaw R, Barnes B, Schlesinger F, Källberg M, et al. Manta: Rapid detection of structural variants and indels for germline and cancer sequencing applications. *Bioinformatics*. 2016;32:1220–2. doi:10.1093/bioinformatics/btv710.
 7. Bickhart DM, Liu GE, Eichler EE, Matukumalli LK, Song J, Van Tassell CP, et al. Copy number variation of individual cattle genomes using next-generation sequencing. *Genome Res*. 2012;22:778–90. doi:10.1101/gr.133967.111.
 8. Genova F, Longeri M, Lyons LAA, Bagnato A, Gandolfi B, Aberdein D, et al. First genome-wide CNV mapping in FELIS CATUS using next generation sequencing data. *BMC Genomics*. 2018;19:1–2. doi:10.1186/s12864-018-5297-2.
 9. Sohrabi SS, Mohammadabadi M, Wu D-D, Esmailizadeh A. Detection of breed-specific copy number variations in domestic chicken genome. *Genome*. 2018;61:7–14. doi:10.1139/gen-2017-0016.
 10. Freedman AH, Gronau I, Schweizer RM, Ortega-Del Vecchyo D, Han E, Silva PM, et al. Genome Sequencing Highlights the Dynamic Early History of Dogs. *PLoS Genet*. 2014;10:e1004016. doi:10.1371/journal.pgen.1004016.
 11. Stella A, Nicolazzi EL, Van Tassell CP, Rothschild MF, Colli L, Rosen BD, et al. AdaptMap: exploring goat diversity and adaptation. *Genet Sel Evol*. 2018;50:61. doi:10.1186/s12711-018-

0427-5.

- .2. Teo SM, Pawitan Y, Ku CS, Chia KS, Salim A. Statistical challenges associated with detecting copy number variations with next-generation sequencing. *Bioinformatics*. 2012;28:2711–8. doi:10.1093/bioinformatics/bts535.
- .3. Neupane M, Hoff JL, Taylor JF, Seabury CM, Womack JE, Bovine T, et al. Refinement of the DST locus associated with bovine respiratory disease complex in Holstein calves. *J Anim Sci*. 2016;94 suppl_5:361–361. doi:10.2527/jam2016-0753.
- .4. Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, et al. Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles. *Proc Natl Acad Sci U S A*. 2005;102:15545–50. doi:10.1073/pnas.0506580102.
- .5. Van De Weijer ML, Bassik MC, Luteijn RD, Voorburg CM, Lohuis MAM, Kremmer E, et al. A high-coverage shrna screen identifies TMEM129 as an E3 ligase involved in ER-associated protein degradation. *Nat Commun*. 2014;5:3832. doi:10.1038/ncomms4832.
- .6. Van De Weijer ML, Van Muijlwijk GH, Visser LJ, Costa AI, Wiertz EJHJ, Lebbink RJ. The E3 ubiquitin ligase TMEM129 is a tri-spanning transmembrane protein. *Viruses*. 2016;8. doi:10.3390/v8110309.
- .7. van den Boomen DJH, Timms RT, Grice GL, Stagg HR, Skodt K, Dougan G, et al. TMEM129 is a Derlin-1 associated ERAD E3 ligase essential for virus-induced degradation of MHC-I. *Proc Natl Acad Sci*. 2014;111:11425–30. doi:10.1073/pnas.1409099111.
- .8. Yu F, Shen XY, Fan L, Yu ZC. Genome-wide analysis of genetic variations assisted by Ingenuity Pathway Analysis to comprehensively investigate potential genetic targets associated with the progression of hepatocellular carcinoma. *Eur Rev Med Pharmacol Sci*. 2014.
- .9. Wang K, Hadley D, Liu R, Bucan M, Li M, Glessner J, et al. PennCNV: An integrated hidden Markov model designed for high-resolution copy number variation detection in whole-genome SNP genotyping data. *Genome Res*. 2007;17:1665–74. doi:10.1101/gr.6861907.

10. Stothard P, Choi JW, Basu U, Sumner-Thomson JM, Meng Y, Liao X, et al. Whole genome resequencing of black Angus and Holstein cattle for SNP and CNV discovery. *BMC Genomics*. 2011;12:559. doi:10.1186/1471-2164-12-559.
11. Bickhart DM, Xu L, Hutchison JL, Cole JB, Null DJ, Schroeder SG, et al. Diversity and population-genetic properties of copy number variations and multicopy genes in cattle. *DNA Res*. 2016;23:253–62. doi:10.1093/dnares/dsw013.
12. Våge DI, Boman IA. A nonsense mutation in the beta-carotene oxygenase 2 (BCO2) gene is tightly associated with accumulation of carotenoids in adipose tissue in sheep (*Ovis aries*). *BMC Genet*. 2010;11:10. doi:10.1186/1471-2156-11-10.
13. Borowska A, Szwaczkowski T, Kamiński S, Hering DM, Kordan W, Lecewicz M. Identification of genome regions determining semen quality in Holstein-Friesian bulls using information theory. *Anim Reprod Sci*. 2018;192:206–15. doi:10.1016/j.anireprosci.2018.03.012.
14. Lirón JP, Prando AJ, Fernández ME, Ripoli M V., Rogberg-Muñoz A, Goszczynski DE, et al. Association between GNRHR, LHR and IGF1 polymorphisms and timing of puberty in male Angus cattle. *BMC Genet*. 2012;13:26. doi:10.1186/1471-2156-13-26.
15. Jiang J, Ma L, Prakapenka D, VanRaden PM, Cole JB, Da Y. A large-scale genome-wide association study in U.S. Holstein cattle. *Front Genet*. 2019;10 MAY:412. doi:10.3389/fgene.2019.00412.
16. Yuen WY, Pasmooij AMG, Stellingsma C, Jonkman MF. Enamel defects in carriers of a novel LAMA3 mutation underlying epidermolysis bullosa. *Acta Derm Venereol*. 2012;92:695–6. doi:10.2340/00015555-1341.
17. Onouchi Y. Genetics of Kawasaki disease: - What we know and don't know. *Circ J*. 2012;76:1581–6.
18. Okada J, Yamada E, Saito T, Ozawa A, Nakajima Y, Pessin JE, et al. Analysis of FAM19A2/TAFA-2 function. *Physiol Behav*. 2019;208:112581. doi:10.1016/j.physbeh.2019.112581.

9. Yamano K, Matsuda N, Tanaka K. The ubiquitin signal and autophagy: an orchestrated dance leading to mitochondrial degradation. *EMBO Rep.* 2016;17:300–16. doi:10.15252/embr.201541486.
10. Castillo PE, Younts TJ, Chávez AE, Hashimotodani Y. Endocannabinoid Signaling and Synaptic Function. *Neuron.* 2012;76:70–81. doi:10.1016/j.neuron.2012.09.020.
11. Basavarajappa BS. The endocannabinoid signaling system: a potential target for next-generation therapeutics for alcoholism. *Mini Rev Med Chem.* 2007;7:769–79. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=1975858&tool=pmcentrez&rendertype=abstract>. Accessed 17 Aug 2019.
12. Ishiguro H, Leonard CM, Sgro S, Onaivi ES. Cannabinoid Receptor Gene Variations in Neuropsychiatric Disorders. In: Murillo-Rodríguez E, editor. *Endocannabinoids: Molecular, Pharmacological, Behavioral and Clinical Features*. 1st edition. Baltimore: Bentham Science Publishers; 2013. p. 3–24.
13. Zajkowska ZE, Englund A, Zunszain PA. Towards a personalized treatment in depression: endocannabinoids, inflammation and stress response. *Pharmacogenomics.* 2014;15:687–98. doi:10.2217/pgs.14.40.
14. Silanikove N. The physiological basis of adaptation in goats to harsh environments. *Small Rumin Res.* 2000;35:181–93. doi:10.1016/S0921-4488(99)00096-6.
15. Piccione G, Giannetto C, Casella S, Caola G. Seasonal change of daily motor activity rhythms in *Capra hircus*. *Can J Anim Sci.* 2008;88:351–5. doi:10.4141/CJAS07125.
16. Malan S. The improved Boer goat. *Small Rumin Res.* 2000;36:165–70. doi:10.1016/S0921-4488(99)00160-1.
17. Tosser-Klopp G, Bardou P, Bouchez O, Cabau C, Crooijmans R, Dong Y, et al. Design and characterization of a 52K SNP chip for goats. *PLoS One.* 2014;9:e86227. doi:10.1371/journal.pone.0086227.

8. Bertolini F, Cardoso TF, Marras G, Nicolazzi EL, Rothschild MF, Amills M. Genome-wide patterns of homozygosity provide clues about the population history and adaptation of goats. *Genet Sel Evol.* 2018;50:59. doi:10.1186/s12711-018-0424-8.
9. Cardoso TF, Amills M, Bertolini F, Rothschild M, Marras G, Boink G, et al. Patterns of homozygosity in insular and continental goat breeds. *Genet Sel Evol.* 2018;50:1–11. doi:10.1186/s12711-018-0425-7.
10. Colli L, Milanese M, Talenti A, Bertolini F, Chen M, Crisà A, et al. Genome-wide SNP profiling of worldwide goat populations reveals strong partitioning of diversity and highlights post-domestication migration routes. *Genet Sel Evol.* 2018;50:58. doi:10.1186/s12711-018-0422-x.
11. Li H, Durbin R. Fast and accurate long-read alignment with Burrows-Wheeler transform. *Bioinformatics.* 2010;26:589–95. doi:10.1093/bioinformatics/btp698.
12. Bickhart DM, Rosen BD, Koren S, Sayre BL, Hastie AR, Chan S, et al. Single-molecule sequencing and chromatin conformation capture enable de novo reference assembly of the domestic goat genome. *Nat Genet.* 2017;49:643–50.
13. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence Alignment/Map format and SAMtools. *Bioinformatics.* 2009;25:2078–9. doi:10.1093/bioinformatics/btp352.
14. Faust GG, Hall IM. SAMBLASTER: Fast duplicate marking and structural variant read extraction. *Bioinformatics.* 2014;30:2503–5. doi:10.1093/bioinformatics/btu314.
15. Chiang C, Layer RM, Faust GG, Lindberg MR, Rose DB, Garrison EP, et al. SpeedSeq: Ultra-fast personal genome analysis and interpretation. *Nat Methods.* 2015;12:966–8. doi:10.1038/nmeth.3505.
16. Larson DE, Abel HJ, Chiang C, Badve A, Das I, Eldred JM, et al. Svtools: Population-Scale Analysis of Structural Variation. *Bioinformatics.* 2019. doi:10.1093/bioinformatics/btz492.
17. Quinlan AR, Hall IM. BEDTools: A flexible suite of utilities for comparing genomic features. *Bioinformatics.* 2010;26:841–2. doi:10.1093/bioinformatics/btq033.

8. Liu GE, Bickhart DM. Copy number variation in the cattle genome. *Funct Integr Genomics*. 2012;12:609–24. doi:10.1007/s10142-012-0289-9.
9. R Development Core Team. R: A Language and Environment for Statistical Computing. Vienna, Austria: R Foundation for Statistical Computing; 2016. doi:10.1007/978-3-540-74686-7.
10. Gu Z, Gu L, Eils R, Schlesner M, Brors B. Circlize implements and enhances circular visualization in R. *Bioinformatics*. 2014;30:2811–2. doi:10.1093/bioinformatics/btu393.
11. Liu X, Li A, Xi J, Feng H, Wang M. Detection of copy number variants and loss of heterozygosity from impure tumor samples using whole exome sequencing data. *Oncol Lett*. 2018;16:4713–20. doi:10.3892/ol.2018.9150.
12. Cridland JM, Macdonald SJ, Long AD, Thornton KR. Abundance and distribution of transposable elements in two drosophila QTL mapping resources. *Mol Biol Evol*. 2013;30:2311–27. doi:10.1093/molbev/mst129.
13. Rogers RL, Cridland JM, Shao L, Hu TT, Andolfatto P, Thornton KR. Landscape of standing variation for tandem duplications in drosophila yakuba and drosophila simulans. *Mol Biol Evol*. 2014;31:1750–66. doi:10.1093/molbev/msu124.
14. Mortazavi A, Williams BA, McCue K, Schaeffer L, Wold B. Mapping and quantifying mammalian transcriptomes by RNA-Seq. *Nat Methods*. 2008;5:621–8. doi:10.1038/nmeth.1226.
15. Wang X, Zheng Z, Cai Y, Chen T, Li C, Fu W, et al. CNVcaller: Highly efficient and widely applicable software for detecting copy number variations in large populations. *Gigascience*. 2017;6:1–12. doi:10.1093/gigascience/gix115.
16. Mi H, Poudel S, Muruganujan A, Casagrande JT, Thomas PD. PANTHER version 10: Expanded protein families and functions, and analysis tools. *Nucleic Acids Res*. 2016;44:D336–42. doi:10.1093/nar/gkv1194.
17. Dennis G, Sherman B, Hosack D, Yang J, Gao W, Lane HC, et al. DAVID: Database for Annotation, Visualization, and Integrated Discovery. *Genome Biol*. 2003;4:P3. doi:10.1186/gb-2003-4-9-r60.

18. Huang DW, Sherman BT, Lempicki RA. Bioinformatics enrichment tools: Paths toward the comprehensive functional analysis of large gene lists. *Nucleic Acids Res.* 2009;37:1-13. doi:10.1093/nar/gkn923.
19. Huang DW, Sherman BT, Lempicki RA. Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. *Nat Protoc.* 2009;4:44-57. doi:10.1038/nprot.2008.211.
20. Gene Ontology Consortium. The Gene Ontology (GO) database and informatics resource. *Nucleic Acids Res.* 2004;32:258D – 261. doi:10.1093/nar/gkh036.
21. Ogata H, Goto S, Sato K, Fujibuchi W, Bono H, Kanehisa M. KEGG: Kyoto encyclopedia of genes and genomes. *Nucleic Acids Res.* 1999;27:29-34. doi:10.1093/nar/27.1.29.
22. Mitchell A, Chang HY, Daugherty L, Fraser M, Hunter S, Lopez R, et al. The InterPro protein families database: The classification resource after 15 years. *Nucleic Acids Res.* 2015;43:D213-21. doi:10.1093/nar/gku1243.
23. Boutet E, Lieberherr D, Tognolli M, Schneider M, Bansal P, Bridge AJ, et al. Uniprotkb/swiss-prot, the manually annotated section of the uniprot knowledgebase: How to use the entry view. In: *Methods in Molecular Biology.* Humana Press, New York, NY; 2016. p. 23-54. doi:10.1007/978-1-4939-3167-5_2.

Supplementary Files Legend

Additional files

Additional file 1 (xlsx format): Supplementary Table 1 Genes in CNV with V_{ST} values above the 99th percentile for each comparison. **Supplementary Table 2** List of CNVR by population. **Supplementary Table 3** Summary of CNVR by population. **Supplementary Table 4** List of global CNVR with frequency above 1%. **Supplementary Table 5** Functional annotation and clustering analysis for global CNVR. **Supplementary Table 6** Functional analysis for CNVR private to breeds and to populations. **Supplementary Table 7** List of the samples used in the analysis. **Supplementary Table 8** Sample processing details.

Additional file 2 (docx format): Supplementary Figures 1-10 CNV differentiation between the following populations; respectively: Boer and East African; Boer and Madagascar; Boer and Southern African; Boer and

West African; East African and Madagascar; East African and Southern African; East African and West African; Madagascar and Southern African; Madagascar and West African and Southern African and Western African.

Supplementary Figure 11-39 CNVR for the following goat breeds, respectively: Abergelle (Ethiopia), Androy (Madagascar), Balaka-Ulongwe (Malawi), Boer (Tanzania and Zimbabwe), Dedza (Malawi), Diana (Madagascar), Galla (Kenya), Gogo (Tanzania), Guera (Mali), Gumez (Ethiopia), Keffa (Ethiopia), Landin (Mozambique), Maasai (Tanzania), Malya (Tanzania), Manica (Mozambique), Mashona (Zimbabwe), Matebele (Zimbabwe), Menabe (Madagascar), Naine (Mali), Norwegian (Tanzania), Nsanje (Malawi), Pare White(Tanzania), Saanen (Tanzania), Small East African (Kenya and Mozambique), Sofia (Madagascar), Soudanaise (Mali), SudOuest (Madagascar), Thyolo (Malawi) and Woyito Guji (Ethiopia). **Supplementary Figure 40** Functional analysis of genes in CNVR highly differentiated among and across populations. **Supplementary Figure 41** Functional analysis of genes in CNVR private to breeds and populations.

Figures

Figure 1
Overall numbers of CNV by chromosome and CNV state.

Figure 2

Distribution of the sizes of CNV for each population by CNV state.

Percentile Plot - CNV Length - Gains

QQ Plot

Figure 3

Percentile plots for CNV gains and losses and a QQ plot for CNV losses.

Figure 4

Population CNV differentiation, estimated by V_{ST} computed across all populations, plotted for each chromosome. The dotted line represents the V_{ST} threshold value for this test (0.601).

Figure 5

Location of the CNVR for the 29 autosomes by population. The outermost numbers are the autosomes, and the other numbers are the start and end positions of each autosome.

Figure 6

Distribution of size of CNVR (in kbp) for each population.

Figure 7

Location of the global CNVR across the 29 autosomes. Blue is for loss; red is for gain and green is for both loss and gain.

Figure 8

Distribution of the CNVR. A, B: Number of CNVR found in different numbers of breeds and populations, respectively. C, D: Distribution of CNVR found in only a single breed and only a single population only, respectively.

Figure 9

Functional analysis of genes in CNV that were highly differentiated between populations based on VST analysis. The analysis is based on the *Bos taurus* gene set.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

Additional file 1.xlsx